

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

***N*-Heterocyclic Carbene-Catalyzed Double Acylation of Enones with Benzils**

Cite this: DOI: 10.1039/x0xx00000x

Ken Takaki,* Akira Ohno, Makoto Hino, Takashi Shitaoka, Kimihiro Komeyama and Hiroto Yoshida

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/


Thiazolium carbene-catalyzed reaction of aromatic 1,2-diketones with enones in aprotic solvents gave double acylation products in good yields, whereas hydroacylation products formed by Stetter reaction were not detected at all. These results suggested a generation of aryloxyenamine species from the 1,2-diketones instead of hydroxyenamines (Breslow intermediates).

N-Heterocyclic carbenes (NHCs) have attracted much attentions as organocatalysts in organic synthesis, because they can convert various electrophilic functional groups into nucleophilic ones via Breslow¹ and deoxy-Breslow intermediates.² For example, the Stetter reaction has been most frequently used in many transformations as a method for Michel addition of acyl anions.³ However, this reaction is applicable to aldehydes only, not to ketones and esters in general. From a mechanistic point of view, when aldehydes **I** are substituted by other carbonyl compounds **V**, functional enamines **VI** would be generated by the reaction with NHC **II**, instead of Breslow intermediates **III** (Scheme 1). If the new species **VI** are sufficiently nucleophilic, acyl and functional group **G** could be delivered to β - and α -position of α,β -unsaturated carbonyl compounds, respectively. This process would be expected to extend the scope of Stetter reaction.

On the other hand, few examples of generation of Breslow intermediate **III** from the carbonyl compounds **V** other than aldehydes **I** have been reported. Scheidt et al. employed acylsilanes⁴ and α -keto carboxylates⁵ in the Stetter reaction to avoid self-condensation or benzoin products, giving rise to the hydroacylation products **IV** in high yields through the elimination of alkoxysilanes or CO₂. Recently, Massi et al. demonstrated that aliphatic 1,2-diketones could be used as aldehyde equivalents, wherein one acyl moieties was eliminated by alcoholic solvents.⁶ As a result, even though the functional enamines **VI** could be generated from **V**, they afforded the same products **IV** as those derived from the corresponding aldehydes **I**.

To develop the new double acylation reaction, we investigated the reaction of benzils **V** (G = ArCO) with enones. 1,2-Diketones have been used as acceptors of the umpolung species formed by NHC.⁷

Moreover, It has been reported that aromatic 1,2-diketones did not afford any Stetter-type products, because 1,4-thiazin-3-ones, 1:1 adducts of thiazolium carbenes and the diketones, were formed exclusively.^{8,9} On the other hand, cyanide ion-catalyzed umpolung of benzils has been achieved in the cross-benzoin condensation,¹⁰ which suggests a possibility to retain the two benzoyl groups in the final products. Fortunately, we found that the reaction gave double acylation products **VII** in good yields for the first time.




Scheme 1 Extension of Stetter reaction.

Initially, benzil (**1a**) was treated with one equivalent of phenyl vinyl ketone (**2a**) under various conditions (Table 1). In the presence of the thiazolium salt **A** (20 mol%), DIPEA as base and DMF as solvent gave better yield of the double acylation product **3aa**, than Et₃N-DCE system (entries 1 and 2). Mild heating of the reaction increased the yield from 53% to 94% yield (entry 3). Of the NHC salts tested, thiazolium salts **B** and **C** gave **3aa** in 77% and 73% yields, respectively, along with trace amounts of hydroacylation product **4** (entries 5 and 7). The triazolium and imidazolium salts **D** and **E** gave little or no expected product (entries 8 and 9). Formation of the by-product **4** with **B** and **C** might be caused by the OH group attached to the thiazolium skeleton. In fact, when the reactions with **A** and **B** were carried out in EtOH solvent, **4** was formed in 11% and


9% yields, respectively, though as a minor product (entries 4 and 6). No reaction took place, of course, in the absence of NHC salts.

Table 1 Optimization of the reaction conditions




Entry	NHC salt	Base	Solvent	Temp (°C)	Yield (%) ^a	
					3aa	4
1	A	Et ₃ N	DCE	rt	17	0
2	A	DIPEA	DMF	rt	53	0
3	A	DIPEA	DMF	50	94	0
4	A	DIPEA	EtOH	50	56	11
5	B	DIPEA	DMF	50	77	trace
6	B	DIPEA	EtOH	50	60	9
7	C	DIPEA	DMF	50	73	trace
8	D	DIPEA	DMF	50	9	0
9	E	DIPEA	DMF	50	0	0

^a NMR yield.



The reaction of benzil (**1a**) with various aryl vinyl ketones **2** was carried out under the optimal conditions (Table 2). The substrates **2** having both electron donating and withdrawing substituents on the *p*-position of the phenyl ring gave the products **3** in high yields (entries 4, 6, and 8), except for *p*-cyanophenyl enone **2i** (entry 9). On the contrary, the substituents on the *o*-position disturbed the reaction to afford **3** in decreased yields, probably due to steric hindrance as well as the reaction with mesityl vinyl ketone (**2e**) (entries 2, 5, and 7).¹¹ Similar difference between 1- and 2-naphthyl substrates **2j** and **2k** was observed (entries 10 and 11). Moreover, the present reaction could be applicable to the heteroaromatic and aliphatic substrates **2l**, **2m** and **2n** (entries 12-14).

Table 2 Reaction with various enones




Entry	Enone 2	R	Time (h)	Product 3	Yield (%) ^a
1	2a	Ph	12	3aa	92
2	2b	<i>o</i> -Tolyl	18	3ab	58
3	2c	<i>m</i> -Tolyl	12	3ac	80

4	2d	<i>p</i> -Tolyl	12	3ad	89
5	2e	Mesityl	36	3ae	13
6	2f	<i>p</i> -Anisyl	12	3af	75
7	2g	<i>o</i> -ClC ₆ H ₄	12	3ag	53
8	2h	<i>p</i> -ClC ₆ H ₄	12	3ah	87
9	2i	<i>p</i> -NCC ₆ H ₄	12	3ai	37
10	2j	1-Naphthyl	12	3aj	67
11	2k	2-Naphthyl	12	3ak	87
12	2l	2-Thienyl	12	3al	87
13	2m	3-Thienyl	12	3am	89
14 ^b	2n	Me	20	3an	68

^a Isolated yield. ^b 3 equiv of **2n** were used.

Next we investigated the reaction of symmetrical and unsymmetrical benzil derivatives **1b-1h** with phenyl vinyl ketone (**2a**) (Table 3). Bis-*p*-tolyl and bis-*p*-anisyl diketones **1b** and **1c** gave the product **3ba** and **3ca** in 48% and 16% yields, respectively, whereas 4-chloro and 4-bromo-substituted diketone **1d** and **1e** produced **3da** and **3ea** in better yields (entries 1-4). In the reaction of unsymmetrical diketones **1f** and **1g**, two regioisomers **3** and **3'** were formed as inseparable mixtures, wherein relatively electron deficient aryl groups were introduced to the β -position of the enone **2a**, though with low selectivities (entries 5 and 6). The reaction of 1-phenylpropane-1,2-dione (**1h**) gave the products **3ha** and **3an** in 42% yield with a ratio of 83/17 (entry 7).

Table 3 Reaction of substituted benzils




Entry	Benzil 1		Product		Yield (%) ^a	Ratio (3/3')
	R ¹	R ²	3	3'		
1	1b	<i>p</i> -Tolyl	<i>p</i> -Tolyl	3ba, –	48	–
2	1c	<i>p</i> -Anisyl	<i>p</i> -Anisyl	3ca, –	16	–
3 ^b	1d	<i>p</i> -ClC ₆ H ₄	<i>p</i> -ClC ₆ H ₄	3da, –	70	–
4 ^b	1e	<i>p</i> -BrC ₆ H ₄	<i>p</i> -BrC ₆ H ₄	3ea, –	73	–
5	1f	Ph	<i>p</i> -Tolyl	3fa, 3ad	80	40 / 60
6	1g	Ph	<i>p</i> -ClC ₆ H ₄	3ga, 3ah	85	55 / 45
7	1h	Ph	Me	3ha, 3an	42	83 / 17

^a Total isolated yield. ^b DMF (0.25 M).

As can be seen in Table 3, electron donating groups of the diketones decreased the product yields (entries 1, 2, and 7). Thus, we attempted their improvement by use of Lewis acid co-catalysts.¹² The yields of **3ba**, **3ca** and (**3ha+3an**) actually increased to 90%, 49% and 62% yields, respectively, with 20 mol% of MgCl₂. However, the effect was not always so positive to all products **3** (see ESI).


More importantly, the reaction of unsymmetric benzils **1f-1h** gave no cross-products, for example, neither **3aa** nor **3ba** was formed in the reaction of **1f**. These results suggested that the double acylation would take place intramolecularly. In order to confirm this hypothesis, the reaction of **2a** with an equimolar mixture of symmetrical benzils **1a** and **1e** was carried out under standard conditions (Scheme 2). The two diketones reacted independently

with **2a** to yield the product **3aa** and **3ea** in 91% and 85% yields, respectively, whereas cross-products were not detected.




Scheme 2 Intramolecular double acylation.

Compared to benzil derivatives, aliphatic 1,2-diketones gave the expected product in lower yields (Scheme 3). Thus, the double acylation product **3ia** was obtained in only 16% yield by the reaction of biacetyl (**1i**) with the enone **2a**, but the corresponding hydroacylation product was not formed. Interestingly, the reaction of cyclohexane-1,2-dione (**1j**) gave 8-membered triketone **3ja** in 27% yield. It is noteworthy that this ring-expansion reaction showed a striking contrast to the thiazolium carbene-catalyzed ring-opening hydroacylation of chalcones with **1j** in EtOH.^{6a}



Scheme 3 Reaction of aliphatic 1,2-diketone.



Scheme 4 Proposed Mechanism.

A plausible mechanism of the double acylation is proposed in Scheme 4. Addition of the carbene **A** to benzil (**1**), followed by migration of benzoyl group to alkoxide moiety yields the acylated Breslow intermediate **G**. Michael addition of **G** to enone **2** generates the zwitterion **H**. Then, intramolecular abstraction of the benzoyl group by enolate results in the formation of the product **3** and regeneration of **A**. Although this mechanism is analogous to that of Stetter reaction, suppression of the benzoyl group elimination from the intermediate **F** or **G** in anhydrous aprotic solvent would realize the double acylation. The low yields with aliphatic 1,2-diketones may be caused by inefficient alkanoyl migration in **F** as well as enolization of the diketone.

In summary, we have succeeded in the extension of Stetter reaction, i.e., thiazolium carbene-catalyzed reaction of benzils with enones gave double acylation products in good yields, whereas hydroacylation by classical Stetter reaction was completely excluded. These results provide new potential for NHC-catalyzed reaction, by allowing the insertion of activated carbon-carbon multiple bonds into other acyl compounds than aldehydes.

This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Notes and references

Department of Applied Chemistry, Graduate School of Engineering, Hiroshima University, Kagamiyama, Higashi-Hiroshima 739-8527, Japan, E-mail: ktakaki@hiroshima-u.ac.jp

† Electronic Supplementary Information (ESI) available: Experimental procedures and spectroscopic data.

- For reviews, see: (a) M. H. Hopkinson, C. Richter, M. Schedler and F. Glorius, *Nature*, 2014, **510**, 485-496; (b) J. Izquierdo, G. E. Hutson, D. T. Cohen and K. A. Scheidt, *Angew. Chem. Int. Ed.*, 2012, **51**, 1686-11698; (c) A. Grossmann and D. Enders, *Angew. Chem. Int. Ed.*, 2012, **51**, 314-325; (d) J. Mahatthanachai and J. W. Bode, *Chem. Sci.*, 2012, **3**, 192-197; (e) X. Bugaut and F. Glorius, *Chem. Soc. Rev.*, 2012, **41**, 3511-3522; (f) V. Nair, R. S. Menon, A. T. Biju, C. R. Sinu, R. R. Paul, A. Jose and V. Streekumar, *Chem. Soc. Rev.*, 2011, **40**, 5336-5346; (g) A. T. Biju, N. Kuhl and F. Glorius, *Acc. Chem. Res.*, 2011, **44**, 1182-1195; (h) P.-C. Chiang and J. W. Bode, *TCI MAIL*, 2011, **149**, 2-17; (i) E. M. Phillips, A. Chan and K. A. Scheidt, *Aldrichimica Acta*, 2009, **42**, 55-66; (j) D. Enders, O. Niemeier and A. Henseler, *Chem. Rev.*, 2007, **107**, 5606-5655.
- (a) M. Schedler, N. E. Wurz, C. G. Daniliuc and F. Glorius, *Org. Lett.*, 2014, **16**, 3134-3137; (b) T. Kato, Y. Ota, S. Matsuoka, K. Takagi and M. Suzuki, *J. Org. Chem.*, 2013, **78**, 8739-8747; (c) S. Matsuoka, S. Namera, A. Washio, K. Takagi and M. Suzuki, *Org. Lett.*, 2013, **15**, 5916-5919; (d) B. Maji, M. Horn and H. Mayr, *Angew. Chem., Int. Ed.*, 2012, **51**, 6231-6235; (e) A. T. Biju, M. Padmanaban, N. E. Wurz and F. Glorius, *Angew. Chem., Int. Ed.*, 2011, **50**, 8412-8415; (f) S. Matsuoka, Y. Ota, A. Washio, A. Katada, K. Ichioka, K. Takagi and M. Suzuki, *Org. Lett.*, 2011, **13**, 3722-3725; (g) C. Fischer, S. W. Smith, D. A. Powell and G. C. Fu, *J. Am. Chem. Soc.*, 2006, **128**, 1472-1473.
- (a) Review, see: M. Christmann, *Angew. Chem., Int. Ed.*, 2005, **44**, 2632-2634; (b) H. Stetter and H. Kuhlmann, *Org. React.*, 1991, **40**, 407-496; (c) H. Stetter, *Angew. Chem., Int. Ed.*, 1976, **15**, 639-647; (d) H. Stetter and M. Schreckenberger, *Angew. Chem., Int. Ed.*, 1973, **12**, 81.

- 4 A. E. Mattson, A. R. Bharadwaj and K. S. Scheidt, *J. Am. Chem. Soc.*, 2004, **126**, 2314-2315.
- 5 M. C. Myers, A. R. Bharadwaj, B. C. Milgram and K. A. Scheidt, *J. Am. Chem. Soc.*, 2005, **127**, 14675-14680.
- 6 (a) O. Bortolini, G. Fantin, M. Fogagnolo, P. P. Giovannini, A. Massi and S. Pacifico, *Org. Biomol. Chem.*, 2011, **9**, 8437-8444; (b) O. Bortolini, G. Fantin, M. Fogagnolo, P. P. Giovannini, V. Venturi, S. Pacifico and A. Massi, *Tetrahedron*, 2011, **67**, 8110-8115.
- 7 V. Nair, S. Vellalath, M. Poonoth, R. Mohan and E. Suresh, *Org. Lett.*, 2006, **8**, 507-509.
- 8 V. Bertolasi, O. Bortolini, A. Donvito, G. Fantin, M. Fogagnolo, P. P. Giovannini, A. Massi and S. Pacifico, *Org. Biomol. Chem.*, 2012, **10**, 6579-6586.
- 9 Ring-opening reaction of cyclic 1,2-diketones to yield α,ω -dicarboxylic acid derivatives has been accomplished also by use of triazolium carbene under aerobic conditions, see: S. Gundala, C.-L. Fagan, E. G. Delany and S. J. Connon, *Synlett*, 2013, **24**, 1225-1228.
- 10 A. S. Demir and Ö. Reis, *Tetrahedron*, 2004, **60**, 3803-3811.
- 11 The double acylation reaction seems to compete with some side reactions like oligomerization of the enones **2**, because they were consumed much faster than benzil **1a** in the low yield reactions (entries 2, 5, 7, and 9).
- 12 (a) J. Qi, X. Xie, R. Han, D. Ma, J. Yang and X. She, *Chem. Eur. J.*, 2013, **19**, 4146-4150; (b) J. Dugal-Tessier, E. A. O'Bryan, T. B. H. Schroeder, D. T. Cohen and K. A. Scheidt, *Angew. Chem. Int. Ed.*, 2012, **51**, 4963-4967; (c) D. T. Cohen and K. A. Scheidt, *Chem. Sci.*, 2012, **3**, 53-57; (d) B. Cardinal-David, D. E. A. Raup and K. A. Scheidt, *J. Am. Chem. Soc.*, 2010, **132**, 5345-5347.