



NJC

Selective induced polarization through electron transfer in non-polar substrates

Journal:	<i>New Journal of Chemistry</i>
Manuscript ID:	NJ-ART-04-2014-000679.R1
Article Type:	Paper
Date Submitted by the Author:	23-Jul-2014
Complete List of Authors:	Tewari, Ashish; Banaras Hindu University, Department of Chemistry Singh, Ved; Banaras Hindu University, Chemistry Srivastava, Priyanka; Banaras Hindu University, Chemistry Singh, Praveen; Banaras Hindu University, Chemistry Kumar, Ranjeet; Banaras Hindu University, Chemistry Khanna, Ranjana; Banaras Hindu university, Department of Chemistry Srivastava, Pankaj; Banaras Hindu University, Chemistry Gnanasekaran, 1. Ramachandran ; Institute of Organic and Biochemistry, Academy of Science Hobza, Pavel; Institute of Organic and Biochemistry, Academy of Science

SCHOLARONE™
Manuscripts

Selective induced polarization through electron transfer in acetone and pyrazole ester derivatives via C-H \cdots O=C interaction

Ashish Kumar Tewari,^{1,*} Priyanka Srivastava,¹ Ved P. Singh,¹ Praveen Singh,¹ Ranjeet Kumar,¹ Ranjana S. Khanna,¹ Pankaj Srivastava,¹ Ramachandran Gnanasekaran² and Pavel Hobza²

¹Department of Chemistry, Faculty of Science, Banaras Hindu University, Varanasi-221 005

² Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 16610 Prague, Czech Republic

Email ID: tashish2002@yahoo.com

ABSTRACT

A set of organic compounds (pyrazole ester derivatives, viz. 5-[3-(substituted)-propoxy]-3-methyl-1-phenyl-1H -pyrazole-4-carboxylic acid methyl ester and 5-[2-(substituted)-ethoxy]-3-methyl-1-phenyl-1H -pyrazole-4-carboxylic acid methyl ester) was synthesized and their affinity and stability towards the acetone molecule were tested by NMR. Further, the host-guest complex formed has been studied by cyclic voltammetric titration. Interestingly, the acetone molecule selectively bound to the methyl group of the pyrazole moiety and stabilized the system. The finding is also supported by quantum chemical calculations using the DFT/B-LYP/TZVPP method. Apart from hydrogen bonding, an important role in stabilizing the complex is played also by the C-H \cdots π interaction governed by dispersion energy. The study proves that methyl-substituted pyrazol ester can act as a receptor of acetone.

KEYWORDS: C-H \cdots O hydrogen bond, ¹H NMR titration, cyclic voltammetry, intermolecular interaction, π -stacking, DFT.

INTRODUCTION

The question of Jeffrey and Saenger of whether $\text{C-H}\cdots\text{O}=\text{C}$ interaction can be referred to as a hydrogen bond even though there is every reason to suspect that the carbon atom is not electronegative and may even carry a net positive charge was denied by Pauling and accepted by the Pimentel and McClellan¹. A refinement of the latter definition led to a quantification by Steiner and Saenger, who in 1993 considered a hydrogen bond as ‘any cohesive interaction $\text{X-H}\cdots\text{Y}$ where H carries a positive and Y a negative (partial or full) charge and the charge on X is more negative than on H’^{2,3}. Recently, a more general definition has been given by Arunan et al.⁴, namely ‘The hydrogen bond is an attractive interaction between a hydrogen atom from a molecule or a molecular fragment X-H in which X is more electronegative than H, and an atom or a group of atoms in the same or a different molecule, in which there is evidence of bond formation’.


Molecular recognition processes can be said to hinge on isotropic interactions, the most important among which is the hydrogen bond⁵. An X-ray single crystal structure analysis of three complexes formed between ethylcalix[4]resorcinarene (CECRA) and acetone molecules suggests a recognition mechanism in which an acetone molecule is captured as a guest in the cavity of a CECRA host molecule through $\text{C-H}\cdots\pi$ interactions between the methyl groups of acetone and the phenyl rings of CECRA⁶. In the case of $\text{C-H}\cdots\text{O}$ hydrogen bonds, the strength of the bond depends on the nature of the other groups attached; even weakly polar C-H groups of methyl⁷ may give carbonyl function forming a weak $\text{C-H}\cdots\text{O}$ hydrogen bond. In this continuation, C5-H in 4,6-dimethyl pyrimidine-2-one derivative is a rich donor, whereas O, N, π are acceptors in this model system⁸. The crystal structure of various organic compounds usually shows $\text{C-H}\cdots\text{O}$ weak interactions⁹⁻¹⁴. These bonds also play an important role in native and misfolded proteins¹⁵.

Recently, 1-(2-naphthyl)-3-methyl-5-pyrazolone coupled with capillary electrophoresis with diode array detection for the determination of carbohydrates such as mannose, galacturonic acid, glucuronic acid, rhamnose, glucose, galactose, xylose, arabinose and fucose has been developed¹⁶.

In the present study, we synthesised the derivatives of 3-methyl-1-phenyl-1H-pyrazole-4-carboxylic acid methyl ester and explored its role as a host for the acetone guest molecule through a weak intermolecular C-H \cdots O hydrogen bond with the support of NMR, cyclic voltammetric (CV) and theoretical studies. The varying chemical shift in the ¹H NMR titration of experimental compounds in CDCl₃ solvent with varying concentration of acetone indicated the formation of host–guest complexation. In CH₂Cl₂ solution, the complex formation between guest and host molecules indicated one electron transfer mechanism in CV studies. The best way to understand this complex formation theoretically is by means of analysing the interaction energy, its correlation with bond length and frequency shift with respect to the localised C-H \cdots O hydrogen bonding. In a usual hydrogen-bond linear complex like X-H \cdots Y, the formation of the hydrogen bond makes the covalent X-H bond weaker with a concomitant decrease of the X-H stretch frequency (red shift)¹⁷. Our analysis thus focuses on the study of the interaction energy, frequency shift and C-H \cdots O hydrogen-bond distances in complexes of compounds **1–5** (Table 1) placed in the acetone environment. The aim of the present work is to study the selectivity and stability of the acetone guest molecule upon complex formation by understanding the role of C-H \cdots O hydrogen bond along with other effects such as π -stacking, steric hindrance etc. using experimental (NMR) and theoretical results.

2. The preparation of the compounds

Pyrazole ester derivatives, viz. 5-[3-(substituted)-propoxy]-3-methyl-1-phenyl-1*H*-pyrazole-4-carboxylic acid methyl ester (1 & 2)[Table 1], 5-[2-(substituted)-ethoxy]-3-methyl-1-phenyl-1*H*-pyrazole-4-carboxylic acid methyl ester (3, 4 & 5)[Table 1], have been synthesised first by synthesising dithioacetals of 3-methyl-1-phenyl-pyrazole-5-one¹⁸ by its treatment with carbon disulphide in the presence of a strong base followed by its treatment with an appropriate alkyl halide. Dithioacetals were then hydrolysed in the presence of a strong base in a corresponding alcohol. Pyrazole esters thus formed were treated with 1,*n*-dibromo alkanes (*n*= 2,3) in excess to synthesise the polymethylene halide of pyrazole ester¹⁹; the subsequent treatment of this halide with a corresponding heterocyclic compound in the presence of a weak base, potassium carbonate, in dimethyl formamide, resulted the desired products. The compounds were characterised by spectroscopic techniques.



Scheme 1. Synthesis of ethylene and propylene linked pyridine and phthalimide substituted pyrazole ester derivatives (1-5)

3. Computational methods

All the systems presented in Table 1 were optimised in vacuum (see the Supporting Information) and in implicit acetone solvent ($\epsilon = 20.7$) using the DFT-D method (the B-LYP functional with the TZVPP basis set²⁰ and Grimme's dispersion (D3) correction²¹ included). This method is known to describe different types of noncovalent interactions reasonably well.²²

To understand the nature of the acetone solvent interaction, an acetone molecule was explicitly connected to the hydrogen atom of the methyl group via a weak C-H \cdots O hydrogen bond (Figure 1) and the structure of the 1:1 complex was fully optimised again in vacuum (see the Supporting Information) and implicit acetone solvent. Basis set superposition error (BSSE) were corrected using counterpoise corrections.²³ All these calculations were performed using the Turbomole Version 6.3 software package.²⁴




Figure 1. The chemical structure of the compounds with an acetone molecule used for theoretical study.

4. Results and discussion

4. a. NMR titration studies

In the solid state, 5-[3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-propoxy]-3-methyl-1-phenyl-1H-pyrazole-4-carboxylic acid methyl ester has two conformations, but in solution only one conformation is observed. This is confirmed through low-temperature NMR in acetone, and a shift in the proton of the methyl group at position 3 is observed to be 1.6 ppm (Figure 2).




Figure 2. The result of the low-temperature NMR experiments in acetone solvent.


The experiments were carried out in chloroform as solvent. Although the interaction of chloroform with acetone is well known,²⁵ in the present studies the interaction of the 3-methyl substituent with acetone is observed to be more favorable, because the oxygen of the carbonyl group is not available to interact with chloroform (calculated interaction energy between acetone and chloroform is -0.11 kcal/mol). The dielectric constant of acetone (20.7) is more compared to chloroform (4.8) which favours the interaction of acetone with the systems under our study. The other reason is also due to the presence of the ester group at the carbon atom adjacent to that of the methyl group in the pyrazole-ring system. The presence of the ester group at position 4 of the pyrazole-ring system makes it easier for the 3-methyl group to undergo no bond resonance and thus to interact with acetone (Scheme 2).

NMR studies²⁶⁻²⁸ are powerful enough to understand such type of weak interactions. ¹H NMR titration experiments of compounds **1–5** (Table 1) with acetone yielded a dramatic change in the chemical shift of 3-methyl protons in all the compounds. During the NMR titration experiments of compound **1**, when acetone was added, the chemical shift of the 3-methyl proton was altered while the rest of the peaks remained unaffected (Table 1), indicating the intermolecular interaction of 3-methyl proton and acetone. With increasing concentration of acetone, a change in the chemical shift of the 3-methyl proton of the pyrazole nucleus was observed without any significant change in the chemical shift of any other proton. This change in the chemical shift of the 3-methyl proton in compound **1** was observed to be 0.48 ppm, which is significant in the identification of any kind of weak interactions between the molecules. Titrations were carried out by increasing the amount of acetone in the solution of compound **1** in CDCl₃. With the addition of 0.2 ml of acetone, there was no change in the chemical shift of the respective proton, whereas on further addition of 0.2 ml of acetone (up to a total acetone volume of 0.4 ml), the change in

the chemical shift of the 3-methyl proton was observed to be 0.14 ppm; an addition of another 0.2 ml of acetone (a total acetone volume of 0.6 ml) led to a change in the chemical shift of 0.33 ppm as compared to the chemical shift of 3-methyl proton of compound 1, and this increase in chemical shift change was further continued with the addition of another 0.2 ml of acetone (a total acetone volume of 0.8 ml), where it was 0.45 ppm. The total addition of 1 ml of acetone in the titration mixture increased the chemical shift of the 3-methyl proton of compound 1 by 0.48 ppm. Further addition of acetone to the titration mixture did not cause any significant change in the chemical shift of the respective protons. In order to understand the strength of such interactions, these experiments were carried out on some other similar molecules and yielded parallel results. The ^1H NMR titration of compound 2 with acetone also showed a reasonable amount of change in the chemical shift for the 3-methyl proton of the pyrazole nucleus. The gradual increase in the concentration of acetone in the titration mixture resulted change in the chemical shift of the 3-methyl proton as 0.21 ppm (0.4 ml of acetone), 0.40 ppm (0.6 ml of acetone), 0.47 ppm (0.8 ml of acetone) and 0.50 ppm (1 ml of acetone). No further alteration of the chemical shift was observed with another increase in the amount of acetone. Similar results were observed with the NMR titration of the rest of the pyrazole-based compounds with acetone. In the case of compound 3, even the addition of 0.4 ml of acetone changed the chemical shift of the 3-methyl proton of the pyrazole nucleus by 0.49 ppm, and with a further addition of acetone, there was a moderate change in the value of the chemical shift of the respective proton. After the addition of 1 ml of acetone in the titration mixture, the total change in the value of the chemical shift was 0.53 ppm. In the case of compounds 4 and 5, the changes in the value of the chemical shift with the addition of acetone were 0.34 ppm, 0.19 ppm (0.4 ml of acetone), 0.46 ppm, 0.39 ppm (0.6 ml acetone), 0.50 ppm, 0.47 ppm (0.8 ml acetone) and 0.51 ppm, 0.47 ppm (1 ml

acetone), respectively. The titration experiments clearly indicate the formation of hydrogen-bonded complexes of compounds 1–5 with acetone via the hydrogen atoms of the 3-methyl substituent.

Table 1: ^1H NMR titration data of compounds with acetone.



Compound No.	Z	n	0.2ml Acetone	0.4ml Acetone	0.6ml Acetone	0.8ml Acetone	1.0ml Acetone	$\Delta\delta$
1		3	2.45	2.59	2.78	2.90	2.93	0.48
2		3	2.46	2.67	2.86	2.93	2.96	0.50
3		2	2.42	2.91	2.95	2.95	2.95	0.53
4		2	2.43	2.77	2.89	2.93	2.94	0.51
5		2	2.47	2.66	2.86	2.91	2.94	0.47

4. b. Cyclic Voltammetric study

The electrochemical properties of compounds were investigated by cyclic voltammetry (CV)^{29,30} in dichloromethane solution at scan rate 100 mV/s for understanding the intermolecular C-H \cdots O hydrogen bonding between experimental molecules and acetone. The experiments were performed in a cell having platinum working (disk) / counter (foil) electrodes and an Ag / Ag⁺

reference electrode. The compounds exhibit single irreversible reduction wave in a potential range of $E_{PC} = -1.68$ to -1.75 V (Figure 3).




Figure 3. Cyclic voltammogram of compounds on titration with different concentration of acetone indicates bonding of acetone molecule with compound via $C-H \cdots O$ weak bond.

The change in peak currents and shift in peak potentials as the concentration of acetone increases can be used to evaluate the formation constants if the electrode reaction is reversible and the formation and dissociation of the complex are fast enough to maintain equilibrium.^{31,32} The analysis of the cyclic voltammograms obtained in the presence and absence of acetone clearly show that the above two requirements are met by the acetone-pyrazole ester system. The formation constants for the reduced and oxidized pyrazole ester derivative were calculated using the method reported by Osa and coworkers.^{31,33} The formation constant K_{ox} was calculated using Eq. (1) (Figure 4) and K_{red} was calculated by substituting K_{ox} in Eq. (2).

$$A^2 = \frac{1}{K_{ox}} \frac{(1-A^2)}{[Comp (1-5)]} + \frac{D_c}{D_f} \dots \dots \dots \text{Eq. (1)}$$

$$\left(a_{E_{1/2}} - f_{E_{1/2}} \right) = 0.0295 \log(K_{ox}/K_{red}) \dots \dots \dots \text{Eq. (2)}$$

Where $A = I_p/I_p^0$; I_p and I_p^0 are the peak currents in the presence and absence of acetone, and D_c and D_f are the diffusion coefficients of the complexed and free compound **1-5** respectively, $a_{E_{1/2}}$ and $f_{E_{1/2}}$ are the half-wave potentials of the compound **1-5** in the presence and absence of acetone.









Figure 4. Plot of A^2 vs. $1 - A^2/[Comp (1-5)]$ for 1:1 complex. $A = I_p / I_p^0$.

The increase in the peak currents at lower concentrations of acetone were used to calculate K_{ox} and the decrease in the peak currents observed at higher concentrations of acetone were used to calculate K_{ox} (Eq. (1)). The shift in the peak potentials at lower concentrations of acetone were used to calculate K_{red} by employing Eq. (1). Since there is no observable change in the peak potential E_p at higher concentrations of acetone ($[acetone]/[Pyrazole\ ester\ derivative] > 2.0 + 0.3$) along with the decrease in peak currents, we are unable to calculate K_{red} .

This indicates an electrochemical change coupled with chemical reaction in the experimental molecules with addition of acetone. Positive shifting of the potentials is suggestive of associated electron withdrawing effects. Reduction wave is indicative of gain of electron by the system which is possible according to the supporting scheme **2**. In this scheme, electron is transferred to carbonyl oxygen leading to the formation of radical anion. Carbon radical undergoes resonance with the π electrons of pyrazole system and anion radical is now created at position 5. Nitrogen atom at position 2 of pyrazole ring system now transfers one of its electrons to carbon-5 of pyrazole ring and forms a cross linked bond. Thus radical is created at position 3 and now hydrogen atoms of methyl group system at position 3 do not undergo any bond resonance to interact with oxygen atom of the carbonyl group of acetone.



Scheme 2: One electron transfer reaction in cyclic voltammetric studies

4. c. Computational study

It is evident from the results of the NMR experiment that the acetone molecule associates with the 3-methyl proton of the pyrazole nucleus. Therefore, we placed the acetone molecule hydrogen-bonded to that methyl proton of pyrazole. The other structures of the complex were also investigated but they were found to be considerably less stable (Figure 5). The optimised structures of all of the complexes are visualised in Figure 1 and selected geometrical parameters are collected in Table 2.




Figure 5. Energy of different possible complex structure of compound 5 with acetone.

The optimised structures of all of the complexes are visualised in Figure 1 and selected geometrical parameters are collected in Table 2.

Table 2. Δ C-H distance and stretching frequencies (ν C-H) Vs Interaction Energy (ΔE) (In acetone solvent)

Compound No.	Δ C-H ₁ (Å)	Δ C-H ₂ (Å)	Δ C-H ₃ (Å)	ν C-H cm ⁻¹	ΔE kcal/mol
1	0.000 (0.001)	-0.001(-0.001)	0.000(0)	+14a(+19a)	-5.1790 (-7.864)
2	-0.002 (-0.001)	0.000(0.001)	0.000(0.001)	+15a, -13a (+8s)	-4.0780 (-4.5058)
3	0.002(-0.001)	0.000(0.001)	0.000(0)	-26a (-15a)	-6.4844 (-6.7296)
4	0.000(0)	-0.001(-0.001)	-0.001(0)	+5a (-4a,+18a)	-4.6985 (-6.760)
5	0.003(0.003)	0.001(0.000)	0.001(0)	+6a, -27s (-21a)	-6.5568 (-8.3693)

s = symmetry; a = asymmetry; '+' = blue shift; '-' = red shift

Values mentioned within parenthesis are from the calculation performed in vacuum.

Δ C-H₁ corresponds to the hydrogen bonded methyl C-H to the carbonyl group of acetone.

Inspecting the resulting structures, we found that the initial planar H-bonded structures passed upon optimisation in vacuum as well as in acetone to stacked structures, but the hydrogen bonds (from methyl) C-H \cdots O (from acetone) remained intact. The resulting stabilisation energies as

well as the shift in the C-H stretching frequencies are summarised in Table 2 and the corresponding correlation between the stabilisation energies and C-H \cdots O are visualised in Figures 6 and 7. The formation of hydrogen-bonded complexes is accompanied by red as well as blue shifts of C-H stretching frequencies. Table 2 also shows the stabilisation energies of all the complexes investigated in acetone (continuous solvent) as well as in vacuum (also Table S1 in the Supporting Information). The stabilisation energies in acetone are reduced on average by 20%. The same table shows the changes of C-H bond lengths upon complex formation; evidently, these changes are rather small and do not correlate with the strength of the H-bond. This is probably caused by the fact that the total stabilisation energy is determined not only by H-bonding but also by C-H $\cdots\pi$ interaction (Figure 1) where the dispersion energy shown in Table 3 plays an important role. It should also be noted that BSSE corrections were important not only in vacuum but also in acetone environment. We have calculated the NBO charges as well as charge transfer for C-H and C=O groups (see Table 3). We believe that these information support our arguments.




Figure 6. The interaction Energy Vs C-H \cdots O distances in acetone solvent




Figure 7. The interaction Energy Vs C-H stretching frequency shift in acetone solvent

A detailed analysis of the optimised structure provides more information on the stability of the complex. In all of the compounds, the free rotation of the methyl group is prevented by intermolecular hydrogen bonding from the nitrogen atom at position 2 of the pyrazole ring and also by the oxygen atom of the acetate group. We proved this by analysing the potential energy surface (PES) scan of torsion angle involving methyl C-H as showed in Figure 8. From this we note that the minima at 0° and 120° correspond to the structures where the hydrogens of the methyl group simultaneously interact with O=C of acetone (intermolecularly) and with the nitrogen atom of the system (intramolecularly).




Figure: 8 Potential energy surface scan of the -CH₃ rotation. The calculations were carried out with the system 4. The minima at 0° and 120° corresponds to the structures where the hydrogens of the methyl group makes simultaneous interactions, intermolecularly with O=C< of acetone and intramolecularly with the 'N' atom.

The added acetone molecule stabilises the complex not only by its hydrogen bonding with the methyl group but also by the C-H... π interaction to the pyrazole ring; apart from that, the phenyl and methoxy groups of the pyrazole are vital to this stabilisation effect. Therefore, the hydrogen-bond angle C-H...O is not linear and varies from 110 to 150 degrees in different complexes.

In compound 3, the phenyl group of the pyrazole group shows intermolecular C-H... π interaction with the 'Z'- group (Table 1) and the methyl group of acetate also forms hydrogen bonds to the acetone; in the case of compound 1, however, due to the increment of the ether chain by a methylene group, the intermolecular C-H... π interaction of the phenyl group is absent. Apart from that, we also noted that unlike in compound 3, the freely rotatable methoxy group is away from the acetone molecule. In the case of compound 4, the acetone molecule has a C-H... π interaction effect both with the pyrazole and phenyl groups. The C-H of the phenyl group forms a weak hydrogen bond with C=O of the 'Z' substituent group. In compound 2, there is an increment in the ether chain; nevertheless, the methyl group of the 'Z' substituent still exhibits a C-H... π interaction with the phenyl group. After complexation with acetone, the pyrazole and phenyl rings are not in the same plane (please check the RMSD value from Table 3).

Table 3. RMSD value of compound 1-5.

Compound No.	H...O distance in Å	Dispersion E kcal/mol	BSSE kcal/mol	$\Delta q(\text{C-H})$ e	$\Delta q(\text{O=C})$ e	RMSD (Å)
1	2.5 (2.6)	-8.6 (-9.2)	-0.8 (0.7)	-0.0038	+0.0044	0.26
2	2.8 (2.9)	-6.2 (-6.1)	-0.9 (0.3)	-0.0035	+0.0011	0.19
3	2.6 (2.5)	-10.0(-9.7)	-1.8 (0.6)	-0.0029	+0.0036	0.27
4	2.5 (2.6)	-7.5 (-8.4)	-1.1 (0.4)	-0.0010	+0.0036	0.34
5	2.7 (2.7)	-9.6 (-9.8)	-1.6 (0.5)	-0.0110	+0.0042	0.36

In compound 5, the acetone molecule is more stabilised by the methoxy and phenyl groups of the pyrazole, and the phenyl group is not involved in intermolecular C-H $\cdots\pi$ interaction or in the hydrogen bond effect with the 'Z' substituent. These effects are reflected in the results, namely in the increase in interaction energy (Figure 6 and 7) and the C-H bond distance (Table 2) and in the decrease in the hydrogen-bond (C-H \cdots O) distance. It is also evident from the RMSD value (Table 3) that the planarity between the pyrazole and phenyl is perturbed in the presence of the acetone molecule.

The optimised structures of all the complexes in vacuum were structurally similar to the corresponding ones in the implicit acetone solvent. However, as arises from the Supplementary Information (Figures S1 and S2 and Table S1), there is a slight variation in frequencies (red and blue shifts) and stabilisation energies.

5. Concluding Remarks

The hydrogen atoms of the 3-methyl group in 3-methyl-1-phenyl-pyrazole-5-one were found to interact with the carbonyl group of acetone. This was shown experimentally by performing a ^1H NMR titration and cyclic voltammetric studies. A low-temperature NMR of the compounds investigated was performed in acetone, with the chemical shift of the 3-methyl proton being about 1.6 ppm. These observations have led to the conclusion that the 3-methyl

protons interact with acetone and form a weak C-H \cdots O bond. The structures of the complexes studied were determined theoretically, and besides the formation of weak C-H \cdots O hydrogen bonds, an important role of C-H $\cdots\pi$ interaction was shown. Thus, these weak non-covalent C-H \cdots O interaction between methyl pyrazole ester and acetone, helps to understand the role of polar aprotic solvent in bimolecular nucleophilic substitution reactions. Such type of studies can be used as an important tool for the study of mechanism of such non-covalent interactions in the near future.

ACKNOWLEDGMENT

We acknowledge UGC India Grant F.37-54/2009 (SR) for its financial support of this work. The Department of Chemistry of Banaras Hindu University is acknowledged for departmental facilities. For the calculation of association constant, authors are highly thankful to Dr. Vellaichamy Ganesan.

References

1. G. A. Jeffrey, W. Saenger, *Hydrogen Bonding in Biological Structures*, Springer, Berlin, 1991.
2. T. Steiner, W. Saenger, *J. Am. Chem. Soc.*, 1993, **115**, 4540.
3. G. R. Desiraju, *Angew. Chem. Int. Ed.*, 2011, **50**, 52.
4. E. Arunan, G. R. Desiraju, R. A. Klein, J. Sadlej, S. Scheiner, I. Alkorta, D. C. Clary, R. H. Crabtree, J. J. Dannenberg, P. Hobza, H. G. Kjaergaard, A. C. Legon, B. Mennucci, D. Nesbitt, *J. Pure Appl. Chem.*, 2011, **83**, 1637.
5. G. A. Jeffrey, *An Introduction to Hydrogen Bonding*, Oxford Univ. Press, New York, 1997.
6. Z. Cao, K. Murayama, K. Aoki, *Analytica Chimica Acta*, 2011, **448**, 47.

7. T. Steiner, *New J. Chemistry*, 1998, **22**, 1099.
8. M. Muthuraman, Y. L. Fur, M. B. Beucher, R. Masse, J. F. Nicoud, S. George, A. Nangia, G. R. Desiraju, *J. Solid State Chem.*, 2000, **152**, 221.
9. P. R. Maulik, K. Avasthi, G. Biswas, S. Biswas, D. S. Rawat, S. Sarkhel, T. Chandra, D. S. Bhakuni, *Acta Cryst.*, 1998, **C54**, 275–277.
10. P. Munshi, K. N. Venugopalan, B. S. Jayashree, T. N. Guru Row, *Crystal Growth and Design*, 2004, **4**, 1105.
11. D. Braga, F. Grepioni, K. Biradha, V. R. Pedireddi, G. R. Desiraju, *J. Am. Chem. Soci.*, 1995, **117**, 3156.
12. A. K. Tewari, R. Dubey, *ARKIVOC*, 2009, **(X)**, 283–291.
13. A. K. Tewari, P. Srivastava, *J. Mol. Struc.*, 2009, **921**, 251.
14. K. Avasthi, S. M. Farooq, C. Bal, R. Kumar, A. K. Tewari, P. R. Maulik, *J. Mol. Str.*, 2007, **842**, 100.
15. A. K. Chamberlain, J. U. Bowie, *J. Mol. Biol.*, 2002, **322**, 497.
16. J. You, X. Sheng, C. Ding, Z. Sun, Y. Suo, H. Wang, Y. Li, *Analytica Chimica Acta*, 2008, **609**, 66.
17. W. Zierkiewicz, D. Michalska, Z. Havlas, P. Hobza, *Chem. Phys. Chem.*, 2002, **3**, 511.
18. A. K. Tewari, P. Srivastava, *Synthetic Commun.*, 2009, **39**, 2837.
19. A. K. Tewari, P. Srivastava, V. P. Singh, A. Singh, R. K. Goel, C. Gopimohan, *Chem. Pharm. Bull. Japan*, 2010, **58 (5)**, 634.
20. A. Schafer, C. Huber, R. Ahlrichs, *J. Chem. Phys.*, 1994, **100**, 5829.
21. S. Grimme, J. Jens Antony, S. Ehrlich, H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.

22. (a) R. Sedlak, T. Janowski, M. Pitonak, J. Rezac, P. Pulay, P. Hobza, P. *J. Chem. Theory Comput.*, 2013, **9**(8), 3364-3374. (b) L. Goerigk, H. Kruse, S. Grimme, *Chem. Phys. Chem.*, 2011, **12**, 3421-3433. (c) J. Rezac, K. E. Riley, P. Hobza, *J. Chem. Theory & Compt.*, 2012, **8**, 4285. (d) S. Haldar, M. Kolar, R. Sedlak, P. Hobza, *J. Phys. Chem. C*, 2012, **116**, 25328-25336. (e) J. Trnka, R. Sedlak, M. Kolar, P. Hobza, *J. Phys. Chem. A*, 2013, **117**, 4331-4337.
23. S. F. Boys, F. Bernardi, *Molecular Physics*, 1970, **19**, 553-566.
24. Y. Chen, Y. Morisawa, Y. Futami, M. A. Czarnecki, H. S. Wang, Y. Ozaki, *J. Phys. Chem. A*, 2014, **118**, 2576-2583.
25. A. Ahlrichs, M. Bar, M. Haser, H. Horn, C. Kolmel, *Chem. Phys. Lett.*, 1989, **162**, 165.
26. M. Felemez, B. Spiess, *J. Am. Chem. Soc.*, 2003, **125**, 7768.
27. H. Dodziuk, A. Ejchart, O. Lukin, M. O. Vysotsky, *J. Org. Chem.*, 1999, **64**, 1503.
28. P. L. Wash, S. Ma, U. Obst, R. Rebek Jr., *J. Am. Chem. Soc.*, 1999, **121**, 7973.
29. T. Nakagaki, K. Shin-ichiro, A. Harano, T. Shinmyozu, *Tetrahedron*, 2010, **66**, 976.
30. M. Oelgemoller, A. Haeuseler, M. Schmittel, A. G. Griesbeck, J. Lex, Y. Inoue, *J. Chem. Soc. Perkin Trans.*, 2, 2002, 676.
31. T. Matsue, D. H. Evans, T. Osa, N. Kobayashi, *J. Am. Chem. Soc.*, 1985, **107**, 3411.
32. J. M. Saveant, E. Vianello, *Electrochim. Acta*, 1963, **8**, 905.
33. T. Matsue, T. Osa, D. H. Evans, *J. Inclus. Phenom.*, 1984, **2**, 547.