



NPR

Ecteinascidins. A Review of the Chemistry, Biology and Clinical Utility of Potent Tetrahydroisoquinoline Antitumor Antibiotics

Journal:	<i>Natural Product Reports</i>
Manuscript ID:	NP-REV-04-2014-000051.R1
Article Type:	Review Article
Date Submitted by the Author:	22-Aug-2014
Complete List of Authors:	Williams, Robert; Colorado State University, Department of Chemistry Inai, Makoto; University of Shizuoka, School of Pharmaceutical Sciences Kan, Toshiyuki; University of Shizuoka, School of Pharmaceutical Sciences; University of Shizuoka, Le, Vy; Colorado State University, Chemistry

SCHOLARONE™
Manuscripts

ARTICLE

Ecteinascidins. A Review of the Chemistry, Biology and Clinical Utility of Potent Tetrahydroisoquinoline Antitumor Antibiotics

Cite this: DOI: 10.1039/x0xx00000x

Received ooth April 2014,
Accepted ooth xxx 2014

DOI: 10.1039/x0xx00000x

www.rsc.org/

V. H. Le,^a M. Inai,^b R. M. Williams,^a and T. Kan.^b

The Ecteinascidin family comprises a number of biologically active compounds, containing two to three tetrahydroisoquinoline subunits. Although isolated from marine tunicates, these compounds share a common pentacyclic core with several antimicrobial compounds found in terrestrial bacteria. Among the tetrahydroisoquinoline natural products, Ecteinascidin 743 (Et-743) stands out as the most potent antitumor antibiotics that it is recently approved for treatment of a number of soft tissue sarcomas. In this article, we will review the backgrounds, the mechanism of action, the biosynthesis, and the synthetic studies of Et-743. Also, the development of Et-743 as an antitumor drug is discussed.

1. Isolation, Structure and Biological Activity of Et-743

Ecteinascidin 743 (Et-743, **1**) is a marine natural product possessing remarkable biological activity against several tumor cell lines.¹ It was first isolated in 1990 from a Caribbean tunicate extract *Ecteinascidia turbinata*,² and was later identified to be produced by the bacterial symbiont *Candidatus Endoecteinascidia frumentensis*.³⁻⁵ In 1992, Sakai *et al.* confirmed the structure of Et-743 via the X-ray crystal structure of the derived N12-oxide.⁶ Et-743 contains three tetrahydroisoquinoline (THIQ) units A, B, and C (Figure 1). The connection between the fully functionalized THIQ units A and B forms a rigid pentacyclic core. Unit C is connected to the pentacycle through a 10-membered macrolactone. Unit A and B are believed to promote the alkylation of DNA through hydrogen bonding with the target DNA. Unit C interacts with DNA-binding proteins, resulting in Et-743's antitumor activity.^{7,8}

Among the Et family members, Et-743 and Et-729 show the most potent antitumor activity on several carcinoma cell types (Table 1).⁹ The major emphasis has been on Et-743 because it was the most abundant compound from the tunicate extract.²




Figure 1. Structure of Et-743.

Et-743 has an effect on cell proliferation against a wide range of cancer subtypes at low concentration.^{10, 11} Izbicka *et al.* illustrated that a continuous exposure of the tumor to Et-743 resulted in a significant improvement of the antitumor activity *in vitro* (Table 2).¹⁰ Their results also revealed that breast, lung, melanoma and ovarian cancers are the most suitable targets for Et-743.

	IC ₅₀ (ng/mL)						IC ₅₀ (μg/mL)					μg/dics
	L1210	P388	A549	HT29	MEL28	CV-1	PS	DNA	RNA	DNAP	RNAP	Bs
Et-743	5.0	0.2	0.2	0.5	5.0	1.0	>1	0.1	0.03	2.0	0.1	0.02
Et-729	< 1.0	0.2	0.2	0.5	5.0	2.5	>1	0.2	0.02	1.5	0.05	0.08
Et-815	25	2.5	5.0	5.0	5.0	5.0		>1	0.1		5.0	0.75
Et-759B		5.0	5.0	5.0	10	25	>1	0.7	0.5		>1	3.90
Et-745B	25	5.0	10	10		25		>1	0.5		3.0	
Et-759C		1.0	2.5	2.5		2.5	2.5		>1	0.5	>5	0.1
Et-745		10	20	25	50	50		>1	0.3		5.0	6.50
Et-731		100	100	100	200	200	>1					6.20
Et-736		0.5	1.0	2.5	2.5	2.5	0.5	0.4	0.1		0.5	0.38
Et-722		1.0	1.0	2.0	2.0	5.0	0.9	0.4	0.1	>1	0.5	0.70
Et-594		10	20	25	25	25	0.8	0.5	0.5		1.0	0.37
Et-597		2.0	2.0	2.0	2.0	2.5	0.7	0.08	0.01		0.25	0.14
Et-583		10	10	10	5.0	25	1.0	1.0	0.4		0.5	0.74

Table 1. Bioactivities of the Ets. L1210 = murine lymphocytic leukemia cells; P388 = murine lymphoblastic cells; A549 = human lung carcinoma; HT29 = human colon carcinoma; MEL28 = human melanoma; CV-1 = monkey kidney cells; PS = protein synthesis inhibition; DNA = DNA synthesis inhibition; RNA = RNA synthesis inhibition; DNAP = DNA polymerase inhibition; RNAP = RNA polymerase inhibition; Bs = *Bacillus subtilis*.

Tumor type	0.1 nM		1.0 nM		10 nM		100 nM		1000 nM	
	1HR	CE	1HR	CE	1HR	CE	1HR	CE	1HR	CE
Ovary	0%	0%	11%	22%	20%	45%	0%	58%	0%	67%
Breast	0%	0%	0%	0%	0%	40%	20%	79%	40%	100%
Non-small cell lung	0%	0%	0%	0%	0%	50%	–	69%	–	85%
Colon	0%	0%	0%	0%	0%	50%	–	43%	–	71%
Melanoma	–	–	–	–	–	71%	–	88%	–	86%
Kidney	0%	0%	0%	0%	0%	43%	0%	50%	0%	67%
Sarcoma	0%	0%	0%	0%	100%	75%	–	67%	–	67%
Corpus uteri	–	–	–	–	–	67%	–	67%	–	67%
Peritoneum	0%	0%	0%	0%	50%	100%	–	–	–	–
Head/neck	0%	0%	0%	0%	0%	100%	–	–	–	–
Lymphoma	–	–	–	–	–	0%	–	0%	–	0%
Pancreas	–	–	–	–	–	100%	–	100%	–	100%
Mesothelioma	–	–	–	–	–	100%	–	100%	–	100%
Prostate	–	–	–	–	–	0%	–	0%	–	0%
Stomach	–	–	–	–	–	0%	–	0%	–	100%

Table 2. Antitumor activity of Et-743 against human tumor colonies after one-hour exposure (1HR) versus continuous exposure (CE).¹⁰

2. Mechanism of action of Et-743

At micromolar concentration, Et-743 was found to inhibit a number of DNA-binding proteins, including NF-Y, TATA binding protein, E2F, and SRF/TCF.¹² It also activates the formation of topoisomerase I-mediated cleavage complexes, and perturbs microtubule arrangement at this high range of concentration.¹³⁻¹⁵ These bioactivities were observed at suprapharmacological concentrations of Et-743, rendering these auxiliary effects of Et-743.

Interestingly, the incubation of NF-Y with Et-743 prior to the addition of DNA results in inhibition at a significantly reduced

drug level (at 10 μM range). This result suggested that transcription factor NF-Y is the target of Et-743, not the CCAAT box on which NF-Y acts. Since NF-Y interacts with several essential genes involving DNA metabolism and cell-cycle regulators,¹⁶ the inhibition of this protein can cause a delay in S-phase progression and accumulation of cells in G₂/M phase, which is in agreement with the observation of Simoens and coworkers.¹⁷ The inhibition of transcription factor NF-Y, however, still occurs at high concentration of Et-743, three times higher than the pharmacological concentration when pre-incubating NF-Y with Et-743.

The study on Et-743-induced suppression of the multidrug resistant gene MDR1 on the human colon carcinoma cell line

SW620 by Jin *et al.* supports transcription factor NF-Y as one of Et-743's target in its mechanism of action.¹⁸ In contrast to the *in vitro* study, the inhibition of MDR1 promoter activation occurs at 50 nM; hence this activity may follow a mechanism unrelated to DNA alkylation. Possibly, at its pharmacological concentration, Et-743 interferes with the binding of NF-Y to other co-activators, such as PCAF, which in turn regulates the expression of several critical genes.

To date, the most widely accepted mechanism of action of Et-743 involves double-strand breaks (DSBs) induced by the transcription-coupled nucleotide excision repair (TC-NER) system following the binding of Et-743 to the minor groove of DNA (Figure 2a). Et-743 alkylates DNA at the N2 of a guanine, forming a DNA adduct.^{7, 19, 20} The presence of the adduct widens the minor groove of the target DNA, and bends DNA toward the major groove.^{21, 22} This event is reversible, favoring 5'-CGG, TGG, GGC, AGC sequences, in which the middle bold G is the site of alkylation.²³ Computer-based dynamic simulation studies illustrated that these sequences optimize the hydrogen-bonding network to both strands of the DNA and orient the carbinolamine of Et-743 on top of the center guanine (Figure 2b). Seaman and Hurley proposed that the above-favored sequences optimize the coordination of hydrogen bond 1 (HB1) and HB2. The combination of HB1, HB2, and HB3 positions and stabilizes HB4 at the site of alkylation.²⁴ The protonated amine (N12) also acts as a catalyst in the generation of the active iminium ion (N2), thus promoting the alkylation.²⁰ Despite numerous thermodynamically favored target sequences, the efficiency of DNA alkylation by Et-743 is low, suggesting that other factors can contribute to the alkylation process.¹⁹ Possibly, the protein-DNA complex increases the sequence selectivity of Et-743.²⁵




Figure 2. a. Et-743-DNA complex. b. Hydrogen-bond network of Et-743 to DNA. The arrows represent the direction of the hydrogen bonds from a donor to an acceptor.

MDR1 promoter contains two GC-rich sequences, thus they are a target for Et-743 alkylation. This event leads to the suppression of P-glycoprotein (P-gp) expression and was found to reverse the sensitivity of P-gp/MDR1 overexpressing cells (KB-8-5 and KB-C-2) to doxorubicin and vincristine, the two

known substrates of P-gp/MDR1.²⁶ Interestingly, Et-743 activity is not subdued at the P-gp levels in human tumor cells, but only at high expression of P-gp as seen in LLC-PK1 pig kidney and Madine-Darby Canine kidney cells.²⁷ This discovery suggested that the use of Et-743 in combination with other antitumor drugs may be beneficial in tumor models resistant to conventional treatments.

DNA adducts formed with Et-743 was found to block gene expressions by causing RNA polymerase II arrest during transcription and competing with certain transcription factors, such as Sp1, for the binding site on DNA.²⁸ These events happen preferentially at the binding site of the fusion proteins FUS-CHOP and EWS-FLI1, leading to the displacement of these oncogenic transcription factors from the promoter.^{29, 30} FUS-CHOP and EWS-FLI1 were known for their role in tumor initiation, specifically myxoid/round cell liposarcoma type I and II by FUS-CHOP and Ewing sarcoma by EWS-FLI1, resulting in the tumor selectivity of Et-743 on these sarcoma.³¹⁻³³




Figure 3. Mode of action of Et-743, dependent on DNA replication.

While other known DNA alkylating agents, such as mitomycin C and cisplatin, requires a deficient nucleotide excision repair (NER) system to efficiently exert their biological activity, Et-743 works the best in the presence of a proficient NER system. Cells deficient in NER genes (*XPG*, *XPA*, *XPD*, or *XPF*) are 2- to 8-fold less sensitive to Et-743, and *XPG*-deficient cells are more resistant to the drug than other strains.³⁴⁻³⁶ Moreover, cells deficient in *XPC* (the gene responsible for recognition of damages in global genome NER (GG-NER)) showed no change

in sensitivity to Et-743, elucidating that the toxicity of Et-743 is independent on the GG-NER but is involved the TC-NER machinery.³⁴ Upon formation of a DNA adduct, TC-NER is recruited to the lesion by the use of the *XPA* gene product (Figure 3). *XPD* will then unwind the damaged region, and *XPF* and *XPG* gene products will induce the single-strand break (SSB) on the opposite strand to the alkylation.²⁸ An Et-743-induced adduct traps the *XPG*-DNA complex, generating irreversible SSBs,³⁷ which are transformed into double-strand breaks (DSBs) during transcription.^{35, 38} Homologous recombination (HR) repair can recognize and repair the DSBs induced by Et-743; hence cells lacking HR repair (common in human tumors) are significantly more sensitive to Et-743.^{38, 39} Studies of Zewail-Foote *et al.* on the less complicated bacterial nuclease UvrABC system demonstrated that the incision event induced by Et-743-DNA adduct is more efficient at the less favored sequence 5'-AGT.⁴⁰ Based on this result, they proposed a different mechanism of how Et-743 induces apoptosis (Figure 4). The UvrA protein of the UvrABC machinery recognizes the Et-743-DNA lesion and initiates the repair mechanism by forming a dimer with another UvrA protein, then a heterotrimer UvrA₂B. This event creates a six-base pair bubble around the alkylated guanine, bending DNA toward its major groove, and resulting in the dissociation of the UvrA₂. UvrC is recruited to the UvrB-DNA complex to incise at both ends of the DNA containing the alkylated guanine, thus the lesion can be fixed. Et-743 can also dissociate from the bubble before the incision happens depending on the stability of the Et-743-DNA adduct, which also leads to cell survival. When the repair mechanism occurs at the favored DNA sequences, the incision is not as effective or does not happen at all, generating a toxic lesion and cell death.




Figure 4. Incision effectiveness on unfavored DNA sequences versus favored DNA sequences.

More recently, Guirouilh-Barbat *et al.* discovered a direct DSB pathway mediated by the TC-NER and Mre11-Rad50-Nbs1 (MRN) complex, independent of replication.⁴¹ The authors proposed that the TC-NER-mediated SSBs induced by Et-743 recruits the MRN complex, causing DSBs (Figure 5). DNA-dependent protein kinase (DNAPK) recognizes these DSBs and phosphorylates serine 139 of H2AX protein (γ H2AX) around DSB sites, causing apoptosis. MRN and γ H2AX also activates the ataxia telangiectasia-mutated (ATM) gene, which in turn will phosphorylate more H2AX, thus intensifying the cell death signal. Besides amplifying the effect of Et-743, MRN and ATM are associated with DNA-damage repair as well via activation of DNA-damage checkpoints, which can result in cell survival.⁴²




Figure 5. Et-743-induced DSBs mediated by TC-NER and MRN independent of replication.

A part of Et-743's biological activity was associated with its ability to modulate the tumor microenvironment as a rapid reduction of blood vessels and mononuclear phagocytes, especially tumor-associated macrophages (TAM), was observed around the tumor.⁴³ The reduction of macrophages was resulted from the activation of enzyme caspase-8, which initiates the extrinsic apoptotic pathway in leukocytes. The binding of decoy R3 onto TRAIL receptor inhibits this apoptotic pathway in neutrophils and T cells, and hence their resistance to Et-743.

These studies reveal that the detailed interactions of Et-743 in cells are enormously complex and much may yet be discovered regarding how this drug reacts at the biochemical and cellular levels.

3. Biosynthesis of the Ecteinascidins

The biosynthetic studies of the ecteinascidins face several obstacles due to the fact that the producing bacterial symbiont has yet to be successfully cultured making it extremely difficult to obtain high quality samples of DNA. In addition, the ecteinascidins and their biosynthetic precursors are present in trace amounts in the tunicate extract. Despite these intrinsic challenges, information about the biosynthesis of these alkaloids has been partially obtained, owing largely to the structural similarity between the ecteinascidins, specifically Et-743, and other THIQ family members, including saframycin B (2),⁴⁴ saframycin Mx1 (3),⁴⁵ safracin A (4),⁴⁶ naphthiridinomycin (5),⁴⁷ and quinocarcin (6) (Figure 6).^{48, 49}

An early study from Kerr's research group revealed that tyrosine and cysteine are associated with the biosynthesis of Et-743.⁵⁰ Serine was originally proposed to make up the α -hydroxyethyl moiety (C1-C22) of Et-743 (Figure 7). However, there was no strong evidence to support this hypothesis. Studies by Peng *et al.* elucidated the role of ketoses in the biosynthesis of naphthiridinomycin and quinocarcin, in which the ketose phosphates are converted to α -hydroxyethyl-thiamine pyrophosphate prior to their incorporation into these natural products.⁵¹ Et-743 shares a similar α -hydroxyethyl moiety with these two bacterial metabolites; hence its structure may also arise from a ketose as well.




Figure 6. Structural similarity between Et-743 (1) and saframycin B (2), saframycin Mx1 (3), safracin A (4), naphthiridinomycin (5) and quinocarcin (6).

The addition of ¹⁴C-labeled diketopiperazines (DKPs) 7 and 8 to the cell-free extract of *E. tubinata* afforded radiolabeled Et-743 (Figure 7). These two DKPs were found to incorporate into Et-743 at a significantly higher level than ¹⁴C-labeled tyrosine (9), suggesting that 8 could be the first committed precursor in Et-743 biosynthesis, and that oxidation of 8 to generate 7 could happen before its incorporation into the pentacyclic core of Et-743.⁵²

The challenge of culturing the tunicate bacterial symbiont *Candidatus Endoecteinascidia frumentensis* has been an obstacle to the discovery of the gene cluster responsible for Et-743 biosynthesis. To gain insights of how Et-743 is produced in the Nature, Rath and coworkers utilized metagenomic sequencing methodology on the tunicate consortium, followed by genetic comparison between the resulting DNA pools with those of the known bacterial THIQ biosynthetic genes, specifically of saframycin A (Sfm), saframycin Mx1 (Saf) and safracin (Sac).⁵³⁻⁵⁵ Their studies resulted in the identification of the gene cluster encoding the enzymes associated with the biosynthesis of Et-743 (Etu) (Scheme 1).³

EtuH is a homologue of SfmD, a tyrosine hydroxylase, functioning to hydroxylate the C3 of a tyrosine. EtuM1, a homologue of SacF and SfmM2, and EtuM2, a homologue of SacC and SfmM3, are believed to C- and O-methylate 3-hydroxytyrosine, respectively. The combination of EtuH, EtuM1 and EtuM2 was proposed to produce tyrosine derivative 13 as previously observed in saframycin and safracin biosynthetic pathways (Scheme 1).⁵⁶




Figure 7. The origin of Et-743.

ARTICLE





Figure 8. Gene cluster encoding the biosynthesis of Et-743. A: NRPS; D: DNA processing; F: fatty acid biosynthesis; H: hydrolase; M: methyltransferases; N: amidotransferases; O: monooxygenase; P: pyruvate cassette; R: regulatory enzymes; T: drug transporter; U: unknown function.

The non-ribosomal peptide synthetase (NRPS) responsible for the biosynthesis of the ecteinascidins contains three modules, *EtuA1*, *EtuA2* and *EtuA3* (Figure 8). The common starter module AL-T is located in *EtuA3*, where a long-chain fatty acid is activated as a thioester **14** (Scheme 1). The *EtuA3* domain was predicted to incorporate cysteine to the fatty acid, followed by the elongation of the chain at the *EtuA1* domain with a glycolic acid to provide **16**. Unlike conventional NRPSs, in which all the components are incorporated collinearly, the biosynthesis of the THIQ family was found to be noncollinear.^{57, 58} The C domain of *SfmC* was found to act as a Pictet-Spenglerase to form the mono- and bis-THIQ intermediates in the biosynthesis of saframycin A.⁵⁹ The role of the C domain of the *EtuA2* module may catalyze a similar transformation in the biosynthesis of the ecteinascidins. Hence, after the reductive release of **16**, aldehyde **17** is condensed with the primary amine of **18** followed by ring closure of the resulting imine via Pictet-Spengler reaction to yield the first THIQ intermediate **19**. The sequence is repeated to insert a second tyrosine derivative **13**, followed by reductive release of the resulting bis-THIQ intermediate to generate aldehyde **22**. Internal condensation of the aldehyde with the secondary amine gives the pentacyclic core **23**, which will undergo hydrolysis of

the long chain *N*-acyl moiety to provide free amine **24** under catalysis of *EtuF3* (a homologue of penicillin acylase).⁶⁰ *EtuO* is a homologue of *SfmO2* and *SacJ*, FAD-dependent monooxygenases, and is proposed to act as a catalyst for the insertion of the last hydroxyl group on the pentacyclic core of Et-743.⁵⁵ There is no reference or evidence of the last few steps in Et-743 biosynthesis. However, the presence of other isolated ecteinascidins can serve as hints of how pentacycle **24** can be converted to Et-743. Pentacycle **24** can be transformed into Et-583 (**25**), which is then methylated at the secondary amine to give Et-597 (**26**). Subsequent transamination of Et-597 may happen to provide Et-596 (**27**), which will incorporate the dioxymethylene to yield Et-594 (**28**). Condensation of Et-594 with 3-hydroxy-4-methoxytyrosine (**29**), followed by the Pictet-Spengler reaction of the resulting imine, gives rise to intermediate **30**, which can be converted to Et-743 via an unknown transformation. Although more work will have to be done before a complete biosynthetic pathway for the ecteinascidins can be mapped, the work of Rath and coworkers provided the first molecular genetic basis of how the ecteinascidins are made in Nature.

ARTICLE




Scheme 1. Proposed biosynthetic pathway of Et-743 (1).

4. Synthetic Studies of Et-743

Due to its scarcity, being derived from a marine origin, Et-743 has garnered substantial interest from the synthetic community which has enabled its introduction into clinical use. While the field of THIQ synthesis is quite mature, the complex molecular structure of Et-743, relative to other members of the family, have rendered it a particularly alluring and relevant synthetic target. Currently, Et-743 is manufactured commercially by a semi-synthetic process from cyanosafraicin B and the enabling synthetic technology to convert cyanosafraicin B into commercial grade Et-743 relied on the pioneering synthetic studies reported from the Corey laboratory.⁶¹


4-1. Corey's total synthesis of Et-743

The first, and landmark total synthesis of Et-743 was reported by Corey and Gin *et al.* in 1996.^{62,63} Their synthesis featured a biomimetic condensation of two tyrosine derivatives **36** and **45**. Synthesis of the right hand segment **36** commenced with gallic ester **31**. After protection of **31** as the TBS ether, the methyl ester was converted to the aldehyde **32** by reduction and oxidation. Condensation of aldehyde **32** and monomethylmalonate proceeded smoothly to give **33** as a 1 : 1 mixture. Upon heating **33** and DPPA, isocyanate formation by Curtius rearrangement and subsequent trapping with benzyl alcohol proceeded to give the (*Z*)-enamide ester as a single isomer. With the desired compound **34** in hand, an asymmetric hydrogenolysis reaction was accomplished by treatment with H₂ in the presence of Rh-DiPAMP catalyst. After changing the nitrogen-protecting group of **35** from Cbz to the Alloc group, acidic hydrolysis of the methyl ester provided the carboxylic acid **36**.



Scheme 2. Reagents and conditions (a) TBSCl, Et₃N, DMAP, CH₂Cl₂, 23 °C (98%); (b) DIBAL-H, CH₂Cl₂, -78 °C (quant); (c) PDC, MS4A, CH₂Cl₂, 23 °C (99%); (d) monomethylmalonate, piperidine, AcOH, MS3A, toluene, 23 °C (92%); (e) (PhO)₂P(O)N₃, Et₃N, MS4A, toluene, 70 °C; BnOH, 70 °C (89%); (f) H₂ (45 psi), Rh[(COD)]-(*R,R*)-DiPAMP]⁺BF₄⁻, MeOH, 23 °C (100%, 96% ee); (g) H₂, 10% Pd/C, EtOAc, 23 °C (quant); (h) AllocCl, pyridine, 23 °C (93%); (i) 0.2 M HCl, AcOH, 110 °C; (j) TBSCl, imidazole, DMF, 23 °C (95%).


As shown in Scheme 3, the left-hand segment **45** was also synthesized by a Rh-catalyzed asymmetric hydrogenolysis reaction of a dehydroamino ester derived from **41**. After protection of sesamol **37** as the MOM ether, regioselective lithiation with *n*-BuLi in the presence of TMEDA and following treatment with MeI the desired alkylation reaction proceeded smoothly. Subsequent *ortho*-lithiation of the MOM ether and addition of DMF gave the benzaldehyde derivative **38**. After changing the protecting group of **38** from the MOM ether to the corresponding benzyl ether, Knoevenagel reaction of **39** with malonate **40** and cleavage of the protecting group provided the carboxylic acid **41**. After incorporation of the nitrogen atom by Curtius rearrangement of **41**, Rh-DiPAMP catalyst-mediated asymmetric hydrogenation of the Cbz-protected *Z*-enamide ester proceeded smoothly to provide the desired substance **42** in 95% yield and 96% ee. After deprotection of the dimethyl acetal by treatment with BF₃·OEt₂ in the presence of H₂O, treatment with BF₃·OEt₂ and MS 4A, the desired intramolecular Pictet-Spengler reaction proceeded to give the tricyclic lactone **44** as a single isomer. Subsequent cleavage of the Cbz and Bn groups under hydrogenolysis conditions gave the left-hand segment **45**.



Scheme 3. Reagents and conditions (a) NaH, DMF/Et₂O, 0 °C; MOMBr, 0 °C (90%); (b) *n*-BuLi, TMEAD, hexane 0 °C; MeI, -78 to 23 °C (87%); (c) *n*-BuLi, THF -30 °C; DMF, 0 °C (64%); (d) MeSO₃H, CH₂Cl₂, 0 °C; (e) NaH, 23 °C; BnBr, 23 °C (86% 2steps); (f) **40**, piperidine, AcOH, MS4A, benzene, 23 °C (99%); (g) Et₃N/HCO₂H, Pd(PPh₃)₄, THF, 23 °C (93%); (h) (PhO)₂P(O)N₃, Et₃N, MS4A, toluene, 70 °C; BnOH, 23 °C; (i) H₂ (20 psi), Rh[(COD)]-(*R,R*)-DiPAMP]⁺BF₄⁻, MeOH, CH₂Cl₂, 23 °C (97%, 96% ee); (j) BF₃·OEt₂, H₂O, CH₂Cl₂, 0 °C; (k) BF₃·OEt₂, MS4A, CH₂Cl₂, 23 °C; (73% 2 steps); (l) H₂, 10% Pd/C, EtOAc, 23 °C.

With both desired segments in hand, condensation of **36** and **45** was accomplished by treatment with CIP and HOAt, followed


by protection of the phenolic residue as the *O*- ether furnishing **46** in good yield. After partial reduction of lactone **46** with $\text{LiAlH}_2(\text{OEt})_2$, deprotection of both TBS ethers gave the cyclization precursor **47**. Upon treatment of **47** with $\text{CF}_3\text{SO}_3\text{H}$ in a 3:2 mixture of $\text{H}_2\text{O} : \text{CF}_3\text{CH}_2\text{OH}$, the desired intramolecular Mannich bis-annulation reaction proceeded smoothly through the *N*-acyl-imine to provide the desired pentacyclic intermediate **48**. After partial reduction of lactone **49** with $\text{LiAlH}_2(\text{OEt})_2$ and subsequent treatment with KCN and AcOH, aminonitrile formation of the incipient aminal derivative proceed to afford **50**. After regioselective incorporation of the trifluoromethyl sulfonyl group on the less-hindered phenol of **50**, the resultant primary alcohol and phenol were protected as their TBDPS and MOM ethers respectively. After spontaneous deprotection of the Alloc group and ether by treatment with a Pd-catalyst and Bu_3SnH , reductive amination of the secondary amine with formaldehyde was carried out by treatment with NaBH_3CN and AcOH. The incorporation of the aryl methyl group via the triflate was accomplished by a Stille cross-coupling to give **52**.



Scheme 4. Reagents and conditions (a) 2-chloro-1,3-dimethylimidazolium hexafluorophosphate (CIP), HOAt, Et_3N , $\text{THF}/\text{CH}_2\text{Cl}_2$, 23 °C; (b) bromide, Cs_2CO_3 , DMF, 23 °C (81%, 2 steps); (c) $\text{LiAlH}_2(\text{OEt})_2$, Et_2O , -78 °C (95%); (d) KF, MeOH, 23 °C; (e) 0.6 M TfOH, $\text{H}_2\text{O}/\text{CF}_3\text{CH}_2\text{OH}$, BHT, 45 °C (89%, 2 steps); (f) $\text{LiAlH}_2(\text{OEt})_2$, THF, 0 °C; AcOH, 4.8 M aq KCN, 23 °C (87%); (g) PhNTf_2 , Et_3N , DMAP, CH_2Cl_2 , -30 °C (74%); (h) TBDPSCl, DMAP, CH_2Cl_2 , 23 °C (89%); (i) MOMBr, DIPEA, CH_2Cl_2 , 23 °C (92%); (j) *n*- Bu_3SnH ,

$\text{PdCl}_2(\text{PPh}_3)_2$, AcOH, CH_2Cl_2 , 23 °C (quant); (k) CH_2O , NaBH_3CN , AcOH, MeCN, 23 °C (95%); (l) SnMe_4 , $\text{PdCl}_2(\text{PPh}_3)_2$, LiCl, DMF, 80 °C (83%).

For the final transformation, oxidation of the phenol **52** to the quinone and incorporation of the side chain was accomplished, as shown in Scheme 5. Oxidation of phenol **52** with benzeneseleninic anhydride proceeded to give the tertiary α -hydroxyl ketone derivative. After deprotection of the TBS ether, incorporation of protected cysteine derivative **54** afforded **53**. The crucial step for construction of **57** into the ten-membered lactone was accomplished in the following sequence. First, Swern-type oxidation proceeded by treatment of **53** with Tf_2O and DMSO followed by the addition of the non-nucleophilic base *i*- Pr_2NEt , resulted in the interesting *o*-quinone methide intermediate **55**. Subsequent addition of Barton's base **56**,⁶⁴ deprotection of the Fmoc group and in situ cyclization to construct the ten-membered ring proceeded with final trapping of the incipient phenolate by acetate furnishing **57**. After deprotection of the alloc group, oxidative deamination was accomplished employing Rappaport's conditions to give the keto-lactone **58**. Condensation of amine **59** with **58** in the presence of silica gel, the desired Pictet-Spengler reaction proceeded to give **60** as a single isomer. Finally, deprotection of the MOM ether and conversion of the α -amino-nitrile to the carbinolamine by exposure to AgNO_3 afforded Et-743 (**1**).




Scheme 5. Reagents and conditions (a) $(\text{PhSeO})_2\text{O}$, CH_2Cl_2 , 23 °C, 15 min (82%); (b) TBAF, THF, 23 °C (91%); (c) **54**,

EDCI, DMAP, CH₂Cl₂, 23 °C (91%); (d) Tf₂O, DMSO, DMSO, –40 °C; DIPEA, 0 °C; *t*-BuOH, 0 °C; **56**, 23 °C; Ac₂O, 23 °C; (e) *n*-Bu₃SnH, PdCl₂(PPh₃)₂, AcOH, CH₂Cl₂, 23 °C (84%); (f) [*N*-methylpyridinium-4-carboxaldehyde]⁺Tf⁻, DBU, DMF, CH₂Cl₂, 23 °C (70%); (g) **59**, silica gel, EtOH (58%); (h) CF₃CO₂H, THF/H₂O; (i) AgNO₃, H₂O, (77%, 2 step).

4-2. Fukuyama and Kan's total synthesis of Et-743


In 2002, Fukuyama and Kan reported the total synthesis of Et-743 (**1**).⁶⁵ Their synthesis featured the powerful Ugi four-component condensation and intramolecular Heck reactions that rapidly assembled the tetrahydroisoquinoline ring. As shown in Scheme 6, phenyl glycinol derivative **66** was synthesized by Mannich-type reaction with chiral template **63**, which was developed by the author's group.⁶⁶ The phenol **62**, which was readily obtained from MOM-protected sesamol **61**, was reacted with **63** to provide **64**. After conversion to the triflate, reduction of the lactone ring and protection of the resulting primary alcohol as the TBDPS ether provided **65**. After incorporation of the methyl group,⁶⁷ removal of the chiral auxiliary was performed by Pb(OAc)₄-mediated oxidation of the amino-alcohol and aminolysis of the imine intermediate by treatment with hydroxylamine yielding **66**.



Scheme 6. Reagents and conditions (a) *n*-BuLi, THF, 0 °C to rt; B(OMe)₃, H₂O₂, AcOH, rt (92%); (b) TFA, CH₂Cl₂, –10 °C (89%); (c) Tf₂O, pyridine, CH₂Cl₂, 0 °C (90%); (d) NaBH₄, MeOH, 0 °C (85%); (e) TBDPSCl, imidazole, DMF, rt (91%); (f) MeZnCl, PdCl₂(dppf) (3 mol%), THF, reflux (97%); (g) Pb(OAc)₄, MeCN, 0 °C; (h) NH₂OH·HCl, NaOAc, EtOH, rt (89% in 2 steps).

As shown in Scheme 7, the right-hand segment was synthesized from inexpensive 3-methylcatechol by employing the DuPHOS-mediated asymmetric hydrogenation. Regioselective protection of **67** was performed by treatment with TsCl. Incorporation of a bromine atom was carried out at the *p*-position of the phenol and adjustment of phenolic protecting groups furnished **68**. The bromide **68** was converted to the benzaldehyde by halogen-lithium exchange and subsequent treatment with DMF. Regioselective introduction of the iodo substituent was next achieved by directed ortho-lithiation of the corresponding dimethylacetal followed by quenching with I₂. Simultaneous cleavage of the MOM ether and the


dimethylacetal and subsequent benzylation of the phenol afforded the iodobenzaldehyde **69**, which was then subjected to Horner-Emmons reaction with the phosphonate **70** to give the (*Z*)-dehydrophenylalanine derivative **71**.⁶⁸ Catalytic asymmetric hydrogenation of **71** proceeded smoothly in the presence of DuPHOS catalyst to afford the aminoester and subsequent basic hydrolysis gave the desired carboxylic acid **72**.⁶⁹



Scheme 7. Reagents and conditions (a) TsCl, TEA, CH₂Cl₂, 0 °C (83%); (b) Br₂, AcOH, CH₂Cl₂, rt (97%); (c) MeI, K₂CO₃, acetone, reflux (96%); (d) NaOH, EtOH/H₂O, reflux (97%); (e) MOMCl, DIPEA, CH₂Cl₂, 0 °C (97%); (f) *n*-BuLi, THF, –60 °C; DMF (79%); (g) HC(OMe)₃, CSA, MeOH, rt (94%); (h) *n*-BuLi, Et₂O, 0 °C to rt; I₂; (i) conc. HCl, THF, rt (72%, 2 steps); (j) BnBr, K₂CO₃, CH₃CN, reflux (98%); (k) **70**, TMG, CH₂Cl₂, rt (93%); (l) Rh[(COD)-(S,S)-Et-DuPHOS]⁺OTf⁻, H₂ (500 psi), EtOAc, 50 °C (99%, 94% ee); (m) LiOH, MeOH/H₂O/THF, 0 °C to rt (quant).

With both desired segments **66** and **72** in hand, condensation was next examined a mixture of amine **66**, carboxylic acid **72**, *p*-methoxyphenyl isocyanide (**73**) and acetaldehyde was heated in MeOH, the desired Ugi's four-component condensation reaction proceeded to afford the dipeptide **74**.⁷⁰ After switching from the TBDPS ether to the acetate, simultaneous cleavage of the Boc group and the MOM ether gave the aminophenol, which was readily cyclized to afford **75**. The dioxopiperazine **75** was converted to the cyclization precursor **76** by a four-step sequence involving mesylation, introduction of a Boc group onto the lactam nitrogen, partial reduction of the ring carbonyl group, and dehydration. The crucial Heck reaction of **76** was performed by treatment with a catalytic amount of Pd₂(dba)₃ and P(*o*-tol)₃ to afford the desired tricycle **77** in good yield.⁷¹⁻⁷³ After switching the protecting groups of the amine and the phenol of **77** to the corresponding *N*-Troc-*O*-Ac compound, the enamide was oxidized with DMDO and immediately treated with CSA to afford the corresponding methoxy alcohol. The subsequent acyliminium ion-mediated reduction under acidic conditions afforded alcohol **78** as a single product. After silylation of the alcohol **78**, cleavages of the two acetyl groups and selective benzylation of the phenolic hydroxyl group, the partial reduction of the lactam carbonyl was performed by treatment with Red-Al to afford the corresponding oxazolidine. Cleavage of the oxazolidine with TMSCN and BF₃·OEt₂


afforded the aminonitrile **79** as a single stereoisomer. Conversion from **79** to aldehyde **80** was accomplished by an acetylation of the regenerated hydroxyl group, cleavage of the TBS ether, and oxidation of the resultant alcohol with Dess-Martin periodinane. During the hydrogenolysis reaction of the benzyl ethers **80** a spontaneous cyclization reaction proceeded to give the desired pentacycle **81**.



Scheme 8. Reagents and conditions (a) **73**, MeCHO, MeOH, reflux (90%); (b) TBAF, THF, rt (89%); (c) Ac₂O, pyridine, DMAP, rt (93%); (d) TFA, anisole, CH₂Cl₂, rt; (e) EtOAc, reflux (87%, 2 steps); (f) MeSO₂Cl, pyridine, CH₂Cl₂, 0 °C (91%); (g) (Boc)₂O, DMAP, CH₃CN, rt (97%); (h) NaBH₄, H₂SO₄, EtOH/CH₂Cl₂, 0 °C; (i) CSA, quinoline, toluene, reflux (88% in 2 steps); (j) Pd₂(dba)₃ (5 mol%), P(*o*-tol)₃ (20 mol%), TEA, CH₃CN, reflux (83%); (k) NaOH, MeOH/H₂O, reflux; (l) Ac₂O, pyridine, DMAP, rt (93%, 2 steps); (m) TFA, CH₂Cl₂, rt; (n) TrocCl, aq. NaHCO₃/CH₂Cl₂, rt (74%, 2 steps); (o) DMDO, MeOH/acetone, 0 °C; cat. CSA (90%); (p) NaBH₃CN, TFA/THF, 0 °C (94%); (q) TBSCl, imidazole, DMF, rt (92%); (r) guanidinium nitrate, NaOMe, MeOH/CH₂Cl₂, 40 °C (85%); (s) BnBr, K₂CO₃, CH₃CN, reflux (91%); (t) Red-Al, THF, 0 °C (82%); (u) TMSCN, BF₃·OEt₂, CH₂Cl₂, 0 °C (73%); (v) Ac₂O, pyridine, DMAP, rt (92%); (w) HF, CH₃CN, rt (quant); (x)

Dess-Martin periodinane, CH₂Cl₂, rt (92%); (y) Pd/C, H₂, THF, rt (84%).

As shown in Scheme 9, selective allylation of the phenol, cleavage of the acetyl group, and condensation of the resultant alcohol with L-cysteine derivative **82** furnished ester **83**. Chemoselective hydrazinolysis of the thioacetate gave the thiol, which, upon exposure to TFA, smoothly underwent cyclization to give the ten-membered sulfide. Subsequent acetylation of the resultant phenol gave **84**. Cleavage of the Troc group followed by reductive alkylation afforded the *N*-methyl amine, whose Alloc group and ether were simultaneously cleaved with a palladium catalyst to give the aminophenol. According to the protocol reported by Corey, conversion to keto-lactone **85** and Pictet-Spengler reaction furnished Ecteinascidin 770. Finally, conversion from the aminonitrile to the labile hemiaminal was effected by treatment with AgNO₃ to give Ecteinascidin 743 (**1**).




Scheme 9. Reagents and conditions (a) bromide, DIPEA, CH₂Cl₂, reflux (89%); (b) K₂CO₃, MeOH, rt (99%); (c) **82**, EDCI, DMAP, CH₂Cl₂, rt (94%); (d) H₂NNH₂, CH₃CN, rt (98%); (e) TFA, CF₃CH₂OH, rt; (f) Ac₂O, pyridine, DMAP, rt (71% in 2 steps); (g) Zn, AcOH, Et₂O, rt (92%); (h) HCHO, AcOH, NaBH₃CN, MeOH, rt (96%); (i) PdCl₂(PPh₃)₂, AcOH, *n*-Bu₃SnH, CH₂Cl₂, rt (89%); (j) [N-methylpyridinium-4-carboxaldehyde]⁺I⁻, DMF/CH₂Cl₂, rt; DBU; citric acid (54%); (k) **59**, NaOAc, EtOH, rt (96%); (l) AgNO₃, CH₃CN/H₂O, rt (93%).

4-3. Zhu's total synthesis of Et-743


In 2005, Zhu and co-workers reported the third total synthesis of Et-743.⁷⁴ The left-hand benzyl bromide **88** was synthesized from sesamol **61**, which was converted to the phenol **62** according to Fukuyama's protocol.⁶⁵ Friedel-Crafts type reaction of **62** and ethyl glyoxalate provided α -hydroxy ester **86**. Incorporation of the methyl group on to the phenol **86** was performed by a two-step sequence including chemoselective conversion to the triflate by using the author's nitrophenyl

triflate⁷⁴ and Suzuki-Miyaura cross coupling with boroxine.⁷⁵ Treatment of benzyl alcohol **87** with thionyl bromide and benzotriazole provided the benzyl bromide **88**.



Scheme 10. Reagents and conditions (a) MOMCl, NaH, Et₂O/DMF, 0 °C to rt (96%); (b) *n*-BuLi, B(OMe)₃, THF then AcOH, H₂O₂, 0 °C to rt (95%); (c) LiCl, MS 3A, HFIP/toluene, ethyl glyoxalate, rt (97%); (d) 4-nitrophenyltriflate, K₂CO₃, DMF, rt (94%); (e) trimethyl boroxine, K₃PO₄, Pd(PPh₃)₄, 1,4-dioxane, reflux (93%); (f) SOBr₂, benzotriazole, CH₂Cl₂, rt (91%).


The right-hand segment phenyl alaninol derivative **95** was synthesized from 3-methylcatechol **67** by utilizing an asymmetric alkylation with glycine derivative **92**.⁷⁶ Regioselective incorporation of the Ts group, methyl group and formyl group provided aldehyde **90**. After reduction of **90**, benzyl alcohol was converted to the corresponding bromide **91**. Upon treatment of **91** and glycine derivative **92** in the presence of the Corey's catalyst,⁷⁷⁻⁷⁹ the desired alkylation reaction proceeded smoothly and subsequent acidic hydrolysis of the imine to provide the amino ester **94**. After reduction of the ester moiety of **94** and basic hydrolysis of the Ts group provided the phenyl alaninol derivative **95**.



Scheme 11. Reagents and conditions (a) TsCl, Et₃N, CH₂Cl₂, -70 °C; (b) MeI, K₂CO₃, acetone, 55 °C (84%, 2 steps) (c) TiCl₄ in CH₂Cl₂, Cl₂CHOCH₃, 0 °C to rt (85%); (d) NaBH₄, MeOH/THF/H₂O, 0 °C (quant); (e) PBr₃, toluene/CH₂Cl₂, 0 °C to rt (96%); (f) **92**, cat. **93** (10 mol%), CsOH·H₂O, CH₂Cl₂, -

78 °C, then after work up THF/H₂O/AcOH (85%); (g) LiBH₄, MeOH/Et₂O, rt; (b) 2M NaOH, EtOH, reflux.


Construction of the DE-ring was accomplished by Pictet-Spengler reaction of amino-phenol **95** and Garner's aldehyde (**96**).⁸⁰⁻⁸² Upon treatment of **95** and **96** with AcOH in CH₂Cl₂ and CF₃CH₂OH, the desired tetrahydroisoquinoline formation reaction proceeded to give **97** as a single isomer.⁸³⁻⁸⁵ Employment of these optimized conditions, the cyclization reaction was accomplished with high regio- and diastereoselectivity. After protection of secondary amine with the Alloc group, chemoselective allylation of the phenol and acetylation of resultant primary alcohol provided **98**. Acidic hydrolysis of **98** by TFA provided amino-alcohol **99**.



Scheme 12. Reagents and conditions (a) **96** AcOH, CH₂Cl₂/CF₃CH₂OH (7:1), MS3A, 20 h (84%); (b) AllocCl, NaHCO₃, CH₂Cl₂, rt, 2 h (88%); (c) bromide, Cs₂CO₃, DMF, rt, 3 h (86%); (d) Ac₂O, pyridine, DMAP, CH₂Cl₂, rt, 1 h (92%); (e) TFA, CH₂Cl₂, rt (72%).

With the desired A-ring bromide **89** and DE ring amine **99** in hand, the next challenge in their synthesis was condensation of these two segments. Upon treatment of **89** and **99** with base, secondary amine formation proceeded to provide desired product **100** and its β-isomer in 68% and 23% yield, respectively. The C-ring precursor **100** was prepared by silylation of the alcohol and cleavage of the acetyl group. After oxidation of the primary alcohol with Dess-Martin periodinane, treatment with TMSCN and a catalytic amount of ZnCl₂ afforded the aminonitrile as a single stereoisomer. For construction of the B-ring, the precursor aldehyde was converted by a four-step sequence involving reduction of the ester moiety, introduction of the acetyl group onto the primary alcohol, cleavage of the TBS ether, and Dess-Martin oxidation of the resultant alcohol. Upon treatment of **102** with TFA, cleavage of the MOM ether and subsequent spontaneous cyclization ensued to give the desired pentacyclic intermediate. After cleavage of the acetyl group, regioselective condensation of the primary alcohol with L-cysteine derivative **103** furnished ester **104**. Upon exposure **104** to 1% TFA in CF₃CH₂OH, cleavage of Troc group and cyclization of the ten-membered sulfide ring proceeded; subsequent acetylation of the resultant phenol gave the desired acetate. Simultaneous deprotection of

the Alloc group and ether, followed by reductive *N*-methylation gave **105**. Finally, according Corey's protocol,⁶² key intermediate **105** was readily converted to **1**.




Scheme 13. Reagents and conditions (a) TEA, MeCN, 0 °C (α -isomer: 68%, β -isomer: 23%); (b) TBSCl, imidazole, DMF, rt (97%); (c) K₂CO₃, MeOH, rt (94%); (d) Dess-Martin periodinane, rt; TMSCN, ZnCl₂, rt (78%); (e) LiBH₄, MeOH, THF, 0 °C (80%); (f) Ac₂O, pyridine, DMAP, CH₂Cl₂ (92%); (g) HF, H₂O, MeCN, rt (91%); (h) Dess-Martin periodinane, rt (93%); (i) TFA, CH₂Cl₂, rt (95%); (j) K₂CO₃, MeOH, rt (96%); (k) **103**, EDCI, DMAP, CH₂Cl₂, rt (95%); (l) TFA, TFE, rt; Ac₂O, pyridine, DMAP, CH₂Cl₂ (77%); (m) *n*-Bu₃SnH, PdCl₂(PPh₃)₂, AcOH, CH₂Cl₂, rt (87%); (n) NaBH₃CN, AcOH, HCHO, rt (96%); (o) Zn, AcOH, rt (92%); (p) 4-formyl-1-methylpyridinium benzenesulfonate, DBU, sat. oxalic acid, DMF/CH₂Cl₂, rt (53%); (q) **59**, NaOAc, EtOH, rt (97%); (r) AgNO₃, MeCN/H₂O, rt (92%).

4-4. Danishefsky's formal synthesis of Et-743

In 2006, Danishefsky and co-workers reported a formal total synthesis of Et-743.⁸⁶ The synthesis started from Borchardt catechol, readily obtained by formylation of **107**.⁸⁷ After cleavage of two methyl ethers, mono-bromination was carried out regioselectively. Subsequent methylenation, Bayer-Villiger oxidation, acidic hydrolysis, and incorporation of the TBDPS group onto phenol provided **109**. After lithiation of **109**, treatment with the Weinreb amide derivative gave the ketone


110.⁸⁸ Asymmetric reduction of **110** was accomplished by the combination of Noyori transfer-hydrogenation reaction⁸⁹ and treatment of **111** with DPPA to give azide **112**. After hydrogenation of azide of **112**, reductive amination provided the two-carbon homologated product. Subsequent protecting group swap of the TBDPS group to the *O*-ether gave **113**. Upon treatment of **113** with HCl, hydrolysis of the dimethyl acetal and cyclization reaction proceeded to afford tetrahydroisoquinoline **114**.



Scheme 14. Reagents and conditions (a) Cl₂CHOMe, TiCl₄, CH₂Cl₂; (b) BBr₃, CH₂Cl₂ (c) Br₂, NaOAc, AcOH, 0 to 25 °C; (d) BrCH₂Cl, Cs₂CO₃, DMF, 105 °C (66%, 2 steps); (e) *m*CPBA, 25 to 64 °C; (f) HCl, 0 to 25 °C (78%, 2 steps); (g) TBDPSCl, TEA, DMAP, 25 °C (89%); (h) *n*-BuLi, toluene/THF, -78 °C; neat Me-(MeO)NC(O)CH₂OBn, -78 °C (80 %); (i) RuCl[(*R,R*)-Ts-DPEN](*p*-cymene), HCO₂H, TEA, DMF, 0 to 40 °C (78%, 95% ee); (j) DPPA, DBU, toluene/DMF, 50 °C (89%, 95% ee); (k) H₂ (1 atm), Pd/C, EtOAc, 25 °C (80%); (l) (MeO)₂CHCHO, AcOH, NaBH₃CN, MgSO₄, MeOH, 0 to 65 °C; (m) TBAF, THF, 0 °C (94%, 2 steps); (n) allyl bromide, NaH, DMF, 0 to 25 °C (84%); (o) 7 M HCl, 1,4-dioxane, 0 to 25 °C (90%).

Condensation reaction of **114** and tyrosine derivative **115**⁹⁰ was performed by treatment with BOPCl in the presence of Et₃N to provide **116**. Preparation of the intramolecular Pictet-Spengler cyclization precursor **117** was carried out by a four-step sequence, involving the deprotection of PMB group, dehydration to the enamide, oxidation to the aldehyde, and Pd-mediated cleavage of ether. Upon treatment of **117** with thirty equivalents of difluoroacetic acid, the desired cyclization proceeded to give the pentacycle **118**. Conversion to Fukuyama's synthetic intermediate **126** from phenol **118** was accomplished by the following sequence. After incorporation of the TBS residue to the phenol **118**, conversion of the *N*-Me amine to the Troc-protected species was carried out by treatment with TrocCl in the presence of TBAI. After changing the protecting group of **119** from TBS to the MOM group, hydration of enamide **120** was employed. DMDO oxidation of

120 provided unstable epoxide **121** and without isolation treatment with NaBH_3CN provided benzyl alcohol **122** as a predominant product. After removal of the two benzyl ethers of **122**, treatment with a 1:1 complex of *n*-BuLi and DIBAL, partial reduction of amide **123** proceeded to provide the oxazolidine **124**. After selective protection of phenol **124** with the group, exposure to KCN and acetic acid provided aminonitrile **125**. Finally cleavage of the MOM group of **125** with TFA gave Fukuyama's synthetic intermediate **126**, constituting a formal total synthesis.




Scheme 15. Reagents and conditions (a) BOPCl, TEA, CH_2Cl_2 , 0 to 25 °C (85%); (b) DDQ, $\text{CH}_2\text{Cl}_2/\text{pH}$ 7.0 buffer, 25 °C (90 %); (c) $\text{Cu}(\text{OTf})_2$, benzene, 85 °C (61%); (d) Dess-Martin periodinane, CH_2Cl_2 , 25 °C (94%); (e) $\text{PdCl}_2(\text{PPh}_3)_2$, *n*-Bu₃SnH, AcOH, 0 to 25 °C (93%); (f) $\text{CHF}_2\text{CO}_2\text{H}$, MgSO_4 , benzene, 100 °C, (42-58%); (g) TBSOTf, TEA, CH_2Cl_2 , 0 °C (quant); (h) TrocCl, TBAI, toluene, 110 °C (92%); (i) TBAF, CH_2Cl_2 , 0 °C, 2 min; (j) MOMCl, DIPEA, (79%); (k) DMDO, CH_2Cl_2 , 0 to 25 °C; (l) NaCNBH_3 (78%) (m) H_2 (1 atm), Pd/C,

EtOAc, 25 °C (77%); (n) DIBAL-H/*n*-BuLi, THF, 0 °C (78%); (o) bromide, DIPEA, CH_2Cl_2 , 50 °C (66%); (p) KCN, AcOH, 25 °C (79%); (q) TFA, CH_2Cl_2 , 20 °C (54%).

4-5. Williams' formal synthesis of Et-743


Williams and Fishlock reported the formal total synthesis of **1** in 2008, that relayed into Fukuyama's synthesis.⁹¹ For this effort, they developed a new, radical-based cyclization reaction to form the left-hand tetrahydroisoquinoline. The synthesis commenced with phenol derivative **116**, which was readily obtained from **107** by using a similar procedure employed by Danishefsky.⁸⁶ Upon treatment of phenol **116** with Garner's aldehyde (**117**)⁸⁰⁻⁸² with titanium phenolate, the Friedel-Crafts reaction proceeded smoothly to give the desired *anti*-product **118**. After protection of the phenolic group as the corresponding *O*- ether, hydrolysis of the oxazolidine ring of **119**, formation of the *trans*-acetone and deprotection of the Boc group via Ohfuné's protocol provided amine **120**.⁹² Construction of the B-ring of **121** was accomplished by a novel glyoxal-imine radical cyclization. Condensation between amine **120** and ethyl glyoxalate proceeded smoothly to provide the imine intermediate **121**. Upon treatment of the glyoxal-imine intermediate **121** with slow addition of *n*-Bu₃SnH and AIBN, the desired radical cyclization reaction proceeded smoothly to provide tetrahydroisoquinoline **122** as a single isomer.



Scheme 16. Reagents and conditions (a) Cl_2CHOMe , TiCl_4 , CH_2Cl_2 ; (b) BBr_3 , CH_2Cl_2 ; (c) Br_2 , NaOAc, AcOH/ CH_2Cl_2 (92%); (d) BrCH_2Cl , Cs_2CO_3 , MeCN, 110 °C (69%); (e) *m*CPBA, CHCl_3 , reflux; (f) 4 M HCl, MeOH/ CH_2Cl_2 , rt (73%, 2 steps); (g) *R*-Garner's aldehyde (**117**) $\text{Ti}(\text{O}i\text{-Pr})_4$, toluene, rt; (h) bromide, Cs_2CO_3 , DMF, rt (65%, 2 steps); (i) TsOH·H₂O, MeOH, rt; (j) 2,2-dimethoxypropane, TsOH·H₂O, DMF, rt (84% 2 steps); (k) TBSOTf, 2,6-lutidine; MeOH; $\text{KF}\cdot 2\text{H}_2\text{O}$, MeOH (76%); (l) ethyl glyoxalate, MS4A, toluene; (m) *n*-Bu₃SnH, AIBN, benzene, reflux.

After reduction of the ester group of **122**, subsequent protection of the resulting primary alcohol as a benzyl ether afforded **123**.

Acylation of the tetrahydroisoquinoline amine **123** was achieved by conversion to the acid chloride from **124**⁹³ to give the desired amide **125** without epimerization. After swapping out the amine protecting group from Fmoc to Boc group, treatment of **126** with Dowex in methanol, the desired removal of acetonide group proceeded smoothly with incorporation of methanol at the benzylic position (as an inconsequential 1:1 diastereomer mixture).




Scheme 17. Reagents and conditions (a) LiAlH_4 , THF, 0 °C; (b) NaH, BnBr, DMF/THF, 0 °C (77%, 2 steps); (c) $(\text{COCl})_2$, CH_2Cl_2 , rt; **120**, DMF, rt (70%); (d) Et_2NH , CH_2Cl_2 , rt; (e) Boc_2O , $\text{EtOH}/\text{CH}_2\text{Cl}_2$, rt; (f) Dowex 50W-X8, MeOH (90%); (g) TBAF, THF, rt (95%); (h) Swern Oxidation; (i) TFA, anisole, CH_2Cl_2 , rt (72%, **128** : **129** = 0.72 : 1); (j) TrocCl, pyridine, CH_2Cl_2 , 0 °C; (k) BnBr, K_2CO_3 , TBAI, acetone, rt (85%, 2 steps); (l) $\text{Pd}(\text{PPh}_3)_4$, pyrrolidine, CH_2Cl_2 , rt; (m) MOMBr, DIPEA, CH_2Cl_2 , rt (56%, 2 steps); (n) DMDO; (o) NaBH_3CN .

Deprotection of the phenolic *O*-TBS group and oxidation of the primary alcohol under Swern conditions provided the aminal **127**. Upon treatment of **127** with TFA, deprotection of the Boc


group and Pictet-Spengler reaction proceeded with elimination concomitant elimination of the methoxy group at the benzylic position to afford **128** and **129** as a 1:1 regioisomeric mixture. After separation, the desired pentacycle **128** was converted to the Fukuyama synthetic intermediate **122** also intercepted by Danishefsky.⁶⁵ Protection of the amine with the Troc group and the phenolic group as the benzyl ether provided **130**. Changing the protecting group from the allyl ether of **130** to the corresponding MOM ether afforded **120** which constituted a formal total synthesis. 4-6. Fukuyama's formal synthesis of Et-743

In 2013, Fukuyama and co-workers reported a second-generation total synthesis of Et-743. The A-ring moiety **136** as the precursor for diazonium salt was prepared from known phenol **132**.^{94,95} Oxidation of **132** with $\text{PhI}(\text{OAc})_2$, and treatment with sodium cyanide gave nitrile **134**. After benzylation of the phenolic hydroxy group, the resulting nitrile was hydrolyzed to furnish carboxamide **135**. Hofmann rearrangement followed by hydrolysis afforded amine **136**.



Scheme 18. Reagents and conditions (a) $\text{PhI}(\text{OAc})_2$, MeOH, 0 °C; (b) NaCN, DMF/ H_2O , 0 °C to rt (37%, 2 steps); (c) BnBr, K_2CO_3 , DMF, rt; (d) aq H_2O_2 , K_2CO_3 , DMSO, rt; (e) $\text{PhI}(\text{OAc})_2$, KOH, MeOH, 0 °C; (f) LiOH, $\text{EtOH}/\text{H}_2\text{O}$, reflux (83%, 4 steps).


The construction of the CDE ring by *N*-acyliminium ion-mediated cyclization reaction is detailed in Scheme 19. Perkin condensation of known dioxopiperazine **138**⁹⁶ with aldehyde **139** proceeded smoothly to give **140**. After introduction of a Boc group at the lactam, cleavage of the benzyl group, stereoselective reduction of the double bond, hydrazinolysis of the acetyl group in **141** followed by selective reduction of the imide carbonyl group with sodium borohydride afforded **142**. Upon treatment of **142** with TFA, the *N*-acyliminium ion-mediated cyclization reaction proceeded smoothly, and subjection of the product to PhNTf_2 under basic conditions afforded bis-triflate **143** in 88% yield. Suzuki-Miyaura coupling of **143** with trimethylboroxine took place selectively at the less-hindered triflate to produce **144** in 92% yield. After the Boc group was switched to a methoxycarbonyl group, partial reduction of the ester moiety in **145** with L-Selectride and subsequent dehydration of the resulting hemi-aminal under acidic conditions afforded enamide **146**. The Tf group was replaced with a MOM group in a one-pot process to afford **147**.



Scheme 19. Reagents and conditions (a) Ac_2O , 130 °C (80%); (b) **139**, *t*-BuOK, THF, -78 to 0 °C; DBU, 0 °C; (c) Boc_2O , DMAP, THF, rt (quant, 2 steps); (d) H_2 (750 psi), Pd/C, EtOAc, rt; (e) $\text{H}_2\text{NNH}_2\cdot\text{H}_2\text{O}$, THF, rt; evaporation; NaBH_4 , MeOH, 0 °C (57%, 2 steps); (f) TFA, $\text{CF}_3\text{CH}_2\text{OH}$, rt; evaporation; PhNTf_2 , DMAP, Cs_2CO_3 , MeCN, rt (88%); (g) trimethylboroxine, $\text{Pd}(\text{PPh}_3)_4$, K_3PO_4 , 1,4-dioxane, 100 °C (92%); (h) HCl, EtOAc, rt; ClCO_2Me , NaHCO_3 , H_2O , 0 °C (91%); (i) L-Selectride, THF, -42 °C; (j) CSA, toluene, reflux (55%, 2 steps); (k) aq KOH, 1,4-dioxane, rt; MOMCl, 0 °C (95%).

After treatment of amine **136** with *t*-butyl nitrite and $\text{BF}_3\cdot\text{OEt}_2$,⁹⁷ Heck reaction of the resulting diazonium salt and enamide **147** from the less-hindered face of the enamide produced coupling product **148** with the desired stereo- and regiochemistry. The dihydroxylation of the highly hindered double bond in **148** and subsequent oxidative cleavage of the resulting 1,2-diol with H_5IO_6 formed a dialdehyde, which underwent facile hydration to afford **150**. Hydrogenolysis of the benzyl ether in **150** gave phenol **151**. Heating **151** in *m*-xylene promoted liberation of the dialdehyde, which was trapped intramolecularly by the electron-rich A-ring moiety to furnish **153**. Subsequent reduction of **153** with Red-Al afforded **154** in 76% yield over the two steps. Treatment of **154** with KCN in acetic acid induced cleavage of the oxazolidine ring, forming aminonitrile **155**. Condensation of the primary hydroxyl group in **155** with cysteine derivative **82** followed by selective cleavage of the *S*-acetyl group with hydrazine furnished thiol **156** in good yield. Upon treatment of **156** with TFA, the cyclic sulfide formation occurred presumably via the generation of an

ortho-quinone methide to give, after acetylation of the phenolic hydroxyl group, compound **157** in 55% yield. Sequential cleavage of the MOM and Alloc protecting groups furnished **158**, which was identical to the intermediate of their previous synthesis, and was converted into Ecteinascidin 743 (**1**) via the published three-step sequence.



Scheme 20. Reagents and condition: (a) $\text{BF}_3\cdot\text{OEt}_2$, *t*-BuONO, THF, -15 to 0 °C; **136**, $\text{Pd}_2(\text{dba})_3$, NaOAc, MeCN/THF, 0 °C to rt; (b) OsO_4 , $\text{K}_3[\text{Fe}(\text{CN})_6]$, K_2CO_3 , quinuclidine-HCl, MeSO_2NH_2 , *t*-BuOH, H_2O , rt (93%, 2 steps); (c) H_5IO_6 , THF, 0 °C (87%); (d) H_2 , Pd/C, MeOH, rt; (e) *m*-xylene, 120 °C; Red-Al, -42 to 60 °C (76%, 2 steps); (f) KCN, AcOH, rt (98%); (g) **82**, EDCI, DMAP, CH_2Cl_2 , rt (92%); (h) $\text{H}_2\text{NNH}_2\cdot\text{H}_2\text{O}$, MeCN, rt (85%); (i) TFA, $\text{CF}_3\text{CH}_2\text{OH}$, 25 °C; toluene, evaporation; Ac_2O , pyridine, rt (55%); (j) TFA, CH_2Cl_2 , rt (64%); (k) $\text{PdCl}_2(\text{Ph}_3\text{P})_2$, AcOH, *n*- Bu_3SnH , CH_2Cl_2 , rt (95%).

5. Development of Et-743 (Yondelis®) as a Clinical Drug: Therapeutic Applications

The pharmacological development and commercialization of Et-743 are performed by PharmaMar (Zeltia group) and Ortho Biotech Products (Johnson & Johnson) under the commercial name Trabectedin or Yondelis®. Phase I clinical trials of Trabectedin were performed in the United States, France, and by the European Organisation for the Research and Treatment of Cancer (EORTC).⁹⁸ The most common toxicity observed in these studies was hematological, including grade 3-4 neutropenia, febrile neutropenia and thrombocytopenia. Grade 3-4 transaminitis was observed, and this adverse event is reversible. Nausea/vomiting, fatigue and hyperbilirubinemia were also associated with Trabectedin treatments. In the American study, the maximum tolerated dose (MTD) was recommended to be 1.2 mg/m² as a continuous infusion over 72 hours every three weeks for phase II studies.⁹⁹ The proposed MTD was 1.5 mg/m² as a continuous infusion over 24 hours every three weeks in the French study, and 1 mg/m² over 1-hour infusion or 1.65 mg/m² over 3-hour infusion every three weeks in the EORTC study.^{100, 101}

The safety and toxicity profile of Trabectedin was obtained from phase II studies on a total of 1,132 patients (all had received prior treatments, including surgery, radiotherapy and/or chemotherapy) with a wide variety of malignancies.¹⁰² These patients were administered with Trabectedin according to one of the following schedules: twenty-four-hour infusion every three weeks at 1.5 mg/m², three-hour infusion every three weeks at 1.3 mg/m², or three-hour infusion every week at 0.58 mg/m². Nineteen deaths were reported only during the first two cycles of the treatments. 27.7% of the cycles were delayed over five days, 10.2% of the patients were discontinued, and dose reduction was necessary to 9% of the patients due to severe adverse events. Hematological toxicity was common, with predictable and reversible neutropenia and thrombocytopenia occurring the most frequently. Elevation of liver enzymes (AST/ALT) was the most common non-hematological event; however, liver damage by Trabectedin can be managed by administration of Dexamethasone thirty minutes prior to Trabectedin infusion. Nine patients experienced rhabdomyolysis, leading to neutropenia, sepsis, renal failure and elevated liver enzymes, and this adverse event was the most fatal among all of the drug-related toxicities. Liver damage associated with Trabectedin treatment can be reduced by Dexamethasone pretreatment. Other Trabectedin-related toxicities, including fatigue, nausea, vomiting, alopecia, mucositis/stomatitis, periphery sensory neuropathy, cardiac disorder, and renal toxicity, were also reported in these studies.

In a randomized phase II study of Trabectedin in patients with advanced or metastatic liposarcoma and leiomyosarcoma after failure of standard treatments, a reduction in tumor progression was recognized, and the overall survival was favorable.¹⁰³ Following these trials, Trabectedin was approved for the treatment of advanced or metastatic soft tissue sarcoma after

failure of anthracyclines and ifosfamide, or in patients that are not suited for these treatments in 2007 in Europe.¹⁰²

The combination of Trabectedin with Cisplatin or Doxorubicin is well-tolerated among patients. Also, synergistic activity was observed with little or no cross-resistance of Trabectedin to Cisplatin- or Doxorubicin-resistant models,¹⁰⁴⁻¹¹⁰ supporting the use of Trabectedin in combination with other antitumor agents. A randomized phase III study of Trabectedin/pegylated liposomal Doxorubicin (PLD) combination on recurrent ovarian cancer showed a significant improvement in progression-free survival and overall response rates compared to PLD alone.¹¹¹ Moreover, this combination was well-tolerated among patients with manageable and non-cumulative adverse events, and a reduction in PLD-related toxicity was observed.¹¹² Therefore, in 2009, Trabectedin was granted approval for treatment of relapsed platinum-sensitive ovarian cancer in combination with PLD in Europe.¹¹³

In 2012, the Children's Oncology Group reported a phase II clinical trial of Trabectedin in children with recurrent rhabdomyosarcoma, Ewing sarcoma and non-rhabdomyosarcoma soft tissue sarcoma.¹¹⁴ Fifty patients ranging from four to twenty-four years old were administered Trabectedin as 24-hour infusion every three weeks at 1.5 mg/m² or three-hour infusion every three weeks at 1.3 mg/m².¹¹⁵ Reported adverse events were less serious compared to the adults' studies with neutropenia being the most common toxicity, where grade 3-4 neutropenia and febrile neutropenia were rare. Other adverse events were also observed, including ALT/AST and GGT elevation and hematological toxicity. Despite the less severe toxicity in this group of patients, the development of Trabectedin to treat these malignancies was not continued due to the lack of meaningful response rates.

Trabectedin was not approved for cancer treatment by the FDA due to its hepatotoxicity and not statistically improving survival rate.⁹⁶ However, it is certified as an orphan drug for soft tissue sarcoma and ovarian cancer in the United States.¹¹⁶ Currently, it is in phase II and III clinical trial for treatment of breast cancer, other pediatric tumors, and first-line treatment of soft tissue sarcoma with translocation. In Japan, trabectedin is developing as a novel treatment of malignant soft tissue sarcoma with translocation and is gaining supportive results from a phase II clinical trials.

It is significant that, this drug, which will likely become more widely adopted as on-going clinical studies advance, was made possible through the pioneering synthetic chemistry efforts of the Corey laboratory, enabling the semi-synthetic manufacturing route from Cyanosaffracin B. Emerging studies on the biosynthesis of the Ecteinascidins, along with many of the additional synthetic studies reviewed here, will likely play a role in the evolution of this potent and fascinating marine alkaloid as a life-saving medicine for various forms of refractory cancers. The story of the Ecteinascidins, serves to underscore the tremendous and continuing importance of natural products discovery, total synthesis efforts and engineered biosynthesis as enabling technologies for the discovery, development and advance of powerful new

therapeutic agents. Future chapters in the history of this complex family of alkaloids, will no doubt be exciting.

Notes and references

^a Department of Chemistry, Colorado State University, Fort Collins, Colorado, 80523 USA.

^b Faculty of Pharmaceutical Sciences, University of Shizuoka.

1. K. L. Rinehart, *Med. Res. Rev.*, 2000, **20**, 1-27.
2. K. L. Rinehart, T. G. Holt, N. L. Fregeau, J. G. Stroth, P. A. Keifer, F. Sun, L. H. Li and D. G. Martin, *J. Org. Chem.*, 1990, **55**, 4512-4515.
3. C. M. Rath, B. Janto, J. Earl, A. Ahmed, F. Z. Hu, L. Hiller, M. Dahlgren, R. Kreft, F. A. Yu, J. J. Wolff, H. K. Kweon, M. A. Christiansen, K. Hakansson, R. M. Williams, G. D. Ehrlich and D. H. Sherman, *ACS Chem. Biol.*, 2011, **6**, 1244-1256.
4. C. Moss, D. H. Green, B. Perez, A. Velasco, R. Henriquez and J. D. McKenzie, *Marine Biology*, 2003, **143**, 99-110.
5. A. E. Perez-Matos, W. Rosado and N. S. Govind, *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology*, 2007, **92**, 155-164.
6. R. Sakai, K. L. Rinehart, Y. Guan and A. H. J. Wang, *Proc. Natl. Acad. Sci. U. S. A.*, 1992, **89**, 11456-11460.
7. B. M. Moore, F. C. Seaman and L. H. Hurley, *J. Am. Chem. Soc.*, 1997, **119**, 5475-5476.
8. M. H. David-Cordonnier, C. Gajate, O. Olmea, W. Laine, J. de la Iglesia-Vicente, C. Perez, C. Cuevas, G. Otero, I. Manzanares, C. Bailly and F. Mollinedo, *Chem. Biol.*, 2005, **12**, 1201-1210.
9. *Application: US*
US Pat., 2003-406997
20040059112, 2004.
10. E. Izbicka, R. Lawrence, E. Raymond, G. Eckhardt, G. Faircloth, J. Jimeno, G. Clark and D. D. Von Hoff, *Ann Oncol*, 1998, **9**, 981-987.
11. H. R. Hendriks, H. H. Fiebig, R. Giavazzi, S. P. Langdon, J. M. Jimeno and G. T. Faircloth, *Ann Oncol*, 1999, **10**, 1233-1240.
12. M. Bonfanti, E. La Valle, J. M. Fernandez Sousa Faro, G. Faircloth, G. Caretti, R. Mantovani and M. D'Incalci, *Anticancer Drug Des.*, 1999, **14**, 179-186.
13. E. J. Martinez, T. Owa, S. L. Schreiber and E. J. Corey, *Proc. Natl. Acad. Sci. U. S. A.*, 1999, **96**, 3496-3501.
14. Y. Takebayashi, P. Pourquier, A. Yoshida, G. Kohlhagen and Y. Pommier, *Proc. Natl. Acad. Sci. U. S. A.*, 1999, **96**, 7196-7201.
15. M. Garcia-Rocha, M. D. Garcia-Gravalos and J. Avila, *Br. J. Cancer*, 1996, **73**, 875-883.
16. J. Zwicker and R. Muller, *Trends Genet.*, 1997, **13**, 3-6.
17. C. Simoens, A. E. Korst, C. M. De Pooter, H. A. Lambrechts, G. G. Pattyn, G. T. Faircloth, F. Lardon and J. B. Vermorken, *Br. J. Cancer*, 2003, **89**, 2305-2311.
18. S. Jin, B. Gorfajn, G. Faircloth and K. W. Scotto, *Proc. Natl. Acad. Sci. U. S. A.*, 2000, **97**, 6775-6779.
19. Y. Pommier, G. Kohlhagen, C. Bailly, M. Waring, A. Mazumder and K. W. Kohn, *Biochemistry*, 1996, **35**, 13303-13309.
20. B. M. Moore, F. C. Seaman, R. T. Wheelhouse and L. H. Hurley, *J. Am. Chem. Soc.*, 1998, **120**, 2490-2491.
21. R. Garcia-Nieto, I. Manzanares, C. Cuevas and F. Gago, *J. Am. Chem. Soc.*, 2000, **122**, 7172-7182.
22. M. Zewail-Foote and L. H. Hurley, *J. Med. Chem.*, 1999, **42**, 2493-2497.
23. M. Zewail-Foote and L. H. Hurley, *J. Am. Chem. Soc.*, 2001, **123**, 6485-6495.
24. F. C. Seaman and L. H. Hurley, *J. Am. Chem. Soc.*, 1998, **120**, 13028-13041.
25. R. Garcia-Nieto, I. Manzanares, C. Cuevas and F. Gago, *J. Med. Chem.*, 2000, **43**, 4367-4369.
26. A. Kanzaki, Y. Takebayashi, X. Q. Ren, H. Miyashita, S. Mori, S. Akiyama and Y. Pommier, *Mol. Cancer Ther.*, 2002, **1**, 1327-1334.
27. J. H. Beumer, T. Buckle, M. Ouwehand, N. E. Franke, L. Lopez-Lazaro, J. H. Schellens, J. H. Beijnen and O. van Tellingen, *Invest. New Drugs*, 2006, **25**, 1-7.
28. S. Feuerhahn, C. Giraudon, M. Martinez-Diez, J. A. Bueren-Calabuig, C. M. Galmarini, F. Gago and J. M. Egly, *Chem. Biol.*, 2011, **18**, 988-999.
29. S. Di Giandomenico, R. Frapolli, E. Bello, S. Uboldi, S. A. Licandro, S. Marchini, L. Beltrame, S. Brich, V. Mauro, E. Tamborini, S. Pilotti, P. G. Casali, F. Grosso, R. Sanfilippo, A. Gronchi, R. Mantovani, R. Gatta, C. M. Galmarini, J. M. Sousa-Faro and M. D'Incalci, *Oncogene*, 2013.
30. P. J. Grohar, L. B. Griffin, C. Yeung, Q. R. Chen, Y. Pommier, C. Khanna, J. Khan and L. J. Helman, *Neoplasia*, 2011, **13**, 145-153.
31. I. Sanchez-Garcia and T. H. Rabbitts, *Proc. Natl. Acad. Sci. U. S. A.*, 1994, **91**, 7869-7873.
32. J. Perez-Losada, B. Pintado, A. Gutierrez-Adan, T. Flores, B. Banares-Gonzalez, J. C. del Campo, J. F. Martin-Martin, E. Battaner and I. Sanchez-Garcia, *Oncogene*, 2000, **19**, 2413-2422.
33. L. A. Owen, A. A. Kowalewski and S. L. Lessnick, *PLoS One*, 2008, **3**, e1965.
34. Y. Takebayashi, P. Pourquier, D. B. Zimonjic, K. Nakayama, S. Emmert, T. Ueda, Y. Urasaki, A. Kanzaki, S. I. Akiyama, N. Popescu, K. H. Kraemer and Y. Pommier, *Nat Med*, 2001, **7**, 961-966.
35. A. B. Herrero, C. Martin-Castellanos, E. Marco, F. Gago and S. Moreno, *Cancer Res.*, 2006, **66**, 8155-8162.
36. G. Damia, S. Silvestri, L. Carrassa, L. Filiberti, G. T. Faircloth, G. Liberi, M. Foiani and M. D'Incalci, *Int. J. Cancer*, 2001, **92**, 583-588.
37. Y. Takebayashi, F. Goldwasser, Y. Urasaki, G. Kohlhagen and Y. Pommier, *Clin Cancer Res*, 2001, **7**, 185-191.
38. D. G. Soares, A. E. Escargueil, V. Poindessous, A. Sarasin, A. de Gramont, D. Bonatto, J. A. Henriques and A. K. Larsen, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 13062-13067.
39. M. Tavecchio, M. Simone, E. Erba, I. Chiolo, G. Liberi, M. Foiani, M. D'Incalci and G. Damia, *Eur J Cancer*, 2008, **44**, 609-618.
40. M. Zewail-Foote, V. S. Li, H. Kohn, D. Bearss, M. Guzman and L. H. Hurley, *Chem. Biol.*, 2001, **8**, 1033-1049.
41. J. Guirouilh-Barbat, C. Redon and Y. Pommier, *Mol. Biol. Cell*, 2008, **19**, 3969-3981.
42. Y. Shiloh, *Trends Biochem. Sci.*, 2006, **31**, 402-410.
43. G. Germano, R. Frapolli, C. Belgiovine, A. Anselmo, S. Pesce, M. Liguori, E. Erba, S. Uboldi, M. Zucchetti, F. Pasqualini, M. Nebuloni, N. van Rooijen, R. Mortarini, L. Beltrame, S. Marchini, I. Fuso Nerini, R. Sanfilippo, P. G. Casali, S. Pilotti, C. M. Galmarini, A. Anichini, A. Mantovani, M. D'Incalci and P. Allavena, *Cancer Cell*, 2013, **23**, 249-262.

44. T. Arai, K. Takahashi, S. Nakahara and A. Kubo, *Experientia*, 1980, **36**, 1025-1027.
45. H. Irschik, W. Trowitzsch-Kienast, K. Gerth, G. Hofle and H. Reichenbach, *J Antibiot (Tokyo)*, 1988, **41**, 993-998.
46. Y. Ikeda, Y. Shimada, K. Honjo, T. Okumoto and T. Munakata, *J Antibiot (Tokyo)*, 1983, **36**, 1290-1294.
47. D. Kluepfel, H. A. Baker, G. Piattoni, S. N. Sehgal, A. Sidorowicz, K. Singh and C. Vezina, *J. Antibiot.*, 1975, **28**, 497-502.
48. F. Tomita, K. Takahashi and K. Shimizu, *J. Antibiot.*, 1983, **36**, 463-467.
49. K. Takahashi and F. Tomita, *J. Antibiot.*, 1983, **36**, 468-470.
50. R. G. Kerr and N. F. Miranda, *J. Nat. Prod.*, 1995, **58**, 1618-1621.
51. C. Peng, J. Y. Pu, L. Q. Song, X. H. Jian, M. C. Tang and G. L. Tang, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 8540-8545.
52. S. Jeedigunta, J. M. Krenisky and R. G. Kerr, *Tetrahedron*, 2000, **56**, 3303-3307.
53. L. Li, W. Deng, J. Song, W. Ding, Q. F. Zhao, C. Peng, W. W. Song, G. L. Tang and W. Liu, *J. Bacteriol.*, 2008, **190**, 251-263.
54. A. Pospiech, B. Cluzel, J. Bietenhader and T. Schupp, *Microbiology*, 1995, **141 (Pt 8)**, 1793-1803.
55. A. Velasco, P. Acebo, A. Gomez, C. Schleissner, P. Rodriguez, T. Aparicio, S. Conde, R. Munoz, F. de la Calle, J. L. Garcia and J. M. Sanchez-Puelles, *Mol. Microbiol.*, 2005, **56**, 144-154.
56. C. Y. Fu, M. C. Tang, C. Peng, L. Li, Y. L. He, W. Liu and G. L. Tang, *J Microbiol Biotechnol*, 2009, **19**, 439-446.
57. R. Finking and M. A. Marahiel, *Annu. Rev. Microbiol.*, 2004, **58**, 453-488.
58. M. A. Fischbach and C. T. Walsh, *Chem Rev*, 2006, **106**, 3468-3496.
59. K. Koketsu, A. Minami, K. Watanabe, H. Oguri and H. Oikawa, *Methods Enzymol.*, 2012, **516**, 79-98.
60. M. Arroyo, I. de la Mata, C. Acebal and M. P. Castillon, *Appl. Microbiol. Biotechnol.*, 2003, **60**, 507-514.
61. C. Cuevas, M. Pérez, M. J. Martín, J. L. Chicharro, C. Fernández-Rivas, M. Flores, A. Francesch, P. Gallego, M. Zarzuelo, F. de la Calle, J. García, C. Polanco, I. Rodríguez and I. Manzanares, *Org. Lett.* 2000, **2**, 2545-2548.
62. E. J. Corey, D. Y. Gin and R. S. Kania, *J. Am. Chem. Soc.*, 1996, **118**, 9202-9203.
63. E. J. Martinez and E. J. Corey, *Org. Lett.* 2000, **2**, 993-996.
64. D. H. R. Barton, J. D. Elliott and S. D. Gero, *Journal of the Chemical Society-Perkin Transactions I*, 1982, 2085-2090.
65. A. Endo, A. Yanagisawa, M. Abe, S. Tohma, T. Kan and T. Fukuyama, *J. Am. Chem. Soc.*, 2002, **124**, 6552-6554.
66. S. Tohma, A. Endo, T. Kan and T. Fukuyama, *Synlett*, 2001, 1179-1181.
67. T. Hayashi, M. Konishi, Y. Kobori, M. Kumada, T. Higuchi and K. Hirotsu, *J. Am. Chem. Soc.*, 1984, **106**, 158-163.
68. G. W. Gokel, R. P. Widera and W. P. Weber, *Org. Synth.*, 1988, **50-9**, 232-235.
69. M. J. Burk, J. E. Feaster, W. A. Nugent and R. L. Harlow, *J. Am. Chem. Soc.*, 1993, **115**, 10125-10138.
70. G. Gokel, G. Ludke and I. Ugi, *Isonitrile Chemistry*, Academic Press, New York, 1971.
71. J. T. Link and L. E. Overman, *Metal-Catalyzed Cross-Coupling Reactions*, Wiley-VCH Verlag GmbH, Weinheim, Germany, 1998.
72. M. Shibasaki, C. D. J. Boden and A. Kojima, *Tetrahedron*, 1997, **53**, 7371-7395.
73. S. Brase and A. De Meijere, *Metal-Catalyzed Cross-Coupling Reactions*, Wiley-VCH Verlag GmbH, Weinheim, Germany, 1998.
74. L. Neuville, A. Bigot, M. E. T. H. Dau and J. P. Zhu, *J. Org. Chem.*, 1999, **64**, 7638-7642.
75. N. Miyaura and A. Suzuki, *Chemical Reviews*, 1995, **95**, 2457-2483.
76. M. De Paolis, X. C. Chen and J. P. Zhu, *Synlett*, 2004, 729-731.
77. E. J. Corey, F. Xu and M. C. Noe, *J. Am. Chem. Soc.*, 1997, **119**, 12414-12415.
78. B. Lygo and P. G. Wainwright, *Tetrahedron Lett.*, 1997, **38**, 8595-8598.
79. M. J. Odonnell, W. D. Bennett and S. D. Wu, *J. Am. Chem. Soc.*, 1989, **111**, 2353-2355.
80. P. Garner and J. M. Park, *J. Org. Chem.*, 1987, **52**, 2361-2364.
81. P. Garner and J. M. Park, *J. Org. Chem.*, 1988, **53**, 2979-2984.
82. P. Garner, J. M. Park and E. Malecki, *J. Org. Chem.*, 1988, **53**, 4395-4398.
83. E. D. Cox and J. M. Cook, *Chemical Reviews*, 1995, **95**, 1797-1842.
84. G. Massiot and T. Mulamba, *Journal of the Chemical Society-Chemical Communications*, 1983, 1147-1149.
85. P. D. Bailey, S. P. Hollinshead, N. R. Mclay, K. Morgan, S. J. Palmer, S. N. Prince, C. D. Reynolds and S. D. Wood, *Journal of the Chemical Society-Perkin Transactions I*, 1993, 431-439.
86. S. Zheng, C. Chan, T. Furuuchi, B. J. Wright, B. Zhou, J. Guo and S. J. Danishefsky, *Angew Chem Int Ed Engl*, 2006, **45**, 1754-1759.
87. A. K. Sinhababu, A. K. Ghosh and R. T. Borchardt, *J. Med. Chem.*, 1985, **28**, 1273-1279.
88. R. M. Williams, P. P. Ehrlich, W. X. Zhai and J. Hendrix, *J. Org. Chem.*, 1987, **52**, 2615-2617.
89. A. Fujii, S. Hashiguchi, N. Uematsu, T. Ikariya and R. Noyori, *J. Am. Chem. Soc.*, 1996, **118**, 2521-2522.
90. C. Chan, R. Heid, S. P. Zheng, J. S. Guo, B. S. Zhou, T. Furuuchi and S. J. Danishefsky, *J. Am. Chem. Soc.*, 2005, **127**, 4596-4598.
91. D. Fishlock and R. M. Williams, *J. Org. Chem.*, 2008, **73**, 9594-9600.
92. M. Sakaitani and Y. Ohfuné, *J. Org. Chem.*, 1990, **55**, 870-876.
93. W. Jin and R. M. Williams, *Tetrahedron Lett.*, 2003, **44**, 4635-4639.
94. H. H. Hussain, G. Babic, T. Durst, J. S. Wright, M. Flueraru, A. Chichirau and L. L. Chepelev, *J. Org. Chem.*, 2003, **68**, 7023-7032.
95. X. C. Chen, J. C. Chen, M. De Paolis and J. P. Zhu, *J. Org. Chem.*, 2005, **70**, 4397-4408.
96. M. Weigl and B. Wunsch, *Tetrahedron*, 2002, **58**, 1173-1183.
97. M. P. Doyle and W. J. Bryker, *J. Org. Chem.*, 1979, **44**, 1572-1574.
98. A. Cioffi and A. Italiano, *Expert Opin Drug Metab Toxicol*, 2012, **8**, 113-122.
99. D. P. Ryan, J. G. Supko, J. P. Eder, M. V. Seiden, G. Demetri, T. J. Lynch, A. J. Fischman, J. Davis, J. Jimeno and J. W. Clark, *Clin Cancer Res*, 2001, **7**, 231-242.
100. A. Taamma, J. L. Misset, M. Riofrio, C. Guzman, E. Brain, L. Lopez Lazaro, H. Rosing, J. M. Jimeno and E. Cvitkovic, *J Clin Oncol*, 2001, **19**, 1256-1265.
101. C. Twelves, K. Hoekman, A. Bowman, J. B. Vermorken, A. Anthony, J. Smyth, C. van Kesteren, J. H. Beijnen, J. Uiters, J. Wanders, J. Gomez, C. Guzman, J. Jimeno and A. Hanauske, *Eur J Cancer*, 2003, **39**, 1842-1851.

102. A. Le Cesne, A. Yovine, J. Y. Blay, S. Delalogue, R. G. Maki, J. L. Misset, P. Frontelo, A. Nieto, J. J. Jiao and G. D. Demetri, *Invest. New Drugs*, 2012, **30**, 1193-1202.
103. G. D. Demetri, S. P. Chawla, M. von Mehren, P. Ritch, L. H. Baker, J. Y. Blay, K. R. Hande, M. L. Keohan, B. L. Samuels, S. Schuetze, C. Lebedinsky, Y. A. Elsayed, M. A. Izquierdo, J. Gomez, Y. C. Park and A. Le Cesne, *J Clin Oncol*, 2009, **27**, 4188-4196.
104. N. Takahashi, W. W. Li, D. Banerjee, K. W. Scotto and J. R. Bertino, *Clin Cancer Res*, 2001, **7**, 3251-3257.
105. E. Erba, I. Muratore, G. Tiozzo, P. Ubezio, T. Colombo, I. Nicoletti, R. Giavazzi, M. Zucchetti, G. Damia, D. Meco, R. Riccardi, G. T. Faircloth, J. Jimeno and M. D'Incalci, *Clinical Cancer Research*, 2001, **7**, 3735s-3735s.
106. K. Scotlandi, S. Perdichizzi, M. C. Manara, M. Serra, S. Benini, V. Cerisano, R. Strammiello, M. Mercuri, G. Reverter-Branchat, G. Faircloth, M. D'Incalci and P. Picci, *Clinical Cancer Research*, 2002, **8**, 3893-3903.
107. N. Takahashi, W. W. Li, D. Banerjee, Y. B. Guan, Y. Wada-Takahashi, M. F. Brennan, T. C. Chou, K. W. Scotto and J. R. Bertino, *Cancer Res.*, 2002, **62**, 6909-6915.
108. D. Meco, T. Colombo, P. Ubezio, M. Zucchetti, M. Zaffaroni, A. Riccardi, G. Faircloth, J. Jose, M. D'Incalci and R. Riccardi, *Cancer Chemotherapy and Pharmacology*, 2003, **52**, 131-138.
109. E. Erba, T. Colombo, P. Ubezio, L. Ferrarese, G. Matera, C. Falcioni, G. Faircloth, J. Jimeno and M. D'Incalci, *Clinical Cancer Research*, 2003, **9**, 6264s-6264s.
110. M. D'Incalci, T. Colombo, P. Ubezio, I. Nicoletti, R. Giavazzi, E. Erba, L. Ferrarese, D. Meco, R. Riccardi, C. Sessa, E. Cavallini, J. Jimeno and G. T. Faircloth, *European Journal of Cancer*, 2003, **39**, 1920-1926.
111. B. J. Monk, T. J. Herzog, S. B. Kaye, C. N. Krasner, J. B. Vermorken, F. M. Muggia, E. Pujade-Lauraine, A. S. Lisyanskaya, A. N. Makhson, J. Rolski, V. A. Gorbounova, P. Ghatage, M. Bidzinski, K. Shen, H. Y. Ngan, I. B. Vergote, J. H. Nam, Y. C. Park, C. A. Lebedinsky and A. M. Poveda, *J Clin Oncol*, 2010, **28**, 3107-3114.
112. C. N. Krasner, A. Poveda, T. J. Herzog, J. B. Vermorken, S. B. Kaye, A. Nieto, P. L. Claret, Y. C. Park, T. Parekh and B. J. Monk, *Gynecol Oncol*, 2012, **127**, 161-167.
113. C. Lebedinsky, J. Gomez, Y. C. Park, A. Nieto, A. Soto-Matos, T. Parekh, V. Alfaro, E. Roy, P. Lardelli and C. Kahatt, *Cancer Chemother Pharmacol*, 2011, **68**, 1223-1231.
114. S. Baruchel, A. Pappo, M. Krailo, K. S. Baker, B. Wu, D. Villaluna, M. Lee-Scott, P. C. Adamson and S. M. Blaney, *Eur J Cancer*, 2012, **48**, 579-585.
115. L. Lau, J. G. Supko, S. Blaney, L. Hershon, N. Seibel, M. Krailo, W. Qu, D. Malkin, J. Jimeno, M. Bernstein and S. Baruchel, *Clin Cancer Res*, 2005, **11**, 672-677.
116. *Drugs R D*, 2006, **7**, 317-328.