



Organic & Biomolecular Chemistry

Accepted Manuscript



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

FULL PAPER

In(OTf)₃-Catalysed One-pot Versatile Pyrrole Synthesis through Domino Annulation of α -Oxoketene-*N,S*-acetals with NitroolefinsAbhijeet Srivastava,¹ Gaurav Shukla,¹ Anugula Nagaraju,¹ Girijesh Kumar Verma,¹ Keshav Raghuvanshi,² Raymond C. F. Jones³ and Maya Shankar Singh^{1*}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

In(OTf)₃-catalyzed robust and sustainable one-pot access to previously unknown and synthetically demanding polysubstituted pyrroles via [3+2] annulation of α -oxoketene-*N,S*-acetals with β -nitrostyrenes has been achieved under solvent-free conditions. The merit of this domino Michael addition/cyclization sequence is highlighted by its operational simplicity, short reaction time (5–10 min), good to excellent yields, tolerance of a large variety of functional groups, and efficiency of producing two new (C–C and C–N) bonds and one highly functionalized pyrrole ring in a single operation, which make it an ideal alternative to existing methods.

Introduction

Pyrroles, one of the prominent and ubiquitous privileged *N*-heterocyclic pharmacophore, exhibit diverse biological and therapeutic activities,¹ and constitute the core structural motif of numerous natural products,² medicinal agents,³ catalysts and conducting polymers.⁴ Among the various pharmacological profiles of pyrrole derivatives, their anti-bacterial,^{5a} anti-inflammatory,^{5b,c} anti-oxidant,^{5d,e} anti-tumor,^{5f-h} anti-fungal,^{5i-k} and immune suppressant activities^{5l} seem to be well documented. Chart 1 constitute notable privileged medicinal scaffolds having multiple substituted pyrroles.⁶ Furthermore, pyrrole based derivatives have broad applications in molecular optics,^{7a} electronics,^{7b} and as gas sensors for organic compounds.^{7c} Moreover, they are also widely employed as versatile building blocks in organic synthesis.⁸





Chart 1. Selected examples of pyrrole-based pharmaceutical compounds.


Traditional pyrrole syntheses^{9–11} generally require the initial preparation of an intermediate that subsequently undergoes a cyclization reaction. In general, pyrroles are prepared by cycloaddition ([4+1] and [3+2]) reactions¹² or intramolecular cyclization reactions.¹³ Albeit the reported approaches are useful tools, most of them suffer from significant limitations such as multiple steps, functional group compatibility, pre-functionalized substrates, expensive catalysts/reagents, long reaction times and harsh reaction conditions. Recently, a number of strategies^{14–18} are employed for the synthesis of pyrroles. Rueping *et al.*¹⁹ reported the synthesis of pyrroles through domino reaction of enaminones with β -bromonitrostyrenes (Scheme 1). The above published protocols not only showcase the power of advances in organic methodology but also provide proof of the vibrant activity in this field. In spite of these outstanding efforts, it is still challenging to prepare polysubstituted pyrroles possessing diverse substituents directly from readily available precursors in a single synthetic operation with environmentally benign fashion.



Scheme 1. Synthetic strategies for the construction of pyrroles.


A possible retrosynthetic analysis for the construction of pyrroles is given in Scheme 2, which comprises the use of simple and easily available synthons such as binucleophilic α -oxoketene-*N,S*-acetals and bielectrophilic β -nitrostyrenes. In continuation of our ongoing research for the development of new methodologies for various heterocyclic systems by exploiting the

reactions of ketene-*S,S*-acetals,^{20,21} we envisioned to explore α -oxoketene-*N,S*-acetals for the construction of highly substituted pyrroles for the first time.



Scheme 2. A retrosynthetic disconnection approach for the synthesis of pyrroles.

Herein, we report a rapid and efficient one-pot synthesis of highly substituted pyrroles via coupling of α -oxoketene-*N,S*-acetals with (*E*)- β -nitrostyrenes catalyzed by Lewis acid under solvent-free conditions at moderate temperature (Scheme 3).



Scheme 3. One-pot synthesis of polysubstituted pyrroles.

Results and Discussion

The feasibility of the coupling reaction was examined using α -oxoketene-*N,S*-acetal **1a** and (*E*)- β -nitrostyrene **2a** as the test substrates under different conditions (Table 1). Initially, the equimolar amount of **1a** and **2a** was treated in EtOH at room temperature without any catalyst. The work up of the reaction afforded 3-benzoyl-2-(methylthio)-1,4-diphenylpyrrole **3aa** in 55% yield after 72 h (Table 1, entry 1). An attempt to increase the yield of **3aa**, the above model reaction was further investigated at 55 °C and in refluxing EtOH. The work up of the reaction provided the desired pyrrole **3aa** in 64% and 68% yields respectively, reducing the time significantly (Table 1, entries 2 and 3). A brief screen of other protic solvents like MeOH and H₂O afforded **3aa** in 55% and 50% yields, respectively taking more than double time (Table 1, entries 4 and 5). Because of the increasing public concern for the harmful effects of organic solvents over environment and human body, we carried out the above model reaction under solvent-free conditions at different temperatures. Notably, the reaction proceeded smoothly at 55 °C furnishing the desired pyrroles **3aa** in 55% yield (Table 1, entry 6). Higher temperatures reduced the yield of **3aa** due to decomposition of the substrates (Table 1, entries 7 and 8).

With a view to optimize the reaction conditions for maximum yield and the lesser reaction time, the model reaction was carried out in the presence of Lewis acid (LA) catalysts such as FeCl₃, AlCl₃, InBr₃, CuBr₂, SbCl₃, SnCl₄ and In(OTf)₃ under solvent-free conditions at 55 °C (Table 1, entries 9–15). A radical change

in reaction time was observed while using various Lewis acids, and to our great pleasure 10 mol % of In(OTf)₃ catalyzed the reaction efficiently and afforded the desired pyrrole **3aa** in 92% yield within 5 min (Table 1, entry 15). Next, reduction in the amount of LA catalyst resulted in lower yield of the desired product, and increasing the catalyst loading did not show any improvement (Table 1, entries 16 and 17). Finally, when the model reaction was carried out in refluxing ethanol in the presence of 10 mol % of In(OTf)₃, **3aa** was obtained in 74% yield in 5 h (Table 1, entry 18). Consequently, the optimum reaction condition was achieved by employing equimolar amount of **1a** and **2a** in the presence of 10 mol % of In(OTf)₃ at 55 °C under solvent-free conditions.

Table 1. Optimization studies.

Entry	Solvent	Catalyst (mol%)	T (°C)	Time	Yield ^a (%)
1	EtOH	None	rt	72 h	55
2	EtOH	None	55	11 h	64
3	EtOH	None	reflux	9 h	68
4	MeOH	None	reflux	20 h	55
5	H ₂ O	None	reflux	22 h	50
6	None	None	55	8 h	55
7	None	None	80	6 h	40
8	None	None	100	4 h	15
9	None	FeCl ₃ (10)	55	30 min	80
10	None	AlCl ₃ (10)	55	20 min	55
11	None	InBr ₃ (10)	55	15 min	82
12	None	CuBr ₂ (10)	55	40 min	63
13	None	SbCl ₃ (10)	55	20 min	85
14	None	SnCl ₄ (10)	55	15 min	82
15	None	In(OTf) ₃ (10)	55	5 min	92
16	None	In(OTf) ₃ (5)	55	10 min	76
17	None	In(OTf) ₃ (15)	55	5 min	92
18	EtOH	In(OTf) ₃ (10)	reflux	5 min	74

^aIsolated pure yields.

Having found an optimum reaction condition, we explored the scope of the substrates and functional group tolerance by synthesizing twenty six new tetra- and penta-substituted pyrroles with high functionalities from different α -oxoketene-*N,S*-acetals (**1a-j**) and (*E*)- β -nitrostyrenes (**2a-h**) as shown in Tables 2 & 3 and Schemes 4 and 5. In all products 2-thiomethyl substituent that stem from α -oxoketene-*N,S*-acetals was kept constant. The electron-donating groups at R³ of nitrostyrenes (**2b**, **2c**) gave slightly higher yield of product (Table 2, entries 2, 3, 7, 9, 11, 14, 15, and 18) as compared to electron-withdrawing groups at R³ of nitrostyrenes (**2d**, **2e**) (Table 2, entries 4, 5, 8, 12, and 16). Similarly, the electron-donating group at R¹ of α -oxoketene-*N,S*-acetal (**1c**) provides significantly higher yield as compared with the electron-withdrawing group (**1d**) (Table 2, entry 6 vs. entry 9).

To illustrate the broad synthetic utility and generality of our developed methodology, further we treated α -oxoketene-*N,S*-acetal bearing extended aromatic ring biphenyl (**1e**), and heteroaryl groups like 2-furyl (**1f**) and 2-thienyl (**1g**) with different (*E*)- β -nitrostyrenes separately under the above

optimized conditions to get the corresponding tetrasubstituted pyrroles in 68–87% yields (Table 2, entries 10–18).

Table 2. Scope of substrates for tetrasubstituted pyrroles.

Entry	R ¹	R ³	Product	Time (min)	Yield ^a (%)
1	C ₆ H ₅ (1a)	C ₆ H ₅ (2a)	3aa	5	92
2	C ₆ H ₅ (1a)	4-MeC ₆ H ₄ (2b)	3ab	5	94
3	C ₆ H ₅ (1a)	4-OMeC ₆ H ₄ (2c)	3ac	5	95
4	C ₆ H ₅ (1a)	4-NO ₂ C ₆ H ₄ (2d)	3ad	8	72
5	4-MeC ₆ H ₄ (1b)	4-NO ₂ C ₆ H ₄ (2d)	3bd	7	73
6	4-OMeC ₆ H ₄ (1c)	C ₆ H ₅ (2a)	3ca	5	89
7	4-OMeC ₆ H ₄ (1c)	4-MeC ₆ H ₄ (2b)	3cb	5	90
8	4-OMeC ₆ H ₄ (1c)	4-NO ₂ C ₆ H ₄ (2d)	3cd	7	70
9	4-CF ₃ C ₆ H ₄ (1d)	4-MeC ₆ H ₄ (2b)	3db	10	62
10	4-PhC ₆ H ₄ (1e)	C ₆ H ₅ (2a)	3ea	6	78
11	4-PhC ₆ H ₄ (1e)	4-MeC ₆ H ₄ (2b)	3eb	6	82
12	4-PhC ₆ H ₄ (1e)	4-FC ₆ H ₄ (2e)	3ee	8	70
13	2-furyl(1f)	C ₆ H ₅ (2a)	3fa	8	74
14	2-furyl(1f)	4-MeC ₆ H ₄ (2b)	3fb	6	80
15	2-furyl(1f)	4-OMeC ₆ H ₄ (2c)	3fc	5	87
16	2-furyl(1f)	4-NO ₂ C ₆ H ₄ (2d)	3fd	9	68
17	2-thienyl(1g)	C ₆ H ₅ (2a)	3ga	6	72
18	2-thienyl(1g)	4-MeC ₆ H ₄ (2b)	3gb	6	84

^aIsolated pure yields.


Scope of α -oxoketene-*N,S*-acetal is not limited to R² as phenyl only but various substituents have also been screened as shown in Table 3. Methyl, *n*-butyl and cyclohexyl substituents also provided good yields under similar reaction conditions (Table 3).

Table 3. Scope of *N*-substituted α -oxoketene-*N,S*-acetals.

Entry	R ¹ /R ²	R ³	Product	Time (min)	Yield ^a (%)
1	4-BrC ₆ H ₄ /Me (1h)	C ₆ H ₅ (2a)	3ha	5	82
2	3-OHC ₆ H ₄ / <i>n</i> -butyl (1i)	4-OMeC ₆ H ₄ (2c)	3ic	8	77
3	C ₆ H ₅ /cyclohexyl (1j)	4-OMeC ₆ H ₄ (2c)	3jc	5	84

^aIsolated pure yields.

In order to further derivatise the pyrrole moiety, we prepared functionally rich nitrostyrenes **2f** and **2g** from 2-ethynylbenzaldehyde and *p*-methoxycinnamaldehyde, respectively. Treatment of α -oxoketene-*N,S*-acetals **1c** and **1e** with the nitrostyrene **2f** provided pyrroles **3cf** and **3ef** in 81% and 72% yields, respectively. Treatment of α -oxoketene-*N,S*-acetal **1a** with (*E*)- β -nitrostyrene **2g** gave pyrrole **3ag** in 67% yield (Scheme 4). These three examples also exhibited functional group tolerance towards the Lewis acid catalyst.




Entry	R ¹	R ³	Product	Time	Yield ^a (%)
1	4-OMeC ₆ H ₄ (1c)	2-CH=CC ₆ H ₄ (2f)	3cf	5 min	81
2	4-PhC ₆ H ₄ (1e)	2-CH=CC ₆ H ₄ (2f)	3ef	6 min	72
3	C ₆ H ₅ (1a)	4-OMeC ₆ H ₄ CH=CH (2g)	3ag	8 min	67

Scheme 4. Substrate functionalization on nitrostyrene.

^a Isolated pure yields.

Next, we turned our attention to expand the diversity of this protocol for the synthesis of pentasubstituted pyrroles. Thus, upon treatment of (*E*)- β -methyl- β -nitrostyrene **2h** with α -oxoketene-*N,S*-acetals **1d** and **1f** under previously described one-pot optimised reaction conditions, furnished pentasubstituted pyrroles **3dh** and **3fh** in 66% and 79% yields, respectively (Scheme 5).




Entry	R ¹	R ³	Product	Time	Yield ^a (%)
1	4-CF ₃ C ₆ H ₄ (1d)	4-OMeC ₆ H ₄ (2h)	3dh	8 min	66
2	2-Furyl (1f)	4-OMeC ₆ H ₄ (2h)	3fh	5 min	79

Scheme 5. Synthesis of pentasubstituted pyrroles.

^a Isolated pure yields.

Considering the high electronic efficacy exerted by the various electron-rich and electron-deficient substituents, we performed intermolecular competition experiments with two different α -oxoketene-*N,S*-acetals, first containing electron-donating group **1c** and second electron-withdrawing group **1d** with simple β -nitrostyrene **2a** in an equimolar ratio. It has been observed that



Scheme 6. Intermolecular competition reactions.

the product pyrrole **3ca** derived from α -oxoketene-*N,S*-acetal **1c** was major (70%) and **3da** formed from **1d** was minor (30%) revealing electron-rich substrate to be more readily functionalized (Scheme 6a). In next attempt, we performed the reaction of α -oxoketene-*N,S*-acetal **1a** with two different (*E*)- β -nitrostyrenes **2c** and **2d** in an equimolar ratio (Scheme 6b). Exclusively, we found only one pyrrole **3ac** in 92% yield derived from nitrostyrene with electron-donating group **2c**. Unused starting material **2d** was recovered in 90% yield highlighting the electron-rich (*E*)- β -nitrostyrene to be converted more preferentially.

The structures of all the newly synthesized pyrroles **3** were deduced from their satisfactory spectral (IR, ^1H & ^{13}C NMR, and HRMS) studies and unequivocally established by the X-ray single crystal diffraction analysis of two representative compounds **3ga**¹⁵ and **3eb** (Figure 1).²² The mass spectra of these compounds displayed molecular ion peaks at the appropriate *m/z* values.





Figure 1. ORTEP diagrams of **3ga** and **3eb**.

On the basis of the above experimental results together with the related reports, a plausible reaction scenario for domino [3+2] heteroannulation is outlined in Scheme 7. We speculate that the reaction occurs as a tandem Michael addition/intramolecular ring closure/elimination process. The first step in the mechanism is believed to be the Michael addition of α -oxoketene-*N,S*-acetal **1** through its Lewis acid activated enolate form **A** to β -nitrostyrene **2** to generate open-chain Michael adduct intermediate **B**. Next, the Michael adduct intermediate **B** undergoes successive intramolecular *N*-cyclization followed by elimination of H_2O and nitroxyl (HNO) via intermediate **C** leading to desired pyrrole **3**.



Scheme 7. Plausible mechanism for formation of pyrroles **3**.

Conclusions

In summary, we have designed and developed an operationally simple, highly efficient one-pot direct method for the synthesis of

tetra- and pentasubstituted pyrroles under solvent-free mild reaction conditions. Further, we have shown α -oxoketene-*N,S*-acetals to be suitable partners in (3 + 2) annulation with β -nitrostyrenes. $\text{In}(\text{OTf})_3$ worked very efficiently with various functional group substituted α -oxoketene-*N,S*-acetals and β -nitrostyrenes yielding highly functionalized pyrroles in good to excellent yields. The scope and diversity of the tolerated substrates in this work is rather broad in comparison to the reported ones. These structurally unique molecules synthesized in an environmentally benign manner may reveal potential application in future drug discovery and development. A possible reaction mechanism was proposed to account for the coupling reaction.

Experimental Section

General procedure for synthesis of 3

A 25 mL round bottom flask equipped with a magnetic stirrer was charged with α -oxoketene-*N,S*-acetal **1** (1.0 mmol), β -nitrostyrene **2** (1.0 mmol) and 10 mol% of $\text{In}(\text{OTf})_3$. The reaction mixture was heated on oil bath at 55 °C for the stipulated period of time (monitored through TLC). After completion of the reaction, the reaction mixture was extracted with EtOAc, washed with water (2×20 mL) followed by brine (1×20 mL), and dried over anhydrous sodium sulphate. The solvent was evaporated under vacuum and the crude residue thus obtained was purified by silica gel column chromatography to give pure pyrrole **3**.

60 3-Benzoyl-2-(methylthio)-1,4-diphenylpyrrole (3aa). Yellow sticky solid. $R_f = 0.55$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 7.88$ (d, $J = 7.2$ Hz, 2H), 7.51–7.38 (m, 5H), 7.31–7.04 (m, 9H), 2.02 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 194.1$, 138.8, 138.5, 133.9, 132.6, 129.9, 129.0, 128.2, 128.1, 128.0, 127.6, 126.5, 126.2, 122.8, 20.8. IR (KBr) (ν_{max} /cm $^{-1}$): 2912, 1670, 1222. HRMS (EI) calcd for $\text{C}_{24}\text{H}_{19}\text{NOS}, m/z: 369.1187$, found 369.1193 [M^+].

3-Benzoyl-2-(methylthio)-1-phenyl-4-(*p*-tolyl)pyrrole (3ab). Yellow sticky solid. $R_f = 0.55$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 7.88$ (d, $J = 7.2$ Hz, 2H), 7.51–7.24 (m, 8H), 7.15–7.10 (m, 2H), 6.96 (d, $J = 7.8$ Hz, 2H), 2.23 (s, 3H), 2.01 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 194.3$, 138.9, 138.5, 135.8, 132.5, 130.9, 129.9, 128.9, 128.7, 128.3, 128.08, 128.00, 127.4, 127.0, 126.5, 126.19, 126.11, 125.2, 122.6, 20.9, 20.7. IR (KBr) (ν_{max} /cm $^{-1}$): 2920, 1656, 1595, 1496, 1225. HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{21}\text{NOS}, m/z: 383.1344$, found 384.1399 [$(\text{M}+\text{H})^+$].

3-Benzoyl-4-(4-methoxyphenyl)-2-(methylthio)-1-phenylpyrrole (3ac). Yellow sticky solid. $R_f = 0.48$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 7.92$ (d, $J = 7.2$ Hz, 2H), 7.55–7.45 (m, 6H), 7.35 (dd, $J = 7.8$, 7.2 Hz, 2H), 7.22 (d, $J = 8.7$ Hz, 2H), 7.11 (s, 1H), 6.75 (d, $J = 8.7$ Hz, 2H), 3.74 (s, 3H), 2.06 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 194.3$, 158.1, 138.8, 138.5, 132.5, 129.9, 128.9, 128.7, 127.9, 126.5, 126.1, 125.9, 122.3, 113.7, 55.1, 20.7. IR (KBr) (ν_{max} /cm $^{-1}$): 2922, 1650, 1340, 1229. HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{21}\text{NO}_2\text{S}, m/z: 399.9293$, found 438.1102 [$(\text{M}+\text{K})^+$].

3-Benzoyl-2-(methylthio)-4-(4-nitrophenyl)-1-phenylpyrrole (3ad).

Yellow (crayola) sticky solid. $R_f = 0.50$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 8.03$ (d, $J = 8.4$ Hz, 2H), 7.87 (d, $J = 7.5$ Hz, 2H), 7.52–7.45 (m, 9H), 7.39 – 7.32 (m, 2H), 2.02 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 193.6$, 146.0, 140.8, 138.4, 138.3, 133.1, 129.9, 129.2, 128.7, 128.6, 128.3, 128.1, 127.8, 126.5, 124.2, 124.1, 123.8, 20.6. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 3064, 2922, 1651, 1595, 1513, 1341, 1228. MS calcd for $\text{C}_{24}\text{H}_{17}\text{N}_2\text{O}_3\text{S}$, m/z : 414.1038, found 415.2 [(M+H) $^+$].

10 2-(methylthio)-4-(4-nitrophenyl)-1-phenyl-3-(*p*-tolyl)pyrrole (3bd).

Yellow (crayola) sticky solid. $R_f = 0.6$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 8.03$ (d, $J = 8.4$ Hz, 2H), 7.79 (d, $J = 7.8$ Hz, 2H), 7.52 (m, 6H), 7.39 (d, $J = 8.4$ Hz, 2H), 7.16 (d, $J = 7.5$ Hz, 2H), 2.36 (s, 3H), 2.00 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 193.4$, 145.9, 144.2, 140.8, 138.4, 135.7, 130.1, 129.19, 129.11, 128.9, 128.5, 127.6, 127.5, 126.5, 123.9, 123.8, 21.7, 20.6. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 2931, 1653, 1559, 1495, 1340, 1247. MS calcd for $\text{C}_{25}\text{H}_{20}\text{N}_2\text{O}_3\text{S}$, m/z : 428.1195, found 429.2 [(M+H) $^+$].

20 3-(4-Methoxybenzoyl)-2-(methylthio)-1,4-diphenylpyrrole (3ca).

White solid, mp = 135–137 °C. $R_f = 0.48$ (19:1 hexane/ethyl acetate). ^1H NMR (400 MHz, CDCl_3): $\delta = 7.92$ (d, $J = 8.8$ Hz, 2H), 7.54 – 7.46 (m, 5H), 7.32 – 7.10 (m, 5H), 6.84 (d, $J = 8.8$ Hz, 2H), 3.82 (s, 3H), 2.05 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 193.1$, 163.3, 138.9, 134.0, 132.4, 131.5, 129.1, 129.0, 128.4, 128.1, 127.4, 126.6, 126.3, 125.8, 125.6, 122.7, 113.3, 55.4, 20.8. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 3019, 2920, 1650, 1595, 1494, 1262. HRMS (EI) calcd for $\text{C}_{25}\text{H}_{21}\text{NO}_2\text{S}$, m/z : 399.1293, found 400.1367 [(M+H) $^+$].

30 3-(4-Methoxybenzoyl)-2-(methylthio)-1-phenyl-4-(*p*-tolyl)pyrrole (3cb).

White solid, mp = 109–111 °C. $R_f = 0.35$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 7.89$ (d, $J = 8.7$ Hz, 2H), 7.51 – 7.24 (m, 6H), 7.18 – 7.14 (m, 2H), 7.00 – 6.95 (m, 2H), 6.82 (d, $J = 8.7$ Hz, 2H), 3.80 (s, 3H), 2.24 (s, 3H), 2.00 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 193.2$, 163.3, 139.0, 135.8, 132.4, 131.5, 131.0, 129.1, 129.0, 128.0, 127.3, 126.6, 126.3, 125.8, 125.2, 122.6, 113.6, 113.4, 55.4, 21.1, 20.8. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 3010, 2933, 1650, 1575, 1482, 1254. HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{23}\text{NO}_2\text{S}$, m/z : 413.1449, found 414.1522 [(M+H) $^+$].

3-(4-methoxybenzoyl)-2-(methylthio)-4-(4-nitrophenyl)-1-phenylpyrrole (3cd).

Yellow sticky solid. $R_f = 0.50$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 8.04$ (d, $J = 8.7$ Hz, 2H), 7.87 (d, $J = 8.7$ Hz, 2H), 7.51–7.39 (m, 8H), 6.84 (d, $J = 8.4$ Hz, 2H), 3.83 (s, 3H), 2.02 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 194.3$, 176.7, 173.7, 140.9, 132.3, 129.1, 128.5, 127.5, 126.5, 123.9, 113.6, 55.4, 20.6. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 3005, 2941, 1666, 1580, 1442, 1338, 1262, 1211. HRMS (EI) calcd for $\text{C}_{25}\text{H}_{20}\text{N}_2\text{O}_4\text{S}$, m/z : 444.1144, found 444.1137 [(M $^+$)].

50 2-(Methylthio)-1-phenyl-4-(*p*-tolyl)-3-(4-trifluoromethyl)pyrrole (3db).

White solid, mp = 105–107 °C. $R_f = 0.60$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 8.01$ (d, $J = 8.0$ Hz, 2H), 7.71 (d, $J = 8.0$ Hz, 2H), 7.44–7.28 (m, 6H), 6.94

(d, $J = 8.8$ Hz, 2H), 6.91 (d, $J = 8.0$ Hz, 2H), 2.47 (s, 3H), 2.02 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 184.3$, 168.8, 143.4, 137.7, 132.9, 132.48, 132.0, 131.6, 129.1, 127.3, 126.8, 125.3, 20.8, 14.7. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 3077, 2921, 1557, 1537, 1480, 1386, 1103. HRMS (EI) calcd for $\text{C}_{26}\text{H}_{20}\text{F}_3\text{NOS}$, m/z : 451.1218, found 451.1229 [(M $^+$)].

60 1,4-Diphenyl-2-(methylthio)-3-(4-phenylbenzoyl)pyrrole (3ea).

White crystalline solid, mp = 103–105 °C, $R_f = 0.60$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 7.95$ (d, $J = 8.4$ Hz, 2H), 7.58–7.08 (m, 18H), 2.05 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 193.7$, 145.1, 140.0, 138.8, 137.2, 133.9, 130.6, 129.0, 128.8, 128.3, 128.1, 127.9, 127.6, 127.2, 126.7, 126.5, 126.3, 126.2, 122.8, 20.8. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 3029, 2921, 1651, 1600, 1496, 1228. HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{23}\text{NOS}$, m/z : 445.1500, found 446.1561 [(M+H) $^+$].

2-(Methylthio)-1-phenyl-3-(4-phenylbenzoyl)-4-(*p*-tolyl)pyrrole (3eb).

White crystalline solid, mp = 131–133 °C. $R_f = 0.55$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 7.96$ (d, $J = 8.1$ Hz, 2H), 7.58–7.35 (m, 12H), 7.23–7.12 (m, 3H), 6.98 (d, $J = 7.8$ Hz, 2H), 2.22 (s, 3H), 2.03 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 193.8$, 145.1, 140.0, 138.9, 137.2, 135.9, 130.9, 130.6, 129.0, 128.9, 128.7, 128.0, 127.9, 127.4, 127.2, 126.7, 126.5, 126.1, 122.6, 21.0, 20.8. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 3030, 2926, 1643, 1602, 1495, 1446, 1229. HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{25}\text{NOS}$, m/z : 459.1657, found 460.1733 [(M+H) $^+$].

4-(4-Fluorophenyl)-2-(Methylthio)-1-phenyl-3-(4-phenylbenzoyl)pyrrole (3ee).

Yellow crystalline solid, mp = 149–152 °C. $R_f = 0.55$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 7.94$ (d, $J = 8.4$ Hz, 2H), 7.59–7.36 (m, 12H), 7.25–7.21 (m, 2H), 7.11 (s, 1H), 6.90–6.85 (m, 2H), 2.04 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 193.6$, 184.7, 166.5, 145.4, 140.0, 138.8, 137.1, 130.5, 129.2, 129.0, 128.8, 128.2, 128.0, 127.2, 126.7, 126.5, 125.2, 122.8, 115.4, 115.1, 20.7. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 2920, 1634, 1494, 1461, 1211. HRMS (EI) calcd for $\text{C}_{30}\text{H}_{22}\text{FNOS}$, m/z : 463.1406, found 463.1426 [(M $^+$)].

1,4-Diphenyl-2-(methylthio)-3-(2-furoyl)pyrrole (3fa).

White crystalline solid, mp = 148–150 °C. $R_f = 0.40$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 7.49$ – 7.41 (m, 6H), 7.30 (d, $J = 7.2$ Hz, 2H), 7.24–7.09 (m, 4H), 6.93 (d, $J = 3.3$ Hz, 1H), 6.34 (d, $J = 1.8$ Hz, 1H), 2.10 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 180.7$, 153.4, 146.4, 138.8, 134.1, 129.0, 128.3, 128.2, 127.9, 127.4, 127.2, 126.5, 126.2, 122.8, 119.8, 112.0, 111.9, 20.9. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 3116, 2917, 1633, 1492. HRMS (EI) calcd for $\text{C}_{22}\text{H}_{17}\text{NO}_2\text{S}$, m/z : 359.0980, found 359.0973 [(M $^+$)].

3-(2-Furoyl)-2-(methylthio)-1-phenyl-4-(*p*-tolyl)pyrrole (3fb).

Yellow amorphous, mp = 127–129 °C. $R_f = 0.42$ (19:1 hexane/ethyl acetate). ^1H NMR (300 MHz, CDCl_3): $\delta = 7.50$ – 7.44 (m, 6H), 7.20 (d, $J = 7.8$ Hz, 2H), 7.06–7.01 (m, 3H), 6.93 (d, $J = 3.0$ Hz, 1H), 6.35 (dd, $J = 1.8$, 1.8 Hz, 1H), 2.27 (s, 3H), 2.09 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 180.8$, 153.5, 146.3, 138.8, 135.8, 131.1, 129.0, 128.9, 128.1, 127.3, 126.9, 126.5, 125.9, 122.6, 119.8, 112.0, 111.9, 20.9. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$):

¹): 3099, 2972, 1656, 1442, 1202. HRMS (EI) calcd for $C_{23}H_{19}NO_2S, m/z: 373.1136$, found 373.1139 [M^+].

3-(2-Furoyl)-4-(4-methoxyphenyl)-2-(methylthio)-1-phenyl pyrrole (3fc). White solid, mp = 140–143 °C. $R_f = 0.40$ (19:1 hexane/ethyl acetate). 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.50$ –7.45 (m, 7H), 7.23 (d, $J = 8.7$ Hz, 2H), 6.94 (d, $J = 3.6$ Hz, 1H), 6.77 (d, $J = 8.4$ Hz, 2H), 6.36 (dd, $J = 1.65, 1.8$ Hz, 1H), 3.75 (s, 3H), 2.09 (s, 3H). ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 180.8, 158.2, 153.5, 146.4, 138.8, 129.0, 128.6, 128.1, 126.7, 126.5, 125.6, 122.5, 122.4, 119.9, 113.8, 112.1, 112.0, 55.2, 20.9$. IR (KBr) (ν_{max}/cm^{-1}): 3010, 2982, 1643, 1446, 1236. HRMS (EI) calcd for $C_{23}H_{19}NO_3S, m/z: 389.1086$, found 389.1076 [M^+].

3-(2-Furoyl)-2-(methylthio)-4-(4-nitrophenyl)-1-phenyl pyrrole (3fd). Brown solid, mp = 87–89 °C. $R_f = 0.45$ (19:1 hexane/ethyl acetate). 1H NMR (300 MHz, $CDCl_3$): $\delta = 8.08$ (d, $J = 8.7$ Hz, 2H), 7.51–7.43 (m, 9H), 7.04 (d, $J = 3.3$ Hz, 1H), 6.42 (d, $J = 2.1$ Hz, 1H), 2.08 (s, 3H). ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 180.0, 153.2, 146.7, 145.9, 141.0, 138.2, 129.1, 128.7, 128.6, 127.5, 126.4, 124.0, 123.7, 120.0, 112.3, 20.6$. IR (KBr) (ν_{max}/cm^{-1}): 3100, 2915, 1665, 1555, 1465, 1421, 1224. HRMS (EI) calcd for $C_{22}H_{16}N_2O_4S, m/z: 404.0831$, found 404.0830 [M^+].

1,4-Diphenyl-2-(methylthio)-3-(2-thienoyl)pyrrole (3ga). White crystalline solid, mp = 152–154 °C. $R_f = 0.45$ (19:1 hexane/ethyl acetate). 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.54$ –7.41 (m, 8H), 7.31 (d, $J = 7.2$ Hz, 2H), 7.24–7.18 (m, 2H), 7.11 (s, 1H), 6.90 (m, 1H), 2.08 (s, 3H). ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 186.0, 145.6, 138.8, 134.9, 133.9, 129.0, 128.4, 128.1, 127.7, 127.5, 126.5, 126.3, 125.6, 122.8, 21.0$. IR (KBr) (ν_{max}/cm^{-1}): 3082, 2916, 1616, 1488, 1465, 1405, 1279, 1229. HRMS (EI) calcd for $C_{22}H_{17}NOS_2, m/z: 375.0752$, found 375.0758 [M^+].

2-(Methylthio)-1-phenyl-3-(2-thienoyl)-4-(*p*-tolyl)pyrrole (3gb). White crystalline solid, mp = 157–159 °C. $R_f = 0.40$ (19:1 hexane/ethyl acetate). 1H NMR (400 MHz, $CDCl_3$): $\delta = 7.57$ –7.53 (m, 5H), 7.51–7.46 (m, 2H), 7.25 (s, 1H), 7.13 (s, 1H), 7.06 (d, $J = 8$ Hz, 2H), 6.95 (dd, $J = 4.8, 4.0$ Hz, 1H), 2.28 (s, 3H), 2.10 (s, 3H). ^{13}C NMR (100 MHz, $CDCl_3$): $\delta = 186.3, 145.7, 138.9, 136.0, 135.0, 134.0, 131.0, 129.2, 129.1, 128.9, 128.2, 127.9, 127.4, 126.6, 125.9, 125.6, 122.7, 22.7, 21.1$. IR (KBr) (ν_{max}/cm^{-1}): 3077, 2914, 1617, 1495, 1406, 1230. HRMS (EI) calcd for $C_{23}H_{19}NOS_2, m/z: 389.0908$, found 389.0902 [M^+].

3-(4-Bromobenzoyl)-4-phenyl-2-(methylthio)-1-methylpyrrole (3ha). Brown sticky solid. $R_f = 0.48$ (19:1 hexane/ethyl acetate). 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.61$ (d, 2H, $J = 8.4$ Hz), 7.36 (d, 2H, $J = 8.4$), 7.13–7.08 (m, 5H), 6.92 (s, 1H), 3.80 (s, 3H), 2.30 (s, 3H). ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 192.8, 137.4, 134.1, 131.3, 131.1, 128.2, 127.7, 127.3, 126.1, 125.8, 122.4, 34.0, 21.0$. HRMS (EI) calcd for $C_{19}H_{16}BrNOS, m/z: 385.0136$, found 386.0219 [$(M+H)^+$].

3-(3-Hydroxybenzoyl)-4-(4-methoxyphenyl)-2-(methylthio)-1-(*n*-butyl)pyrrole (3ic). Brown solid, mp = 114–116 °C. $R_f = 0.60$ (19:1 hexane/ethyl acetate). 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.18$ –7.13 (m, 2H), 7.00–6.98 (m, 3H), 6.79–6.74 (m, 2H), 6.60

6.57 (d, 2H, $J = 8.7$ Hz), 4.32 (s, 0.35H), 4.04 (2H, dd, $J = 7.5, 7.2$ Hz), 3.61 (s, 3H), 2.19 (s, 3H), 1.73–1.71 (m, 2H), 1.37–1.18 (m, 2H), 0.93–0.80 (m, 3H). ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 194.8, 157.9, 156.7, 155.7, 140.1, 132.1, 130.0, 129.0, 128.7, 127.0, 126.2, 125.7, 123.0, 122.3, 120.7, 119.9, 116.0, 114.4, 113.6, 55.1, 46.7, 33.6, 21.5, 19.9, 13.6, 0.9$. HRMS (EI) calcd for $C_{23}H_{25}NO_3S, m/z: 395.1555$, found 418.1480 [$(M+Na)^+$].

60 3-Benzoyl-4-(4-methoxyphenyl)-2-(methylthio)-1-cyclohexyl pyrrole (3jc). Brown solid. m. p. = 111–113 °C. $R_f = 0.60$ (19:1 hexane/ethyl acetate). 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.84$ –7.76 (m, 2H), 7.40–7.35 (m, 1H), 7.27–7.22 (m, 2H), 7.11–7.08 (m, 2H), 6.94 (s, 1H), 6.66 (d, 2H, $J = 8.1$ Hz), 3.69 (s, 3H), 2.26 (s, 3H), 2.07–1.91 (m, 4H), 1.80–1.56 (m, 2H), 1.52–1.20 (m, 4H). ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 194.5, 157.8, 138.8, 132.2, 129.9, 128.8, 128.6, 127.8, 127.2, 126.9, 125.4, 124.8, 117.7, 117.6, 113.5, 55.3, 34.7, 25.8, 25.3, 21.7$. HRMS (EI) calcd for $C_{25}H_{27}NO_2S, m/z: 405.1762$, found 428.1689 [$(M+Na)^+$].

70 4-(2-Ethynylphenyl)-2-(methylthio)-3-(4-methoxyphenyl)-1-phenylpyrrole (3cf). Yellow (crayola) sticky solid. $R_f = 0.45$ (19:1 hexane/ethyl acetate). 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.72$ (d, $J = 7.8$ Hz, 2H), 7.42–6.94 (m, 10H), 6.65 (d, $J = 7.8$ Hz, 2H), 3.66 (s, 3H), 3.01 (s, 1H), 1.98 (s, 3H). ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 191.9, 162.8, 138.8, 136.7, 133.3, 132.2, 131.3, 129.9, 129.5, 128.8, 128.4, 127.9, 126.4, 126.1, 126.0, 124.79, 124.72, 123.8, 120.2, 112.9, 112.8, 83.5, 80.4, 55.2, 20.7$. IR (KBr) (ν_{max}/cm^{-1}): 3279, 2921, 1643, 1596, 1497, 1461, 1253. HRMS (EI) calcd for $C_{27}H_{21}NO_2S, m/z: 423.1293$, found 422.1182 [$(M-H)^+$].

4-(2-Ethynylphenyl)-2-(methylthio)-1-phenyl-3-(4-phenylbenzoyl)pyrrole (3ef). Yellow (crayola) sticky solid. $R_f = 0.45$ (19:1 hexane/ethyl acetate). 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.87$ (d, $J = 8.1$ Hz, 2H), 7.51–7.27 (m, 14H), 7.16–7.09 (m, 2H), 7.01 (d, $J = 7.2$ Hz, 1H), 3.11 (s, 1H), 2.09 (s, 3H). ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 192.6, 144.5, 140.0, 138.8, 137.2, 136.8, 133.3, 130.5, 130.0, 129.2, 128.9, 128.8, 128.6, 128.5, 128.0, 127.8, 127.0, 126.4, 126.2, 126.1, 124.7, 124.2, 120.4, 83.5, 80.5, 20.7$. IR (KBr) (ν_{max}/cm^{-1}): 3281, 3058, 2921, 1645, 1599, 1497, 90 1227. HRMS (ESI) calcd for $C_{32}H_{23}NOS, m/z: 469.1500$, found 470.1572 [$(M+H)^+$].

3-Benzoyl-4-((E)-(4-methoxystyryl))-2-(methylthio)-1-phenyl pyrrole (3ag). Brown sticky solid. $R_f = 0.45$ (19:1 hexane/ethyl acetate). 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.93$ (d, $J = 7.5$ Hz, 2H), 7.53–7.41 (m, 9H), 7.17 (d, $J = 8.7$ Hz, 2H), 6.76 (d, $J = 8.7$ Hz, 2H), 6.72 (d, $J = 10.2$ Hz, 2H), 3.74 (s, 3H), 1.93 (s, 3H). ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 193.8, 158.7, 139.2, 138.7, 132.5, 130.3, 129.9, 128.9, 128.1, 127.1, 127.0, 126.5, 123.7, 121.9, 118.1, 113.8, 55.1, 20.5$. IR (KBr) (ν_{max}/cm^{-1}): 2921, 1596, 1509, 1249. MS calcd for $C_{27}H_{23}NO_2S, m/z: 425.1449$, found 424.1 [M^+].

4-(4-Methoxyphenyl)-5-methyl-2-(methylthio)-1-phenyl-3-(4-trifluoromethyl)pyrrole (3dh). Brown solid, mp = 164–166 °C. $R_f = 0.55$ (19:1 hexane/ethyl acetate). 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.83$ (d, $J = 8.1$ Hz, 2H), 7.53–7.46 (m, 5H), 7.35 (d,

$J = 6.6$ Hz, 2H), 7.02 (d, $J = 8.4$ Hz, 2H), 6.69 (d, $J = 8.4$ Hz, 2H), 3.69 (s, 3H), 2.05 (s, 3H), 2.02 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 192.6, 158.0, 141.8, 137.7, 133.6, 133.2, 132.8, 132.3, 130.7, 130.0, 129.9, 129.1, 128.7, 128.5, 126.7, 126.4, 125.5, 125.1, 124.6, 123.0, 113.5, 55.1, 21.1, 12.0$. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 3058, 3043, 2921, 1643, 1557, 1482, 1386, 1212, 1172. HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{22}\text{F}_3\text{NO}_3\text{S}$, m/z : 481.1323, found 481.1326 [(M+Na) $^+$].

3-(2-Furoyl)-4-(4-methoxyphenyl)-5-methyl-2-(methylthio)-1-phenylpyrrole (3fh). White solid, mp = 185–188 °C. $R_f = 0.38$ (19:1 hexane/ethyl acetate). ^1H NMR (400 MHz, CDCl_3): $\delta = 7.53\text{--}7.49$ (m, 3H), 7.39 (dd, $J = 8.4, 6.8$ Hz, 3H), 7.18 (d, $J = 8.8$ Hz, 2H), 6.93 (d, $J = 3.2$ Hz, 1H), 6.81 (d, $J = 8.4$ Hz, 2H), 6.35 (dd, $J = 3.4, 1.6$ Hz, 1H), 3.77 (s, 3H), 2.09 [s, 6H, {3H each of 5-methyl and 2-methylthio}]. ^{13}C NMR (100 MHz, CDCl_3): $\delta = 180.9, 157.8, 153.6, 146.1, 137.8, 130.5, 129.8, 129.1, 128.69, 128.64, 128.0, 127.2, 125.6, 122.3, 119.4, 113.6, 111.9, 55.2, 21.3, 12.2$. IR (KBr) ($\nu_{\text{max}}/\text{cm}^{-1}$): 2920, 1629, 1471, 1249. HRMS (EI) calcd for $\text{C}_{24}\text{H}_{21}\text{NO}_3\text{S}$, m/z : 403.1242, found 403.1250 [M $^+$].

Copies of ^1H , ^{13}C and MS/HRMS are listed in Supporting Information.

Acknowledgement

We gratefully acknowledge the generous financial support from the Science and Engineering Research Board (SB/S1/OC-25/30/2013), the Council of Scientific and Industrial Research (02(0072)/12/EMR-II), and the UGC-UKIERI Partnership-2013 (184-16/2013(IC), New Delhi. K. R. is highly grateful to DAAD and George-August-Universität-Göttingen for financial support as DAAD fellowship.

Notes and references

^{1*} Department of Chemistry, Faculty of Science, Banaras Hindu University, Varanasi-05, UP, India. Fax: +91-542-2368127; Tel: +91-542-6702502; E-mail: mssinghbhu@yahoo.co.in

² Institut für Organische und Biomolekulare Chemie, Georg-August-Universität, Tammannstr. 2, 37077, Göttingen, Germany; E-mail: raghuvanshikeshav@gmail.com

³ Department of Chemistry, Loughborough University, Loughborough, Leics, LE11 3TU, UK; E-mail: r.c.f.jones@lboro.ac.uk

⁴⁰ † Electronic Supplementary Information (ESI) available:
See DOI: 10.1039/b000000x/

- (a) V. Estévez, M. Villacampa and J. C. Menéndez, *Chem. Soc. Rev.*, 2010, **39**, 4402; (b) V. Estévez, M. Villacampa and J. C. Menéndez, *Chem. Soc. Rev.*, 2014, doi: 10.1039/c3cs60015g; (c) H. Fan, J. Peng, M. T. Hamann and J.-F. Hu, *Chem. Rev.*, 2008, **108**, 264; (d) A. Hall, S. Atkinson, S. H. Brown, I. P. Chessell, A. Chowdhury, G. M. P. Giblin, P. Goldsmith, M. P. Healy, K. S. Jandu, M. R. Johnson, A. D. Michel, A. Naylor and J. A. Sweeting, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 1200; (e) C. V. Galliford and K. A. Scheidt, *Angew. Chem., Int. Ed.*, 2007, **46**, 8748.
- (a) D. L. Boger, C. W. Boyce, M. A. Labroli, C. A. Sehon and Q. Jin, *J. Am. Chem. Soc.*, 1999, **121**, 54; (b) A. Grube and M. Köck, *Org. Lett.*, 2006, **8**, 4675; (c) K. A. McArthur, S. S. Mitchell, G. Tsueng, D. J. White, J. Grodberg, K. S. Lam and B. C. Potts, *J. Nat. Prod.*, 2008, **71**, 1732.

- (a) J. Lehuédé, B. Fauconneau, L. Barrier, M. Ourakow, A. Piriou and J.-M. Vierfond, *Eur. J. Med. Chem.*, 1999, **34**, 991; (b) A. Fürstner, *Angew. Chem.*, 2003, **115**, 3706; *Angew. Chem., Int. Ed.*, 2003, **42**, 3582.
- (a) A. Berlin, B. Vercelli and G. Zotti, *Polym. Rev.*, 2008, **48**, 493; (b) I. F. Gimenez and O. L. Alves, *J. Braz. Chem. Soc.*, 1999, **10**, 167; (c) J. R. Reynolds, P. A. Poropatic and R. L. Toyooka, *Macromolecules* 1987, **20**, 958.
- (a) R. W. Bürl, D. McMinn, J. A. Kaizerman, W. Hu, Y. Ge, Q. Pack, V. Jiang, M. Gross, M. Garcia, R. Tanaka and H. E. Moser, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 1253; (b) V. J. Demopoulos and E. Rekka, *J. Pharm. Sci.*, 1995, **84**, 79; (c) J. M. Muchowski, *Adv. Med. Chem.*, 1992, **1**, 109; (d) A. S. Demir, I. M. Akhmedov and O. Sesenoglu, *Tetrahedron*, 2002, **58**, 9793; (e) D. Wang, X. Hu and G. Zhao, *Int. J. Food Sci. Technol.*, 2008, **43**, 1880; (f) J. T. Gupton, *Top. Heterocycl. Chem.*, 2006, **2**, 53; (g) N. Amishiro, S. Nagamura, E. Kobayashi, A. Okamoto, K. Gomi and H. Saito, *Chem. Pharm. Bull.*, 1999, **47**, 1393; (h) N. Amishiro, S. Nagamura, E. Kobayashi, A. Okamoto, K. Gomi, M. Okabe and H. Saito, *Bioorg. Med. Chem.*, 2000, **8**, 1637; (i) H. M. Meshram, B. R. V. Prasad and D. A. Kumar, *Tetrahedron Lett.*, 2010, **51**, 3477; (j) M. Del Poeta, W. A. Schell, C. C. Dykstra, S. Jones, R. R. Tidwell, A. Czarny, M. Bajic, A. Kumar, D. Boykin and J. R. Perfect, *Antimicrob. Agents Chemother.*, 1998, **42**, 2495; (k) M.-Z. Wang, H. Xu, T.-W. Liu, Q. Feng, S.-J. Yu, S.-H. Wang and Z.-M. Li, *Eur. J. Med. Chem.*, 2011, **46**, 1463; (l) F. A. Davis, K. Bowen, H. Xu and V. Velvadapu, *Tetrahedron*, 2008, **64**, 4174.
- (a) Y. Arikawa, H. Nishida, O. Kurasawa, A. Hasuoka, K. Hirase, N. Inatomi, Y. Hori, J. Matsukawa, A. Imanishi, M. Kondo, N. Tarui, T. Hamada, T. Takagi, T. Takeuchi and M. Kajino, *J. Med. Chem.*, 2012, **55**, 4446; (b) M. Menichincheri, C. Albanese, C. Alli, D. Ballinari, A. Bargiotti, M. Caldarelli, A. Ciavolella, A. Cirila, M. Colombo, F. Colotta, V. Croci, R. D'Alessio, M. D'Anello, A. Ermoli, F. Fiorentini, B. Forte, A. Galvani, P. Giordano, A. Isacchi, K. Martina, A. Molinari, J. K. Moll, A. Montagnoli, P. Orsini, F. Orzi, E. Pesenti, A. Pillan, F. Roletto, A. Scolaro, M. Tato, M. Tibolla, B. Valsasina, M. Varasi, P. Vianello, D. Volpi, C. Santocanale and E. Vanotti, *J. Med. Chem.*, 2010, **53**, 7296; (c) B. D. Roth, *Prog. Med. Chem.*, 2002, **40**, 1.
- (a) R. E. Nizurski-Mann and M. P. Cava, *Heterocycles*, 1992, **34**, 2003; (b) C. D'Silva and D. A. Walker, *J. Org. Chem.*, 1998, **63**, 6715; (c) B. P. J. de Lacy Costello, P. Evans, N. Guernion, N. M. Ratcliffe, P. S. Sivanand and G. C. Teare, *Synth. Met.*, 2000, **114**, 181.
- (a) T. P. Toube, B. A. Trofimov, G. Cirrincione, A. M. Almerico, E. Aiello, G. Dattolo, H. McNab and L. C. Monahan in *Chemistry of Heterocyclic Compounds: Pyrroles, Part II, The Synthesis, Reactivity and Physical Properties of Substituted Pyrroles*, Vol. 48 (Ed.: R. A. Jones), John Wiley & Sons Inc., New York, 2008, pp. 1; (b) E. T. Pelkey in *Prog. Heterocycl. Chem.* Vol. **17** (Eds.: G. W. Gribble and J. A. Joule), Elsevier, S. A., 2005, pp. 109.
- (a) D. Akbaslar, O. Demirkol and S. Giray, *Synth. Commun.*, 2014, **44**, 1323; (b) A. Devi, Shallu, M. L. Sharma and J. Singh, *Synth. Commun.*, 2012, **42**, 1480; (c) G. E. Veitch, K. L. Bridgwood, K. Rands-Trevor and S. V. Ley, *Synlett*, 2008, 2597; (d) C. Paal, *Ber. Dtsch. Chem. Ges.*, 1885, **18**, 367; (e) L. Knorr, *Ber. Dtsch. Chem. Ges.*, 1884, **17**, 1635; (f) Z. H. Zhang, J. J. Li and T. S. Li, *Ultrason. Sonochem.*, 2008, **15**, 673; (g) J. M. Manley, M. J. Kalman, B. G. Conway, C. C. Ball, J. L. Havens and R. Vaidyanathan, *J. Org. Chem.*, 2003, **68**, 6447.
- (a) V. Estévez, M. Villacampa and J. C. Menéndez, *Chem. Commun.*, 2013, **49**, 591; (b) A. Herath and N. D. P. Cosford, *Org. Lett.*, 2010, **12**, 5182; (c) A. Hantzsch, *Ber. Dtsch. Chem. Ges.*, 1890, **23**, 1474; (d) A. W. Trautwein, R. D. Süßmuth and G. Jung, *Bioorg. Med. Chem. Lett.*, 1998, **8**, 2381; (e) L. Calvo, A. González-Ortega and M. C. Sañudo, *Synthesis*, 2002, 2450.
- (a) H. J. Deng, Y. J. Fang, M. C. Chen, M. C. Liu, H. Y. Wu and J. X. Chen, *Appl. Organomet. Chem.*, 2012, **26**, 164; (b) N. Ono, *Heterocycles*, 2008, **75**, 243.
- (a) N. C. Misra, K. Panda, H. Ila and H. Junjappa, *J. Org. Chem.*, 2007, **72**, 1246; (b) S. Toshiaki, S. Takuya, I. Reiko, S. Norio and K.

- Takeo, *Eur. J. Org. Chem.*, 2010, **22**, 4237; (c) E. P. J. Ng, Y.-F. Wang and S. Chiba, *Synlett*, 2011, 783.
- 13 (a) S. D. Joshi, U. A. More, V. H. Kulkarni and T. M. Aminabhavi, *Curr. Org. Chem.*, 2013, **17**, 2279; (b) V. F. Ferreira, M. C. B. V. de Souza, A. C. Cunha, L. O. R. Pereira and M. L. G. Ferreira, *Org. Prep. Proced. Int.*, 2001, **33**, 411; (c) C. Schmuck and D. Rupprecht, *Synthesis*, 2007, 3095.
- 14 (a) F. Tripoteau, L. Eberlin, M. A. Fox, B. Carboni and A. Whiting, *Chem. Commun.*, 2013, **49**, 5414.
- 15 (a) M. P. Huestis, L. Chan, D. R. Stuart and K. Fagnou, *Angew. Chem., Int. Ed.*, 2011, **50**, 1338; (b) S. Rakshit, F.W. Patureau and F. Glorius, *J. Am. Chem. Soc.*, 2010, **132**, 9585.
- 16 (a) X. Xin, D. Wang, X. Li and B. Wan, *Angew. Chem., Int. Ed.*, 2012, **51**, 1693; (b) Y. Jiang, W. C. Chan and C. M. Park, *J. Am. Chem. Soc.*, 2012, **134**, 4104; (c) M. Egi, K. Azechi and S. Akai, *Org. Lett.*, 2009, **11**, 5002; (d) A. V. Kel'in, A.W. Sromek and V. Gevorgyan, *J. Am. Chem. Soc.*, 2001, **123**, 2074.
- 17 (a) G. Balme, *Angew. Chem., Int. Ed.*, 2004, **43**, 6238; (b) M. Zhang, H. Neumann and M. Beller, *Angew. Chem., Int. Ed.*, 2013, **52**, 597;
- 20 (c) J. Zeng, Y. Bai, S. Cai, J. Ma and X.-W. Liu, *Chem. Commun.*, 2011, **47**, 12855; (d) X. Lin, Z. Mao, X. Dai, P. Lu and Y. Wang, *Chem. Commun.*, 2011, **47**, 6620.
- 25 18 (a) E. Lourdusamy, L. Yao and C.-M. Park, *Angew. Chem., Int. Ed.*, 2010, **49**, 7963; (b) D. Srimani, Y. Ben-David and D. Milstein, *Angew. Chem., Int. Ed.*, 2013, **52**, 4012.
- 25 19 M. Rueping and A. Parra, *Org. Lett.*, 2010, **12**, 5281.
- 20 (a) G. K. Verma, R. K. Verma, G. Shukla, N. Anugula and M. S. Singh, *Tetrahedron*, 2013, **69**, 6612; (b) G. K. Verma, R. K. Verma and M. S. Singh, *RSC Adv.*, 2013, **3**, 245; (c) T. Chanda, R. K. Verma and M. S. Singh, *Chem. Asian J.*, 2012, **7**, 778; (d) R. K. Verma, G. K. Verma, G. Shukla and M. S. Singh, *RSC Adv.*, 2012, **2**, 2413; (e) M. S. Singh, G. C. Nandi and S. Samai, *Green Chem.*, 2012, **14**, 447.
- 30 21 (a) G. Shukla, R. K. Verma, G. K. Verma, K. Raghuvanshi, A. Nagaraju and M. S. Singh, *RSC Adv.*, 2013, **3**, 13811; (b) S. Koley, S. Chowdhury, T. Chanda, B. J. Ramulu and M. S. Singh, *Tetrahedron*, 2013, **69**, 8013; (c) S. Samai, T. Chanda, H. Ila and M. S. Singh, *Eur. J. Org. Chem.*, 2013, 4026; (d) M. S. Singh, A. Nagaraju, G. K. Verma, G. Shukla, R. K. Verma, A. Srivastava and K. Raghuvanshi, *Green Chem.*, 2013, **15**, 954; (e) R. K. Verma, G. K. Verma, G. Shukla, A. Nagaraju and M. S. Singh, *ACS Comb. Sci.*, 2012, **14**, 224; (f) G. C. Nandi, M. S. Singh, H. Ila, and H. Junjappa, *Eur. J. Org. Chem.*, 2012, 967; (g) G. C. Nandi, S. Samai and M. S. Singh, *J. Org. Chem.*, 2011, **76**, 8009; (h) S. Chowdhury, G. C. Nandi, S. Samai and M. S. Singh, *Org. Lett.*, 2011, **13**, 3762.
- 40 22 CCDC 906650 (**3ga**), 946357 (**3eb**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.