

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

Synthesis of β -damascone derivatives with lactone ring and their feeding deterrent activity against aphids and lesser mealworm

Cite this: DOI: 10.1039/x0xx00000x

A. Gliszczynska,^{*a} K. Dancewicz,^b M. Hnatejko,^c M. Szczepanik,^c B. Gabryś,^bReceived 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Starting from β -damascone six new lactones were obtained. The Claisen-Johnson rearrangement of allylic alcohol and halolactonization of γ,δ -unsaturated acid was the key step of presented synthesis. The structures of new derivatives were determined by spectroscopic data. The antifeedant activity of β -damascone towards two insect species with different feeding habits and food preferences, i.e. peach potato aphid *Myzus persicae* (Sulz.) and lesser mealworm, *Alphitobius diaperinus* Panzer was studied as well as the biological consequences of structural modification of the starting substrate. The most successive structural modifications of β -damascone in terms of antifeedant activity towards *M. persicae* were the incorporation of lactone moiety and concomitant presence of bromine in the side chain. All β -damascone derivatives with lactone moiety deterred the feeding of adults and larvae of *A. diaperinus*. Halo- δ -lactones were more active than halo- γ -lactones and *A. diaperinus* adults were more sensitive to the compounds studied than larvae.

^a Department of Chemistry, Wrocław University of Environmental and Life Sciences, Norwida 25, 50-375 Wrocław, Poland, e-mail: anna.gliszczynska@wp.pl, Fax: +48 71 3284124, Tel: +48 71 3205257.

^b Department of Biology and Ecology, University of Zielona Góra, Szafrana 1, 65-516 Zielona Góra, Poland.

^c Nicolaus Copernicus University, Department of Invertebrate Zoology, Lwowska 1, 87-100, Toruń, Poland.

Introduction

β -Damascone (**1**) belongs to the group of norisoprenoids defined as rose ketones, because they were discovered in the 1960s during a quest to identify the characteristic smell of Bulgarian rose oil^{1,2}. This compound has been identified not only as component of rose oil but also creates tea aroma, occurs in some types of tobacco, wine and whiskey³⁻⁷. β -Damascone (**1**) has a pleasant blackcurrant-plum note and is widely used in perfume compositions. Apart from this commercial importance, β -damascone has been shown to have a variety of biological activities. For example, it was discovered that this compound possesses cancer chemopreventive potential⁸ and appeared to be toxic towards three species of mosquitoes, *Aedes aegypti* L., *Ae. albopictus* (Skuse) and *Anopheles quadrimaculatus* Say, the housefly, *Musca domestica* L., the stable fly, *Stomoxys calcitrans* L., and the sand fly, *Lutzomyia shannoni* (Dyar)⁹⁻¹¹. Apart from that, information on the effect of β -damascone (**1**) on arthropods and their behavior is very scarce. Our interests in the synthesis of isoprenoid lactones with damascone skeleton were motivated by the search for new biologically active compounds that can reduce the population of insect pests and can be useful in protection of crops. Insects are responsive to many plant lower terpenoids and their synthetic derivatives. For example, the repellent properties of linalool and α -terpineol to the peach potato aphid

Myzus persicae were reported by Hori^{12,13} and (*S*)-limonene restrained phloem sap ingestion and had other negative effects on the behaviour of this aphid¹⁴. Citral, linalool, (*S*)-limonene, α -ionone, and camphene reduced the total and mean probing time of aphids and their settling on the leaves¹⁵. Following these studies, several analogues of natural terpenoids, including the lactones, have been synthesized and their biological activity examined. We discovered active feeding deterrents among lactones derived from natural isoprenoids: pulegone^{16,17}, piperitone¹⁸, farnesol¹⁹. It appeared that the antifeedant activity of synthetic analogues was highly enantiospecific and depended on various substituents and functional groups^{14,18,20}. Chemical transformation of the piperitone molecule by the introduction of a lactone moiety and a halogen atom strongly changed its antifeedant properties against *M. persicae* and the lesser mealworm *Alphitobius diaperinus*^{18,21}.

In the present study, we concentrated on two insect pests with different feeding habits and food preferences, i.e. peach potato aphid *Myzus persicae* (Sulz.) (Hemiptera: Aphididae) and lesser mealworm, *Alphitobius diaperinus* Panzer (Coleoptera: Tenebrionidae). Aphids are responsible for at least 2% of all losses attributed to insect feeding²². Moreover, the indirect damage caused by aphids due to virus transmission exceeds their direct impact on crops²³. *M. persicae* alone can infest plants of over 40 different families and it is able to transmit over 100 plant viruses²⁴. At the same time, it developed clones resistant to one or more insecticides²³. The lesser mealworm is a cosmopolitan pest inhabiting chicken and broiler houses in vast numbers. In poultry houses the beetles consume manure, spilled feed, dead birds and other organic materials^{25,26}. At the same time, the adults and larvae have been reported as potential vectors for the transfer of *Campylobacter jejuni* and *Salmonella enterica* between successive broiler flocks²⁷ and other pathogens²⁸.


Chemicals targeting insect taste receptors are considered potential bioinsecticides to protect crops²⁹. Aphids possess the piercing-sucking mouthparts and they feed on the phloem sap of the living plants, while the lesser mealworm is a general stored products pest that feeds using the chewing mouthparts. Moreover, aphids lack external taste receptors, which makes the preingestional rejection or acceptance of the food impossible³⁰. In contrast, the lesser mealworm's gustatory receptors are located on their mouthparts, which allows the preingestional response to food quality. The different food preferences, modes of food uptake, and ability to preingestional evaluation of food chemistry may determine different behavioural responses of these insects. In the present study, we present the synthesis and structural modifications of β -damascone (**1**)-derived compounds, including the lactones, as well as their biological activities expressed as antifeedant properties to *M. persicae* and *A. diaperinus*.

Results and discussion

Chemical synthesis

Six new isoprenoid lactones **6-11** were obtained in a six-step synthesis (Scheme 1) from naturally occurring ketone - β -damascone (**1**). The damascone carbon skeleton consists of a trimethyl substituted cyclohexane ring with a four-carbon α,β -unsaturated ketone side chain. The starting material was commercially available β -damascone (**1**).

The first step of the synthesis was the reduction of double bond in side chain of compound (**1**) with lithium aluminium hydride according to the standard procedure. The reaction course and purity of the product were monitored by GC on a capillary column (HP-5). Known dihydro- β -damascone (**2**) which was identified in tobacco³¹ and synthesized previously by Mori et al. in two-steps chemical synthesis from trimethylcyclohexanone³² was obtained in high 99% yield. The structure of **2** was confirmed by the comparison its spectral data with literature one.



Scheme 1. Synthesis of lactones from β -damascone. Reagents (i) LiAlH_4 ; (ii) $\text{CH}_3\text{C}(\text{OEt})_3$, CH_3COOH , 138°C ; (iii) 1. KOH , EtOH , 2. HCl ; (iv) NBS/NCS , THF ; (v) DBU

Ketone (**2**) was next transformed in high 89% yield into corresponding allylic alcohol – dihydro- β -damascol (**3**) by treatment with LiAlH_4 . Preparation of dihydro- β -damascol (**3**) was patented for the first time in 1973³³. The crude alcohol (**3**), without further

purification (97% purity according to the GC analysis) was subjected to the orthoacetate modification of Johnson-Claisen rearrangement³⁴. The mechanism involves the esterification of alcohol followed by subsequent elimination of ethanol to form ketene acetal. The latter is rearranged to the γ,δ -unsaturated esters in the [3.3] sigmatropic shift during heating. In our case, the reaction afforded new compound ethyl 2-(2-butylidene-1,3,3-trimethylcyclohexyl)-acetate (**4**) in 98% yield. Absorption band at 1743 cm^{-1} of carbonyl group and characteristic quartet in the ^1H NMR spectrum from two protons at 4.05 ppm confirmed the presence of carboethoxy group in ester (**4**).

The ester (**4**) was subsequently hydrolyzed in ethanolic KOH solution to the corresponding acid (**5**) in 85% yield. This compound has not been obtained before. In the next step acid (**5**) was subjected to halolactonization reactions giving access to bromo- and chlorolactones (Scheme 1). The bromolactonization of acid (**5**) was carried out with *N*-bromosuccinimide (NBS) in tetrahydrofuran, affording a mixture of two new products in ratio 45%:55% (according to GC). We separate them using column chromatography and established their structure on the basis of spectroscopic data. The products of cyclization were δ -bromo- γ -lactone (**6**) (minor, 17% yield) and γ -bromo- δ -lactone (**7**) (major, 27% yield). The reaction of γ,δ -unsaturated acid (**5**) with *N*-chloro-succinimide (NCS) was carried out to obtain chlorolactones. Using chloride as an electrophilic agent we observed a similar situation as in the process of bromolactonization. Two new lactones were formed as products of cyclization - δ -chloro- γ -lactone (**8**) and γ -chloro- δ -lactone (**9**). According to GC a mixture consists of 41% of γ -lactone (**8**) and 59% δ -lactone (**9**), which were obtained with 18 and 17% yield respectively. The doublet of doublets protons H-8 in γ -lactone (4.24 ppm) and δ -lactone (4.74 ppm) were found in the ^1H NMR spectra and also proved intermolecular cyclization.

The dehydro-halogenation reaction of δ -halo- γ -lactones (**6**), (**8**) and γ -halo- δ -lactone (**7**), (**9**) with 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) gave new cyclic lactones (**10**), (**11**) in good yields (47-57%) as the only products (Scheme 1). The formation of unsaturated lactone (**10**) as the only product of the dehydrohalogenation of lactones (**6**), (**8**) is a result of E2 elimination. In the IR spectrum of unsaturated lactone (**10**) the absorption bands for stretching vibrations for carbonyl group at 1778 cm^{-1} were observed. The location of double bond in the side chain of bicyclic lactone (**10**) between C-8 and C-9 was confirmed by the signals from olefin protons in the ^1H NMR spectrum. The doublets of triplets from H-8 and H-9 protons, at 5.60 ppm and 5.77 ppm respectively indicate the presence of double bond. Moreover the coupling constant $J=15.5\text{ Hz}$ between these protons proved their *trans* orientation. The shift of the absorption band of $\text{C}=\text{O}$ band in lactone moiety from the range of $1716\text{--}1733\text{ cm}^{-1}$ for δ -lactones to higher frequency 1762 cm^{-1} indicates that in the case of product (**11**) the conversion of δ -lactone ring into γ -lactone occurred. In the ^1H NMR spectrum signal from proton H-8 appeared as a doublet of doublets. It was coupled with protons at C-9 at 4.25 ppm with coupling constant $J=10.9$ and 2.2 Hz . These coupling are clearly observed in 2D COSY spectrum (Fig. 1). This type of cyclization is the result of E2 elimination. The lack of signal from one of $\text{CH}_2\text{-3}$ protons as well as any signals characteristic for olefinic protons and finally shape of the signal from H-8 proton indicated the presence of cyclopropane ring. Its formation can be explained by expulsion of proton at C-3 by DBU leading to carbanion stabilized by enolate anion. The second step, cyclization, is the results of the attack of electron pair as a nucleophile on the C-7a carbon atom with simultaneous removing of halogen³⁵.




Fig.1. COSY spectrum of lactone 11.

Biological studies

Feeding deterrent activity against *Myzus persicae*

The potency and mechanism of antifeedant activity of the compounds studied varied depending on the compound structure and the duration of exposure of aphids to β -damascone (**1**) and its derivatives. On control leaves, aphids did not leave the substrate (=leaf) during the no-choice 15-minute experiment and the total probing time was approximately 70% of the time spent on the leaf. Aphids started probing immediately after exposure (the delay was 14 seconds on average) and the average probe was relatively long, i.e., approximately 3.5 minutes (Table 1). In the choice-test, there were no significant differences in aphid preferences to settle on either the untreated or the control (i.e., ethanol-treated) leaf, irrespective of the exposure time (Fig. 2). β -Damascone (**1**) appeared a weak attractant during the initial contact with the treated leaves (no-choice test): although the first probe was delayed in comparison to control, further probing was rarely interrupted. There were twice as few probes and the probes were twice as long in comparison to aphids on control leaves (Table 1). However, this effect did not translate into aphid preferences during settling: no significant differences in the number of aphids on treated and untreated leaves were found during the 24-hour experiment (Fig. 2).




Fig. 2. Effect of β -damascone and its derivatives on settling preferences of *Myzus persicae* in the choice test. Data are expressed as values of indices of deterrence (DI). The standard error is indicated on the bar. * $P < 0.05$; *** $P < 0.001$; (Student t -test).

The compound synthesized in the first step in the β -damascone modification, the dihydro- β -damascone (**2**) did not evoke any changes in aphid behaviour during initial contact with the treated leaves (no-choice test) as well as in the long-term experiment showing aphid preferences during settling (choice-test). Similar results were found after the application of dihydro- β -damascol (**3**), except that the probes were significantly, four times shorter in comparison to control. However, aphids did not restrain from probing – the total probing time was comparable to control, which finally caused no discrimination between treated and untreated leaves during the 24-hour settling experiment. The β -damascone-derived ethyl 2-(2-butyldiene-1,3,3-trimethylcyclohexyl)-acetate (**4**) did not cause significant differences in aphid responses to plants during initial 15 minutes after exposure except considerable delay before the first probe (Table 1).

Table 1. Modification of *Myzus persicae* behaviour by β -damascone and its derivatives in no-choice tests^a.

Compound	Time spent on the leaf (s)	Total probing time (s)	Time to first probe (s)	Number of probes	Mean probing time (s)
Control	900.0 \pm 0.0	608.6 \pm 51.8	13.7 \pm 6.3	5.4 \pm 0.9	209.8 \pm 69.7
1	892.5 \pm 7.5	640.6 \pm 44.1	123.1 \pm 37.2*	2.4 \pm 0.5*	435.7 \pm 90.1*
2	844.5 \pm 47.9	575.8 \pm 79.3	51.5 \pm 28.5	6.2 \pm 1.3	194.8 \pm 65.9
3	809.5 \pm 51.3	457.3 \pm 68.0	14.8 \pm 3.0	7.7 \pm 0.9	77.7 \pm 22.5*
4	867.2 \pm 30.0	467.5 \pm 78.7	154.2 \pm 50.0*	4.1 \pm 0.9	204.2 \pm 66.6
5	892.8 \pm 7.2	585.8 \pm 68.4	12.5 \pm 5.0	5.8 \pm 1.0	189.6 \pm 76.7
6	539.0 \pm 117.0*	256.6 \pm 80.8*	104.7 \pm 64.4	4.5 \pm 1.0	43.9 \pm 15.8*
7	900.0 \pm 0.0	707.5 \pm 36.4	25.7 \pm 7.8	4.3 \pm 0.7	249.3 \pm 70.8
8	799.8 \pm 68.6	545.6 \pm 76.8	27.4 \pm 7.2	4.3 \pm 0.8	264.5 \pm 93.2
9	887.2 \pm 9.1	635.4 \pm 68.0	22.7 \pm 6.1	5.4 \pm 1.0	208.9 \pm 47.5
10	888.6 \pm 10.4	640.1 \pm 102.1	26.9 \pm 11.6	3.3 \pm 0.6	335.8 \pm 102.7

^aValues represent means of $n=12$ replicates \pm SE. An asterisk within a column denotes statistically significant differences in relation to control ($P < 0.05$).

However, from the second hour after exposure onwards, aphids showed significant preference for untreated leaves. The indices of deterrence reached relatively high values of 0.5 and 0.6 after 2 and 24 hours, respectively (Fig. 2). Further molecular modification, the synthesis of 2-(2-butyldiene-1,3,3-trimethylcyclohexyl) acetic acid (**5**) did not cause any change in the biological activity towards *M. persicae* initially, but after 24-hour exposure, aphid settling was significantly hindered ($ID=0.2$). In contrast, the incorporation of the lactone moiety and the halogen atoms into the molecule did have significant impact on aphid behaviour. However, the four halolactones obtained (**6-9**) differed in activity depending on the size of the lactone ring and the kind and position of the halogen atoms in the molecule. In the behavioural no-choice test, the application of δ -bromo- γ -lactone (**6**) caused a significant decrease in the total time spent on the treated leaves and total and mean probing time, while the γ -bromo- δ -lactone (**7**) did not (Table 1). Consequently, the exposure to δ -bromo- γ -lactone (**6**) caused the avoidance of treated leaves by freely moving aphids in choice-test, from the beginning until the end of the experiment in contrast to γ -bromo- δ -lactone (**7**) (Fig. 2). The replacement of the bromine atom by chlorine atom to synthesize the respective δ -chloro- γ -lactone (**8**) and γ -chloro- δ -lactone (**9**) did not cause changes in aphid behaviour during initial contact with the studied compounds (Table 1). However, long-term exposure to γ -chloro- δ -lactone (**9**) impeded aphid settling significantly (Fig. 2). Similar effects on aphid settling and behaviour were caused by the unsaturated bicyclic γ -lactone (**10**) and bicyclic δ -lactone (**11**): aphid initial responses on treated leaves did not differ from those on control leaves in the no-choice experiment (Table 1) but the settling of aphids was significantly impeded after longer

exposure times. The settling-deterrent effect of lactones (**10**) and (**11**) increased in potency in the course of time, with indices of deterrence reaching 0.7 and 0.3 after 24 hours for lactones (**10**) and (**11**), respectively (Figure 2).

In summary, the antifeedant activity of β -damascone-derived compounds was manifested both, as immediate and delayed effects on aphid behaviour. The immediate effects during initial contacts with the allelochemical were expressed as the reduction of total time spent on the leaf, total probing time (δ -bromo- γ -lactone (**6**)), delayed time to the first probe (β -damascone (**1**) and ester (**4**)), and the reduction of mean probing time (dihydro- β -damascol (**3**) and δ -bromo- γ -lactone (**6**)). Considering average durations of probes (0.7 – 1.2 minutes), aphid stylets on (**3**) and (**6**)-treated leaves did not penetrate beyond epidermis³⁶. The delayed effects were expressed after longer times of aphid exposure to the compounds studied. The strongest and the most durable effects on aphid settling were evoked by the application of ethyl 2-(2-butyldiene-1,3,3-trimethyl-cyclohexyl)-acetate (**4**), δ -bromo- γ -lactone (**6**), and unsaturated bicyclic γ -lactone (**10**). Moreover, the deterrent effect increased in potency in the course of time. With the exception of δ -bromo- γ -lactone (**6**), the effects of ester (**4**) and lactone (**10**) were probably only postingestional because the settling impediment occurred later than two hours after exposure. δ -Bromo- γ -lactone (**6**) had probably partially preingestional and ingestional/postingestional deterrent activity: although aphids were discouraged from probing immediately after they had gained access to plants, the probing was not entirely eliminated, which most likely allowed the consumption of plant sap.

Feeding deterrent activity against *Alphitobius diaperinus*

The compounds studied showed varying antifeedant activity, which depended on the structure of the compound and the developmental stage of the lesser mealworm. Starting β -damascone (**1**) was a moderate feeding deterrent for both developmental stages, especially in no-choice test. The food consumed by the larvae and adults in this test represented 59.26 and 51.5% of the consumption in the control, respectively (Fig. 3). The introduction of a lactone moiety into a β -damascone molecule changed its antifeedant properties. Unsaturated γ -lactone (**10**) and bicyclic δ -lactone (**11**) were very good feeding deterrents, but only against adults. In bioassays with larvae a strong activity in the choice test was observed. Unfortunately, in the no choice tests larvae were eating intensively the treated food as in the case of unsaturated lactone (**10**) (Table 2).




Fig. 3 Effect of β -damascone and its derivatives on feeding of *Alphitobius diaperinus* in the no-choice test. Data are expressed as percentages of control consumption. The standard error is indicated on the bar. * $P < 0.05$; *** $P < 0.001$; NS: not significant (Student t -test).

Table 2. Feeding deterrent activity of the studied compounds in choice and no-choice tests against *A. diaperinus*

Compound	Deterrence coefficients \pm SE ^a					
	Larvae			Adults		
	A	R	T	A	R	T
Dose	1 %					
1	30.70 \pm 15.94 abc	55.59 \pm 3.41 a	86.30 \pm 17.37 a	34.74 \pm 12.00 a	56.94 \pm 4.94 ab	91.68 \pm 10.52 ab
2	25.30 \pm 6.80 ab	70.69 \pm 4.46 b	95.99 \pm 9.24 a	57.22 \pm 2.05 ab	63.20 \pm 5.93 ab	120.42 \pm 6.68 abc
3	62.41 \pm 9.69 c	76.71 \pm 3.60 bc	139.12 \pm 4.6 a	75.47 \pm 4.12 b	59.34 \pm 9.29 ab	134.81 \pm 12.81 bcd
4	33.90 \pm 1.20 abc	81.74 \pm 4.55 bcde	115.64 \pm 3.32 ab	46.53 \pm 15.52 a	65.59 \pm 11.02 b	112.13 \pm 16.07 abc
5	8.47 \pm 4.85 ab	90.42 \pm 1.88 cdef	98.89 \pm 3.84 a	50.74 \pm 7.85 a	81.05 \pm 17.72 b	131.79 \pm 4.59 bcd
6	33.58 \pm 2.69 c	77.80 \pm 3.11 bcd	111.38 \pm 3.67 ab	50.34 \pm 1.63 a	23.20 \pm 0.30 a	73.54 \pm 20.70 a
7	82.45 \pm 11.50 d	80.19 \pm 1.17 bcde	162.64 \pm 1.83 b	90.13 \pm 7.63 b	91.05 \pm 1.38 b	181.17 \pm 1.64 d
8	67.04 \pm 4.36 c	97.77 \pm 0.53 f	164.81 \pm 10.56 b	72.94 \pm 1.29 b	89.48 \pm 10.36 b	162.42 \pm 8.72 cd
9	85.71 \pm 5.30 d	98.29 \pm 2.96 f	184 \pm 4.29 b	78.70 \pm 3.82 b	64.69 \pm 8.23 b	143.38 \pm 10.34 bcd
10	-2.08 \pm 4.74 a	91.43 \pm 1.07 def	89.35 \pm 7.42 a	71.24 \pm 2.43 b	77.45 \pm 2.60 b	148.69 \pm 9.58 cd
11	36.01 \pm 2.07 b	93.77 \pm 2.77 ef	129.79 \pm 5.21 ab	72.24 \pm 4.60 b	81.27 \pm 4.13 b	153.5 \pm 2.87 cd
Dose	0.5 %					
7	29.78 \pm 14.53 a	84.61 \pm 4.97 b	114.39 \pm 16.06 ab	82.45 \pm 8.84 ab	66.95 \pm 2.44 a	149.40 \pm 11.06 a
8	29.99 \pm 6.83 a	84.22 \pm 6.03 ab	114.21 \pm 8.75 a	85.53 \pm 1.02 b	61.61 \pm 8.90 a	147.14 \pm 9.46 a
9	57.12 \pm 2.47 a	65.62 \pm 11.96 a	122.74 \pm 12.22 b	64.28 \pm 3.10 a	63.24 \pm 8.86 a	127.53 \pm 9.43 a
Dose	0.1 %					
7	NT	NT	NT	51.91 \pm 10.71 a	50.67 \pm 14.70 a	102.58 \pm 20.2 a
8	NT	NT	NT	65.71 \pm 24.34 a	31.99 \pm 8.74 a	97.70 \pm 27.43 a
9	NT	NT	NT	47.56 \pm 5.07 a	62.65 \pm 14.87 a	110.2 \pm 17.46 a

^a Values are the means of the four replicates, each set up with ten larvae or adults ($n=40$). A: absolute coefficients; R: relative coefficients; T: total coefficients. NT: Not tested. Means followed by the same letters within each column and for each concentration are not significantly different (one-way ANOVA and Tukey's test ($P < 0.05$)).

A significant decrease in feeding of both developmental stages of *A. diaperinus* in the presence of halolactones was observed. Activity of bromolactones was related to the size of the lactone ring. A much stronger antifeedant for both developmental stages of the lesser mealworm was γ -bromo- δ -lactone (**7**) in comparison with the δ -bromo- γ -lactone (**6**). Bromolactone with a larger ring was the most potent antifeedant among the studied compounds (only 5.22% consumption compared with the control) for adults. Its activity is comparable with the most known active antifeedant, azadirachtin¹⁷. This compound also strongly reduced the larvae feeding (consumption in relation to the control was 9.69% in no-choice test, Fig. 3). Replacement of a bromine atom by a chlorine atom leads to an increase of activity of obtained δ -chloro- γ -lactone (**8**). This chlorolactone with a smaller ring reduced three times the larvae feeding and two times the adults feeding in comparison with the δ -bromo- γ -lactone (Fig. 3). Generally, chlorolactones (**8**, **9**) strongly reduced insects feeding, but for larvae the best antifeedant was the one with a larger ring. Its activity was similar to the activity of azadirachtin. In the case of adults this compound (**9**) was a little weaker as feeding deterrent in comparison with δ -chloro- γ -lactone (**8**), but these differences were not significant, especially in the no-choice test (Table 2). Considering the results of the no-choice tests the strongest antifeedants for both developmental stages were halolactones with a larger ring (Fig. 3). These particularly active compounds i.e. γ -bromo- δ -lactone (**7**) and both chlorolactones (**8**, **9**) were also very strong antifeedants against adults at the lower dose i.e. 0.5%. For larvae their activity except for chlorolactone (**9**) in the no-choice test was poor. These compounds used as 0.1% solutions also affected feeding of adults. According to the classification of Nawrot et al. they can be considered as good antifeedants³⁷. These studies show that the halogen atoms are responsible for the high antifeedant activity of lactone derivatives of β -damascone (**1**). A dramatic reduction of consumption of food treated with halolactones obtained from piperitone was also observed. In the presence of the above-mentioned compounds, the food consumed by the adults of the lesser mealworm in the no-choice test represented only 2.16–2.46% of the consumption in the control²¹. Halogen atoms modify also antifeedant activity of the other monoterpenes. For instance, the number and kind of halogen (Cl or Br) substituents in the cyclohexane ring of cyclic monoterpenes isolated from *Plocamium cartilagineum* significantly affects the activity of obtained derivatives towards chrysomelid, *Leptinotarsa*

decemlineata L. and two aphid species *Myzus persicae* Sulz and *Ropalosiphum padi* L.³⁸. The menthol derivatives with halogen substituents show a much stronger activity against mosquitoes than starting product³⁹. Among intermediate products of synthesis of lactones from β -damascone (**1**) a good antifeedant was allylic alcohol - dihydro- β -damascol (**3**) which reduced feeding in the no-choice tests by 75.52 (larvae) and 85.83% (adults) (**Fig. 3**). Adult feeding reduction of over 60% was observed also in the presence of the ester (**4**) and acid (**5**). The results showed a significantly higher sensory sensitivity of adults of *A. diaperinus* in comparison with larvae. Taking into account the total deterrence coefficients among the 11 compounds, only bromolactone (**6**) was clearly weaker deterrent for adults (**Table 2**).

Conclusion

The deterrent activity of individual β -damascone derivatives varied in potency, the time of expression, and the duration of the effect, depending on spatial structure of the lactone and species/developmental stage of the insect.

The most successive structural modifications of β -damascone in terms of antifeedant activity towards *M. persicae* were the incorporation of lactone moiety and concomitant presence of bromine in the side chain. δ -Bromo- γ -lactone (**6**) was the most potent antifeedant at pre-, post- and intraingestional levels. Ester (**4**) and unsaturated γ -lactone (**10**) were active at postingestional levels by impeding aphid settling after longer exposure times.

All β -damascone derivatives with lactone moiety deterred the feeding of adults and larvae of *A. diaperinus*. The potency of the activity was related to the size of the ring and the incorporation of chlorine or bromine into the molecule. Generally, halo- δ -lactones were more active than halo- γ -lactones and the adults were more sensitive to the compounds studied than larvae. Nevertheless, the strongest antifeedants for both developmental stages of the lesser mealworm were γ -bromo- δ -lactone (**7**), δ -chloro- γ -lactone (**8**), and γ -chloro- δ -lactone (**9**).

Experimental section

Reagents

β -Damascone (90% purity), N-bromosuccinimide (99% purity) and N-chlorosuccinimide (98% purity) were purchased from Aldrich, triethylorthoacetate from Fluka.

General Procedures

Analytical TLC was performed on silica gel (Kieselgel 60 F₂₅₄, Merck) with mixture of hexane, acetone and diethyl ether in various ratios as developing systems. Compounds were detected by spraying the plates with solution of Ce(SO₄)₂ (1 g), H₃[P(Mo₃O₁₀)₄] (2 g) in 10% H₂SO₄, followed by heating to 120-200°C.

Column chromatography was performed on silica gel (Kiesel gel 60, 230-400 mesh ASTM, Merck) with a mixture of hexane, acetone and diethyl ether (in various ratios) as eluents.

Gas chromatography was performed on Agilent Technologies 6890N Network GC instrument equipped with autosampler, split injection (20:1) and FID detector using HP-5 column (30 m x 0.32 mm x 0.25 μ m) with hydrogen as carrier gas. The temperature programme was as follows: injector 250°C, detector (FID) 250°C,

column temperature: 100°C, 100-300°C (rate 30°/min), 300°C (hold 2 min).

¹H NMR, ¹³C NMR, DEPT 135, HMQC and ¹H-¹H COSY spectra were recorded in CDCl₃ solutions on a Bruker Avance AMX 300 spectrometer.

IR spectra were determined using Mattson IR 300 Thermo-Nicolet spectrophotometer using KBr pellets or as neat.

Melting points (uncorrected) were determined on Boetius apparatus.

The indexes of refraction were measured on Carl Zeiss Jena refractometer.

Chemistry

Synthesis and separation of compounds

Synthesis of dihydro- β -damascone (**2**)

A solution of β -damascone **1** (4 g, 20.8 mmol) in anhydrous diethyl ether (10 mL) was added dropwise to the stirring solution of LiAlH₄ (0.4 g) in dry diethyl ether (20 mL). After 5 h (GC, TLC) water was added and the product was extracted with diethyl ether. The organic layer was washed with brine, dried over anhydrous MgSO₄ and filtered. The solvent was evaporated in vacuo and pure ketone (**2**) was obtained (3.9 g, 99% yield), with the physical and spectral data consistent with literature report³².

Synthesis of dihydro- β -damascol (**3**)

Using the same procedure of reduction reaction alcohol **3** was obtained (3.6 g, 89% yield) from the ketone (**2**), with the physical and spectral data consistent with literature report³³.

Synthesis of ethyl 2-(2-butylidene-1,3,3-trimethylcyclohexyl)-acetate (**4**)

A mixture of alcohol (**3**) (3.6 g, 18.4 mmol), triethylorthoacetate (21.6 mL, 115.2 mmol) and catalytic amount of propionic acid (1 drop) was heated at 138°C for 10 h under the conditions for distillative removal of ethyl alcohol. When the reaction was completed (GC, TLC), the mixture was chromatographed on silica gel. Elution with hexane/acetone (19:1) gave the pure ester (**4**) (4.7 g, 98% yield); (oily liquid); ¹H NMR (CDCl₃), δ : 0.89 (t, *J* = 7.3 Hz, 3H, CH₃-10), 1.17 (s, 3H, one of the (CH₃)₂C<), 1.20 (s, 3H, one of the (CH₃)₂C<), 1.22 (t, *J* = 7.4 Hz, 3H, CH₃-17), 1.27-1.65 (m, 11H, CH₂-9, CH₂-4, CH₂-5, CH₂-6, CH₃-13), 2.14 (m, 2H, CH₂-8), 2.37 and 2.48 (two d, *J* = 13.7 Hz, 2H, CH₂-14), 4.05 (q, *J* = 7.4 Hz, 2H, CH₂-16), 5.20 (t, *J* = 7.0 Hz, 1H, H-7). ¹³C NMR (CDCl₃), δ : 13.96 and 14.30 (C-10 and C-17), 17.59 (C-5), 23.76 (C-9), 30.51 and 30.64 (C-11 and C-12), 31.73 (C-13), 32.31 (C-8), 35.33 (C-1), 36.33 (C-6), 39.63 (C-3), 41.21 (C-4), 47.70 (C-14), 59.75 (C-16), 126.82 (C-7), 149.07 (C-2), 172.19 (C-15). IR (film, cm⁻¹): 2959 (s), 2932 (s), 2870 (m), 1743 (s), 1193 (w).

Synthesis of 2-(2-butylidene-1,3,3-trimethylcyclohexyl)acetic acid (**5**)

Ester **4** (4 g, 15 mmol) was heated under reflux for 3 h in 10% ethanol solution of KOH (40 mL). After evaporation of ethanol in vacuo the residue was diluted with water and organic impurities were removed by the extraction with diethylether (3 x 40 mL). The water layer was acidified with 0.1 M HCl and the product was extracted with diethyl ether (3 x 40 mL). The ethereal fraction was washed with brine and dried over anhydrous MgSO₄. Evaporation of solvent in vacuo afforded pure acid **5** (3 g, 85% yield); (oily liquid); ¹H NMR (CDCl₃), δ : 0.87 (t, *J* = 6.3 Hz, 3H, CH₃-10), 1.19 (s, 6H, (CH₃)₂C<), 1.23 (s, 3H, CH₃-13), 1.29-1.62 (m, 9H, CH₂-9, CH₂-4, CH₂-5, CH₂-6), 2.11-2.21 (m, 2H, CH₂-8), 2.41 and 2.55 (two d, *J* =

13.8 Hz, 2H, CH₂-14), 5.26 (t, *J* = 7.0 Hz, 1H, H-7), 9.1 (s, 1H, -COOH). ¹³C NMR (CDCl₃), δ: 13.88 (C-10), 17.55 (C-5), 23.63 (C-9), 30.52 i 30.58 (C-11 and C-12), 31.62 (C-13), 32.23 (C-8), 35.29 (C-1), 36.36 (C-6), 39.43 (C-3), 41.29 (C-4), 47.54 (C-14), 127.12 (C-7), 148.88 (C-2), 178.32 (C-15). IR (film, cm⁻¹): 2958 (s), 1706 (s), 1463 (m).

Preparation of bromolactones (6), (7)

A solution of acid (1.86 g, 7.8 mmol) and *N*-bromosuccinimide (7.8 mmol) in THF (30 mL) and a drop of acetic acid were stirred at room temperature for 24 h. The mixture was diluted with diethyl ether, washed with saturated NaHCO₃ solution and brine. The organic layer was dried over anhydrous MgSO₄ and the solvent was evaporated in vacuo. The mixture of products was subjected to column chromatography (hexane/acetone, 3:1). Data of isolated lactones are given below:

7a-(1-Bromobutyl)-3a,7,7-trimethylhexahydrobenzofuran-2-one

(6) (0.42 g, 17% yield); (oily liquid); ¹H NMR (CDCl₃), δ: 0.94 (t, *J* = 7.3 Hz, 3H, CH₃-11), 1.10 (s, 3H, CH₃-14), 1.34 and 1.40 (2s, 6H, (CH₃)₂C<), 1.50-2.33 (m, 10H, CH₂-4, CH₂-5, CH₂-6, CH₂-9, CH₂-10), 2.50 and 2.73 (two d, *J* = 18.1 Hz, 2H, CH₂-3), 4.28 (dd, *J* = 10.3 and 1.2 Hz, 1H, H-8). ¹³C NMR (CDCl₃), δ: 13.36 (C-11), 17.27 (C-10), 22.62 (C-5), 27.03 and 28.11 (C-12 and C-13), 30.11 (C-14), 36.29 (C-9), 38.75 (C-4), 38.97 (C-6), 40.88 (C-3a), 44.30 (C-7), 45.27 (C-3), 59.13 (C-8), 91.91 (C-7a), 175.90 (C-2). IR (film, cm⁻¹): 2958 (m), 2874 (m), 1776 (s), 1463 (w), 1226 (w).

7a-Bromo-3a,7,7-trimethyl-8-propyloctahydroisochromen-3-one

(7) (0.67 g, 27% yield); (colourless crystal); mp: 95-104°C; ¹H NMR (CDCl₃), δ: 0.98 (t, *J* = 7.0 Hz, 3H, CH₃-11), 1.16-1.27 (m, 2H, CH₂-10), 1.32 (s, 3H, CH₃-3a), 1.40 and 1.45 (two s, 6H, (CH₃)₂C<), 1.52-1.97 (m, 8H, CH₂-6, CH₂-5, CH₂-4, CH₂-9), 2.44 and 3.15 (two d, *J* = 19.6 Hz, 2H, CH₂-3), 4.93 (d, *J* = 10.6 Hz, 1H, H-8). ¹³C NMR (CDCl₃), δ: 13.69 (C-11), 18.24 (C-10), 21.22 (C-5), 24.08 (C-14), 28.48 (C-3a), 33.62 and 35.67 (C-12 and C-13), 35.24 (C-9), 35.67 (C-4), 40.70 (C-6), 41.05 (C-3), 84.89 (C-7a), 84.97 (C-8), 170.90 (C-2). IR (KBr, cm⁻¹): 3007 (s), 2918 (s), 1716 (s), 1460 (s).

Preparation of chlorolactones (8), (9)

A solution of acid (1.87 g, 7.8 mmol) and *N*-chlorosuccinimide (7.8 mmol) in THF (30 mL) and a drop of acetic acid were stirred at room temperature for 24 h. The mixture was diluted with diethyl ether, washed with saturated NaHCO₃ solution and brine. The organic layer was dried over anhydrous MgSO₄ and the solvent was evaporated in vacuo. The mixture of products was subjected to column chromatography (hexane/diethyl ether, 3:1). Data of isolated lactones are given below:

7a-(1-Chlorobutyl)-3a,7,7-trimethylhexahydrobenzofuran-2-one

(8) (0.39 g, 18% yield); (oily liquid); ¹H NMR (CDCl₃), δ: 0.94 (t, *J* = 7.3 Hz, 3H, CH₃-11), 1.11 (s, 3H, CH₃-3a), 1.33 and 1.41 (two s, 6H, (CH₃)₂C<), 1.50-2.24 (m, 10H, CH₂-4, CH₂-5, CH₂-6, CH₂-9, CH₂-10), 2.52 and 2.62 (two d, *J* = 18 Hz, 2H, CH₂-3), 4.24 (dd, *J* = 10.6 and 1.6 Hz, 1H, H-8). ¹³C NMR (CDCl₃), δ: 13.43 (C-11), 17.18 (C-10), 21.90 (C-5), 26.61 and 28.27 (C-12 and C-13), 29.60 (C-14), 36.84 (C-9), 37.91 (C-6), 38.82 (C-4), 40.10 (C-3a), 43.96 (C-7), 45.50 (C-3), 66.74 (C-8), 92.63 (C-7a), 175.95 (C-2). IR (film, cm⁻¹): 1464 (s), 1775 (s), 2875 (s), 2959 (s).

7a-Chloro-3a,7,7-trimethyl-8-propyloctahydroisochromen-2-one

(9) (0.36 g, 17% yield); (colourless crystal); mp: 81-96°C; ¹H NMR (CDCl₃), δ: 0.97 (t, *J* = 7.3 Hz, 3H, CH₃-11), 1.27, 1.31 and 1.35 (three s, 9H, (CH₃)₂C<, CH₃-3a), 1.51-2.16 (m, 10H, CH₂-4, CH₂-5, CH₂-6, CH₂-9, CH₂-10), 2.38 and 3.11 (two d, *J* = 19.5 Hz, 2H,

CH₂-3), 4.74 (dd, *J* = 10.3 and 1.6 Hz, 1H, H-8). ¹³C NMR (CDCl₃), δ: 13.72 (C-11), 18.15 (C-10), 21.11 (C-5), 24.63 (C-14), 31.10 and 33.02 (C-12 and C-13), 34.31 (C-9), 35.07 (C-4), 39.95 (C-6), 40.02 (C-7), 40.22 (C-3a), 41.58 (C-3), 80.64 (C-7a), 84.57 (C-8), 175.95 (C-2). IR (KBr, cm⁻¹): 2968 (w), 2931 (w), 2869 (w), 1733 (s), 1463 (s).

Dehydrohalogenation procedure

1,8-Diazabicyclo[5.4.0]undec-7-ene (6 mmol) was added to a solution of the corresponding halolactone (3 mmol) in dry methylene chloride (20 mL) at room temperature. When the reaction was completed (GC, TLC), the mixture was concentrated in vacuo to remove methylene chloride. The residue was diluted with diethyl ether. The ethereal solution was washed with brine, dried (MgSO₄) and evaporated in vacuo. The residue was chromatographed on silica gel. Elution with hexane/acetone (6:1) gave lactones (10) and (11) (47-57% yield) with following data:

7a-((E)-But-1-enyl)-3a,7,7-trimethylhexahydrobenzofuran-2-one

(10) (oily liquid); ¹H NMR (CDCl₃), δ: 0.92 (s, 3H, one of the (CH₃)₂C<), 1.01 (t, *J* = 7.5 Hz, 3H, CH₃-11), 1.03 (s, 3H, one of the (CH₃)₂C<), 1.25 (s, 3H, CH₃-3a), 1.40-1.62 (m, 6H, CH₂-4, CH₂-5, CH₂-6), 1.97 and 2.42 (two d, *J* = 16.3 Hz, 2H, CH₂-3), 2.11 (dq, *J* = 7.5, 6.3 and 1.3 Hz, 2H, CH₂-10), 5.60 (dt, *J* = 15.5 and 1.3 Hz, 1H, H-8), 5.77 (dt, *J* = 15.5 and 6.3 Hz, 1H, H-9). ¹³C NMR (CDCl₃), δ: 13.80 (C-11), 17.75 (C-5), 21.66 and 24.99 (C-12 and C-13), 25.66 (C-10), 26.47 (C-14), 36.25 (C-3a), 36.49 (C-4), 36.57 (C-6), 41.75 (C-7), 45.91 (C-3), 91.98 (C-7a), 125.01 (C-9), 134.33 (C-8), 176.73 (C-2). IR (film, cm⁻¹): 2964 (s), 2932 (s), 1778 (m), 1462 (m), 1289 (m), 1125 (m), 971 (m).

3a,7,7-Trimethyl-8-propylhexahydrocyclopropa[1,2]benzofuran-2(3H)-one (11)

(11) (oily liquid); ¹H NMR (CDCl₃), δ: 0.96 (t, 3H, *J* = 7.3 Hz, CH₃-11), 1.14 and 1.18 (two s, 6H, (CH₃)₂C<), 1.21 (s, 3H, CH₃-3a), 1.29-2.11 (m, 11H, CH₂-6, CH₂-5, CH₂-4, CH₂-9, CH₂-10 and H-3), 4.25 (dd, *J* = 10.9 and 2.2 Hz, 1H, H-8). ¹³C NMR (CDCl₃), δ: 13.92 (C-11), 17.73 (C-10), 17.86 (C-14), 20.45 (C-5), 26.13 (C-12), 28.04 (C-3a), 29.50 (C-13), 30.52 (C-7), 33.11 (C-9), 34.12 (C-3), 37.27 (C-6), 41.51 (C-4), 45.59 (C-7a), 82.73 (C-8), 175.13 (C-2). IR (film, cm⁻¹): 2936 (s), 2873 (s), 1762 (s), 1466 (m), 1327 (m), 1188 (s), 976 (s).

Bioassays

Myzus persicae

Cultures of aphids and their settling (choice-test) were described earlier by Grudniewska et al¹⁸.

Aphid initial behavioural responses (no-choice test).

Aphid behaviour during initial contact with the studied compound was measured by direct monitoring of the freely moving aphids on a treated leaf, using a video recording. This bioassay allows studying the preingestional (activating olfactory receptors) and ingestional (activating gustatory receptors) effects of a chemical, which means that the results may show the mode of action of the compound – whether it is active at plant surface or within plant tissues. The experiment was carried out for 15 min (12 aphids/compound). The following parameters were derived from data obtained in this experiment: total time of aphid presence on the leaf, total time of probing, number of probes, and mean duration of a probe. The time spent on the leaf and the duration of probing were recorded basing on the relationship between antennal and body movements and penetration of the stylets as described by Hardie et al., who

associated styled penetration with the position of antennae parallel to the abdomen and the cessation of body movements⁴⁰.

Statistical analysis. The data derived from choice-test (aphid settling) were analysed using Student t-test. If aphids showed clear preference for the leaf treated with the tested compound ($P < 0.05$), the compound was described as having **attractant** properties. If aphids settled mainly on the control leaf ($P < 0.05$), the compound studied in the respective choice-test was stated a **deterrent**. From the data thus obtained the relative index of deterrence (*DI*) was calculated using the equation:

$$DI = (C - T)/(C + T)$$

where *C* and *T* are the numbers of aphids settled on control and treated leaves, respectively. The value of *DI* ranged between plus 1 (ideal deterrent) and minus 1 (ideal attractant).

The data derived from no-choice-test (aphid behavioural responses) were analyzed by means of one-way analysis of variance (ANOVA) followed by comparison of deterrence coefficients by Tukey's test ($P < 0.05$).

Alphitobius diaperinus

The insect culture of *Alphitobius diaperinus* culture was described earlier by Szczepanik et al²¹.

To investigate the antifeedant activity of compounds studied choice and no-choice tests were used according to the procedure described previously¹⁷. The choice test is very sensitive, but insects could easily avoid treated food. Feeding-deterrent activity of chemical compounds observed in a choice test thus needs to be confirmed in no-choice test. The application of these two types of tests also allowed us to evaluate the mode of action of the compounds studied: did they act on the taste organs only (according to the definition of antifeedants) or were they toxic to the insects?

For the feeding assays, 1 ml of acetone solution of the test compounds at a concentration of 1% were applied to the 1 g of oat flakes (Melvit SA Warsaw, Poland) using a micropipette. After evaporation of acetone, weighed flakes were placed in Petri dishes and offered to 10 larvae or 10 unsexed adults during the following three-day period. All trials were kept in an incubator at $29 \pm 1^\circ\text{C}$ in the dark. The amount of food eaten was the basis for calculating the deterrence coefficients (relative R and absolute A) according to the formulae^{36,41}.

$$R = (C - E) / (C + E) \times 100 \text{ (choice test)}$$

$$A = (CC - EE) / (CC + EE) \times 100 \text{ (no-choice test)}$$

where C, CC and E, EE are the weights of the control and the treated food consumed by the insects in the choice and no-choice tests, respectively.

The total deterrence coefficient, which ranged from -200 to +200, served as an activity index. A compound for which both indices reach a value of 100 and the sum 200 is called an ideal deterrent. Activity of the tested compounds was evaluated according to the classification of Nawrot et al. and Koul^{16,42}

The compounds with the highest activity (7, 8, 9) were used for further studies as 0.5 and 0.1% solutions. Because 0.5% solution moderately decreased feeding of larvae, the lowest concentration of these halolactones was used only against adults. The research

methodology was the same as in the case of 1% solutions. Four replicates of both tests for each compound were conducted on each insect life stage.

From a practical point of view, it is important to evaluate the level of insect feeding in a no-choice situation. To estimate and compare larval and adult feeding levels in the no-choice tests, the amount of treated food consumed was expressed as a percentage of the consumption in the control according to the formula:

$$(EE/CC) \times 100$$

where *EE* and *CC* are the weights of the treated and control food consumed in no-choice tests, respectively.

Statistical analysis. Antifeedant activity was analyzed by means of one-way analysis of variance (ANOVA) followed by comparison of deterrence coefficients by Tukey's test ($P < 0.05$) using PAST (Paleontological Statistics) software²⁶. The Student's t-test (Microsoft Office 2010, Excel) was used to compare the mean values of consumption by the larvae and adults in the no-choice test.


Acknowledgements

This Project was financed by European Union from the European Regional Development Found Grant No. POIG.01.03.01-00-158/09-03 and partly by Nicolaus Copernicus University, Rector's grant no. 502-B/2013.

References

1. E. Demole, P. Enggist, U. Sauberli, M. Stoll, E.Sz. Kovats, *Helv. Chim. Acta.*, 1970, 53, 541-551.
2. F.C. Huang, G. Horváth, P. Molnár, E. Turcsi, J. Deli, J. Schrader, G. Sandmann, H. Schmidt, W. Schwab, *Phytochemistry*, 2009, 70, 457-464.
3. W. Kawasaki, K. Matsui, Y. Akakabe, N. Itai, T. Kajiwar, *Phytochemistry*, 1998, 49, 1565-1568
4. H.Z. Shi, G.S. Liu, Z.F. Xie, C. Li, Y.W. Wei, F. Wang, *Acta Tobacaria Sin*, 2008, 14, 23-27.
5. L. Poisson, P. Schieberle, *J. Agric. Food Chem.*, 2008, 56, 5813-5819.
6. E. Boido, A. Lloret, K. Medina, L. Farina, F. Carrau, G. Versini, E. Dellacassa, *J. Agric. Food Chem.*, 2003, 51, 5408-5413.
7. K. Bauer, D. Garbe, H. Surburg, *Common Fragrance and Flavor Materials*, WILEY-VCH, 1997, 65-66.
8. C. Gerhauser, K. Klimo, W. Hümmer, J. Hölzer, A. Petermann, A. Garreta-Rufas, F-D. Böhmer, P. Schreier, *Mol. Nutr. Food Res.*, 2009, 53, 1237-1244, DOI 10.1002/mnfr.200800492.
9. P.E. Kaufman, R.S. Mann, J.F. Butler, *Pest Manag. Sci.*, 2010, 67, 26-35.
10. J.F. Butler, *Insect Repellents: Principles, Methods and Use*, ed. By Debboun S., Frances S.P., Strickmann D., CRC Press, Taylor and Francis Group, New York, NY, 2006, 161-194.
11. L.R. Askham, C.F. Dunham, L.D. Felix, 2005, US Patent 6958146.
12. M. Hori, *J. Chem. Ecol.*, 1998, 24:1425-1432.

13. M. Hori, *Appl. Entomol. Zool.*, 1999, 34: 113–118.
14. A. Halarewicz-Pacan, B. Gabryś, K. Dancewicz, C. Wawrzęńczyk, *J. Plant Protect. Res.*, 2003, 43 (2): 133–142.
15. Gabrys, B., Dancewicz, K., Halarewicz-Pacan, A., Janusz, E. 2005. *IOBC/WPRS Bull.* 28:29–34.
16. J. Nawrot, I. Dams, C. Wawrzęńczyk, *J. Stored Prod. Res.*, 2009, 45, 221–225.
17. M. Szczepanik, I. Dams, C. Wawrzęńczyk, *Entomol. Exp. Appl.*, 2008, 128, 337–345.
18. Grudniewska, K. Dancewicz, A. Białońska, Z. Ciunik, B. Gabryś, C. Wawrzęńczyk, *RSC Advances*, 2011, 1, 498–510.
19. Gliszczynska, D. Masłowiec, M. Szczepanik, J. Nawrot, C. Wawrzęńczyk, *Przemysł Chemiczny*, 2011, 90, 759–763.
20. K. Dancewicz, B. Gabryś, I. Dams, C. Wawrzęńczyk, *Progress in Plant Protection*, 2008, 48: 817–821.
21. M. Szczepanik, A. Grudniewska, B. Zawitowska, C. Wawrzęńczyk, *Pest Manag. Sci.*, 2014, 70, 953–958.
22. P. W. Wellings, S. A. Ward, A. F. G. Dixon, R. Rabbinge, *Crop Loss Assessment*, 1989, 49–64.
23. Ch-A. Dedryver, A. Le Ralec, F. F. Dedryver, 2010. *Comptes Rendus Biologies*, 2010, 539–553
24. R. L. Blackman, V. F. Eastop, *Aphids on the World's Crops: An Identification Guide*. John Wiley and Sons. Chichester, 1985, New York, Brisbane, Toronto, Singapore.
25. R. W. Miller, R. E. Redfern, *J. Econ. Entomol.*, 1988, 81, 1137–1139.
26. I. M. Rueda, R. C. Axtell, *J. Agric. Entomol.*, 1997, 14, 81–91.
27. W.C. Hazeleger, N. M. Bolder, R. R. Beumer, W. F. Jacobs-Reitsma, *Appl. Environ. Microbiol.*, 2008, 74, 6887–6891.
28. M. Szczepanik, A. Szumny, A. Grudniewska, C. Wawrzęńczyk, *Pesticides*, 2005, 4, 25–32.
29. O. Koul, G. S. Dhaliwal, *Phytochemical biopesticides*. CRC Press. 2003, 236.
30. J. Pettersson, W. F. Tjallingii, J. Hardie, *Host plant selection and feeding*. In: Van Emden H.F., Harrington R. *Aphids as crop pests*. CAB International, 2007, 87–114.
31. J.S. Zhang, L. Ma, R.Y. Xu, E.M. Yao, G.M. Yang, *Yanacao Keji*, 2007, 2, 26–29.
32. K. Mori, M. Ohki, K. Okada, Y. Takei, M. Matsui, *Agr. Biol. Chem.*, 1973, 37, 2907–2911.
33. K.H. Schulte-Elte, *Ger. Offen.*, 1973, DE 2244680 A1 19730322.
34. W.S. Johnson, L. Werthemann, W.R. Bartlett, T.J. Brocksom, T.T. Li, D.J. Faulkner, M.R. Petersen, *J. Am. Chem. Soc.*, 1970, 92, 741–743.
35. R. Obara, A. Szumny, A. Żołnierczyk, T. Olejniczak, A. Białońska, Z. Ciunik, C. Wawrzęńczyk, *Polish J. Chem.*, 2005, 79, 981–993.
36. H. A. van Hoof, *An investigation of the biological transmission of a non-persistent virus*, 1958, 112
37. J. Nawrot, E. Błoszyk, J. Harmatha, L. Novotny, Drożdż B, *Acta Entomol. Bohemos.*, 1986, 83, 327–335.
38. V. H. Argandoña, J. Roviroso, A. San- Martín, A. Riqueleme, A. R. Díaz-Marrero, M. Cueto, J. Darías, O. Santana, A. Guadaño, A. Gonzáles-Coloma. *J. Agric. Food Chem.*, 2002, 50, 7029–7033.
39. R. Samarasekera, I. S. Weerasinghe, K. D. P. Hemalal. *Pest Manag. Sci.*, 2008, 64, 290–295.
40. J. Hardie, M. Holyoak, N. J. Taylor, D. C. Griffiths, *Entomol. Exp. Appl.*, 1992, 62, 233–239.
41. P. Harrewijn, Resistance mechanisms of plant genotypes to various aphid species. In *Aphid - Plant Genotype Interactions*, Campbell, R. K.; Eikenbary, R. D., Eds. Elsevier: Amsterdam, 1990, 117–130.
42. O. Koul, *Insect Antifeedants*, 2004, CRC Press, London/New York/Washington, pp. 25–42.



We report the chemical modification of *t*-β-damascone (**1**) into new lactones derivatives and their biological activity towards aphids and lesser mealworm.