

# RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.


You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## Copper-catalyzed carbonylative Suzuki coupling of aryl iodides with arylboronic acids under ambient pressure of carbon monoxide

Laijin Cheng, Yanzhen Zhong, Zhuchao Ni, Hongyan Du, Fengli Jin, Qi Rong, and Wei Han\*

– Table of Contents Entry –



# Copper-catalyzed carbonylative Suzuki coupling of aryl iodides with arylboronic acids under ambient pressure of carbon monoxide<sup>‡</sup>

Laijin Cheng,<sup>a</sup> Yanzhen Zhong,<sup>a</sup> Zhuchao Ni,<sup>a</sup> Hongyan Du,<sup>a</sup> Fengli Jin,<sup>a</sup> Qi Rong,<sup>a</sup> and Wei Han<sup>\*a</sup>

Received (in XXX, XXX) Xth XXXXXXXXX 200X, Accepted Xth XXXXXXXXX 200X

First published on the web Xth XXXXXXXXX 200X

DOI: 10.1039/b000000x

An efficient and ligandless nanocopper-catalyzed carbonylative cross-coupling of aryl iodides with arylboronic acids at ambient CO pressure in poly(ethylene glycol), has been developed. This protocol is general, practical, and recyclable, and offers an attractive alternative to the corresponding palladium-mediated carbonylative Suzuki process for the construction of biaryl ketone scaffolds.


Cross-coupling of aryl halides, arylboronic acids and carbon monoxide (the carbonylative Suzuki reaction<sup>1</sup>) has been developed as a powerful tool for the construction of biaryl and heterobiaryl ketones. Moreover, these structural motifs are core scaffolds that are found in a tremendous range of pharmaceuticals, advanced organic materials, and photosensitizers.<sup>2</sup> Consequently, establishing a mild, practical, economical, and efficient access to biaryl and heterobiaryl ketones is of great significance.

Since the first example of carbonylative Suzuki reaction was reported by Suzuki et al.<sup>1</sup> extensive work has been done to make the transformation more efficient and to widen its substrate scope.<sup>3</sup> However, the overwhelming majority of the reported methods requires a high pressure (often  $\geq 5$  bar). Although few examples of carbonylative Suzuki coupling of aryl iodides under ambient pressure of carbon monoxide have been reported<sup>4</sup>, palladium remains the metal of choice in almost all cases, causing several shortcomings, particularly from the perspective of its pragmatic application because of the high cost, toxicity of the metal, and the necessity of utilizing expensive ligands. To address these issues, the development of more sustainable, economical, and eco-friendly protocol for the aforementioned transformation is particularly desirable. Very recently, we first reported iron replaced palladium to catalyze carbonylative Suzuki reaction and satisfactory results can be obtained.<sup>5</sup> However, this protocol is inefficient to catalyzed successive carbonylation of aryl diiodides.

Copper has a strong advantage in terms of cost and toxicity compared to palladium. And copper has been widely applied in organic synthesis and oftentimes emerges as an alternative to conduct established palladium-catalyzed cross-coupling reactions.<sup>6</sup> Copper-based catalysts have been broadly employed in oxidative carbonylation of alcohols with carbon monoxide.<sup>7</sup> Comparatively, copper-catalyzed carbonylation of aryl halide is quite limited. Moreover, the limited examples of aminocarbonylation<sup>8</sup> and carbonylative sonogashira<sup>9</sup> of aryl iodide require high pressures of carbon monoxide and ligands to ensure efficient catalysis. Copper-catalyzed carbonylation

of diphenyliodonium tetrafluoroborate with arylboronic acids for the synthesis of biaryl ketones was reported by Kang and co-workers.<sup>10</sup> Since diphenyliodonium tetrafluoroborate is made from phenyl iodide, it would be advantageous to directly carbonylate aryl iodides. Unfortunately, until to date no general copper-catalyzed carbonylative Suzuki reaction of aryl halide exists. We report here first example of copper-catalyzed carbonylative Suzuki reaction of aryl iodide with arylboronic acid under atmospheric pressure of carbon monoxide. Our previous work demonstrated palladium nanoparticles catalyzed carbonylative Suzuki reaction of aryl

**Table 1** Nanocopper-catalyzed carbonylative Suzuki reaction of **1a** with **2a**<sup>a</sup>



Ent ry	acid	base	solvent	yield of <b>3aa</b> (%)	yield of <b>3a'a'</b> (%)
1	-	K <sub>3</sub> PO <sub>4</sub>	PEG-400	73	14
2	<i>t</i> -BuCOOH	K <sub>3</sub> PO <sub>4</sub>	PEG-400	90(75) <sup>b</sup>	< 5
3	CH <sub>3</sub> COOH	K <sub>3</sub> PO <sub>4</sub>	PEG-400	25	< 5
4	CF <sub>3</sub> COOH	K <sub>3</sub> PO <sub>4</sub>	PEG-400	43	trace
5	<i>t</i> -BuCOOH	K <sub>2</sub> CO <sub>3</sub>	PEG-400	55	43
6	<i>t</i> -BuCOOH	Na <sub>2</sub> CO <sub>3</sub>	PEG-400	82	8
7	<i>t</i> -BuCOOH	KF	PEG-400	72	trace
8	<i>t</i> -BuCOOH	TBAF	PEG-400	5	-
9	<i>t</i> -BuCOOH	AcOK	PEG-400	21	trace
10	<i>t</i> -BuCOOH	K <sub>3</sub> PO <sub>4</sub>	PEG-200	82	10
11	<i>t</i> -BuCOOH	K <sub>3</sub> PO <sub>4</sub>	PEG-2000	5	-
12	<i>t</i> -BuCOOH	K <sub>3</sub> PO <sub>4</sub>	Ethylene glycol	10	89
13	<i>t</i> -BuCOOH	K <sub>3</sub> PO <sub>4</sub>	PEGDM-190	8	< 5
14 <sup>c</sup>	<i>t</i> -BuCOOH	K <sub>3</sub> PO <sub>4</sub>	PEG-400	82	< 5
15 <sup>d</sup>	<i>t</i> -BuCOOH	K <sub>3</sub> PO <sub>4</sub>	PEG-400	45	trace
16 <sup>e</sup>	<i>t</i> -BuCOOH	K <sub>3</sub> PO <sub>4</sub>	PEG-400	47	trace

<sup>a</sup>Reaction conditions (unless otherwise stated): **1a** (0.5 mmol), **2a** (0.75 mmol), CO (balloon), base (1.0 mmol), acid (0.25 mmol), nanocopper (20 mol %), solvent (2.0 mL), 80 °C. <sup>b</sup>Nanocopper (10 mol %). <sup>c</sup>Cu<sub>2</sub>O (20 mol %). <sup>d</sup>CuCl (20 mol %). <sup>e</sup>CuBr (20 mol %).


iodide with a high level of efficacy even in the absence of a ligand in PEG-400 [poly(ethylene glycol) with an average molecular weight of 400 Da].<sup>4a</sup> This work prompts us to search for nanocopper-catalyzed carbonylative Suzuki reaction without the assistance of an additional ligand.

Initially, we employed copper nanoparticles for the catalysis of carbonylative Suzuki reaction of 4-iodonitrobenzene with phenylboronic acid and carbon monoxide (1 atm) to identify optimum reaction conditions. Selected results are summarized in Table 1. On the basis of our previous work of developing PEG-based ligand-free transition metal-catalyzed cross coupling reactions,<sup>5,4a,11</sup> we performed the model reaction in PEG-400 with K<sub>3</sub>PO<sub>4</sub> as a base. Consequently, the desired product **3aa** can be obtained in 73% yield, whereas the side product from Suzuki coupling was also formed in 14% yield (Table 1, entry 1). Very recently, we reported pivalic acid effectively suppressed Suzuki coupling and favored carbonylative Suzuki coupling.<sup>4a</sup> As expected, in the presence of the pivalic acid an excellent yield of **3aa** was generated with a little amount of the side product (< 5%) (Table 1, entry 2). Other acids such as CH<sub>3</sub>COOH and CF<sub>3</sub>COOH just gave poor results (Table 1, entries 3 and 4). Successively, other various bases were tested. Na<sub>2</sub>CO<sub>3</sub> and KF showed superior results compared with other bases such as K<sub>2</sub>CO<sub>3</sub>, TBAF, and AcOK (Table 1, entries 5–9), albeit with less efficiency in terms of yields than K<sub>3</sub>PO<sub>4</sub>. Of the solvents evaluated, PEG-400 gave the best results (Table 1, entry 2). Notably, ethylene glycol and poly(ethylene glycol) dimethyl ether (PEGDM-190) were found to be ineffective (Table 1, entries 12 and 13), suggesting that the unique structure of PEG-400 bearing both hydroxyl groups and a polyether chain played a critical role in this reaction. Other catalyst, such as Cu<sub>2</sub>O, CuCl, and CuBr were examined and were observed to be less effective than nanocopper catalyst (Table 1, entries 14–16).

Next, we investigated the first copper-catalyzed method for carbonylative Suzuki reactions of various aryl iodides. However, we observed that more unactivated substrates compared with 4-iodonitrobenzene were unsuitable to the above-established reaction conditions. In the table 1, KF as a base gave a better selectivity than K<sub>3</sub>PO<sub>4</sub>, albeit with moderate efficiency (entry 3). After a concise reevaluation, we found out that the addition of KF (0.5 equiv) can ensure effective catalysis at an elevated temperature (100 °C) (Table 2). The carbonylative Suzuki reactions of aryl iodides with phenylboronic acid demonstrated that electron-deficient as well as electron-rich aryl iodides worked well. And the electronic nature of the aryl iodides had a minor impact on the observed yields. Fluoro- and trifluoromethyl-substituted aryl iodides were reactive and afforded the corresponding products in 75–91% yields (Table 2, entries 2, and 4–6), which are potentially useful for the synthesis of pharmaceuticals and advanced materials.<sup>12</sup> Sterically hindered 2-iodotoluene also proceeded smoothly (Table 2, entry 9). We were pleased to find that a heterocycle 4-iodo-3,5-dimethylisoxazole (**1o**) and 1-iodonaphthalene were suitable substrates and provided the expected products in 90% and 87%, respectively (Table 2,

entries 15 and 14). Intriguingly, 1,4-diiodobenzene (**1p**) and 1,3-diiodobenzene (**1q**) could undergo successive carbonylation effectively and furnish corresponding diketones that are important intermediates for advanced functional materials,<sup>2a,2d</sup> in 87% and 75% yields, respectively (Scheme 1).

**Table 2** Nanocopper-catalyzed carbonylative Suzuki reaction of **2a** with various aryl iodides<sup>a</sup>




Entry	aryl iodide	product	time (h)	yield (%) <sup>b</sup>
-------	-------------	---------	----------	------------------------

1 <sup>c</sup>			5	90
2			9	75
3			12	89
4			12	91
5			12	90
6			4	91
7			9	80
8			5	95
9			24	93
10			9	81
11			9	91
12			9	92
13			12	83
14			12	87
15			9	90


<sup>a</sup>Reaction conditions(unless otherwise stated): **1a** (0.5 mmol), **2a** (0.75 mmol), CO (balloon), K<sub>3</sub>PO<sub>4</sub> (0.5 mmol), *t*BuCOOH (0.25 mmol), nanoCu (20 mol%), PEG-400 (2.0 mL), 100 °C. <sup>b</sup>Yield of isolated product after column chromatography. <sup>c</sup> **1a** (0.5 mmol), **2a** (0.75 mmol), CO (balloon), K<sub>3</sub>PO<sub>4</sub> (1.0 mmol), *t*BuCOOH (0.25 mmol), nanocopper (20 mol %), PEG-400 (2.0 mL), 80 °C.

To further explore the scope, we extensively surveyed the carbonylative Suzuki reactions of various aryl iodides with representative arylboronic acids under the optimized conditions (Scheme 2). In general, unsubstituted, *para*-, *meta*-, and *ortho*-substituted arylboronic acids afforded desired products in good to excellent yields. The electronic nature of the arylboronic acids was observed to be loosely correlated with their reactivity. The reaction conditions were compatible with methyl, methoxy, aldehyde, cyano, fluoro, and chloro groups. Treatment of naphthalen-2-ylboronic acid (**2k**) with **1n** produced desired product (**3nk**) in 82% yield. To our delight, a heterocyclic boronic acid **2m** was quite effective


and provided desired product **3km** in 90% yield [eqn (1)].



**Scheme 1** Successive carbonylative Suzuki reactions of diiodobenzenes




**Scheme 2** Carbonylative Suzuki reactions of aryl iodides with arylboronic acids. *Reaction conditions* (unless otherwise stated): **1** (0.5 mmol), **2** (0.75 mmol), CO (balloon), K<sub>3</sub>PO<sub>4</sub> (0.5 mmol), KF (0.25 mmol), *t*BuCOOH (0.25 mmol), nano-Cu (20 mol %), PEG-400 (2.0 mL), 100 °C. Yield of isolated product after column chromatography was given. <sup>a</sup>**1g** (0.5 mmol), **2g** (0.75 mmol), CO (balloon), K<sub>3</sub>PO<sub>4</sub> (1.0 mmol), *t*BuCOOH (0.25 mmol), nanocopper (20 mol %), PEG-400 (2.0 mL), 80 °C.




To check the reusability of PEG-400 and nanocopper catalyst, the model reaction of 4-iodonitrobenzene (**1a**) with phenylboronic acid (**2a**) and carbon monoxide (1 atm) was tested (Figure 1). The reaction mixture was extracted with diethyl ether, and then, the nanocopper–PEG-400 proceeded to the second run by charging with the same substrates under carbon monoxide (1 atm). Gratifyingly, the catalytic system can be recycled up to ten times with good yields (Figure 1). And Cu leaching in the organic phase (using ICP-MS) was observed to be 1.2% of the starting Cu.





**Figure 1** Reusability of catalyst. *Reaction conditions:* **1a** (0.5 mmol), **2a** (0.75 mmol), CO (balloon),  $K_3PO_4$  (1.0 mmol),  $tBuCOOH$  (0.25 mmol), nanocopper (20 mol %), PEG-400 (2.0 mL), 80 °C.

Although commercially available nanocopper was directly employed to catalyze the carbonylative Suzuki reactions, we still wondered whether the catalysis occurred on the cluster surface or by leached copper species.<sup>13</sup> A mercury additive was used to test nano-metal catalyst.<sup>14</sup> Control experiments were conducted, as shown in Scheme 3. The reaction of **1k** with **2a** under carbon monoxide (1 atm) was almost completely inhibited when 200 equiv of Hg(0) with respect to copper was added. Thus, it is likely that copper nanoparticles are the real working catalyst.



**Scheme 3** The effect of the presence of Hg additive

In conclusion, we have developed the first copper-catalyzed carbonylative Suzuki reactions of aryl iodides with arylboronic acids, which affords the best product yields in the absence of a ligand and under atmospheric pressure of carbon monoxide. The transformation tolerates a variety of functional groups on both coupling partners. Diiodobenzenes such as 1,4-diiodobenzene (**1p**) and 1,3-diiodobenzene (**1q**) can also undergo successive carbonylative Suzuki reactions. Notably, the catalytic system is based on environmentally benign solvent and can be recycled up to ten times with satisfactory yields. Control experiments suggest that copper nanoparticles likely catalyzed the reactions.

## Notes and references

<sup>a</sup> Jiangsu Collaborative Innovation Center of Biomedical Functional Materials, Jiangsu Key Laboratory of Biofunctional Materials, Key Laboratory of Applied Photochemistry, School of Chemistry and Materials Science, Nanjing Normal University, Wenyuan Road NO.1, 210023 Nanjing (China). Fax: (+) 86-(0)25-8589-1455  
E-mail: whhanwei@gmail.com

† Electronic Supplementary Information (ESI) available. See DOI: 10.1039/b000000x/

‡ The work was sponsored by the Natural Science Foundation of China (21302099), the Natural Science Foundation of Jiangsu Province (BK2012449), the Natural Science Foundation of Jiangsu Provincial Colleges and Universities (12KJB150014), the Scientific Research Start-up Foundation of Nanjing Normal University (2011103XGQ0250), the SRF for ROCS, SEM, and the Priority Academic Program Development of Jiangsu Higher Education Institutions.

- (a) T. Ishiyama, H. Kizaki, N. Miyaura and A. Suzuki, *Tetrahedron Lett.*, 1993, **34**, 7595.
- (a) C. Y. K. Chan, Z. J. Zhao, J. W. Y. Lam, J. Z. Liu, S. M. Chen, P. Lu, F. Mahtab, X. J. Chen, H. H. Y. Sung, H. S. Kwok, Y. G. Ma, I. D. Williams, K. S. Wong and B. Z. Tang, *Adv. Funct. Mater.*, 2012, **22**, 378; (b) K. Maeyama, K. Yamashita, H. Saito, S. Aikawa and Y. Yoshida, *Polym. J.*, 2012, **44**, 315; (c) A. Wen, Z. Wang, T. Hang, Y. Jia, T. Zhang, Y. Wu, X. Gao and Z. Yang, *J. Chromatogr. B*, 2007, **856**, 348; (d) W. L. Zhao and E. M. Carreira, *Org. Lett.*, 2006, **8**, 99; (e) A. L. Ong, A. H. Kamaruddin and S. Bhatia, *Process Biochem.*, 2005, **40**, 3526; (f) M. Furusawa, Y. Ido, T. Tanaka, T. Ito, K. Nakaya, I. Ibrahim, M. Ohyama, M. Inuma, Y. Shirataka and Y. Takahashi, *Helv. Chim. Acta.*, 2005, **88**, 1048; (g) F. Bosca, M. A. Miranda, *J. Photochem. Photobiol. B*, 1998, **43**, 1; (h) N. De Kimpe, M. Keppens and G. Froncz, *Chem. Commun.*, 1996, **5**, 635; (i) G. Dorman and G. D. Prestwich, *Biochemistry*, 1994, **33**, 5661.
- For some recent reviews on Pd-catalyzed carbonylations of aryl halides, see: (a) S. T. Gadge and B. M. Bhanage, *RSC Adv.*, 2014, **4**, 10367; (b) X.-F. Wu, H. Neumann and M. Beller, *Chem. Rev.*, 2013, **113**, 1; (c) X.-F. Wu, H. Neumann and M. Beller, *Chem. Soc. Rev.*, 2011, **40**, 4986; (d) R. Grigg and S. P. Mutton, *Tetrahedron*, 2010, **66**, 5515; (e) A. Brennfürher, H. Neumann and M. Beller, *Angew. Chem. Int. Ed.*, 2009, **48**, 4114.
- (a) Q. Zhou, S. H. Wei and W. Han, *J. Org. Chem.*, 2014, **79**, 1454; (b) M. Z. Cai, J. Peng, W. Y. Hao and G. D. Ding, *Green Chem.*, 2011, **13**, 190; (c) H. L. Li, M. Yang, Y. X. Qi and J. J. Xue, *Eur. J. Org. Chem.*, 2011, 2662; (d) M. Z. Cai, G. M. Zheng, L. F. Zha and J. Peng, *Eur. J. Org. Chem.*, 2009, 1585; (e) G. M. Zheng, P. P. Wang and M. Z. Cai, *Chin. J. Chem.*, 2009, **27**, 1420; (f) B. M. O'Keefe, N. Simmons and S. F. Martin, *Org. Lett.*, 2008, **10**, 5301; (g) S. Z. Zheng, L. W. Xu and C. G. Xia, *Appl. Organometal. Chem.*, 2007, 772; (h) P. Prediger, A. V. Moro, C. W. Nogueira, L. Savegnago, P. H. Menezes, J. B. T. Rocha and G. Zeni, *J. Org. Chem.*, 2006, **71**, 3786; (i) M. J. Dai, B. Liang, C. H. Wang, Z. J. You, J. Xiang, G. B. Dong, J. H. Chen and Z. Yang, *Adv. Synth. Catal.*, 2004, **346**, 1669; (j) B. M. O'Keefe, N. Simmons and S. F. Martin, *Tetrahedron*, 2001, **67**, 4344; (k) T. Ishiyama, H. Kizaki, T. Hayashi, A. Suzuki and N. Miyaura, *J. Org. Chem.*, 1998, **63**, 4726-4731; (l) T. Ishiyama, H. Kizaki, N. Miyaura and A. Suzuki, *Tetrahedron Lett.*, 1993, **34**, 7595.
- Y. Z. Zhong and W. Han, *Chem. Commun.*, 2014, **50**, 3874.
- For special reviews on copper-catalyzed cross-couplings, see: (a) J. Magano and J. R. Dunetz, *Chem. Rev.*, 2011, **111**, 2177; (b) H. H. Rao and H. Fu, *Synlett*, 2011, 745; (c) D. S. Surry and S. L. Buchwald, *Chem. Sci.*, 2010, **1**, 13; (d) G. Evano, N. Blanchard and M. Toumi, *Chem. Rev.*, 2008, **108**, 3054; (e) F. Monnier and M. Taillefer, *Angew. Chem. Int. Ed.*, 2008, **47**, 3096; (f) J. P. Corbet and G. Mignani, *Chem. Rev.*, 2006, **106**, 2651; (g) I. P. Beletskaya, and A. V. Chepurkov, *Coord. Chem. Rev.*, 2004, **248**, 2337; (h) S. V. Ley and A. W. Thomas, *Angew. Chem. Int. Ed.*, 2003, **42**, 5400; (i) J. Hassan, M. Sevignon, C. Gozzi, E. Schulz and M. Lemaire, *Chem. Rev.*, 2002, **102**, 1359; (j) P. Siemsen, R. C. Livingston and F. Diederich, *Angew. Chem. Int. Ed.*, 2000, **39**, 2632.
- For selective examples of copper-catalyzed oxidative carbonylation of alcohols, see: (a) M. Casiello, A. Monopoli, P. Cotugno, A. Milella, M. M. Dell'Anna, F. Ciminale and A. Nacci, *Journal of Molecular Catalysis A: Chemical*, 2014, **381**, 99; (b) H. Xiong, W. L. Mo, J. L. Hu, R. X. Bai and G. X. Li, *Ind. Eng. Chem. Res.*, 2009, **48**, 10845; (c) S. A. Anderson and T. W. Root, *J. Mol. Catal. A: Chem.*, 2004, **220**, 247; (d) V. Raab, M. Merz and J. Sundermeyer, *J. Mol. Catal. A: Chem.*, 2001, **175**, 51; (e) Y. Sato, M. Kagotani, T. Yamamoto, Y. Souma, *Appl. Catal. A Gen.*, 1999, **185**, 219; (f) R. Tesel, M. M. Mauri and P. Rebora, *Ind. Eng. Chem. Prod. Res. Dev.*, 1980, **19**, 396.

- 8 J. M. Liu, R. Z. Zhang, S. F. Wang, W. Sun and C. G. Xia, *Org. Lett.*, 2009, **11**, 1321.
- 9 P. Tambade, Y. Patil, N. Nandurkar and B. Bhanage, *Synlett.*, 2008, **6**, 886.
- 5 10 S.-K. Kang, T. Yamaguchi, T.-H. Kim and P.-S. Ho, *J. Org. Chem.*, 1996, **61**, 9082.
- 11 (a) L. F. Yao, Q. Zhou, W. Han and S. H. Wei, *Eur. J. Org. Chem.*, 2012, 6856; (b) W. Han, N. Liu, C. Liu and Z. L. Jin, *Chin. Chem. Lett.*, 2010, **21**, 1411; (c) W. Han, C. Liu and Z. L. Jin, *Adv. Synth. Catal.*, 2008, **350**, 501; (d) W. Han, C. Liu and Z. L. Jin, *Org. Lett.*, 2007, **9**, 4005.
- 12 (a) R. B. Appell, L. T. Boulton, E. D. Daus, M. Hansen, C. H. Hanson, J. Heinrich, C. Kronig, R. C. Lloyd, D. Louks, M. Nitz, C. Praquin, J. A. Ramsden, H. Samuel, M. Smit and M. Willets, *Org. Process Res. Dev.*, 2013, **17**, 69; (b) Z. Y. Yang, Z. G. Chi, T. Yu, X. Q. Zhang, M. N. Chen, B. J. Xu, S. W. Liu, Y. Zhang and J. R. Xu, *J. Mater. Chem.*, 2009, **19**, 5541.
- 15 13 (a) L. D. Pachón and G. Rothenberg, *Appl. Organomet. Chem.*, 2008, **22**, 288; (b) D. Astruc, *Inorg. Chem.*, 2007, **46**, 1884; (c) M. B. Thathagar, J. E. Elshof and G. Rothenberg, *Angew. Chem. Int. Ed.*, 2006, **45**, 2886; (d) M. B. Thathagar, J. Beckers and G. Rothenberg, *Adv. Synth. Catal.*, 2003, **345**, 979.
- 20 14 J. A. Widegren and R. G. Finke, *J. Mol. Catal. A: Chem.*, 2003, **198**, 317.