

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Structural basis for Na⁺-sensitivity in dopamine D2 and D3 receptors

Cite this: DOI: 10.1039/x0xx00000x

M. Michino,^{ab} R. B. Free,^c T. B. Doyle,^c D. R. Sibley^c and L. Shi^{*abc}Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

To understand the structural basis for the Na⁺-sensitivity of ligand binding to dopamine D2-like receptors, using computational analysis in combination with binding assays, we identified interactions critical in propagating the impact of Na⁺ on receptor conformations and on the ligand-binding site. Our findings expand the pharmacologically-relevant conformational spectrum of these receptors.

The phenomenon of Na⁺-sensitivity is conserved among many members of class A G-protein coupled receptors, such as the adrenergic², dopaminergic³, adenosine⁴, opioid⁵, and neurotensin⁶ receptors. The dopamine D2-like receptors, consisting of the D2, D3, and D4 receptors (D2R, D3R, and D4R), are coupled to the G-protein α subunits (Gi/o) that inhibit adenylyl cyclase⁷. The binding property of some ligand classes of these receptors is known to be sensitive to sodium ion (Na⁺)^{3,8} – a physiologically relevant level of Na⁺ (~150 mM) has been shown to decrease the affinity of agonists including the endogenous agonist dopamine, while enhancing the affinity for some antagonists^{3,9}, compared to those in the absence of Na⁺. Although this phenomenon of Na⁺-sensitivity has been experimentally known over many years, the structural basis for the allosteric effects of Na⁺ on agonist and antagonist binding has not been elucidated.

In D2R, the mutation of Asp80(2.50)[†] to Ala or Glu was shown to abolish Na⁺-sensitivity⁸, and it was proposed that the residues near Asp80(2.50) form a square pyramidal Na⁺ binding site in the D2R¹⁰. In recent years, several ultra-high-resolution crystal structures of class A G-protein coupled receptors have revealed that the Na⁺-binding site indeed involves the residue Asp(2.50) (reviewed in Katritch *et al.*¹¹). Both the 1.8 Å-resolution adenosine A_{2A} receptor structure and 2.1 Å-resolution β_1 adrenergic receptor (β_1 AR) structure show that a Na⁺ ion is coordinated by side chain oxygen atoms of Asp(2.50), Ser(3.39), and three water molecules in the middle of a water-filled channel within the transmembrane (TM) domain¹². The allosteric effects of Na⁺ on ligand binding in D2R have previously been studied by computational simulations¹³. Based on the structures of bovine rhodopsin and β_2 AR, Ericksen *et al.* modeled the Na⁺-induced conformations of D2R using normal mode

analysis to rationalize the enhanced binding of substituted benzamides and 1,4-disubstituted piperidines/piperazines (1,4-DAPs)^{13a}. Based on the structure of β_2 AR, Selent *et al.* modeled the allosteric effects of Na⁺ in the *apo* state of D2R using microsecond scale all-atom molecular dynamics (MD) simulations and showed that Na⁺ enters the receptor from the extracellular side, binds at Asp(2.50), and locks the rotamer toggle switch Trp(6.48) to the inactive state^{13b}. Recently, Filizola and colleagues carried out MD simulations in three subtypes of opioid receptors, and revealed important dynamic nature of Na⁺ binding¹⁴.

To better understand the structural basis of the effect of Na⁺ on ligand binding affinity in D2R and D3R, in combination with experimental binding assays, we carried out molecular modeling and simulation analysis of the receptors in complex with ligands whose binding are either sensitive or insensitive to Na⁺.

We first investigated the ability of physiological concentrations of Na⁺ to modulate the binding of these antagonist ligands (Fig. S1) to the D2R and D3R. The affinities of the two substituted benzamides, eticlopride and sulpiride, for the D2R are increased (~3-fold for eticlopride and ~23-fold for sulpiride) in the presence of Na⁺ (Fig. 1A,B). This effect of Na⁺ on sulpiride binding to the D2R is similar to that previously observed^{3,8}. In contrast, the presence or absence of Na⁺ does not affect the affinity of the butyrophenones, spiperone (Fig. 1C) or methylspiperone (data not shown), for the D2R. Interestingly, we found that the affinity of the tricyclic antipsychotic, zotepine, for the D2R is decreased by ~7-fold in the presence of Na⁺ (Fig. 1D). Similar results were observed with the D3R in that the binding of spiperone was not affected by Na⁺, whereas the binding of sulpiride and eticlopride were increased, and the binding of zotepine was decreased by Na⁺ (Fig. S2). These results illustrate how the Na⁺ bound state of the D2R and D3R can differentially affect the binding of different ligands to the receptors.

To computationally investigate atomistic details of the Na⁺ effect on the ligand binding modes, these ligands were docked to equilibrated D2R and D3R models based on the D3R crystal structure^{1,15}. The initial ligand poses were selected from top-scoring poses, taking into account of the implications from available mutagenesis experimental data (see SI). Briefly, for sulpiride, a pose similar to that of

eticlopride in the D3R crystal structure was selected, with the tertiary amine in the ethyl-pyrrolidine moiety forming a salt bridge to the Asp(3.32) side chain, and the methoxy-substituent of the benzamide moiety forming an intramolecular hydrogen bond with the amide-NH. For spiperone, a pose with the butyrophenone moiety in the orthosteric binding site (OBS) was selected, in which π - π interactions are formed with aromatic residues in TM6, in agreement with site-directed mutagenesis data, and structure-activity relationship data of spiperone and its analogs, N-methylspiperone and NAPS¹⁶. For zotepine, the pose in which the tricyclic moiety is bound in the OBS was selected.




Fig 1 Experimental binding affinity curves for the D2R with and without Na⁺ for eticlopride (A), sulpiride (B), spiperone (C), and zotepine (D). Radioligand binding assays with D2R containing membranes were performed as described in the Methods section. Membranes were incubated with 0.5 nM [³H]-methylspiperone and the indicated concentrations of competing ligand in the absence or presence of 144 nM Na⁺. The data are expressed as a percentage of the control [³H]-methylspiperone binding observed in the absence of competing ligand. The curves represent an average of three independent experiments. K_i values were calculated from the IC₅₀ values using the Cheng-Prusoff equation¹⁷ and are as follows: eticlopride: 0.67 ± 0.11 nM (-Na⁺), 0.27 ± 0.03 nM (+Na⁺); sulpiride: 379 ± 141 nM (-Na⁺), 16.7 ± 1.6 nM (+Na⁺); spiperone: 0.77 ± 0.12 nM (-Na⁺), 0.83 ± 0.26 nM (+Na⁺); zotepine: 5.8 ± 0.5 nM (-Na⁺), 38.4 ± 5 nM (+Na⁺).

We then performed extensive MD simulations of the resulting receptor-ligand complexes either in the presence or absence of Na⁺ bound in the Na⁺-binding site (Table S1). Throughout the simulations, similar to that observed in the crystal structures, the bound Na⁺ is stably coordinated by the side chain oxygen atoms of Asp(2.50), Ser(3.39), and three water molecules (Fig. S3). Other residues including Leu(2.46), Ala(2.49), Met(3.35), Asn(7.45), and Ser(7.46) are within 5 Å and contribute to forming the interaction

network that support the Na⁺ binding and propagate its impact (Fig. S3).

The salt bridge interactions between the protonated amines of sulpiride and eticlopride with Asp(3.32) are maintained in both Na⁺-bound and -unbound conditions in D2R and D3R. However, in the absence of Na⁺, the ethyl-pyrrolidine moieties shift down in the OBS (Fig. 2B,C), resulting in altered interactions between the sulfonamide N and the side chains of Ser(5.42) and Ser(5.43) in sulpiride, and the benzamide moiety tilting away from the vertical orientation in eticlopride. On the other hand, the binding mode of spiperone in D3R remains largely the same whether in the presence of absence of Na⁺ (Fig. 2D). In both conditions, the phenyl-substituted spiro moiety of spiperone makes more extensive interactions with TM2, TM3 and TM7 compared to sulpiride or eticlopride, which may be masking the effect of Na⁺ on the ligand binding mode. Interestingly, unlike for sulpiride, eticlopride, and spiperone, the binding mode for zotepine in D2R forms the salt bridge interaction with Asp114(3.32) only in the presence of Na⁺. In the absence of Na⁺, this interaction is lost, and the protonated amine and the ethoxy O atom form an optimal intramolecular interaction, which may contribute to higher binding affinity of zotepine in the Na⁺-unbound condition (Fig. 2E). To further evaluate the role of this intramolecular interaction of zotepine, we characterized the conformational energetics of zotepine and its analog in which the O6 atom is replaced by a C atom, and found that the intramolecular interaction would stabilize zotepine in its lowest-energy conformers, whereas the analog prefers an extended conformation (Fig. S4). Furthermore, MD simulations of D2R in complex with the zotepine-analog showed that the salt bridge interaction with Asp114(3.32) is maintained in the Na⁺-unbound condition (Fig. S5). Thus, the binding modes of Na⁺-sensitive ligands sulpiride, eticlopride, and zotepine are dependent on the presence of Na⁺, while that of Na⁺-insensitive ligand spiperone is unaffected by the absence Na⁺ (Fig. 2B-E).

To correlate the differential binding modes of Na⁺-(in)sensitive ligands with experimentally observed binding affinities, we calculated the MM/GBSA receptor-ligand binding energy for the frames of the equilibrated portions of the MD trajectories. In agreement with the experimental findings, for sulpiride and eticlopride, the binding energy values were lower (higher affinity) for the ligand poses in the Na⁺-bound condition than the -unbound condition (Fig. 3A,B); while for spiperone, the binding energy values with and without Na⁺ were comparable (Fig. 3C). For zotepine, the binding energy values are lower in the Na⁺-unbound condition, consistent with the experimentally observed slightly enhanced affinity in the absence of Na⁺ (Fig. 3D).

To characterize the allosteric impact of Na⁺ on the receptor conformations that might have contributed to the differential binding modes of Na⁺-sensitive ligands in the presence and absence of Na⁺, we analyzed and compared the changes in the interaction network




Fig. 2 Ligand binding mode of Na⁺-sensitive and -insensitive ligands. (A) The relative locations of the Na⁺-binding site (blue rectangle) and ligand binding site (violet rectangle) are shown in the D3R-eticlopride structure¹. The Na⁺ modeled into the structure is coordinated by the side chain oxygen atoms of Asp(2.50) and Ser(3.39), shown in blue sticks. The Na⁺ is shown as yellow sphere. The ligand binding modes of the Na⁺-sensitive ligands, eticlopride (B), sulpiride (C) and zotepine (E), differ in the Na⁺-bound (green) vs. Na⁺-unbound (gray) conditions, while that of Na⁺-insensitive ligand, spiperone (D), are similar in both conditions.

between the Na⁺-bound vs -unbound conditions. Our analysis for D3R and D2R in complex with sulpiride showed that the presence of Na⁺ strengthens interactions not only in the vicinity of the Na⁺-binding site amongst TMs 2, 3, 6, and 7 (see below), but the impact is also propagated to the extracellular ends of TMs 1, 2, and 7, and to the intracellular ends of TMs 3 and 5 (Fig. 4A). This change in interaction network correlates well with the principal mode of motion calculated by the principal component analysis of trajectories. In the Na⁺-unbound condition, the primary motion can be characterized by an outward movement of the extracellular segments of TMs 1, 2, 7, whereas in the Na⁺-bound condition, the motion is smaller and in the inward direction, consistent with the strengthened interactions in this region in the presence of Na⁺ (Fig. 4B). By contrast, similar analysis for the D3R-sulpirone complex did not show marked difference between the Na⁺-bound vs -unbound trajectories.




Fig. 3 Ligand binding energy and receptor conformation. The MM/GBSA ligand binding energy are plotted against the Asp(3.32)-Tyr(7.43) distances in the eticlopride (A), sulpiride (B), spiperone (C), and zotepine (D) complexes. The Na⁺-bound condition is in blue, and the -unbound condition is in gray. The binding energy and distance values are calculated for the frames at 6ns-interval in the equilibrated portions of the MD trajectories.

Specifically, in the Na⁺ binding site, the rotamer state of Ser(3.39) is maintained near $\chi_1 = -60$ degrees in the Na⁺-bound condition, similarly to that in the D3R structure¹, while in the Na⁺-unbound condition, the rotamer fluctuates among multiple rotamer states (Fig. S6). In the ligand binding site, a hydrogen bond interaction between the side chains of Asp(3.32) and Tyr(7.43) located above the Na⁺-binding site is easily broken in the absence of Na⁺ – the distance between the carboxylate O of Asp(3.32) side chain and the hydroxyl O of Tyr(7.43) side chain is ~ 3 Å in the Na⁺-bound trajectories, compared to >5 Å in the Na⁺-unbound trajectories in simulations with Na⁺-sensitive ligands sulpiride, eticlopride, and zotepine (Fig. S7A). The breaking of this hydrogen bond may contribute to the altered binding mode of these ligands in the Na⁺-unbound condition. Interestingly, our comparative analysis of the eticlopride and sulpiride simulations indicate eticlopride is more rigid in the binding site, which may facilitate Asp(3.32) to interact with Tyr(7.43) (Fig. S8). Thus, eticlopride would rely less on the bound-Na⁺ to stabilize the Asp(3.32)-Tyr(7.43) interaction, consistent with the less Na⁺ sensitivity of eticlopride than sulpiride (Fig. 1).

In the region adjacent to both the ligand and Na⁺-binding sites, the interactions among TMs 3 and 6, specifically between Cys(3.36) and Phe(6.44), are less frequently formed in the Na⁺-unbound conditions (Fig. S7B). Interestingly, Phe(6.44) is part of the so-called “P-I-F” motif which reconfigures between the active and inactive states of the β_2 AR¹⁸. The rearrangement observed between TM3 and TM6 from the Na⁺-bound to -unbound condition is in the same direction as that from the inactive to active state (Fig. S7). At the extracellular ends of TMs 1, 2, and 7, the TM1-TM2 and TM1-TM7 interfaces are each strengthened by two pairwise residue interactions in the Na⁺-bound condition (Fig. 4A).

Although the crystal structure of D3R was not resolved with a bound Na⁺, based on the differed rotamer states of Ser(3.39) in the presence and absence of Na⁺, our analysis strongly supports the presence of Na⁺ at Asp(2.50), consistent with a retrospective analysis of this medium resolution structure that showed electron densities compatible with Na⁺ in close proximity to Asp(2.50)¹¹. Our findings also indicate that the presence of Na⁺ has an allosteric impact on the ligand binding site, consequently altering the ligand binding mode and modulating the binding affinity of Na⁺-sensitive ligands. In particular, Na⁺ binding is associated with a critical interaction between the side chains of Asp(3.32) and Tyr(7.43) located in between the Na⁺-binding site and the ligand binding site. In the Na⁺-unbound condition, the less than optimal binding modes of Na⁺-sensitive ligands, eticlopride and sulpiride, are correlated with the weakening or breaking of this hydrogen bond interaction, whereas the binding modes of the Na⁺-insensitive ligand spiperone appear to mask the impact of the bound Na⁺.




Fig. 4 Allosteric impact of Na⁺ on TM domain. (A) The Na⁺-modulated interaction network commonly found in D3R-sulpiride and D2R-sulpiride complexes is shown in black dotted lines drawn between residue pairs with significantly different interaction frequencies in the Na⁺-bound condition compared to the -unbound condition. The residues shown are for D3R. The Na⁺-coordinating residues are in blue sticks; the residues that may involve in ligand binding¹⁹, are in violet sticks; the remaining residues in the network are in cyan sticks. Sulpiride is in green spheres. (B) The principal mode of motion calculated by the principle component analysis is shown in yellow arrows for the D3R-sulpiride complex in the Na⁺-bound (blue) and -unbound (gray) conditions.

Intriguingly, the critical role of Asp(3.32)-Tyr(7.43) interaction we identified in this study for the binding Na⁺-sensitive ligands is consistent with the results from a previous study²⁰, in which the mutation of Tyr(7.43) to Cys decreases the binding of sulpiride 185 fold, whereas it reduces the binding of Na⁺-insensitive methylspiperone only 3 fold.

In addition to the effect on ligand binding affinity, Na⁺-binding has also been implicated in modulating signaling efficacy. The comparison of the Na⁺-bound inactive-state structure to the active-state A_{2A} adenosine receptor structure showed the Na⁺/water pocket in the active state collapses in volume from ~200 to 70 Å³ due to movements of TM helices and would not allow coordination for Na⁺^{12a, 21}. The collapse of the Na⁺-binding site is thus suggested to correlate with the unbinding of Na⁺ and contribute to the negative allosteric effect of Na⁺ on agonist-binding²². Furthermore, Ala mutations of residues Asp(2.50), Asn(7.49), Ser(3.39), Trp(6.48), and Asn(7.45) were shown to either abrogate or reduce the agonist-stimulated receptor signaling²³. In addition, Na⁺ dissociation in the active state may shift the protonation state of Asp(2.50), and be related to the pH-dependence of activation in β₂AR²⁴. In this paper, however, we characterize an Na⁺-unbound inactive state that is yet to be revealed by crystallography – in contrast to all the crystal structures of aminergic receptors¹⁹, the Asp(3.32)-Tyr(7.43) interaction is dissociated in this state, which can be stabilized by a bulky Na⁺-insensitive ligand like spiperone.

Conclusions

The results of our comparative analysis of the MD simulations in the presence or absence of bound Na⁺ reveal an allosteric interaction network from the Na⁺-binding site to both the extracellular and intracellular sides of the TM domain and underlies the Na⁺-sensitivity of ligands. Thus, we establish the allosteric mechanism by which ligand binding affinity is modulated in the presence of bound Na⁺ at Asp(2.50). The distinct conformations of the receptor bound to Na⁺-sensitive vs Na⁺-insensitive antagonists studied herein represent sub-states of the inactive state. These conformations will aid in providing a pharmacologically relevant ensemble of receptor conformations in the structure-based virtual high-throughput screening for novel ligand discovery²⁵.

Notes and references

^a Department of Physiology and Biophysics, Weill Medical College of Cornell University, New York, NY

^b Computational Chemistry and Molecular Biophysics Unit, National Institute on Drug Abuse – Intramural Research Program, National Institutes of Health, Baltimore, MD

^c Molecular Neuropharmacology Section, National Institute of Neurologic Disorders and Stroke, National Institutes of Health, Bethesda, MD

^d Institute for Computational Biomedicine, Weill Medical College of Cornell University, New York, NY.

† The number in parenthesis indicates the Ballesteros and Weinstein index of the residue²⁶.

Electronic Supplementary Information (ESI) available: experimental and computational details, supplemental table and figures. See DOI: 10.1039/c000000x/

- 1 E. Y. Chien, W. Liu, Q. Zhao, V. Katritch, G. W. Han, M. A. Hanson, L. Shi, A. H. Newman, J. A. Javitch, V. Cherezov and R. C. Stevens, *Science*, 2010, **330**, 1091.
- 2 L. E. Limbird, J. L. Speck and S. K. Smith, *Molecular pharmacology*, 1982, **21**, 609.
- 3 K. A. Neve, *Molecular pharmacology*, 1991, **39**, 570.

- 4 Z. G. Gao, S. K. Kim, A. S. Gross, A. Chen, J. B. Blaustein and K. A. Jacobson, *Molecular pharmacology*, 2003, **63**, 1021.
- 5 C. B. Pert, G. Pasternak and S. H. Snyder, *Science*, 1973, **182**, 1359.
- 6 S. Martin, J. M. Botto, J. P. Vincent and J. Mazella, *Molecular pharmacology*, 1999, **55**, 210.
- 7 R. Fredriksson, M. C. Lagerstrom, L. G. Lundin and H. B. Schiöth, *Molecular pharmacology*, 2003, **63**, 1256.
- 8 K. A. Neve, B. A. Cox, R. A. Henningsen, A. Spanoyannis and R. L. Neve, *Molecular pharmacology*, 1991, **39**, 733.
- 9(a) K. A. Neve, R. A. Henningsen, J. M. Kinzie, T. De Paulis, D. E. Schmidt, R. M. Kessler and A. Janowsky, *The Journal of pharmacology and experimental therapeutics*, 1990, **252**, 1108; (b) J. A. Schetz and D. R. Sibley, *The Journal of pharmacology and experimental therapeutics*, 2001, **296**, 359.
- 10 K. A. Neve, M. G. Cumbay, K. R. Thompson, R. Yang, D. C. Buck, V. J. Watts, C. J. DuRand and M. M. Teeter, *Molecular pharmacology*, 2001, **60**, 373.
- 11 V. Katritch, G. Fenalti, E. E. Abola, B. L. Roth, V. Cherezov and R. C. Stevens, *Trends in biochemical sciences*, 2014, **39**, 233.
- 12(a) W. Liu, E. Chun, A. A. Thompson, P. Chubukov, F. Xu, V. Katritch, G. W. Han, C. B. Roth, L. H. Heitman, A. P. Ijzerman, V. Cherezov and R. C. Stevens, *Science*, 2012, **337**, 232; (b) J. L. Miller-Gallacher, R. Nehme, T. Warne, P. C. Edwards, G. F. Schertler, A. G. Leslie and C. G. Tate, *PLoS One*, 2014, **9**, e92727.
- 13(a) S. S. Ericksen, D. F. Cummings, H. Weinstein and J. A. Schetz, *The Journal of pharmacology and experimental therapeutics*, 2009, **328**, 40; (b) J. Selent, F. Sanz, M. Pastor and G. De Fabritiis, *PLoS computational biology*, 2010, **6**, e1000884.
- 14 Y. Shang, V. LeRouzic, S. Schneider, P. Bisignano, G. W. Pasternak and M. Filizola, *Biochemistry*, 2014, **53**, 5140.
- 15 M. Michino, P. Donthamsetti, T. Beuming, A. Banala, L. Duan, T. Roux, Y. Han, E. Trinquet, A. H. Newman, J. A. Javitch and L. Shi, *Molecular pharmacology*, 2013, **84**, 854.
- 16(a) D. O. Kiesewetter, W. C. Eckelman, R. M. Cohen, R. D. Finn and S. M. Larson, *International journal of radiation applications and instrumentation. Part A, Applied radiation and isotopes*, 1986, **37**, 1181; (b) F. Boeckler, H. Lanig and P. Gmeiner, *J Med Chem*, 2005, **48**, 694; (c) L. Albizu, M. Cottet, M. Kralikova, S. Stoev, R. Seyer, I. Brabet, T. Roux, H. Bazin, E. Bourrier, L. Lamarque, C. Breton, M. L. Rives, A. Newman, J. Javitch, E. Trinquet, M. Manning, J. P. Pin, B. Mouillac and T. Durroux, *Nature chemical biology*, 2010, **6**, 587.
- 17 Y. Cheng and W. H. Prusoff, *Biochemical pharmacology*, 1973, **22**, 3099.
- 18 S. G. Rasmussen, H. J. Choi, J. J. Fung, E. Pardon, P. Casarosa, P. S. Chae, B. T. Devree, D. M. Rosenbaum, F. S. Thian, T. S. Kobilka, A. Schnapp, I. Konetzki, R. K. Sunahara, S. H. Gellman, A. Pautsch, J. Steyaert, W. I. Weis and B. K. Kobilka, *Nature*, 2011, **469**, 175.
- 19 M. Michino, T. Beuming, P. Donthamsetti, A. H. Newman, J. A. Javitch and L. Shi, *Pharmacol Rev*, 2015, **67**, 198.
- 20 D. Fu, J. A. Ballesteros, H. Weinstein, J. Chen and J. A. Javitch, *Biochemistry*, 1996, **35**, 11278.
- 21 V. Katritch, V. Cherezov and R. C. Stevens, *Annual review of pharmacology and toxicology*, 2013, **53**, 531.
- 22 H. Gutierrez-de-Teran, A. Massink, D. Rodriguez, W. Liu, G. W. Han, J. S. Joseph, I. Katritch, L. H. Heitman, L. Xia, A. P. Ijzerman, V. Cherezov, V. Katritch and R. C. Stevens, *Structure*, 2013, **21**, 2175.
- 23 A. Massink, H. Gutierrez-de-Teran, E. B. Lenselink, N. V. Ortiz Zacarias, L. Xia, L. H. Heitman, V. Katritch, R. C. Stevens and A. P. Ijzerman, *Molecular pharmacology*, 2015, **87**, 305.
- 24(a) A. Ranganathan, R. O. Dror and J. Carlsson, *Biochemistry*, 2014, **53**, 7283; (b) P. Ghanouni, H. Schambye, R. Seifert, T. W. Lee, S. G. Rasmussen, U. Gether and B. K. Kobilka, *J Biol Chem*, 2000, **275**, 3121.
- 25 D. J. Osguthorpe, W. Sherman and A. T. Hagler, *The journal of physical chemistry. B*, 2012, **116**, 6952.
- 26 J. A. Ballesteros and H. Weinstein, *Methods in Neurosciences*, 1995, **Volume 25**, 366.