

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Organic & Biomolecular Chemistry

COMMUNICATION

Peptide 2-formylthiophenol esters do not proceed through Ser/Thr ligation pathway, but participate in a peptide aminolysis to enable peptide condensation and cyclization

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Chun Ling Tung^{† a,b}, Clarence T. T. Wong^{† a,b}, and Xuechen Li^{* a,b}

Peptide thiol salicylaldehyde (SAL) ester unexpectedly do not follow a Ser/Thr ligation pathway to react with peptides containing N-terminal Ser/Thr, but proceed towards a peptide aminolysis in DMSO. The reaction takes place even at a low substrate concentration (1mM). The method has been successfully used to synthesize several natural cyclic peptides, with high ratio of monocyclic to dimeric products.

Chemoselective peptide ligation enables two side chain unprotected peptide segments to be joined together at the termini with the generation of the natural peptidic linkage at the ligation site, which has dramatically advanced the protein chemical synthesis.¹ Alternatively, some partially chemoselective aminolysis-based methods, which require the internal lysine side chain to be protected, have also been used in the convergent synthesis of peptides and proteins.² These aminolysis-mediated peptide condensation methods are often limited to condense the peptides at the C-terminal glycine and proline sites, due to the proneness to epimerization with other amino acids. Recently, our laboratory has developed a chemoselective Ser/Thr ligation (STL),³ in which one side chain unprotected peptide segment with a C-terminal salicylaldehyde (SAL) ester reacts with another side chain unprotected peptide segment with N-terminal serine or threonine to form an *N,O*-benzylidene acetal intermediate, followed by acidolysis to reveal the natural Xaa-Ser/Thr linkage (Figure 1). This method has been successfully applied in protein synthesis, protein semi-synthesis and cyclic peptide synthesis.^{3,4,5}

A logical inquiry along the line of STL chemistry is to use peptide thiol-salicylaldehyde (2-formylthiophenol) esters under the Ser/Thr ligation condition and is the subject of this communication. From the point of physical organic chemistry, the thiolester is a better leaving group than the O-ester counterpart, thus we expect that a 1, 5 *S* to *N* acyl transfer would be more feasible than a 1, 5 *O* to *N* acyl transfer (Figure 1). To the end, we first tested this idea by using Fmoc-Ala thiol-SAL ester to react with a serine derivative. These two components were dissolved in pyridine acetate buffer (AcOH:Pyr 1:1, mol: mol), and indeed the expected *N,O*-benzylidene acetal product was obtained, which upon acidolysis gave rise to Fmoc-Ala-Ser-Phe-OMe tripeptide, as analyzed by LC-MS (Figure 2, a). However, when a heptapeptide (Ac-SPKMQG thiol-SAL ester)

was reacted with a hexapeptide (NH₂-SFAVGA-CO₂H) with N-terminal Ser under the same condition as above, we were surprised to see that no ligated product was observed and the starting material remained intact as analyzed by LC-MS (Figure 2, b). We have varied the peptide sequences for the ligation study, none of which gave the ligated product. The underlying reason is unclear to us yet.




Figure 1. Peptide (thiol) salicylaldehyde ester-mediated reactions.

In the 1970s, Kemp has reported that aminolysis of 8-acetoxy-1-naphthaldehyde or 2-acetoxybenzaldehyde with primary amines generated amides. Under the condition, an initial hemiaminal formation took place, followed by intramolecular 1,5 *O* to *N* acyl transfer to generate the amide.⁶ Following this strategy, Ito and coworker utilized 2-formyl-4-nitrothiophenol esters to facilitate the amide formation by hemiaminal-mediated acyl transfer.⁷ These results were very promising, however, all these studies were limited to simple substrates (Scheme 1). These precedent examples prompted us to explore the utility of peptide thiol SAL esters for

^aDepartment of Chemistry, State Key Laboratory of Synthetic Chemistry, The University of Hong Kong.

^bChong Yuet Ming Chemistry Building, Pokfulam Road, Hong Kong.

[†] Authors share equal contributions

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

Communication

peptide condensation via hemiaminal formation-mediated aminolysis.

Initially, when peptide **7** (Ac-VGDTYA thiol-SAL ester) and peptide **8** (NH₂-VEGFA-CO₂H) were dissolved in DMSO (10 mM) with 2 equiv. of DIEA, a peptide condensation was indeed observed to give **9** within 6 h (Figure, 2c).⁶ As a comparison, Agrigento et al have used peptides with C-terminal *p*-chloro or *p*-nitro thiophenyl ester for the peptide aminolysis (DIEA, DMSO), in which the reaction needed 20-96 h at the substrate concentration of 60 mM.^{2f} Thus, the hemiaminal formation-mediated reaction help expedite the aminolysis. Next, we evaluated the epimerization issue under this condition. Peptide **10** (Ac-FVGFSDTYGA thiol-SAL ester) and peptide **11** (Ac-FVGFSDTYGa thiol-SAL ester) were synthesized using the epimerization-free peptide hydrazine displacement approach.⁸ These two peptide thiol SAL esters were then reacted with peptide **12** (H₂N-AVYAAPYLAGG-CO₂H), respectively. The ligation could go to completion within few hours. However, we observed the significant epimerized product, which was proved by comparing the retention time of C-terminal L- and D-alanine peptides (Figure 3). Thus, like other aminolysis-mediated peptide condensation^{2a,2c}, this method will be limited to condense the peptide segment at the site with the C-terminal glycine or proline residue. In addition, this method will not tolerate the unprotected lysine and cysteine residues in the peptide sequence.





Figure 2. Thiol-SAL ester mediated reactions.



Scheme 1. Previous examples using a hemiaminal-mediated acyl transfer strategy.

Organic & Biomolecular Chemistry




Figure 3. RP-HPLC profile of epimerization determination between the ligation reaction of Ac-FVGFSDTYGA thiol-SAL ester (**10**)/ Ac-FVGFSDTYGa thiol-SAL ester (**11**) with NH₂-AVYAAPYLAGG-CO₂H (**12**). A. Co-injection of ligated products. B. Ligation products from using C-terminal L-Ala thiol-SAL ester and C. Ligation products from using C-terminal D-Ala thiol-SAL ester.

To further probe the optimal conditions, Ac-AFQIG thiol-SAL ester (**13**) was allowed to react with NH₂-GLVYA-CO₂H (**14**) under different conditions at a substrate concentration of 1 mM. Under neutral or basic aqueous solution, the hydrolysis product predominated (Entry 2-3, Table 1).⁹ Organic solvents (DMSO, DMF and 1-methyl-2-pyrrolidone) were more suitable. Increase of the amount of DIEA to 10 equiv. caused side reactions. In addition to DIEA, other bases including DBU and DEA were found to be unsuitable. Therefore, the optimal condition is to use 1-3 equiv. of DIEA in DMSO. Under the same condition, a peptide thiol SAL ester (**15**) reacted much faster than its peptide thiophenyl ester counterpart (**17**) (Figure 4).

Table 1. Optimization of the reaction conditions.

Entry	Solvent	Base	Conversion
1	PBS (pH 6)	/	<5%
2	PBS (pH 7.4)	/	<5%
3	PBS (pH 8.5)	/	<5%
4	DMSO	10 eq. DIEA	69%
5	DMSO	10 eq. Imid.	67%
6	DMSO	3 eq. DIEA	91%
7	DMSO-NMM	3 eq. DIEA	90%
8	DMSO	1 eq. DIEA	90%
9	DMSO	3 eq. DBU	<5%
10	DMSO	3 eq. DEA	<5%
11	DMF	1 eq. DIEA	82%
12	DMF	3 eq. DIEA	80%
13	DMF	10 eq. DIEA	38%
14	1-methyl-2-pyrrolidone	3 eq. DIEA	77%
15	50% DMSO/ H ₂ O	3 eq. DIEA	22%




Figure 4. Comparison of the aminolysis rate between a peptide thiol SAL ester (**15**) and peptide thiophenyl ester (**17**). Reaction **A** was performed for 5 h at 10 mM while **B** for 20 h at 10 mM.

As the C-terminal residue was limited to the glycine or proline residue, we next explored the scope and limitations on the N-terminal residues. The synthesis of peptides containing C-terminal Gly/Pro thiol SAL esters was straightforward using the direct coupling condition. We elected to use the peptide thiol-SAL ester of Ac-ITGEFNAX (X = Gly or Pro) (**18a-b**) as the N-terminal peptide and the peptide NH₂-ZGVFA-CO₂H (Z = amino acid) (**19a-j**) as the C-terminal peptide. The two peptide segments were dissolved at 1 mM concentration in DMSO with 3 equiv. DIEA. The reaction progress was analyzed by LC-MS at 4 h and 8 h. As seen in Table 2, the reactions with the peptide C-terminal Gly thiol-SAL ester could be completed or nearly completed within 4 h, while the peptide C-terminal Pro thiol-SAL ester required 8 h. We also observed a small amount of the imine product for the peptide C-terminal proline thiol-SAL ester, which was consistent with Kemp's observation that, certain substrates stopped at the imine stage without inducing acyl transfer.

Table 2. Examples of thiol SAL-mediated aminolysis of C-terminal Gly and Pro thiol-SAL ester with different N-terminal amino acids.

Entry	X	Z	Product	Conversion	
				4h	8h
1 ^a	Gly	Asp	12a	77	78
2 ^a	Gly	Glu	12b	80	89
3 ^a	Gly	Pro	12c	75	77
4 ^a	Gly	Ser	12d	82	82
5 ^a	Gly	Gly	12e	85	85
6 ^b	Pro	Asp	12f	65	83
7 ^b	Pro	Glu	12g	60	90
8 ^b	Pro	Pro	12h	61	71

9 ^b	Pro	Ala	12i	79	85
10 ^b	Pro	Gly	12j	90	98

^aReaction conditions: peptide thiol-SAL ester (**18a-b**) (1 equiv.), C-terminal peptides (**19a-j**) (1.2-1.3 equiv.), DIEA (3.0 equiv.), DMSO at a final conc. of 1 mM, r.t.; Conversion was calculated by the ratio of the product area over the product and the hydrolyzed peptide.

^bReaction condition: peptide thiol-SAL ester (**18a-b**) (1.5 equiv.), C-terminal peptides (**19a-j**) (1 equiv.), DIEA (3.0 equiv.), DMSO at a final conc. of 1 mM, r.t.; Conversion was calculated by the ratio of the product area over the product and the remaining C-terminal peptide.

To demonstrate the effectiveness of this method, we next applied it for the synthesis of the natural cyclic peptides. Many natural cyclic peptides exhibit broad biological activities, including anti-cancer, anti-bacteria, and anti-virus; thus cyclic peptides serve as a useful scaffold for the development of new therapeutic agents.¹⁰ As such, the development of different methods for the synthesis of natural and unnatural cyclic peptides is attracting much attention.¹¹

Table 3. Synthesis of natural cyclic peptides via peptide thiol-SAL ester-mediated cyclization

Entry	Name	Sequence	Ring size	M:D	Yield ^a
1	Phakellistain -13	Cyclo-[FGPTLWP]	7	8:1	41%
2	Cyclosquamosin D	Cyclo-[GVVSYYPG]	8	99:1	49%
3	Dichotomin G	Cyclo-[LPSTFPPIPI]	9	9:1	50%
4	Antamanide	Cyclo-[AFFPPFFVPP]	10	99:1	48%
5	Stalladelin D	Cyclo-[VPSYFPAAIG]	11	99:1	45%

^aIsolated by HPLC. M: monocyclic product. D: dimerized product.

We selected several natural cyclic peptides with ring sizes made of 7 to 11 amino acid residues, containing Gly or Pro in the sequence. These peptides include phakellistain-13¹², cyclosquamosin D¹³, Dichotomin G¹⁴, Antamanide¹⁵ and Stalladelin D¹⁶. The cyclization was performed at Pro-Phe, Gly-Gly, Pro-Leu, Pro-Leu, Pro-Ala and Gly-Gly sites, respectively, via the thiol-SAL ester-mediated peptide cyclization. The cyclizations were performed at 1 mM in DMSO, which proceeded smoothly with completion in 4-8 h. The monocyclic products were obtained as major or sole products (Table 3, Figure 5). Unfortunately, highly constrained tetrapeptides could not be cyclized using this method, in which only the dimerized cyclic products were observed.




Figure 5. RP-HPLC profiles of the crude reaction mixture of the thiol SAL-mediated cyclization. * = partial removal of Trp(Boc) protecting group.

Conclusions

In conclusion, the peptide thiol-SAL ester surprisingly could not participate in Ser/Thr ligation, but could undergo a direct aminolysis, likely through a hemiaminal-mediated acyl transfer.¹⁷ This thiol-SAL ester mediated aminolysis method can serve as an alternative approach for the convergent synthesis of peptides and cyclic peptides. We have demonstrated the effectiveness of this method in the synthesis of several natural cyclic peptides. The application of this method towards protein chemical synthesis is ongoing.

Acknowledgements

This work was supported by the National Basic Research Program of China (973 program, 2013CB836900), the General Research Fund of the Research Grants Council of Hong Kong (HKU702813P), AoE/M-12/06 and Peacock Program-Project Development Fund (KQC201109050074A).

Notes and references

- (a) V. R. Pattabiraman, J. W. Bode, *Nature* 2011, **480**, 471-479; (b) C. P. R. Hackenberger, D. Schwarzer, *Angew. Chem. Int. Ed.* 2008, **47**, 10030-10047; (c) L. Raibaut, N. Ollivier, O. Melnyk, *Chem. Soc. Rev.* 2012, **41**, 7001-7015; (d) C. T. T. Wong, C. L. Tung, X. Li, *Mol. Biosyst.* 2013, **9**, 826-833; (e) P. E. Dawson, T. W. Muir, I. Clark-Lewis, S. B. H. Kent, *Science* 1994, **266**, 776-779.
- (a) S. Aimoto, *Curr. Org. Chem.* 2001, **5**, 45-87; (b) R. J. Payne, S. Ficht, W. A. Greenberg, C. H. Wong, *Angew. Chem. Int. Ed.* 2008, **47**, 4411-4415; (c) G. Chen, Q. Wan, Z. Tan, C. Kan, Z. Hua, K. Ranganathan, S. J. Danishefsky, *Angew. Chem. Int. Ed.* 2007, **46**, 7383-7387; (d) D. Crich, I. Sharma, *Angew. Chem. Int. Ed.* 2009, **48**, 2355-2358; (e) B. L. Wilkinson, C. K. Chun, R. J. Payne, *Biopolymers* 2011, **96**, 137-146; (f) P. Agrigento, F. Albericio, S. Chamoine, I. Dacquignis, H. Koc, M. Eberle, *Org. Lett.* 2014, **16**, 3922-3925; (g) S. Popovic, H. Bieräugel, R. J. Detz, A. M. Kluwer, J. A. A. Koole, D. E. Streefkerk, H. Hiemstra, J. H. van Maarseveen, *Chem. Eur. J.* 2013, **19**, 16934-16937; (h) H. Hojo, Y. Murasawa, H. Katayama, T. Ohira, Y. Nakahara, Y. Nakahara, *Org. Biomol. Chem.* 2008, **6**, 1808-1813.
- (a) X. Li, H. Y. Lam, Y. Zhang, C. K. Chan, *Org. Lett.* 2010, **12**, 1724-1727; (b) Y. Zhang, C. Xu, H. Y. Lam, C. L. Lee, X. Li, *Proc. Natl. Acad. Sci. U.S.A.*, 2013, **110**, 6657-6662; (c) C. T. Wong, T. Li, H. Y. Lam, Y. Zhang, X. Li, *Front. Chem. Biol.* 2014, **2**, 28.
- (a) C. L. Lee, X. Li, *Curr. Opin. Chem. Biol.* 2014, **22**, 108-114; (b) H. Y. Lam, Y. Zhang, H. Liu, J. Xu, C. T. Wong, C. Xu, X. Li, *J. Am. Chem. Soc.* 2013, **135**, 6272-6279; (c) C. T. T. Wong, H. Y. Lam, T. Song, G. Chen, X. Li, *Angew. Chem. Int. Ed.* 2013, **52**, 10212-10215; (d) C. Xu, H. Y. Lam, Y. Zhang, X. Li, *Org. Commun.* 2013, **49**, 6200-6202; (e) Y. Zhang, T. Li, X. Li, *Org. Biomol. Chem.* 2013, **11**, 5584-5587.
- (a) P. M. Levine, T. W. Craven, R. Bonneau, K. Kirshenbaum, *Org. Lett.* 2014, **16**, 512-515; (b) P. M. Levine, T. W. Craven, R. Bonneau, K. Kirshenbaum, *Chem. Commun.* 2014, **50**, 6909-6912; (c) Levine, P. M.; Craven, T. W.; Bonneau, R.; Kirshenbaum, K. *Org. Biomol. Chem.* 2013, **11**, 4142-4146.
- D. S. Kemp, F. Vellaccio Jr. *J. Org. Chem.* 1975, **40**, 3003-3004.
- A. Ishiwata, T. Ichiyangaji, M. Takatani, Y. Ito, *Tetrahedron Lett.* 2003, **44**, 3187-3190. In this study, the substrate was limited to the Fmoc-Val-SAr.
- G. M. Fang, Y. M. Li, F. Shan, Y. C. Huang, J. B. Li, Y. Lin, H. K. Cui, L. Liu, *Angew. Chem. Int. Ed.* 2011, **50**, 7645-7649.
- Different from the result of reference 6, in which aqueous buffers could be used.
- (a) A. T. Bockus, C. M. McEwen, R. S. Lokey, *Curr. Top. Med. Chem.* 2013, **13**, 821-836; (b) S. H. Joo, *Biomol. Ther.* 2012, **20**, 19-26.
- (a) C. J. White, A. K. Yudin, *Nat. Chem.* 2011, **3**, 509-524; (b) J. P. Tam, C. T. Wong, *J. Biol. Chem.* 2012, **287**, 27020-27025.
- W. L. Li, Y. H. Yi, H. M. Wu, Q. Z. Xu, H. F. Tang, D. Z. Zhou, H. W. Lin, Z. H. Wang, *J. Nat. Prod.* 2003, **66**, 146-148.
- A. Dellai, I. Maricic, V. Kumar, S. Arutyunyan, A. Bouraoui, A. Nefzi, *Bioorg. Med. Chem. Lett.* 2010, **20**, 6020-6023.
- H. Morita, A. Shishido, T. Kayashita, K. Takeya, H. C. Itokawa, *J. Nat. Prod.* 1997, **60**, 404-407.
- I. L. Karle, T. Wieland, D. Schermer, H. C. Ottenheym, H. C. J. *Proc. Natl. Acad. Sci. USA.* 1979, **76**, 1532-1536.
- Y. R. Zhou, J. Zhou, X. K. Wang, X. L. Huang, H. M. Wu, X. Cheng, *Chin. Chem. Lett.* 1996, **7**, 237-240.
- We do not have a direct evidence for the formation of a hemiaminal.