


Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Rawal's catalyst as an effective stimulant for the highly asymmetric Michael addition of β -keto esters to functionally rich nitro-olefins†

A. Suresh Kumar, T. Prabhakar Reddy, R. Madhavachary, and Dhevalapally B. Ramachary*

Receipt/Acceptance Data [DO NOT ALTER/DELETE THIS TEXT]

Publication data [DO NOT ALTER/DELETE THIS TEXT]

DOI: 10.1039/b000000x [DO NOT ALTER/DELETE THIS TEXT]

A general approach to asymmetric synthesis of highly substituted dihydroquinolines was achieved through neighboring *ortho*-amino group engaged sequential Michael/amination/dehydration reactions on (*E*)-2-(2-nitrovinyl)anilines with cyclic and acyclic β -keto esters in the presence of a catalytic amount of Rawal's quinidine-NH-benzyl squaramide followed by TFA.

Recently organocatalytic domino/cascade reactions have emerged as powerful strategy for the complex molecular synthesis by employing simple and readily available precursors in a simple operational manner.¹ In this scenario, one of the important protocol for the C–C bond formation is asymmetric Michael reaction, which has been extensively used in cascade reactions.² Wherein, the Michael reaction of β -keto esters to nitro-olefins is a fully atom-economic transformation with prominent synthetic potential as the resulting functional groups will be in high demand.³ To control the reactivity and selectivity of this Michael reaction, in 2003 Takemoto and co-workers identified the bifunctional thiourea-tertiary amine as an efficient catalyst.⁴ A milestone report by Rawal's group showed that replacement of the thiourea function by a squaramide catalyst as the hydrogen-bond donor allows a dramatic decrease in the catalyst loading and also increase in the reaction rate/selectivity.^{5,6} The pioneering work of Rawal's and other groups from the past few years has shown the importance of chiral hydrogen-bonding catalysis through squaramide catalyst.^{5,6}

Nowadays, bio-mimetic one-pot synthesis of the highly functionalized chiral Michael adducts for synthetic applications are in high demand. Recently, we and others have reported the natural products inspired dihydrocoumarins synthesis through a Michael addition of 1,3-dicarbonyls to 2-(2-nitrovinyl)phenol and subsequent intramolecular hemiacetalization.⁷ In this reaction, the *ortho*-hydroxyl group participates as neighboring group both during and after the reaction and makes this sequence as a powerful tool among the existing annulation methods. With this inspiration, herein we describe an unusual efficient neighboring *ortho*-amino group engaged asymmetric organocatalytic sequential one-pot reaction for the synthesis of drug-like chiral dihydroquinolines from simple substrates and catalysts by using sequential Michael/amination/dehydration reactions.

Dihydroquinolines are versatile synthetic intermediates in organic synthesis and also prevalent unit in biological and pharmaceutical substances (Figure 1).⁸ Even though numerous synthetic methods have been known for their synthesis, there is an urgent requirement

to develop an efficient asymmetric protocol due to their many applications.⁹




Figure 1 Biologically active molecules containing 1,4-dihydroquinoline.

Recently, we and Kim group independently reported the asymmetric synthesis of 2,3,4-trisubstituted 1,4-dihydroquinolines by the cascade Michael/amination/dehydration reaction of aldehydes or ketones with (*E*)-2-(2-nitrovinyl)anilines through enamine-catalysis with excellent selectivity.¹⁰ However, there is no suitable asymmetric method to synthesise highly functionalized 1,4-dihydroquinolines in optically pure form from (*E*)-2-(2-nitrovinyl)anilines with β -keto esters through enolate-catalysis.¹¹ In continuation of our research interest in this area, we envisaged functionally rich (*E*)-2-(2-nitrovinyl)anilines and β -keto esters as the potential substrates for the newly designed Michael/amination/dehydration reaction sequence (Scheme 1).

To find the best catalytic conditions, we screened a number of emerging hydrogen-bond-donating organocatalysts for the reaction of *N*-Boc-(*E*)-2-(2-nitrovinyl)aniline **1a** with 1.3 equiv. of ethyl 2-oxocyclopentanecarboxylate **2a** (Table 1). Reaction of **1a** with 1.3 equiv. of **2a** under 10 mol% of quinidine-NH-thiourea **3a**-catalysis in toluene at 25 °C for 4 h furnished the **4aa** in 70% yield with 58% *ee* and >99% *de* (Table 1, entry 1). The same reaction in DCM for 4 h furnished the product **4aa** with *ee* increased to 75% with 68% yield and >99% *de* (Table 1, entry 2). Reaction under the hydroquinidine-NH-thiourea **3b**-catalysis furnished **4aa** with increased yield (82%) and no change in *ee* and *de* (Table 1, entry 3). With these moderate results, we moved to investigate this reaction with other set of hydrogen-bond-donating catalysts **3c-e** based on the chiral squaramide derivatives discovered by Rawal's group. Recently Rawal's squaramide catalysts have been found to be more powerful hydrogen-bond-donating catalysts compared to

School of Chemistry, University of Hyderabad, Hyderabad-500 046, India. E-mail: ramsc@uohyd.ernet.in; Fax: +91-40-23012460


† Electronic supplementary information (ESI) available: Experimental procedures and analytical data (¹H NMR, ¹³C NMR and HRMS) for all new compounds. See DOI: 10.1039/xxxxxxx

Communication


www.rsc.org/[JOURNAL] | [JOURNAL NAME HERE]

their corresponding thiourea analogues in both catalytic activity and selectivity.^{5,6} Therefore, herein we shown interest to investigate Rawal's catalysts **3c-e** for the high asymmetric induction.


(*E*)-2-(2-Nitrovinyl)phenol in Michael reactions through enolate-catalysis: Ramachary, Enders and Hong groups



(*E*)-2-(2-Nitrovinyl)aniline in Michael reactions through enamine-catalysis: Kim and Ramachary groups



(*E*)-2-(2-Nitrovinyl)aniline in Michael reactions through enolate-catalysis: This work



Scheme 1 Reaction design for the 1,4-dihydroquinoline synthesis.

Unfortunately, reaction of **1a** with **2a** under 5 mol% of quinine-NH-squaramide **3c** in DCM at 25 °C for 6 h gave the product **4aa** in 71% yield with poor *ee* (18%) and good *de* (Table 1, entry 4). After this poor selectivity, we shown interest to investigate the electronic and steric factors of *N*-aryl group in the squaramide catalyst, we used quinine-NH-benzyl squaramide catalyst **3d** for the high selectivity. Surprisingly, the Michael reaction of **1a** with **2a** under 5 mol% of **3d**-catalysis furnished the product (-)-**4aa** in 71% yield with 97% *ee* and >99% *de* within 5 h (Table 1, entry 5). This result is indicating that the presence of benzyl group in the squaramide catalyst has significant effect on the outcome of selectivity. In a similar manner, the Michael reaction of **1a** with **2a** under the 5 mol% of quinidine-NH-benzyl squaramide **3e**-catalysis furnished the opposite enantiomer (+)-**4aa** in 77% yield with 98% *ee* and >99% *de* within 5 h (Table 1, entry 6). Whereas 10 mol% of **3e**-catalysis furnished the product (+)-**4aa** in 77% yield with decreased *ee* of 83% (entry 7), which indicates that the catalyst loading has a significant effect on the selectivity of the reaction. Use of brine instead of DCM as solvent with 2 mol% of catalyst **3e** the aforesaid product was obtained in 48% yield with 91% *ee* and >99% *de* in 16 h (Table 1, entry 8). To obtain the highest selectivity of the product **4aa**, we thought of using sterically hindered isopropyl 2-oxocyclopentanecarboxylate **2b** with **1a** under **3d/3e**-catalysis. Surprisingly, the reaction of **1a** with **2b** under 5 mol% of **3e**-catalysis in DCM at RT for 6 h furnished the product (+)-**4ab** in 80% yield with >99% *ee* and *de* (Table 1, entry 10). Similar reaction under **3d**-catalysis furnished the opposite enantiomer (-)-**4ab** in 75% yield with >99% *ee* and *de* (Table 1, entry 11). To further investigate the reaction conditions, we carried out the Michael reactions of other *N*-protected-(*E*)-2-(2-nitrovinyl)anilines **1b/1c** with **2b** under **3d**- or **3e**-catalysis in DCM at RT. It has shown that other carbamate protecting groups (Cbz and CO₂Et) were tolerated and the desired products were obtained in moderate yields (42–65%) with excellent *ee*'s (96 to >99%) and *de*'s (97 to >99%) within 8 h (Table 1, entries 12–14). Finally we envisioned the optimized condition to be 25 °C in DCM under 5 mol% of Rawal's catalysts **3d** or **3e** to furnish the Michael adduct **4ab** in 75–80% yield with >99% *ee* and >99% *de* (Table 1, entry 10–11).


Table 1 Reaction optimization.^a

Entry	Pg	R	Catalyst 3 (5 mol%)	Solvent (0.3 M)	Time (h)	Yield (%) ^b	<i>ee</i> (%) ^c	<i>de</i> (%) ^c
1 ^d	1a: Boc	2a: Et	3a	Toluene	4	4aa (70)	58	>99
2 ^d	1a: Boc	2a: Et	3a	DCM	4	4aa (68)	75	>99
3 ^d	1a: Boc	2a: Et	3b	DCM	4	4aa (82)	74	>99
4 ^d	1a: Boc	2a: Et	3c	DCM	6	4aa (71)	18	>99
5	1a: Boc	2a: Et	3d	DCM	5	4aa (71)	97	>99
6	1a: Boc	2a: Et	3e	DCM	5	4aa (77)	98	>99
7 ^d	1a: Boc	2a: Et	3e	DCM	5	4aa (77)	83	>99
8 ^d	1a: Boc	2a: Et	3e	Brine	16	4aa (48)	91	>99
9 ^d	1a: Boc	2b: ⁱ Pr	3e	DCM	6	4ab (75)	95	>99
10	1a: Boc	2b: ⁱ Pr	3e	DCM	6	4ab (80)	>99	>99
11	1a: Boc	2b: ⁱ Pr	3d	DCM	6	4ab (75)	>99	>99
12	1b: Cbz	2b: ⁱ Pr	3e	DCM	8	4bb (65)	>99	>99
13	1b: Cbz	2b: ⁱ Pr	3d	DCM	8	4bb (63)	98	97
14	1c: CO ₂ Et	2b: ⁱ Pr	3e	DCM	8	4cb (42)	96	>99

^a Unless otherwise mentioned, all reactions were carried out with **1** (0.2 mmol), **2** (0.26 mmol), catalyst **3** (5 mol%) in DCM (0.3 M) at rt. ^b Yield refers to the column purified product. ^c *Ee* and *de* was determined by CSP HPLC analysis. ^d 10 mol% of catalyst **3** was used. ^e Reaction performed with 2 mol% of **3e**.

With the optimized conditions in hand, the scope of the Rawal's quinidine-NH-benzyl squaramide **3e**-catalyzed asymmetric Michael reaction was investigated. A series of substituted *N*-Boc-(*E*)-2-(2-nitrovinyl)anilines **1d-g** were reacted with 1.3 equiv. of cyclic β-keto esters **2a-f** catalyzed by 5 mol% of **3e** at 25 °C in DCM for 8–96 h to furnish the highly substituted chiral Michael adducts **4db-af** in 40–95% yields with excellent *ee*'s and *de*'s (Table 2). Electronic and steric nature of the *N*-Boc-(*E*)-2-(2-nitrovinyl)anilines were investigated with β-keto ester **2b**. Halogen substituted *N*-Boc-(*E*)-2-(2-nitrovinyl)anilines **1d-e** reacted well with **2b** and furnished the products (+)-**4db** and (+)-**4eb** in excellent yields (90 and 95%) and *ee*'s (99%) with good to moderate *de*'s (97 and 70%) within 8 h respectively. Methyl substituted *N*-Boc-(*E*)-2-(2-nitrovinyl)anilines **1f-g** gave the desired products (-)-**4fb** and (+)-**4gb** in moderate to good yields (40 and 70%) with excellent *ee*'s (>99%) and moderate *de*'s (47 and 88%) through **3e**-catalysis for 96 and 72 h, respectively. We also tested the other cyclic β-keto esters to investigate the generality of this asymmetric reaction. Methyl 2-oxocyclopentanecarboxylate **2c** with **1a** under the **3e**-catalysis furnished the product (-)-**4ac** in 81% yield with 94% *ee* and 95% *de* within 8 h. Intriguingly, six-membered cyclic β-keto ester **2d** gave the product (+)-**4ad** only in 40% yield with 94% *ee* and 84% *de* for 96 h; but simple ethyl 2-oxocyclohexanecarboxylate **2d'** gave the Michael product (+)-**4ad'** in 65% yield with >99% *ee* and >99% *de* for within 40 h (Table 2). Heterocyclic β-keto ester **2e** with **1a** under **3e**-catalysis furnished the product (+)-**4ae** in 60% yield with >99% *ee* and *de* for 24 h. In a similar manner, 2-acetylcyclopentanone **2f** with **1a** under **3e**-catalysis furnished the product (+)-**4af** in 62% yield with 94% *ee* and 79% *de* within 8 h.


It was observed that compared to five membered cyclic β -keto esters, six-membered cyclic β -keto esters gave the corresponding products in low yield. The structure and absolute stereochemistry of the Michael products **4** were confirmed by NMR analysis and also finally confirmed by X-ray structure analysis on (+)-**4ae** as shown in Fig. S1 (Supporting Information-1).¹²

Table 2 Reaction scope.^{a-c}

^a Yield refers to the column purified product. ^b Ee and de were determined by CSP HPLC analysis. ^c In parenthesis values refers to minor ee.

In order to expand the reaction scope, we explored the utilization of different acyclic β -keto esters **2g-l**. First the reaction of **1a** with 1.3 equiv. of ethyl acetoacetate **2g** under the catalysis of **3e** at 25 °C for 8 h in DCM furnished the product **4ag** in 76% yield with 1:1 *dr*. For the clear understanding of selectivity and also for the HPLC separation, we transformed the product **4ag** in situ into easily separable and also very important 1,4-dihydroquinoline (+)-**5ag** in 72% yield with 98% *ee* through TFA-mediated cascade amination/dehydration in DCM at 0-25 °C for 6 h in one-pot manner (Table 3). We further investigated the substrate scope for Rawal's quinidine-NH-benzyl squaramide **3e**-catalyzed and TFA-promoted asymmetric Michael/amination/dehydration reaction sequence with **1a-g** and **2h-l** for the synthesis of chiral 1,4-dihydroquinolines (Table 3). Methyl and *t*-butyl-3-oxobutanoates

2h-i with **1a** under **3e**-catalysis in DCM followed by TFA-mediated one-pot amination/dehydration reaction generated the cyclised products (+)-**5ah-ai** in 68% and 65% yields with 98% and 99% *ee*, respectively (Table 3). Likewise, ethyl 3-oxopentanoate and 3-oxohexanoate **2j-k** generated the expected cyclised products (+)-**5aj** and (+)-**5ak** in 65% and 60% yields with excellent *ee*'s

Table 3 Reaction application.^{a-c}

^a Yield refers to the column purified product. ^b Ee and de were determined by CSP HPLC analysis. ^c Only Micheal product obtained. ^d In parenthesis values refers to minor ee. ^e Reaction of (+)-**4al** with TFA gave ethyl 2-phenylquinoline-3-carboxylate in 65% yield (see SI).

(99% and 97%) after 36 h respectively (Table 3). In the case of ethyl 3-oxo-3-phenylpropanoate **2l**, the Michael product (+)-**4al** was obtained in 79% yield with 90% *ee* and 55% *de* at 25 °C for 8 h; but when in-situ treatment of (+)-**4al** with TFA at 0-25 °C for 24 h furnished the unexpected by-product ethyl 2-phenylquinoline-3-carboxylate in 65% yield may be due to the electronic/steric hindrance of phenyl group (Table 3 and see Supporting Information-I). Reaction of substituted *N*-Boc-(*E*)-2-(2-nitrovinyl)anilines **1d-e** with **2g** under **3e**-catalysis followed by in situ treatment with TFA furnished the cyclised products (+)-**5dg** and (+)-**5eg** in good yields (62 and 60%) with excellent *ee*'s (99 and 98%) at 25 °C for 12 h, respectively (Table 3). Surprisingly, the reaction of methyl substituted *N*-Boc-(*E*)-2-(2-nitrovinyl)aniline **1g** with **2g** under **3e**-catalysis followed by in situ treatment with

Communication

www.rsc.org/[JOURNAL] | [JOURNAL NAME HERE]

TFA furnished the only Michael product (+)-**4gg** in 76% yield with 94/99% *ee* and 1:1 *dr* at 25 °C for 48 h without cyclization may be due to the steric hindrance of methyl groups (Table 3).¹³ After these interesting results, TFA-mediated cascade amination/dehydration protocol was applied to the many of Table 2 compounds in DCM at 0-25 °C; for within 1 h compounds were decomposed.





Figure 2 Proposed transition state for the asymmetric Michael reactions.

With controlled experimental data in hand, herein we firmly elucidate the mechanism of the asymmetric Michael reaction through double hydrogen-bonding assembly by **3e**-catalysis and propose that the reaction most likely proceeds via **TS-1** mechanism (Figure 2). In the case of the addition of β -keto-esters **1** to substituted 2-(2-nitrovinyl)anilines **2** via Rawal's **3e**-catalysis, we can rationalize the observed stereochemistries through a favoured double hydrogen-bonding assembly transition state where the less hindered *si*-face of **2** approaches the *si*-face of the *in situ* generated enol as shown in **TS-1**. Outcome of decent selectivity and reactivity for the β -keto-ester **1** addition to the 2-(2-nitrovinyl)anilines **2** could be explained by soft involvement due to the steric hindrance of neighboring group Ar-NHBoc as shown in Figure 2. Fascinatingly, the Michael reaction of simple (*E*)-(2-nitrovinyl)benzene **1h** and *N*-Boc-(*E*)-3-(2-nitrovinyl)aniline **1i** with **2b** via **3e**-catalysis in DCM at 25 °C for 6 h furnished the expected products (-)-**4hb** in 63% yield with 98% *ee* and >99% *de*; and (-)-**4ib** in 71% yield with 99% *ee* and >99% *de* respectively as shown in eq. (1). This result gives evident for the fact that there is not much of neighboring *ortho*-amino group participation in the reaction pre-transition state and also Rawal's catalyst **3e** is sufficient enough to activating both the substrates in the pre-transition state along with steric hindrance of neighboring group Ar-NHBoc to control the selectivity.



In summary, we have developed Rawal's quinidine-NH-benzyl squaramide **3e**-catalyzed and TFA-promoted asymmetric sequential Michael/amination/dehydration reaction of *N*-protected-(*E*)-2-(2-nitrovinyl)anilines **1** with β -keto esters **2** for the synthesis of enantioenriched and highly functionalized 1,4-dihydroquinolines. Further work is in progress to utilize 1,4-dihydroquinoline derivatives for biological studies.

Acknowledgements


This work was made possible by a grant from the Department of Science and Technology (DST), SERB, New Delhi [Grant No.: EMR/2015/000860]. ASK thanks University Grants Commission (UGC)-Dr. D. S. Kothari Postdoctoral Fellowship (DSKPDF) Scheme, New Delhi for his postdoctoral research fellowship. We thank Dr P. Raghavaiah for his help in X-ray structural analysis.

Notes and references

- For selected reviews on asymmetric organocatalytic sequential one-pot reactions, see: (a) D. Enders, C. Grondal and M. R. M. Huettl, *Angew. Chem. Int. Ed.*, 2007, **46**, 1570-1581; (b) A. Erkkila, I. Majander and P. M. Pihko, *Chem. Rev.*, 2007, **107**, 5416-5470; (c) S. Bertelsen and K. A. Jorgensen, *Chem. Soc. Rev.*, 2009, **38**, 2178-2189; (d) B. Westermann, M. Ayaz and S. S. van Berkel, *Angew. Chem. Int. Ed.*, 2010, **49**, 846-849; (e) C. Grondal, M. Jeanty and D. Enders, *Nat. Chem.*, 2010, **2**, 167-178; (f) D. B. Ramachary and S. Jain, *Org. Biomol. Chem.*, 2011, **9**, 1277-1300; (g) M. Rueping, J. Dufour and F. R. Schoepke, *Green Chem.*, 2011, **13**, 1084-1105; (h) D. B. Ramachary and Y. V. Reddy, *Eur. J. Org. Chem.*, 2012, 865-887; (i) J. Yu, F. Shi and L. -Z. Gong, *Acc. Chem. Res.*, 2011, **44**, 1156-1171; For selected papers on asymmetric organocatalytic sequential one-pot reactions, see: (j) N. S. Chowdari, D. B. Ramachary and C. F. Barbas III, *Org. Lett.*, 2003, **5**, 1685-1688; (k) D. B. Ramachary and C. F. Barbas III, *Chem. Eur. J.*, 2004, **10**, 5323-5331; (l) J. W. Yang, M. T. Hechavarria Fonseca and B. List, *J. Am. Chem. Soc.*, 2005, **127**, 15036-15037; (m) Y. Huang, A. M. Walji, C. H. Larsen and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2005, **127**, 15051-15053; (n) M. Marigo, S. Bertelsen, A. Landa and K. A. Jorgensen, *J. Am. Chem. Soc.*, 2006, **128**, 5475-5479; (o) D. Enders, M. R. M. Huettl, C. Grondal and G. Raabe, *Nature* 2006, **441**, 861-863; (p) W. Wang, H. Li, J. Wang and L. Zu, *J. Am. Chem. Soc.*, 2006, **128**, 10354-10355; (q) M. Rueping, A. P. Antonchick and T. Theissmann, *Angew. Chem. Int. Ed.*, 2006, **45**, 3683-3686; (r) J. Wang, H. Li, H. Xie, L. Zu, X. Shen and W. Wang, *Angew. Chem. Int. Ed.*, 2007, **46**, 9050-9053; (s) A. Carlone, S. Cabrera, M. Marigo and K. A. Jorgensen, *Angew. Chem. Int. Ed.*, 2007, **46**, 1101-1104; (t) J. Zhou and B. List, *J. Am. Chem. Soc.*, 2007, **129**, 7498-7499; (u) O. Penon, A. Carlone, A. Mazzanti, M. Locatelli, L. Sambri, G. Bartoli and P. Melchiorre, *Chem. Eur. J.*, 2008, **14**, 4788-4791; (v) D. Enders, C. Wang and J. W. Bats, *Angew. Chem. Int. Ed.*, 2008, **47**, 7539-7542; (w) D. Zhu, M. Lu, P. J. Chua, B. Tan, F. Wang, X. Yang and G. Zhong, *Org. Lett.*, 2008, **10**, 4585-4588; (x) H. Ishikawa, T. Suzuki and Y. Hayashi, *Angew. Chem. Int. Ed.*, 2009, **48**, 1304-1307; (y) M. Rueping, A. Kuenkel, F. Tato and J. W. Bats, *Angew. Chem. Int. Ed.*, 2009, **48**, 3699-3702; (z) R. Imashiro, H. Uehara and C. F. Barbas III, *Org. Lett.*, 2010, **12**, 5250-5253; (aa) K. Jiang, Z. -J. Jia, L. Wu and Y. -C. Chen, *Org. Lett.*, 2010, **12**, 2766-2769; (ab) H. Ishikawa, M. Honma and Y. Hayashi, *Angew. Chem. Int. Ed.*, 2011, **50**, 2824-2827.
- (a) C. Min, X. Han, Z. Liao, X. Wu, H.-B. Zhou and C. Dong, *Adv. Synth. Catal.*, 2011, **353**, 2715; (b) A. Dondoni and A. Massi, *Angew. Chem., Int. Ed.*, 2008, **47**, 4638; (c) D. W. C. MacMillan, *Nature*, 2008, **455**, 304; (d) C. F. Nising and S. Bräse, *Chem. Soc. Rev.*, 2008, **37**, 1218; (e) S. S. Mossé and A. Alexakis, *Chem. Commun.*, 2007, 3123; (f) D. Almaşi, D. A. Alonso and C. Nájera, *Tetrahedron: Asymmetry*, 2007, **18**, 299; (g) S. B. Tsogoeva, *Eur. J. Org. Chem.*, 2007, 1701; (h) B. List and J. W. Yang, *Science*, 2006, **313**, 1584; (i) R. Ballini, G. Bosica, D. Fiorini, A. Palmieri and M. Petrini, *Chem. Rev.*, 2005, **105**, 933; (j) L.-W. Xu and C.-G. Xia, *Eur. J. Org. Chem.*, 2005, 633; (k) W. Notz, F. Tanaka and C. F. Barbas III, *Acc. Chem. Res.*, 2004, **37**, 580; (l) J. Christoffers and A. Baro, *Angew. Chem., Int. Ed.*, 2003, **42**, 1688; (m) O. M. Berner, L. Tedeschi and D. Enders, *Eur. J. Org. Chem.*, 2002, 1877.
- (a) Z. Dong, X. Jin, P. C. Wang, C. Min, J. Zhang, Z. Chen, H.-B. Zhou and C. Dong, *ARKIVOC*, 2011, (ix), 367; (b) R. Manzano, J. M. Andrés, M. D. Muruzábal and R. Pedrosa, *Adv. Synth. Catal.*, 2010, **352**, 3364; (c) S. Ban, D.-M. Du, H. Liu and W. Yang, *Eur. J. Org. Chem.*, 2010, 5160; (d) T. Tsubogo, Y. Yamashita and S. Kobayashi, *Angew. Chem. Int. Ed.*, 2009, **48**, 9117; (e) B. Tan, X. Zhang, P. J. Chua and G. F. Zhong, *Chem. Commun.*, 2009, 779; (f) W.-M. Zhou, H. Liu and D.-M. Du, *Org. Lett.*, 2008, **10**, 2817; (g) P. Gao, C. Wang, Y. Wu, Z. Zhou and C. Tang, *Eur. J. Org. Chem.*,

- 2008, 4563; (h) N. Mase, K. Watanabe, H. Yoda, F. Tanaka and C. F. Barbas III, *J. Am. Chem. Soc.*, 2006, **128**, 4966; (i) W. Wang, J. Wang and H. Li, *Angew. Chem., Int. Ed.*, 2005, **44**, 1369; (j) C. Czekelius and E. M. Carreira, *Angew. Chem., Int. Ed.*, 2005, **44**, 612; (k) N. Mase, R. Thayumanavan, F. Tanaka and C. F. Barbas III, *Org. Lett.*, 2004, **6**, 2527; (l) T. Okino, Y. Hoashi and Y. Takemoto, *J. Am. Chem. Soc.*, 2003, **125**, 12672.
- 4 (a) T. Okino, Y. Hoashi, T. Furukawa, X. Xu and Y. Takemoto, *J. Am. Chem. Soc.*, 2005, **127**, 119; (b) T. Okino, Y. Hoashi and Y. Takemoto, *J. Am. Chem. Soc.*, 2003, **125**, 12672.
- 5 (a) K. S. Yang, A. E. Nibbs, Y. E. Turkmen and V. H. Rawal, *J. Am. Chem. Soc.*, 2013, **135**, 16050; (b) Y. Zhu, J. P. Malerich and V. H. Rawal, *Angew. Chem., Int. Ed.*, 2010, **49**, 153; (c) Y. Qian, G. Ma, A. Lv, H.-L. Zhu, J. Zhao and V. H. Rawal, *Chem. Commun.*, 2010, **46**, 3004; (d) H. Konishi, T. Y. Lam, J. P. Malerich and V. H. Rawal, *Org. Lett.*, 2010, **12**, 2028; (e) J. P. Malerich, K. Hagihara and V. H. Rawal, *J. Am. Chem. Soc.*, 2008, **130**, 14416.
- 6 (a) H. Y. Bae and C. E. Song, *ACS Catal.*, 2015, **5**, 3613; (b) P. Chauhan, S. Mahajan, U. Kaya, D. Hack and D. Enders, *Adv. Synth. Catal.*, 2015, **357**, 253; (c) D.-F. Chen, F. Zhao, Y. Hu and L.-Z. Gong, *Angew. Chem., Int. Ed.*, 2014, **53**, 10763; (d) L. Albrecht, G. Dickmeiss, F. C. Acosta, C. R. E. Schrich, R. L. Davis and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2012, **134**, 2543; (e) R. I. Storer, C. Aciro and L. H. Jones, *Chem. Soc. Rev.*, 2011, **40**, 2330; (f) H.-L. Song, K. Yuan and X.-Y. Wu, *Chem. Commun.*, 2011, **47**, 1012; (g) H. Y. Bae, S. Some, J. H. Lee, J.-Y. Kim, M. J. Song, S. Lee, Y. J. Zhang and C. E. Song, *Adv. Synth. Catal.*, 2011, **353**, 3196; (h) J. Alemln, A. Parra, H. Jiang and K. A. Jørgensen, *Chem. Eur. J.*, 2011, **17**, 6890; (i) H. Jiang, M. W. Paixão, D. Monge and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2010, **132**, 2775; (j) L. Dai, S.-X. Wang and F.-E. Chen, *Adv. Synth. Catal.*, 2010, **352**, 2137; (k) J. W. Lee, T. H. Ryu, J. S. Oh, H. Y. Bae, H. B. Jang and C. E. Song, *Chem. Commun.*, 2009, 7224.
- 7 For selected examples of organocatalytic asymmetric syntheses of chromans and 3,4-dihydrocoumarins, see: (a) P. Kotame, B. -C. Hong and J. -H. Liao, *Tetrahedron Lett.*, 2009, **50**, 704-707; (b) L. Zu, S. Zhang, H. Xie and W. Wang, *Org. Lett.*, 2009, **11**, 1627-1630; (c) A. E. Nibbs, A. -L. Baize, R. M. Herter and K. A. Scheidt, *Org. Lett.*, 2009, **11**, 4010-4013; (d) H. Lv, L. You and S. Ye, *Adv. Synth. Catal.*, 2009, **351**, 2822-2826; (e) D. B. Ramachary and R. Sakthidevi, *Chem. Eur. J.*, 2009, **15**, 4516-4522; (f) B. -C. Hong, P. Kotame, C. -W. Tsai and J. -H. Liao, *Org. Lett.*, 2010, **12**, 776-779; (g) J. -L. Li, S. -L. Zhou, B. Han, L. Wu and Y. -C. Chen, *Chem. Commun.*, 2010, **46**, 2665-2667; (h) K. -S. Choi and S. -G. Kim, *Tetrahedron Lett.*, 2010, **51**, 5203-5206; (i) X. -F. Wang, Q. -L. Hua, Y. Cheng, A. -L. An, Q. -Q. Yang, J. -R. Chen and W. -J. Xiao, *Angew. Chem. Int. Ed.*, 2010, **49**, 8379-8383; (j) D. B. Ramachary and R. Sakthidevi, *Org. Biomol. Chem.*, 2010, **8**, 4259-4265; (k) D. Enders, C. Wang, X. Yang and G. Raabe, *Adv. Synth. Catal.*, 2010, **352**, 2869-2874; (l) D. Lu, Y. Li and Y. Gong, *J. Org. Chem.*, 2010, **75**, 6900-6907; (m) B. -C. Hong, P. Kotame and J. -H. Liao, *Org. Biomol. Chem.*, 2011, **9**, 382-386; (n) D. B. Ramachary, M. S. Prasad and R. Madhavachary, *Org. Biomol. Chem.*, 2011, **9**, 2715-2721; (o) Q. -L. Hua, C. Li, X. -F. Wang, L. -Q. Lu and J. -R. Chen, *ACS Catal.*, 2011, **1**, 221-226; (p) D. Enders, X. Yang, C. Wang, G. Raabe and J. Runsik, *Chem. Asian. J.*, 2011, **6**, 2255-2259; (q) B. -C. Hong, P. Kotame and G. -H. Lee, *Org. Lett.*, 2011, **13**, 5758-5761.
- 8 (a) J. D. Shields, D. T. Ahneman, T. J. A. Graham and A. G. Doyle, *Org. Lett.*, 2014, **16**, 142; (b) C. Luo and Y. Huang, *J. Am. Chem. Soc.*, 2013, **135**, 8193; (c) W. Yang, H.-X. He, Y. Gao and D.-M. Du, *Adv. Synth. Catal.*, 2013, **355**, 3670; (d) S. N. Mistry, C. Valant, P. M. Sexton, B. Capuano, A. Christopoulos and P. J. Scammells, *J. Med. Chem.*, 2013, **56**, 5151; (e) V. Sridharan, P. A. Suryavanshi and J. C. Menéndez, *Chem. Rev.*, 2011, **111**, 7157; (f) S. Heeb, M. P. Fletcher, S. R. Chhabra, S. P. Diggle, P. Williams and M. Cámara, *FEMS Microbiol. Rev.*, 2011, **35**, 247; (g) J. Fotie, M. Kaiser, D. A. Delfin, J. Manley, C. S. Reid, J.-M. Paris, T. Wenzler, L. Maes, K. V. Mahasenan, C. Li and K. A. Werbovets, *J. Med. Chem.*, 2010, **53**, 966; (h) X. Liu and Y. Lu, *Org. Biomol. Chem.*, 2010, **8**, 4063; (i) L. Foucoute, F. Gourand, M. Dhilly, P. Bohn, G. Dupas, J. Costentin, A. Abbas, F. Marsais, L. Barré and V. Levacher, *Org. Biomol. Chem.*, 2009, **7**, 3666; (j) J. P. Michael, *Nat. Prod. Rep.*, 2008, **25**, 166.
- 9 For selected synthesis of chiral 1,4-dihydroquinolines, see: (a) X. Zhang, X. Song, H. Li, S. Zhang, X. Chen, X. Yu and W. Wang, *Angew. Chem., Int. Ed.*, 2012, **51**, 7282; (b) F. Rezgui, P. Mangeney and A. Alexakis, *Tetrahedron Lett.*, 1999, **40**, 6241; (c) X. Liu and Y. Lu, *Org. Lett.*, 2010, **12**, 5592.
- 10 (a) D. B. Ramachary, K. S. Shruthi and R. Madhavachary, *Eur. J. Org. Chem.*, 2015, 6413; (b) Y. Lee and S.-G. Kim, *J. Org. Chem.*, 2014, **79**, 8234.
- 11 During the preparation of this manuscript, Kim group reported asymmetric synthesis of 1,4-dihydroquinolines with lower selectivity (<90% ee) and more catalyst loading, see: H. Kim and S.-G. Kim, *Tetrahedron Lett.*, 2015, **56**, 4819.
- 12 CCDC-1432348 for (+)-**4ae** contains the supplementary crystallographic data for this paper. This data can be obtained free of charge via <http://www.ccdc.cam.ac.uk/conts/retrieving.html> or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB21EZ, UK; fax: (+44)1223-336-033; or <mailto:deposit@ccdc.cam.ac.uk>. See supporting information for crystal structure.
- 13 For organocatalytic high-yielding synthesis of racemic products **4** and **5** from **1** and **2** through [Q + QD]-catalysis followed by TFA, see Table S1 and S2 in Supporting Information-I. We have also done racemic products **4** synthesis from **1** and **2** through DBU or DABCO-catalysis, but reaction conversions are very low even after 24 h at 25 °C.

Graphical Abstract for Table of Contents:



Short Statement

A general approach to the asymmetric synthesis of highly substituted dihydroquinolines was achieved through neighboring *ortho*-amino group engaged sequential Michael/amination/dehydration reactions on (*E*)-2-(2-nitrovinyl)anilines with cyclic and acyclic β -keto esters in the presence of a catalytic amount of Rawal's quinidine-*NH*-benzyl squaramide followed by TFA (see Scheme).