

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



RSC Advances

ARTICLE

Copper-catalyzed rapid C-H nitration of 8-aminoquinolines by using sodium nitrite as the nitro source under mild conditions

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x


www.rsc.org/

Xiaolei Zhu,^a Li Qiao,^a Pingping Ye,^a Beibei Ying,^a Jun Xu,^a Chao Shen^b and Pengfei Zhang^{*a}

A mild, rapid and efficient method for copper-catalyzed nitration of quinolines at C5 or C7 position was reported firstly by using sodium nitrite as the nitro source. A series of nitrated quinoline derivatives were achieved in moderate to good yields via remote C-H activation. The method overcomes some shortcomings reported in previous nitration strategies, such as employing expensive metal catalysts or using toxic and unstable nitrating agents. Furthermore, some of products can be obtained through filtration which displayed the operational simplicity of this methodology. Importantly, this protocol may provide a practical synthetic tool for the introduction of nitro group into drug molecules.

Introduction

Organic nitro compounds are important raw materials in the synthesis of dyes, plastics, explosives, perfumes and pharmaceuticals.¹ The traditional routes to aromatic nitro compounds, such as the electrophilic aromatic nitration, usually require harsh conditions (HNO₃/H₂SO₄), and suffer from regioselectivity issues as well as the unsatisfactory functional group tolerance (Scheme 1 eq 1).² To overcome these problems, recently several strategies of regiospecific synthesis of nitroarenes, including the ispo-nitration of aryl boronic acids,³ aryl halides, pseudohalides,⁴ and aryl carboxylic acids⁵ have been developed (Scheme 1 eq 2). Alternatively, ispo-oxidation of aryl amines or azides also provided a nitration protocol to give the nitroarenes indirectly.⁶ However, these promising approaches are low atom economy. Furthermore, related wastes are produced.



Scheme 1 A series of strategies leading to nitroarenes

In view of the encouraging achievements made in recent transition-metal-catalyzed auxiliary-assisted C-H functionalization which successfully construct C-C, C-O, C-S, C-N and C-halogen bonds,⁷ a similar strategy would be appealing also for the direct C-H nitration (Scheme 1 eq 3). Specially, liu et al. has shown a method for palladium-catalyzed *ortho* nitration of (hetero)arenes at 130 °C.⁸ Bi's copper-catalyzed *ortho* nitration of (hetero)arenes also required high temperature and used expensive AgNO₂ as the nitro source.⁹ The latest radical C-H nitration of arenes by Jiao and co-workers performs at 80 °C, but employed expensive Pd catalyst.¹⁰ Obviously, these protocols should abide a number of limitations, mainly using expensive metal catalysis, choosing costly silver(I) salt as the nitrating agent or requiring harsh reaction conditions.¹¹ Therefore, mild and efficient strategies for selective nitration is still a great need. In 2014, the carretero group have disclosed a mild and efficient copper-catalyzed nitration of aromatic derivatives but using unstable nitric acid as the nitrating agent at 100 °C.¹² Just recently, the Ribas group reported a mild protocol for remote C-H nitration of 8-aminoquinoline amides using inexpensive cobalt nitrate hexahydrate as catalyst and *tert*-butyl nitrate (TBN) as nitro source.¹³ However, the lack of selectivity for the unsymmetrical

^a College of Material, Chemistry and Chemical Engineering, Hangzhou Normal University, Hangzhou 310036 (China). E-mail: chxyzpf@hotmail.com

^b College of Biology and Environmental Engineering, Zhejiang Shuren University, Hangzhou 310015 (China).

† Electronic Supplementary Information (ESI) available: Detailed experimental procedures and analytical data. CCDC 1444178 for **3a**. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/x0xx00000x

COMMUNICATION

Journal Name

quinoline rings (Ratio C5:C7: 3:1), the excess use of unstable and toxic nitrating agent and long reaction times (18h) have limited the synthesis value of this methodology. Herein, we report a copper-catalyzed mild and efficient method for the remote C-H nitration of 8-aminoquinoline amides using cheap and stable sodium nitrite as the nitro source (Scheme 1 eq 4). It is noteworthy that the reaction was completed within one minute at room temperature.

Results and discussion


Our original goal was to achieve the copper-catalysed 8-aminoquinoline directed *ortho* C-H nitration of benzoic acid derivatives. We started our study by choosing the reaction of benzamide **1a** with sodium nitrite as the model (Table 1). To our surprise, the nitration product at the *ortho* position of benzoic acid derivatives was not observed. Instead, in the presence of $\text{Cu}(\text{NO}_3)_2$ (10 mol%), PivOH (20 mol%), and $\text{PhI}(\text{OAc})_2$ (2.0 equiv.) at 70 °C in CH_3CN overnight, unexpected nitrated product **2a** was obtained in 70% yield by filtration (Table 1, entry 15). Encouraged by this result, we further investigated the reaction parameters from catalyst, oxidant, and solvent. Gratifyingly, the reaction proceeded rapidly at room temperature in higher yield when $\text{PhI}(\text{TFA})_2$ was used as an oxidant, and the additive PivOH was indispensable (Table 1, entry 9). The transformation could not take place in the presence of other oxidants, such as TBHP and $\text{K}_2\text{S}_2\text{O}_8$ (Table 1, entries 16-17). The

Table 1 Screening of reaction conditions for nitration of 8-aminoquinoline amides^{a,b}

Entry	Catalyst	Oxidant	Solvent	Yield (%) ^b
1	CuI	$\text{PhI}(\text{TFA})_2$	CH_3CN	0
2	PdCl_2	$\text{PhI}(\text{TFA})_2$	CH_3CN	0
3	AgNO_3	$\text{PhI}(\text{TFA})_2$	CH_3CN	0
4	$\text{Fe}(\text{NO}_3)_3$	$\text{PhI}(\text{TFA})_2$	CH_3CN	0
5	$\text{Co}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	CH_3CN	0
6 ^c	$\text{Cu}(\text{OAc})_2$	$\text{PhI}(\text{TFA})_2$	CH_3CN	76
7	$\text{Cu}(\text{NO}_3)_2$	none	CH_3CN	0
8	none	$\text{PhI}(\text{TFA})_2$	CH_3CN	0
9	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	CH_3CN	86
10 ^e	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	CH_3CN	80
11 ^{d,e}	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	CH_3CN	30
12 ^f	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	CH_3CN	73
13 ^g	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	CH_3CN	85
14 ^h	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	CH_3CN	83
15	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{OAc})_2$	CH_3CN	trace (70) ^e
16	$\text{Cu}(\text{NO}_3)_2$	TBHP	CH_3CN	0
17	$\text{Cu}(\text{NO}_3)_2$	$\text{K}_2\text{S}_2\text{O}_8$	CH_3CN	0
18	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	DMF	trace
19	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	CH_3NO_2	trace
20	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	dioxane	trace
21	$\text{Cu}(\text{NO}_3)_2$	$\text{PhI}(\text{TFA})_2$	DCE	trace

^a Reaction conditions: **1a** (0.20 mmol), sodium nitrite (1.1 equiv), catalyst (10 mol %), oxidant (1.5 equiv), PivOH (20 mol %), solvent (2.0 mL); ^b Isolated yield of **2a**. ^c The reaction was completed in 5 min; ^d No PivOH was added; ^e The reaction was carried out at 70 °C; ^f The reaction time was 30 min; ^g Under N_2 ; ^h Under O_2 .


Table 2 Substrate scope of Cu(II)-catalysed nitration of C5-H of 8-aminoquinoline amides^{a,b}



^a Reaction conditions: **1** (0.20 mmol), sodium nitrite (0.22 mmol), $\text{Cu}(\text{NO}_3)_2$ (10 mol %), $\text{PhI}(\text{TFA})_2$ (1.5 equiv.), PivOH (20 mol %), CH_3CN (2.0 mL); ^b Isolated yield.

yield of product **2a** was declined slightly with reaction temperature increasing (Table 1, entry 10). Instead of $\text{Cu}(\text{NO}_3)_2$, $\text{Cu}(\text{OAc})_2$ resulted in lower yield and longer reaction time (Table 1, entry 6). Other metal salts such as CuI , PdCl_2 , AgNO_3 , $\text{Fe}(\text{NO}_3)_2$, $\text{Co}(\text{NO}_3)_2$ proved ineffective for this reaction (Table 1, entries 1-5). A screening of different solvents showed that CH_3CN was still the optimum solvent and no reaction occurred in other solvents (Table 1, entries 18-21). When the reaction time was extended to 30 min, however, the nitration product just be obtained in lower yield (Table 1, entry 12). Impressively, control experiments revealed that atmosphere has no effect on the reaction outcome, which highlighted the operational simplicity of this method (Table 1, entries 13 and 14). Furthermore, the reaction did not work in

Table 3 Substrate scope of Cu(II)-catalysed nitration of C7-H of 8-aminoquinoline amides^{a,b}




^a Reaction conditions: **3** (0.20 mmol), sodium nitrite (0.22 mmol), $\text{Cu}(\text{NO}_3)_2$ (10 mol %), $\text{PhI}(\text{TFA})_2$ (1.5 equiv), PivOH (20 mol %), CH_3CN (2.0 mL); ^b Isolated yield.


absence of either $\text{Cu}(\text{NO}_3)_2$ or $\text{PhI}(\text{TFA})_2$ (entries 7-8).

With the optimized conditions in hand, the scope of this reaction was investigated (Table 2). The results showed that a wide range of 8-aminoquinoline amides reacted smoothly with sodium nitrite to generate the corresponding nitrated products (**2a-2k**). 8-aminoquinolinebenzamide was successfully converted into the desired product **2a** in an excellent yield of 86% under the standard conditions. The substrate bearing Me group on the phenyl ring could afford the nitrated product **2b** in 84% yield. Moreover, aliphatic amides, including those with methyl, cyclopropyl, cyclohexyl and *tert*-butyl could also be employed to afford the target compounds **2c-2f** in moderate to good yields. Aliphatic amides with oxygen heteroatom **1g**, **1i** and **1j** were also found apply to this transformation. In addition, the furancarboxylic acid derived substrate **1h** was suitable for this reaction. Especially, the substrate of Ph group on the quinoline ring was also been well tolerated, giving the desired product **2k** in 84% yield.


Encouraged by the wide functional group tolerance of this C5-H nitration, we wondered whether this protocol could be applied to the nitration of quinoline ring at C7 position. To our delight, the reaction exclusively occurred at the C7 position of quinoline ring under the typical reaction conditions if the C5 position was brominated (table 3).^{14a} Various C5-brominated aminoquinoline amides reacted with sodium nitrite to obtain the desired product **4a-4f** in moderate to good yields. The structure of the product **4a** was confirmed by X-ray crystallography (1444178, CCDC NO.), as shown in the supporting information.



Scheme 2 Gram scale reaction




Scheme 3 Application



Scheme 4 control experiments

Scheme 5 plausible mechanism



Given the easy available of the raw materials and the operational simplicity of this method, we performed on a gram scale reaction (Scheme 2), and the nitrated product **2a** was obtained in 82% yield. The obtained product could be further transformed into *N*-(5-aminoquinolin-8-yl)benzamide **5** in 88% yield by Pd/C hydrogenation.¹⁵ These two transformations demonstrated that the nitration product is also provided with synthetic usefulness. Importantly, quinolone derivate **1l** was easily converted to C5 nitrated product **2l** under standard conditions which might be further used to synthesize some interesting compounds of biological and pharmacological activities (Scheme 3).¹⁶

In order to obtain the mechanistic information about this transformation, some control experiments were carried out. Substrate **6** (1-aminonaphthalene derived benzoic acid) did not give any nitrated product under standard conditions (Scheme 3, a). No desired product was obtained in the reaction when the substrate was changed into *N*-methyl aminoquinoline **7** or 8-acyloxyquinoline **8** (Scheme 4, a). These reactions showed that the formation of the bidentate structure was crucial to the remote C-H nitration.¹⁷ To figure out whether the radical progress would be involved in the reaction, a radical inhibition experiment of **1a** with the addition of radical scavenger TEMPO (2,2,6,6-tetramethyl-1-piperidinyloxy) or hydroquinone was further performed under the standard reactions (Scheme 4, b). Completely suppression of the reaction suggested that a radical pathway might be responsible for this transformation. In addition, capturing of radical test was carried out to prove the feasibility of the radical progress and the result revealed that the nitro radical ($\text{NO}_2^\bullet$) was existent in the reaction process (Scheme 4, c). The influence of the quantity of $\text{PhI}(\text{TFA})_2$ and reaction atmosphere were further investigated (Scheme 4, d; Table 1, entries 11 and 12). The results explained that reducing the oxidant loading would apparently affect the reaction efficiency. This phenomenon illustrated that the $\text{PhI}(\text{TFA})_2$ plays a dual role in the reaction, generating nitro radical and oxidating the intermediate in the reaction process.

Based on above experimental results and our previous studies,¹⁴ a plausible mechanism was depicted in Scheme 5. First, the coordination of substrate **1a** with $\text{Cu}(\text{II})$ produced chelated complex

A. Afterwards, the deprotonation of the amide group leads to formation of the complex **B**. Then, nitro radical($\text{NO}_2\bullet$) was released in the presence of $\text{PhI}(\text{TFA})_2$ and NaNO_2 . Subsequently, the single electron transfer (SET) occurred between the nitro radical($\text{NO}_2\bullet$) and $\text{Cu}(\text{II})$ complex **B** to afford the $\text{Cu}(\text{I})$ intermediate **C**, followed by the formation of complex **D** through oxidation. Finally, the intermediate **E** generated through proton transfer process, and the desired product **2a** was obtained via ligand dissociation, along with the regeneration of the $\text{Cu}(\text{II})$ species to complete the catalytic cycle.

Conclusions

In summary, we have developed a novel copper-catalyzed highly regioselective direct C-H nitration of quinolines with cheap sodium nitrite, which afforded good yields of C5-H or C7-H nitrated quinoline derivatives. This transformation featured a mild reaction system, wide substrate scope as well as the short reaction times. Importantly, this protocol may provide a green and practical approach to construct nitrated quinoline derivatives, which were the key intermediates in the drug synthesis.

Experimental section

General information

Unless indicated, all reagents were obtained commercially and used without further purification. Flash chromatography was carried out with silica gel (200-300 mesh). ^1H and ^{13}C NMR spectras were recorded in CDCl_3 or $\text{DMSO}-d_6$ on a Bruker Advance 500 spectrometer at ambient temperature with CDCl_3 or $\text{DMSO}-d_6$ as solvent and tetramethylsilane (TMS) as the internal standard. ^1H NMR data were reported as follows: chemical shift (δ ppm), multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet), coupling constant (J values, Hz). ^{13}C NMR data were reported in terms of chemical shift (δ ppm). Melting points were determined on an X-5 Data microscopic melting point apparatus. The small-angle X-ray diffraction(SAXRD) data was performed at room temperature using German Bruker D4 X-ray diffractometer. Analytical thin layer chromatography (TLC) was performed with Merk precoated TLC (silica gel 60 F254) plates. High resolution mass spectra (HRMS) were recorded using Agilent 6530 QTOF mass spectrometer. **1a-1k** were obtained according to literature reported¹⁴, **3a-3f** were formed according to literature reported.^{14a}

General procedure

General method for Copper(II)-catalyzed C5-H and C7-H nitration of 8-aminoquinoline amides. (**2a**, **2b**, **2k**, **4a** and **4b**): A mixture of amide (0.2 mmol), NaNO_2 (15.0 mg, 0.22 mmol, 1.1 equiv), $\text{PhI}(\text{TFA})_2$ (129.0 mg, 0.3 mmol, 1.5 equiv), $\text{Cu}(\text{NO}_3)_2$ (3.8 mg, 0.02 mmol, 10 mol %), PivOH (4.0 mg, 0.04 mmol, 20 mol %), and CH_3CN (2.0 mL) were stirred at room temperature for 1 min. After completion (monitored by TLC), the mixture was filtered through a celite pad, then washed with CH_3CN and dried to give the corresponding product. (**2c-2j**, **4c-4f**): A mixture of amide (0.2 mmol), NaNO_2 (15.0 mg, 0.22 mmol, 1.1 equiv), $\text{PhI}(\text{TFA})_2$ (129.0 mg, 0.3 mmol, 1.5 equiv), $\text{Cu}(\text{NO}_3)_2$ (3.8 mg, 0.02 mmol, 10 mol %),

PivOH (4.0 mg, 0.04 mmol, 20 mol %), and CH_3CN (2.0 mL) were stirred at room temperature for 1 min. After completion (monitored by TLC), the solvent was removed under reduced pressure and the residue was purified by silica gel column chromatography to give the corresponding product.

Pd/C hydrogenation of 2a. Compound **2a** (60.0 mg, 0.2 mmol), and Pd/C (1.2 mg, 20%) were taken in a dried schlenk tube with a magnetic stir bar under hydrogen, then MeOH (3.0 mL) was added with a syringe and the resulting mixture was stirred for 6 h at room temperature. After completion (monitored by TLC), the mixture was filtered through a celite pad, and the solvent was removed under reduced pressure. The residue was purified by silica gel column chromatography to give the reduction product.

Acknowledgements

Financial support from the National Natural Science Foundation of China (No. 21376058, 21302171), Changjiang Scholars and Innovative Research Team in Chinese University (IRT 1231), Zhejiang Provincial Natural Science Foundation of China (No. LZ13B020001).

Notes and references

- (a) R. Ballini, L. Barboni, F. Fringuelli, A. Palmieri, F. Pizzo and L. Vaccaro, *Green Chem.*, 2007, **9**, 823; (b) W.-P. Hou, Q. Wei, G.-S. Liu, J. Chen, J. Guo and Y.-G. Peng, *Org. Lett.*, 2015, **17**, 4870; (c) N. Ono, The Nitro Group in Organic Synthesis, Wiley-VCH, Weinheim, 2001; (d) M. V. Papadopolou, B. B. Trunz, W. D. Bloomer, C. McKenzie, S. R. Wilkinson, C. Prasittichai, R. Brun, M. Kaiser and E. Torreele, *J. Med. Chem.*, 2011, **54**, 8214.
- (a) K. Schofield, Aromatic Nitration, Cambridge University Press, Cambridge, 1980; (b) E. Mitscherlich, *AnInPhys.Chem.*, 1834, **31**, 625; (c) E. Mitscherlich, *AnIn Pharm.*, 1834, **12**, 305; (d) M. Feldman and J. W. Wheeler, *J. Chem. Educ.*, 1967, **44**, 464; (e) G. A. Olah, R. Malhotra and S. C. Narang, Nitration: Methods and Mechanisms, VCH: New York, 1989; (f) K. Schofield, Aromatic Nitration, Cambridge University Press: Cambridge, 1980.
- (a) R. R. Yadav, R. A. Vishwakarma and S. B. Bharate, *Tetrahedron Lett.*, 2012, **53**, 5958; (b) X.-F. Wu, J. Schranck, H. Neumann and M. Beller, *Chem. Commun.*, 2011, **47**, 12462; (c) M. Jiang, H.-J. Yang, Y. Li, Z.-Y. Jia and H. Fu, *RSC Adv.*, 2013, **3**, 25602; (d) S. Manna, S. Maity, S. Rana, S. Agasti and D. Maiti, *Org. Lett.*, 2012, **14**, 1736; (e) G. K. S. Prakash, C. Panja, T. Mathew, V. Surampudi, N. A. Petasis and G. A. Olah, *Org. Lett.*, 2004, **6**, 2205.
- (a) S. Saito and Y. Koizumi, *Tetrahedron Lett.*, 2005, **46**, 4715; (b) S. Paik and M. G. Jung, *Bull. Korean. Chem. Soc.*, 2012, **33**, 689.
- (a) D. Baruah, P. Pahari and D. Konwar, *Tetrahedron Lett.*, 2015, **56**, 2418; (b) P. Natarajan, R. Chaudhary and P. Venugopalan, *J. Org. Chem.*, 2015, **80**, 10498.
- (a) V. V. Patil and G. S. Shankarling, *J. Org. Chem.*, 2015, **80**, 7876; (b) H. Firouzabadi, N. Iranpoor and K. Amani, *Green Chemistry*, 2001, **3**, 131; (c) S. Rozen and M. Carmeli, *J. Am. Chem. Soc.*, 2003, **125**, 8118; (d) M. Carmeli and S. Rozen, *J.*

Org. Chem., 2006, **71**, 4585; (e) K. R. Reddy, C. U. Maheswari and M. Venkateshwar, *Adv. Synth. Catal.*, 2009, **351**, 93.

7 (a) J. Yi, L. Yang, C.-G. Xia and F.-W. Li, *J. Org. Chem.*, 2015, **80**, 6213; (b) H.-W. Liang, K. Jiang, W. Ding, Y. Yuan, L. Shuai, Y.-C. Chen and Y. Wei, *Chem. Commun.*, 2015, **51**, 16928; (c) C.-C. Xia, Z.-J. Wei, C. Shen, J. Xu, Y. Yang, W.-K. Su and P.-F. Zhang, *RSC Adv.*, 2015, **5**, 52588; (d) L.-Z. Zhu, R.-H. Qiu, X. Cao, S. Xiao, X.-H. Xu, C.-T. Au and S.-F. Yin, *Org. Lett.*, 2015, **17**, 5528; (e) H.-J. Qiao, S.-Y. Sun, F. Yang, Y. Zhu, W.-G. Zhu, Y.-X. Dong, Y.-S. Wu, X.-T. Kong, L. Jiang and Y.-J. Wu, *Org. Lett.*, 2015, **17**, 6086; (f) Z. Wang, Y. Kuninobu and M. Kanai, *Org. Lett.*, 2014, **16**, 4790; (g) C. Shen, P.-F. Zhang, Q. Sun, S.-Q. Bai, T. S. A. Hor and X.-G. Liu, *Chem. Soc. Rev.*, 2011, **40**, 2761; (h) A. M. Suess, M. Z. Ertem, C. J. Cramer and S. S. Stahl, *J. Am. Chem. Soc.*, 2013, **135**, 9797; (i) G. G. Pawar, A. Brahmanandan and M. Kapur, *Org. Lett.*, 2016, **18**, 448.

8 L. Zhang, Z.-H. Liu, H.-Q. Li, G.-C. Fang, B. D. Barry, T. A. Belay, X.-H. Bi and Q. Liu, *Org. Lett.*, 2011, **13**, 6536.

9 W. Zhang, S.-J. Lou, Y.-K. Liu and Z.-Y. Xu, *J. Org. Chem.*, 2013, **78**, 5932.

10 Y.-F. Liang, X.-Y. Li, X.-Y. Wang, Y.-P. Yan, P. Feng and N. Jiao, *ACS Catal.*, 2015, **5**, 1956.

11 (a) Y.-K. Liu, S.-J. Lou, D.-Q. Xu, and Z.-Y. Xu, *Chem. -Eur. J.*, 2010, **16**, 13590; (b) D. Katayev, K. F. Pfister, T. Wendling and L. J. Gooben, *Chem. -Eur. J.*, 2014, **20**, 9902; (c) P. Sadhu, S. K. Alla and T. Punniyamurthy, *J. Org. Chem.*, 2015, **80**, 8245; (d) F. Xie, Z.-S. Qi and X.-W. Li, *Angew. Chem. Int. Ed.*, 2013, **52**, 11862; (e) W. Zhang, S.-B. Ren, J. Zhang and Y.-k. Liu, *J. Org. Chem.*, 2015, **80**, 5973.

12 E. Hernando, R. R. Castillo, N. Rodriguez, R. G. Arrayas and J. C. Carretero, *Chem. Eur. J.*, 2014, **20**, 13854.

13 C. J. Whiteoak, O. Planas, A. Company and X. Ribas, *Adv. Synth. Catal.*, 2016, **358**, 1679.

14 (a) J. Xu, X.-L. Zhu, G.-B. Zhou, B.-B. Ying, P.-P. Ye, L.-Y. Su, C. Shen and P.-F. Zhang, *Org. Biomol. Chem.*, 2016, **14**, 3016; (b) J. Xu, C. Shen, X.-L. Zhu, P.-F. Zhang, M. J. Ajitha, K.-W. Huang, Z.-F. An and X.-G. Liu, *Chem. Asian J.*, 2016, **11**, 882.

15 (a) K. Murugan, A. V. Raichurkar, F. R. N. Khan and P. S. Iyer, *Bioorg. Med. Chem. Lett.*, 2015, **25**, 1100; (b) H. Esaki, T. Hattori, A. Tsubone, S. Mibayashi, T. Sakata, Y. Sawama, Y. Monguchi, H. Yasuda, K. Nosaka and H. Sajiki, *ChemCatChem*, 2013, **5**, 3629.

16 (a) C. Manera, M. G. Cascio, V. Benetti, M. Allara, T. Tuccinardi, A. Martinelli, G. Saccomanni, E. Vivoli, C. Ghelardini, V. D. Marzob and P. L. Ferrarini, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 6505; (b) C.-H. Han, A. Chatterjee, M. J. Noetzel, J. D. Panarese, E. Smith, P. Chase, P. Hodder, C. Niswender, P. J. Conn, C. W. Lindsley and S. R. Stauffer, *Bioorg. Med. Chem. Lett.*, 2015, **25**, 384; (c) R. D. R. Reis, E. C. Azevedo, M. C. B. V. D. Souza, V. F. Ferreira, R. C. Montenegro, A. J. Araújo, C. Pessoa, L. V. C. Lotufo, M. O. D. Moraes, J. D. B. M. Filho, A. M. T. D. Souza, N. C. D. Carvalho, H. C. Castro, C. R. Rodrigues and T. R. A. Vasconcelos, *Eur. J. Med. Chem.*, 2011, **46**, 1448; (d) B. Medapi, P. Suryadevara, J. Renuka, J. P. Sridevi, P. Yogeeswari and D. Sriram, *Eur. J. Med. Chem.*, 2015, **103**, 1; (e) Y.-Q. Zhang, W. A. Guiguemde, M. Sigal, F.-Y. Zhu, M. C. Connelly, S. Nwaka and R. K. Guy, *Bioorg. Med. Chem.*, 2010,

18, 2756; (f) Y.-Y. Zhang, J. A. Clark, M. C. Connelly, F.-Y. Zhu, J. Min, W. A. Guiguemde, A. Pradhan, L. Iyer, A. Furimsky, J. Gow, T. Parman, F. E. Mazouni, M. A. Phillips, D. E. Kyle, J. Mirsalis and R. K. Guy, *J. Med. Chem.*, 2012, **55**, 4205; (g) N. Klein, I. Senkovska, I. A. Baburin, R. Grunker, U. Stoeck, M. Schlichtenmayer, B. Streppel, U. Mueller, S. Leoni, M. Hirscher and S. Kaskel, *Chem. -Eur. J.*, 2011, **17**, 13007.

17 (a) H. Tang, B. Zhou, X.-R. Huang, C. Wang, J. Yao and H. Chen, *ACS Catal.*, 2014, **4**, 649; (b) Y. Aihara and N. Chatani, *J. Am. Chem. Soc.*, 2014, **136**, 898.

Copper-catalyzed rapid C-H nitration of 8-aminoquinolines by using sodium nitrite as the nitro source under mild conditions

Xiaolei Zhu,^a Li Qiao,^a Pingping Ye,^a Beibei Ying,^a Jun Xu,^a Chao Shen^b and Pengfei Zhang^{*a}

We report the first example of copper(II) catalyzed remote C-H nitration of 8-aminoquinoline amides by using sodium nitrite as nitration reagent under mild conditions in 1 minute which undergoes single electron process.

