



**Environmental
Science**
Processes & Impacts

**PFAS soil and groundwater contamination via industrial
airborne emission and land deposition in SW Vermont and
Eastern New York State, USA**

Journal:	<i>Environmental Science: Processes & Impacts</i>
Manuscript ID	EM-ART-10-2020-000427.R1
Article Type:	Paper

SCHOLARONE™
Manuscripts

PFAS soil and groundwater contamination via industrial airborne emission and land deposition in SW Vermont and Eastern New York State, USA

Tim Schroeder¹ (tschroeder@bennington.edu)

David Bond¹ (dbond@bennington.edu)

Janet Foley¹ (jfoley@bennington.edu)

¹Bennington College, 1 College Drive, Bennington, VT 05201

Abstract

In order to understand the extent to which airborne PFAS emission can impact soil and groundwater, we conducted a sampling campaign in areas of conserved forest lands near Bennington, VT/Hoosick Falls, NY. This has been home to sources of PFAS air-emissions from Teflon-coating operations for over 50 years. Since 2015, the Vermont and New York Departments of Environmental Conservation have documented ~1,200 residential wells and two municipal water systems across a 200 km² area contaminated with perfluorooctanoic acid (PFOA). Given the large aerial extent of the plume, and the fact that much of the contaminated area lies up-gradient and across rivers from manufactures, we seek to determine if groundwater contamination could have resulted primarily from air-emission, land deposition, and subsequent leaching to infiltrating groundwater. Sampling of soils and groundwater in the Green Mountain National Forest (GMNF) downwind of factories shows that both soil and groundwater PFOA contamination extend uninterrupted from inhabited areas into conserved forest lands. Groundwater springs and seeps in the GMNF located 8 km downwind, but >300 meters vertically above factories, contain up to 100 ppt PFOA. Our results indicate that air-emitted PFAS can contaminate groundwater and soil in areas outside of those normally considered down-gradient of a source with respect to regional groundwater flow.

Environmental Significance Statement

Poly- and perfluoroalkyl substances (PFAS) are a class of widely used chemicals of emerging environmental concern. Most instances of PFAS contamination have resulted from direct land applications at industrial sites, locations of fire-fighting foam use, or biosolids application. In these cases, groundwater contamination typically migrates in the direction of regional groundwater flow. Where PFAS contamination results from airborne emission, contamination needs to be investigated in areas outside of those normally considered at-risk relative to the location of manufacturing facilities. In our study area, >200 km² of upland regions may have been contaminated by airborne PFAS emission.

Introduction

Poly- and perfluoroalkyl-substances (PFAS) are contaminants of emerging widespread concern. They have been used to apply non-stick, water repellent, and stain resistant coatings to a wide range of manufactured products since the 1950's.¹ PFAS have been found to be bioaccumulative, and have been linked by epidemiological and animal-based studies to a wide

1
2
3 range of health issues, including kidney and testicular cancers,^{2,3} ulcerative colitis,⁴ thyroid
4 disease,^{5,6} and immunological problems.⁷ PFAS are present in the blood serum of nearly all
5 people and animals.⁸⁻¹⁴ Blood serum levels in people are highly variable and dependent on
6 proximity to manufacturers, military bases, or other point sources.^{11,15} Primary exposure
7 pathways are through drinking water, eating contaminated food, food packaging, exposure to
8 other products manufactured using PFAS, and/or occupational exposure.¹⁶⁻²¹ PFAS are highly
9 persistent in the environment and have been found in soil, lakes, rivers, and oceans in all
10 corners of the globe, including polar regions.²²⁻³⁰ PFAS are not broken down by any known
11 natural process, and can be transported as anions in surface water or groundwater, through the
12 atmosphere as dust and/or aerosols, and may be mobilized as aerosols from sea spray.^{3,31} The
13 qualities of many PFAS - bioaccumulative, highly persistent, and mobile - pose new questions to
14 environmental science and regulation.
15
16
17

18 This study seeks to test the hypothesis that airborne PFAS emissions from manufactures in
19 southwest Vermont and eastern New York State contaminated soil and groundwater at
20 significant distances (>8 km) from emission sources. Air emission of PFAS has been shown to
21 have impacted water from manufactures in West Virginia^{32,33} and North Carolina³⁴. However, the
22 scale at which impacts can occur is debated. Early modelling and sampling studies indicated
23 limited PFAS deposition more than a few kilometers from the source^{32,35}, while more recent
24 sampling has suggested that air deposition can impact water >10 km from sources^{34,36}.
25 Addressing this question is difficult because of the pervasive nature of PFAS contamination and
26 the multiple pathways by which it can be introduced to the environment. Our study area is an
27 ideal case to address this question because of the large areas of conserved lands in the Green
28 Mountain National Forest downwind of emissions sources, within which point-source
29 contamination is very unlikely to have occurred. We collected soil and groundwater samples
30 from regions with specific spatial relations to air-emission sources and ran pairwise statistical
31 tests on data to examine spatial patterns of contamination and delineate any signature of an air-
32 deposition plume.
33
34
35
36
37

38 **Site History and Description**

39 The Bennington/Hoosick region's industrial history stretches back to the 1800's, when mills
40 were built in river valleys. The surrounding uplands were developed largely for agricultural use.
41 Several of the riverside industrial sites were repurposed as plastics manufacturers in the second
42 half of the 20th Century. The most significant of these are the McCaffrey Street factory in
43 Hoosick Falls, NY, which operated from 1956 to present, ChemFab in North Bennington, VT,
44 which operated from 1978 until 2002, and Taconic Plastics in Petersburg, NY, which has
45 operated from 1961 to present. All three factories utilized dispersions containing ammonium
46 perfluorooctanoate, which was vaporized during a baking step at ~300°C and released from
47 smokestacks as PFOA and other PFAS. Many of the region's PFAS-contaminated wells are in
48 the uplands surrounding the factories in river valleys. Because the factories all lie in valley
49 bottoms (groundwater discharge points), it is extremely unlikely for contamination to migrate
50 from factory sites to uplands via groundwater flow.
51
52
53
54
55
56
57
58
59
60

1
2
3 PFOA contamination was discovered in the Hoosick Falls village water supply wells in 2015 at
4 concentrations ranging from 400 to 600 ppt. Subsequent investigations conducted by the New
5 York and Vermont Departments of Environmental Conservation have now discovered PFOA in
6 over 1200 residential wells over an area of >200 km² across the region (Fig. 1).
7

8 **Geologic Setting**

9
10 Bennington and Hoosick Falls both lie within river valleys to the west of the Green Mountains
11 (Figure 1). The valley bottoms are filled with variable-thickness alluvial and fluvial gravels, while
12 valley sides and hilltops are generally covered by variable-thickness (0-50 m thick) glacial till.³⁷
13 Bedrock beneath unconsolidated materials in the valleys and hills of Bennington/Hoosick region
14 is dominantly a mixture of dolomitic limestone, phyllite, with minor quartzite. All bedrock
15 formations are highly fractured, faulted, and folded.³⁸ Thus bedrock permeability is highly
16 dependent on local structures, and is extremely heterogeneous and non-isotropic.³⁸ Bedrock
17 beneath the Green Mountains is highly deformed quartzite and gneiss, the permeability of which
18 is highly dependent on local fractures. Bedrock in the Green Mountains is covered by a very thin
19 (0-5 m thick) layer of sandy, rocky till.
20
21
22
23

24 Most private wells in the PFAS-impacted region are completed within the dolomite and/or
25 phyllite bedrock units. However, several private wells and the Village of Hoosick Falls public
26 wells are completed within the alluvial/fluvial gravel formations in valley bottoms. The degree of
27 PFAS contamination in wells is highly variable spatially, with uncontaminated wells occurring
28 within 200 meters of wells with over 1,000 ppt PFOA (Figure 1). Kim et al.³⁹ have hypothesized
29 that the variability is in part due to the fracture-controlled nature of groundwater flow in the
30 bedrock aquifer and possible groundwater recharge to some rock units occurring from distant
31 sources along very long flow paths.
32
33

34 **Methods**

35 *Study Design and Sampling*

36
37 The ubiquitous nature of PFAS contamination raises the question: how does one differentiate
38 locally air-emitted PFAS contamination from “baseline” PFAS levels caused by long-range air
39 transport, while also considering the possibility of undocumented point sources? This question
40 has significant legal importance in the Bennington/Hoosick Falls area, as regulators and courts
41 are assessing responsibility and liability for the water contamination. We designed a sampling
42 strategy to take advantage of the large areas of conserved forestland surrounding Bennington;
43 many of these lands, including the Green Mountain National Forest have been conserved since
44 prior to the first synthesis of PFAS. Prevailing wind patterns are dominantly west-to-east with a
45 secondary peak wind direction out of the southeast (Fig 1 inset). We divided our sample
46 collection sites into five regions relative to these wind patterns:
47
48
49
50

- 51 1. Bennington Local – Water sources and soil near PFAS industrial sites in Bennington;
52 hypothesized to be impacted by air emission
- 53 2. Downwind – samples from the Green Mountain National Forest directly east of emitters
54 in Bennington and Hoosick Falls; hypothesized to be impacted to by air emission
55
56
57

3. North of Main Wind Pattern – samples from the Green Mountain National Forest and other conserved forestlands that are north of the west-to-east wind pattern from the known emitters; hypothesized to be not impacted by air emission (i.e. background)
4. Local Upwind – samples from the Pittsfield State Forest, directly upwind of known emitters in Bennington and Hoosick Falls; hypothesized to be not impacted to by air emission (i.e. background)
5. Far Afield – samples collected from conserved forestlands more than 50 km distant from known industrial sites; hypothesized to be not impacted to by air emission (i.e. background)

We selected soils sampling sites in order to minimize the chance that any PFAS could have been applied directly to land by human activities. All sampling sites were either within the Green Mountain National Forest, New York DEC forest preserve or state forest land, or other protected tracts of private or state-owned land in VT and NY. Sample sites were accessed via foot and are located at least 200 meters from a road or 50 meters from a hiking trail. Sample locations were selected in clusters in order to achieve representative coverages from specific regions.

Soil samples were collected with a one-inch-diameter, 16-inch long, stainless steel push sampler. At each site, the sampler was driven to refusal depth (generally 12 to 16 inches) at four to eight locations within a five-square-meter area. The soil was then thoroughly homogenized in a stainless-steel bowl before being transferred into sample containers provided by Eurofins labs.

Groundwater seeps were sampled by driving a 0.5 cm diameter stainless steel mini-well into the center of the seep, then drawing water from the well with a 250 ml polyethylene syringe. The water was then decanted from the syringe into two 250 ml Trimza-treated sample bottles provided by Eurofins labs. Springs and surface water samples were collected from the source with a 500 ml polyethylene beaker, and then transferred to two 250 ml Trimza-treated sample bottles provided by Eurofins labs.

Full details of sampling and analysis procedures are included in ESI-1.

Data Analysis/Statistics

Interpretation of our analytical results hinges on determining if there is a statistically significant difference between PFAS concentrations in our five sampling regions. We conducted statistical analysis of PFOA and PFOS. Analysis results returned below detection limits were not included in statistical comparisons. Significant outliers were also excluded from statistical analysis so that sample groups would better approximate a normal distribution. For analytical results that were below the 95% confidence quantification limit, the estimated value from the lab is used in our statistical comparisons. In comparing PFOA and PFOA concentrations between sampling regions, we first ran a one-way ANOVA (analysis of variance) test to determine if there is a statistically significant difference between the mean values of any of the groups. If the ANOVA test rejected the null hypothesis, meaning that there is a statistically significant difference between groups, we proceeded to run a Tukey-HSD (Honestly Significant Difference) pairwise test, which compares sample-group means with a studentized range with a 95% confidence

1
2
3 interval for the entire sample set. All statistical analyses were conducted using the “Statsmodel”
4 package in Python. All code and output are included in ESI-3
5
6
7

8 **Data Quality**

9

10 *Duplicate Sampling:* Over the course of this study, we collected five duplicate water samples
11 and one duplicate soil sample in order to test the consistency of laboratory results. Duplicate
12 water samples were collected by simply filling additional sample bottles from the same source.
13 Duplicate soil samples were collected by filling additional samples bottles from the stainless-
14 steel bowl following homogenization. Full data from the duplicate sampling is included in
15 supplementary information ESI-1.
16
17

18 *Repeat Water Sampling:* Monitoring wells on the Bennington College campus and several
19 residential wells were sampled multiple times over this course of this investigation to study the
20 temporal variability of PFAS concentrations. The results also provide a reasonable test of the
21 quality of our sampling techniques and the consistency of laboratory results. Results were very
22 consistent between sampling events, with the PFOA concentration of water sources varying less
23 than 20% between most sampling events; see data and graphs in ESI-5.
24
25

26 *Sequence analysis of soil samples:* One concern with our study design is the possibility of
27 cross-contamination between soil samples because we used the same equipment at each
28 sample site. Though we carefully cleaned the equipment between each site, influence of prior
29 samples on each result is still a concern. To test for this effect, we plotted the PFOA and PFOS
30 concentrations in the sequence that they were collected with color codes for the sampling region
31 (See supplemental information ESI-2 Figure 1). Visual examination of these plots does not
32 suggest any influence of a prior sample on the result of subsequent samples. There is no
33 smearing effect apparent after high- or low-concentration samples, and the dominant influence
34 on the PFOA concentration is the region from which the sample was collected.
35
36
37

38 *Comparison between multiple soils studies:* We have compiled analysis data from additional
39 PFAS soil studies conducted in the Bennington, VT region to test that data collected in this
40 study is consistent with the results of other sampling and analysis techniques. These include:
41
42

- 43 1. Samples collected by the Vermont Dept. of Environmental Conservation (VT-DEC)
44 around the North Bennington area impacted by the ChemFab factory contamination
45 (<https://dec.vermont.gov/commissioners-office/pfoa>)
46
- 47 2. Samples collected for preparation of the *Draft Conceptual Site Model Site Investigation*
48 *Report* prepared by Barr Engineering on behalf of St. Gobain Performance Plastics;
49 samples collected around the Bennington region impacted by ChemFab
50 (<https://anrweb.vt.gov/PubDocs/DEC/PFOA/Conceptual%20Site%20Model%20Site%20Investigation/DRAFT-CSM-Site-Investigation-Report-text-only-FEB2018.pdf>)
51
52
- 53 3. Samples collected in a forested region of Bennington by a contractor for a solar
54 developer as part of the permitting process for a solar farm on the site (
55 <https://epuc.vermont.gov/?q=node/64/127312/FV-PFEXAFF-PTL>)
56
57

- 1
2
3 4. PFAS soil background study across Vermont commissioned by VT-DEC
4 ([https://anrweb.vt.gov/PubDocs/DEC/PFOA/Soil-Background/PFAS-Background-](https://anrweb.vt.gov/PubDocs/DEC/PFOA/Soil-Background/PFAS-Background-Vermont-Shallow-Soils-03-24-19.pdf)
5 [Vermont-Shallow-Soils-03-24-19.pdf](https://anrweb.vt.gov/PubDocs/DEC/PFOA/Soil-Background/PFAS-Background-Vermont-Shallow-Soils-03-24-19.pdf)).
6
7

8 The samples from these studies were collected in vertical profiles at varying depths at each site.
9 We have averaged results from the top 18 inches in each depth profile in order to provide the
10 most direct comparison to the results of our sampling technique. We have also divided our
11 samples into sets of the impacted areas (Bennington Local and Downwind regions), and the
12 sampling regions that we hypothesize to be not impacted by local air deposition (North of Wind
13 Pattern, Upwind, and Far-Afield).
14

15 With the exception of the Barr Site Investigation⁴⁰, our data is consistent with these other
16 investigations. A Tukey-Kramer pairwise test between the sample groups (See ESI-2 Figs. 4
17 and 5) shows that dry soil PFOA concentrations from impacted areas of our study are not
18 significantly different from those of the initial VT-DEC samples or the sampling done on the
19 proposed Apple-Hill solar farm, but they are significantly different those of the Barr Site
20 Investigation. The Barr samples are also significantly lower than those of the VT-DEC and
21 Apple-Hill studies. Dry soil PFOA concentrations in the peripheral (non-impacted) samples of
22 our study are not significantly different from those in the Vermont Soil Background study, or
23 those of the Barr Site Investigation. Dry soil perfluorooctane-sulfonate (PFOS) concentrations
24 were not significantly different between any of the studies.
25
26
27
28

29 The lower dry soil PFOA concentrations from the Barr Site Investigation relative to those of our
30 study and the other two studies may be due to sampling techniques and/or the land cover of the
31 sampling sites. Our study, the VT-DEC samples, and the Solar-Farm samples were collected
32 with hand-operated soil probes, while the Barr samples were collected from boreholes during
33 drilling with a casing-advance rotary technique that involved significant amounts of water
34 pumped down the hole. The sample sites from the Barr Investigation were also almost entirely
35 in areas of developed land cover, with disturbed soils, while the other studies' sites were more
36 commonly located in forestland or grassland. These differences are further addressed in the
37 Discussion section below.
38
39

40 **Results/Discussion**

41 *PFAS Distribution in Surface Soils*

42
43
44 Soil sampling results (included in ESI-4) support our hypothesis that the Bennington Local and
45 Downwind regions were impacted by air deposition from local PFAS air emitters. Dry soil PFOA
46 concentrations mapped in Figures 2 and 3 and graphed in Figure 4 exhibit distinctly higher
47 concentrations in the Bennington Local and Downwind regions relative to the three peripheral
48 sampling regions. The mean values of soil PFOA concentration in the Local (5.6 ppt) and
49 Downwind (4.3 ppt) regions is higher than the mean soil PFOA in the other three sampling
50 regions (North = 1.33 ppb, Upwind = 1.16 ppb, Far = 0.62 ppb).
51
52

53
54 One-way ANOVA showed a statistically significant difference between soil dry PFOA
55 concentration in the five sampling areas; $F(4,60) = 7.383$, $p = 0.000038$. The Tukey-HSD
56
57

1
2
3 pairwise test indicates that the mean soil PFOA concentration in the Local and Downwind
4 regions is statistically significantly higher than that of the North of Wind Pattern and Far-Afield
5 regions, though not significantly higher than the Upwind region (See ESI-4 Fig. 12), even though
6 the mean PFOA concentration in the Upwind region is much lower than that of the North region.
7 We were only able to collect four samples in the Upwind area, and thus the confidence range
8 around the Upwind PFOA mean is much wider, which likely contributed to the failed test for
9 statistical difference.
10
11

12 The mean values of soil PFOS concentration (Fig. 4) in the Local (1.8 ppt) and Downwind (0.66
13 ppt) regions is not apparently different than the mean soil PFOS in the other three sampling
14 regions (North = 0.76 ppb, Upwind = 0.55 ppb, Far = 0.61 ppb). One-way ANOVA showed that
15 there is not a statistically significant difference between soil dry PFOS concentration in the five
16 sampling areas; $F(4,60) = 0.355$, $p = 0.839$. Because PFOS was claimed to have not been used
17 by local industries, this pattern further supports our hypothesis of soil impacts from local air
18 emitters.
19
20

21 PFHpA was detected in 74% of samples in the downwind area, but only 29% of samples from
22 other areas, and PFHxA was detected in 53% of the downwind samples, but only 10% of the
23 samples from the other regions.
24
25

26 Our Local and Downwind samples have an average PFOA:PFNA ratio of 21.9 and 22.8
27 respectively, while average ratios in the other three sampling areas are much lower; North
28 PFOA:PFNA = 5.8, Upwind PFOA:PFNA = 6.2, and Far Afield PFOA:PFNA = 2.2 (Fig 2).
29 Globally, the PFAS air deposition on soils likely occurs via a combination of direct PFAS
30 deposition⁴¹, and long-range-transport of volatile precursor compounds, including fluorotelomer
31 alcohols (FTOH) followed by oxidation to PFAS in aquatic, atmospheric, or biologic media.⁴²⁻⁴⁵
32 Data compiled by Rankin et al.²⁵ from other sources suggests that that the range of FTOH
33 oxidation reactions should produce PFOA:PFNA ratios between 1:1 and 6:1, while direct air
34 deposition of PFOA may produce PFOA:PFNA ratios greater than 8:1 because PFOA was used
35 much more commonly in industrial processes than PFNA. The higher PFOA:PFNA ratio in the
36 Local and Downwind regions relative to other sampling regions further supports our hypothesis
37 that local industrial emissions source of direct air deposition in the region downwind of industrial
38 emitters in Bennington and Hoosick Falls.
39
40
41
42

43 An area of at least 200 km² has been contaminated to a level of between 5 and 6 ppb dry soil
44 PFOA by industrial emitters in the Bennington/Hoosick Falls area, which corresponds to roughly
45 1,000 kg of PFOA sequestered in soils. This likely represents only a small fraction of the total
46 PFOA used during the region's manufacturing history. Our results represent a minimum extent
47 of the plume; additional work is needed to fully delineate its full extent. The size of the
48 Bennington/Hoosick soil plume is roughly similar to the reported size of a soil and groundwater
49 PFOA plume near Merrimack, New Hampshire, which is where ChemFab relocated to from
50 North Bennington in 2002⁴⁶.
51
52
53
54

55 *Groundwater PFAS Distribution*

56
57
58
59
60

1
2
3 PFOA concentrations in water samples through the Bennington valley (ESI-5) are highly
4 variable, ranging from non-detect to 3100 ppt. Both the range and variability are consistent with
5 water sampling of domestic wells performed by VT-DEC (Fig 1). Groundwater samples in the
6 Downwind region of the Green Mountain National Forest ranged from non-detect in several
7 springs to 100 ppt from a groundwater seep located ~6.5 km from ChemFab (Figure 3). Several
8 other groundwater seeps and springs located between 6 and 8 km downwind of ChemFab
9 ranged in concentration from 6 to 41 ppt. This area on the west-facing slope of the Green
10 Mountains contains no roads or human habitation, and is 200 to 800 vertical meters above (up-
11 gradient) known PFOA emission sources. Thus, the only reasonable explanation for water
12 contamination there is PFOA leaching from soil that was contaminated by air deposition.
13
14
15

16 A cluster of water samples collected from the North-of-Wind-Pattern region of the Green
17 Mountain National Forest was below detection limits for PFOA and all other PFAS analytes.
18 This is consistent with lower soil PFOA levels detected in this area, and thus provides a control
19 that further supports the hypothesis that groundwater contamination in the Downwind region is a
20 result of air deposition from the local industries.
21
22

23 PFOA water contamination appears to extend less far (~8 km) from the air emitters than the
24 anomalously high soil concentrations (Figures 3 and 5). We detected PFOA in several springs
25 and seeps beyond the westernmost ridge of the Green Mountains, but the values are relatively
26 low. The extent and scope of air-deposited PFAS in our study appears to be less than that
27 documented by Galloway et al.³⁶ from the Washington Works plant in West Virginia. These
28 investigators found PFOA in surface waters >10 ppt at distances up to 30 km from the emission
29 source. The primary difference between these sites may be in the mass of PFOA vapors
30 emitted per year. The Washington Works factory was DuPont's primary Teflon production
31 center; the Bennington/Hoosick manufacturers are relatively small by comparison.
32
33
34

35 *Groundwater Contamination via Airborne PFAS deposition*

36
37 Drinking water is an important human exposure pathway for PFAS^{21,47}, thus we are primarily
38 concerned with the transference of air-deposited PFAS from soil to groundwater. While our
39 results indicate that air-emitted PFAS can impact groundwater, a fundamental question remains
40 for environmental professionals: what soil concentration is necessary for PFOA to begin
41 leaching from soil to infiltrating groundwater? Our dataset does not offer a clear-cut answer to
42 this question. The degree of soil PFOA contamination in the Green Mountain National Forest is
43 on the same order as that in the Bennington Valley. However, groundwater PFOA
44 concentrations fall to non-detect values with greater distance east in the mountains where the
45 soil PFOA concentration is still in the 4-6 ppb range (Figures 4 & 5).
46
47
48

49 Insights may lie in the nature of the soils themselves. The undisturbed forest soils in the Green
50 Mountains have significantly higher organic carbon content than the human-disturbed soils in
51 the Bennington Valley (see data in ESI-4). Experimental studies suggest that higher organic
52 carbon content causes higher PFAS retention in soils via electrostatic interactions.⁴⁸⁻⁵⁵ Miao et
53 al.⁵³ and Li et al.⁵⁵ both demonstrate positive, though not particularly strong, correlations
54 between organic carbon content and PFAS soil retention, with the effect being strongest for
55 longer chain PFAS. Some of the highest PFOA soil concentrations found in our study come
56
57
58
59
60

1
2
3 from very organic-rich soils on the west slope of the Green Mountains, ~6.5 km to the east and
4 1,000 vertical feet above the ChemFab factory. This includes significant outlier soil PFOA
5 concentrations of 23 ppb, collected in 2018, and 96 ppb when we resampled the same site in
6 2019. When soil and water PFOA level are plotted vs. distance from ChemFab (Fig 5a), it is
7 apparent that water PFOA concentration decreases dramatically with distance from the factory,
8 while soil PFOA remains high to much larger distances (into the Green Mountains). If soil PFOA
9 is normalized to total organic carbon (Fig 5b), a more regular linear relationship between
10 distance and concentration emerges.
11
12

13
14 Soil pH is also a possible factor in PFAS retention, with adsorption decreasing at higher pH due
15 to the surface charge on soils becoming less positive under alkaline conditions.^{55,56} These
16 laboratory results are consistent with our observation of high PFOA retention in the Green
17 Mountain soils, which are above quartzo-feldspathic bedrock (low pH), while the valley soils are
18 above carbonate bedrock (high pH).
19

20
21 The type of land cover/use may also have an effect on retention vs. leaching of PFAS from soil.
22 In order to examine this relationship, we intersected soil sample location points from our study
23 and the three other soil studies conducted in the Bennington area with the 2016 National Land
24 Cover Dataset (NLCD), and binned the NLCD categories into “Forest”, “Grassland/Pasture”, and
25 “Developed”, the last of which includes residential, commercial, and tilled agricultural land (Fig
26 6). Soils in developed land areas are generally low in organic carbon, and may be subject to
27 regular disturbance (i.e., tilling & mowing), which could increase the tendency for them to leach
28 rather than retain PFAS. The results of our sampling and the other three studies in the
29 Bennington area show highest PFOA in areas of forest cover, and lowest PFOA in developed
30 land soils, with grassland/pasture being intermediate between the two. A Tukey-HSD pairwise
31 test of these land use groups indicates that soils in forested land have statistically significant
32 higher PFOA concentrations than those on developed land (See SEI-3 Figs 9 and 10).
33
34
35

36
37 Taken together, our observations indicate that multiple factors need to be considered when
38 assessing the potential for PFAS to leach from soil to groundwater. It is likely that soil PFOA
39 levels were higher in the Bennington valley while the ChemFab factory was active, and that a
40 portion of that soil PFOA has subsequently leached to groundwater. In the Green Mountains
41 and other forested areas, a higher fraction of the PFOA deposited on soils has been retained
42 rather than leached.
43

44 *Baseline soil PFAS contamination in the northeastern United States*

45

46
47 Soil results from our sampling sites that are not directly downwind of the Bennington/Hoosick
48 PFAS emitters may be indicative of baseline level of soil PFAS contamination in the northeast
49 United States caused by long-distance air transport of PFAS and precursor chemicals. The
50 mean values of major PFAS compounds from these sites are PFOA = 1.12 ppb, PFOS = 0.61
51 ppb. The mean values of our sites that are more remote from the Bennington/Hoosick emitters
52 are PFOA = 0.62 ppb, PFOS = 0.53 ppb.
53

54
55 There are only a few published analyses with which to compare our soil results. Rankin et al.²⁵
56 analyzed PFAS soil levels in sites globally. The nearest soil sample site to the
57
58
59
60

1
2
3 Bennington/Hoosick area this study is from Holderness, NH, which contained 1.2 ppb PFOA.
4 This is broadly similar to many of our background results (average 1.12 ppb), but the PFOS
5 concentration in this sample (1.8 ppb) is considerably higher than our downwind and
6 background results. Other results reported by Rankin et al. for sites near Cortland, NY and
7 Penns Grove, NJ are generally similar to the Holderness site. Mejia-Avendano et al.⁵⁷ reported
8 on soils in Lac Megantic, Quebec that were contaminated with PFAS by AFFF application. This
9 study included analyses of background soils away from the fire-fighting sites: PFOA = 0.250
10 ppb, and PFOS = 0.160 ppb. This area of southern Quebec is somewhat more remote from
11 industrial regions of the northeastern US, which could account for the lesser contaminated soils
12 there. Discrepancies in baseline levels between these studies could also be due to variations in
13 local soil conditions (i.e. sorption capacity), local/regional contamination sources, or analytical
14 techniques.
15
16
17

18 **Conclusions and Implications**

19
20 PFAS soil sampling and analysis reveal a statistically significant enrichment of PFOA in areas
21 local to, and downwind of factories that emitted PFAS to the atmosphere during baking of wet
22 dispersions. Our results are further supported by higher PFOA:PFNA ratios in the local and
23 downwind regions relative to peripheral areas. The presence of elevated PFOA in groundwater
24 seeps and springs within the Green Mountain National Forest in the downwind region supports
25 our hypothesis that air-emitted PFAS can cause groundwater contamination over large areas
26 downwind of emission sites. Environmental investigations related to such manufacturing
27 facilities need to expand in scope to consider regions downwind of emission sites, rather than
28 being limited to areas in the down-gradient direction of groundwater flow.
29
30
31

32 **Conflicts of Interest**

33
34 This research was supported by National Science Foundation grants #1711637 and #1634237.
35 The authors have no conflicts of interest in this work.
36
37

38 **Acknowledgements**

39
40 The authors would like to thank the Green Mountain National Forest, Merck Forest and
41 Farmland Center, Smokey House Preserve, and New York State Forest Preserve for allowing
42 soil and water samples to be collected from their properties. The authors would also like to
43 thank the Vermont Department of Environmental Conservation for helpful collaboration during
44 this project.
45
46
47

48 **Figures and Captions**

49
50 **Figure 1.** Map of PFAS-impacted region surrounding Bennington, VT and Hoosick Falls, NY.
51 Locations and ranges of groundwater PFOA concentrations are indicated for samples analyzed
52 for this study, along with analyses compiled from the Vermont Department of Environmental
53 Conservation and Site Investigation Reports prepared for St. Gobain Performance Plastics. Few
54 analytical results are publically available on the New York side of the plume because of privacy
55
56
57

concerns raised by New York state government. The boundary of the plume within New York is estimated from general area maps provided by the NY Department of Environmental Conservation. In all, approximately 1,200 private and public wells have been contaminated over an area exceeding 200 km². Inset Wind Rose diagram is based on daily two-minute maximum wind direction recorded at the Morse State Airport in Bennington for the period January 1 1998 through July 31 2018. This data includes a daily maximum two-minute average wind speed and direction (7162 records). This two-minute average maximum wind is taken to represent the wind during any given day that is most likely to transport PFAS aerosols from points of emission. The wind speed and directions were analyzed with the “Openair” package within the “R” environment.³⁸ The dominant aerosol transport direction should be from west to east, with some secondary transport from south-southeast to north-northwest.




Figure 2. Map showing all dry soil PFOA concentrations selected water concentrations analyzed for this study. Sample sites are divided into regions in relation to manufacturers that emitted airborne PFOA in Bennington and Hoosick Falls: 1) Bennington Local, 2) Downwind, 3) North of Wind Pattern, 4) Upwind, and 5) Far-afield. All samples were collected from forest landscapes on conserved lands with minimal human disturbance.




Figure 3. Detail map and profile section of soil and water PFOA concentrations in the downwind-of-emitters sampling areas within the Green Mountain National Forest. Line of profile on map is shown in yellow. The profile shows elevated soil and water PFOA levels in regions up-gradient, though down-wind, of possible PFOA point sources.




Figure 4. Box plots of dry soil PFOA and PFOS concentrations; subdivided by the sampling regions delineated in Figure 2. Soil PFOA concentrations are distinctly higher in the downwind from emitters area than in the other three sample regions, while the soil PFOS concentrations are broadly similar across all sample regions.




Figure 5. Graphs of PFOA water and soil concentrations vs. sample location distance from ChemFab source of air emission. Water concentration from this study are plotted as triangles and those from VT-DEC domestic well sampling are plotted as "X's". Soil PFOA concentration on the right graph have been normalized to Total Organic Carbon, which shows a more pronounced decrease with distance from ChemFab.





Figure 6: Box-plot and jitter-plot of soil PFOA concentrations in the Bennington region divided by the land cover/use for each sampling point. Forest soils contain more PFOA than Grasslands or areas of developed/tilled land, indicating higher PFAS soil-retention and or air-deposition in forested regions.



References Cited

- 1 Z. Wang, J. C. Dewitt, C. P. Higgins and I. T. Cousins, A Never-Ending Story of Per- and Polyfluoroalkyl Substances (PFASs)?, *Environ. Sci. Technol.*, 2017, **51**, 2508–2518.
- 2 V. Barry, A. Winqvist and K. Steenland, Perfluorooctanoic Acid (PFOA) Exposures and Incident Cancers among Adults Living Near a Chemical Plant, *Environ. Health Perspect.*,

- 2013, **121**, 1313–8.
- 3 W. Nicole, PFOA and cancer in a highly exposed community: new findings from the C8
4 science panel., *Environ. Health Perspect.*, 2013, **121**, A340.
- 5
6 4 K. Steenland, L. Zhao, A. Winquist and C. Parks, Ulcerative Colitis and Perfluorooctanoic
7 Acid (PFOA) in a Highly Exposed Population of Community Residents and Workers in the
8 Mid-Ohio Valley, *Environ. Health Perspect.*, 2013, **121**, 900–905.
- 9
10 5 A. Winquist and K. Steenland, Perfluorooctanoic Acid Exposure and Thyroid Disease in
11 Community and Worker Cohorts, *Epidemiology*, 2014, **25**, 255–264.
- 12 6 S. Shrestha, M. S. Bloom, R. Yucel, R. F. Seegal, Q. Wu, K. Kannan, R. Rej and E. F.
13 Fitzgerald, Perfluoroalkyl substances and thyroid function in older adults, *Environ. Int.*,
14 2015, **75**, 206–214.
- 15 7 B. Granum, L. S. Haug, E. Namork, S. B. Stølevik, C. Thomsen, I. S. Aaberge, H. van
16 Loveren, M. Løvik and U. C. Nygaard, Pre-natal exposure to perfluoroalkyl substances
17 may be associated with altered vaccine antibody levels and immune-related health
18 outcomes in early childhood, *J. Immunotoxicol.*, 2013, **10**, 373–379.
- 19 8 K. Kato, L.-Y. Wong, L. T. Jia, Z. Kuklennyik and A. M. Calafat, Trends in Exposure to
20 Polyfluoroalkyl Chemicals in the U.S. Population: 1999–2008 †, *Environ. Sci. Technol.*,
21 2011, **45**, 8037–8045.
- 22 9 L. W. Y. Yeung and S. A. Mabury, Are humans exposed to increasing amounts of
23 unidentified organofluorine?, *Environ. Chem.*, 2016, **13**, 102–110.
- 24 10 E. Sinclair, D. T. Mayack, K. Roblee, N. Yamashita and K. Kannan, Occurrence of
25 perfluoroalkyl surfactants in water, fish, and birds from New York State, *Arch. Environ.*
26 *Contam. Toxicol.*, 2006, **50**, 398–410.
- 27 11 Y. Gao, J. Fu, H. Cao, Y. Wang, A. Zhang, Y. Liang, T. Wang, C. Zhao and G. Jiang,
28 Differential Accumulation and Elimination Behavior of Perfluoroalkyl Acid Isomers in
29 Occupational Workers in a Manufactory in China, *Environ. Sci. Technol.*, 2015, **49**, 6953–
30 6962.
- 31 12 C. E. Müller, A. O. De Silva, J. Small, M. Williamson, X. Wang, A. Morris, S. Katz, M.
32 Gamberg and D. C. G. Muir, Biomagnification of Perfluorinated Compounds in a Remote
33 Terrestrial Food Chain: Lichen–Caribou–Wolf, *Environ. Sci. Technol.*, 2011, **45**, 8665–
34 8673.
- 35 13 J. Bytingsvik, S. P. J. van Leeuwen, T. Hamers, K. Swart, J. Aars, E. Lie, E. M. E. Nilsen,
36 Ø. Wiig, A. E. Derocher and B. M. Jenssen, Perfluoroalkyl substances in polar bear
37 mother–cub pairs: A comparative study based on plasma levels from 1998 and 2008,
38 *Environ. Int.*, 2012, **49**, 92–99.
- 39 14 E. Zafeiraki, D. Costopoulou, I. Vassiliadou, L. Leondiadis, E. Dassenakis, R. L. A. P.
40 Hoogenboom and S. P. J. van Leeuwen, Perfluoroalkylated substances (PFASs) in home
41 and commercially produced chicken eggs from the Netherlands and Greece,
42 *Chemosphere*, 2016, **144**, 2106–2112.
- 43 15 G. W. Olsen, J. M. Burris, D. J. Ehresman, J. W. Froehlich, A. M. Seacat, J. L. Butenhoff
44 and L. R. Zobel, Half-life of serum elimination of
45 perfluorooctanesulfonate, perfluorohexanesulfonate, and perfluorooctanoate in retired
46 fluorochemical production workers., *Environ. Health Perspect.*, 2007, **115**, 1298–305.
- 47 16 S. A. Tittlemier, K. Pepper, C. Seymour, J. Moisey, R. Bronson, X. L. Cao and R. W.
48 Dabeka, Dietary exposure of Canadians to perfluorinated carboxylates and
49 perfluorooctane sulfonate via consumption of meat, fish, fast foods, and food items
50 prepared in their packaging, *J. Agric. Food Chem.*, 2007, **55**, 3203–3210.
- 51 17 K. Hoffman, T. F. Webster, S. M. Bartell, M. G. Weisskopf, T. Fletcher and V. M. Vieira,
52 Private drinking water wells as a source of exposure to perfluorooctanoic acid (PFOA) in
53 communities surrounding a fluoropolymer production facility., *Environ. Health Perspect.*,
54 2011, **119**, 92–7.
- 55
56
57
58
59
60

- 1
2
3 18 R. R. Worley, X. Yang and J. Fisher, Physiologically based pharmacokinetic modeling of
4 human exposure to perfluorooctanoic acid suggests historical non drinking-water
5 exposures are important for predicting current serum concentrations, *Toxicol. Appl.*
6 *Pharmacol.*, 2017, **330**, 9–21.
- 7 19 U. Eriksson and A. Kärrman, World-Wide Indoor Exposure to Polyfluoroalkyl Phosphate
8 Esters (PAPs) and other PFASs in Household Dust, *Environ. Sci. Technol.*, 2015, **49**,
9 14503–14511.
- 10 20 L. A. Schaider, S. A. Balan, A. Blum, D. Q. Andrews, M. J. Strynar, M. E. Dickinson, D. M.
11 Lunderberg, J. R. Lang and G. F. Peaslee, Fluorinated Compounds in U.S. Fast Food
12 Packaging, *Environ. Sci. Technol. Lett.*, 2017, **4**, 105–111.
- 13 21 X. C. Hu, C. Dassuncao, X. Zhang, P. Grandjean, P. Weihe, G. M. Webster, F. Nielsen
14 and E. M. Sunderland, Can profiles of poly- and Perfluoroalkyl substances (PFASs) in
15 human serum provide information on major exposure sources?, *Environ. Heal. A Glob.*
16 *Access Sci. Source*, 2018, **17**, 11.
- 17 22 B. Boulanger, J. Vargo, J. L. Schnoor and K. C. Hornbuckle, Detection of perfluorooctane
18 surfactants in Great Lakes water, *Environ. Sci. Technol.*, 2004, **38**, 4064–4070.
- 19 23 J. Armitage, I. T. Cousins, R. C. Buck, K. Prevedouros, M. H. Russell, M. MacLeod and
20 S. H. Korzeniowski, Modeling Global-Scale Fate and Transport of Perfluorooctanoate
21 Emitted from Direct Sources, *Environ. Sci. Technol.*, 2006, **40**, 6969–6975.
- 22 24 C. M. Butt, U. Berger, R. Bossi and G. T. Tomy, Levels and trends of poly- and
23 perfluorinated compounds in the arctic environment, *Sci. Total Environ.*, 2010, **408**,
24 2936–2965.
- 25 25 K. Rankin, S. A. Mabury, T. M. Jenkins and J. W. Washington, A North American and
26 global survey of perfluoroalkyl substances in surface soils: Distribution patterns and mode
27 of occurrence, *Chemosphere*, 2016, **161**, 333–341.
- 28 26 P. Casal, Y. Zhang, J. W. Martin, M. Pizarro, B. Jiménez and J. Dachs, Role of Snow
29 Deposition of Perfluoroalkylated Substances at Coastal Livingston Island (Maritime
30 Antarctica), *Environ. Sci. Technol.*, 2017, **51**, 8460–8470.
- 31 27 G. Sammut, E. Sinagra, R. Helmus and P. de Voogt, Perfluoroalkyl substances in the
32 Maltese environment – (I) surface water and rain water, *Sci. Total Environ.*, 2017, **589**,
33 182–190.
- 34 28 J. Meng, T. Wang, S. Song, P. Wang, Q. Li, Y. Zhou and Y. Lu, Tracing perfluoroalkyl
35 substances (PFASs) in soils along the urbanizing coastal area of Bohai and Yellow Seas,
36 China, *Environ. Pollut.*, 2018, **238**, 404–412.
- 37 29 U. K. Vedagiri, R. H. Anderson, H. M. Loso and C. M. Schwach, Ambient levels of PFOS
38 and PFOA in multiple environmental media, *Remediation*, 2018, **28**, 9–51.
- 39 30 J. H. Johansson, Y. Shi, M. Salter and I. T. Cousins, Spatial variation in the atmospheric
40 deposition of perfluoroalkyl acids: source elucidation through analysis of isomer patterns,
41 *Environ. Sci. Process. Impacts*, 2018, **20**, 997–1006.
- 42 31 F. Wania*, A Global Mass Balance Analysis of the Source of Perfluorocarboxylic Acids in
43 the Arctic Ocean, , DOI:10.1021/ES070124C.
- 44 32 C. A. Barton, L. E. Butler, C. J. Zarzecki, J. Flaherty and M. Kaiser, Characterizing
45 perfluorooctanoate in ambient air near the fence line of a manufacturing facility:
46 comparing modeled and monitored values., *J. Air Waste Manag. Assoc.*, 2006, **56**, 48–
47 55.
- 48 33 K. L. Davis, M. D. Aucoin, B. S. Larsen, M. A. Kaiser and A. S. Hartten, Transport of
49 ammonium perfluorooctanoate in environmental media near a fluoropolymer
50 manufacturing facility, *Chemosphere*, 2007, **67**, 2011–2019.
- 51 34 M. Sun, E. Arevalo, M. Strynar, A. Lindstrom, M. Richardson, B. Kearns, A. Pickett, C.
52 Smith and D. R. U. Knappe, Legacy and Emerging Perfluoroalkyl Substances Are
53 Important Drinking Water Contaminants in the Cape Fear River Watershed of North
54
55
56
57
58
59
60

- 1
2
3 Carolina, *Environ. Sci. Technol. Lett.*, 2016, **3**, 415–419.
- 4 35 C. A. Barton, C. J. Zarzecki and M. H. Russell, A site-specific screening comparison of
5 modeled and monitored air dispersion and deposition for perfluorooctanoate, *J. Air Waste*
6 *Manag. Assoc.*, 2010, **60**, 402–411.
- 7 36 J. E. Galloway, A. V. P. Moreno, A. B. Lindstrom, M. J. Strynar, S. Newton, A. A. May, A.
8 A. May, L. K. Weavers and L. K. Weavers, Evidence of Air Dispersion: HFPO-DA and
9 PFOA in Ohio and West Virginia Surface Water and Soil near a Fluoropolymer
10 Production Facility, *Environ. Sci. Technol.*, 2020, **54**, 7175–7184.
- 11 37 D. J. DeSimone, *No Title Surficial Geology of the Bennington Area, Vermont: VGS Open*
12 *File report VG2017-1*, 2017.
- 13 38 M. Kim, J.J., Ryan, P., Schroeder, T., Romanowicz, E., Boutt, D., Belaval, in *Geological*
14 *Society of America Abstracts with Programs. Vol. 51, No. 1*, Geological Society of
15 America, 2019.
- 16 39 M. Kim, J.J., Ryan, P., Schroeder, T., Romanowicz, E., Boutt, D., Belaval, in *Geological*
17 *Society of America Abstracts with Programs*, Geological Society of America], 2019.
- 18 40 B. Engineering, *Conceptual Site Model Site Investigation Report: Bennington VT March*
19 *2018*, 2018.
- 20 41 K. Prevedouros, I. T. Cousins, R. C. Buck and S. H. Korzeniowski, Sources, Fate and
21 Transport of Perfluorocarboxylates, *Environ. Sci. Technol.*, 2006, **40**, 32–44.
- 22 42 E. Webster and D. A. Ellis, *Environ. Toxicol. Chem.*, 2010, **29**, 1703–1708.
- 23 43 D. A. Ellis, J. W. Martin, A. O. De Silva, S. A. Mabury, M. D. Hurley, M. P. Sulbaek
24 Andersen and T. J. Wallington, Degradation of Fluorotelomer Alcohols: A Likely
25 Atmospheric Source of Perfluorinated Carboxylic Acids, *Environ. Sci. Technol.*, 2004, **38**,
26 3316–3321.
- 27 44 B. F. Scott, C. Spencer, S. A. Mabury and D. C. G. Muir, Poly and Perfluorinated
28 Carboxylates in North American Precipitation †, *Environ. Sci. Technol.*, 2006, **40**, 7167–
29 7174.
- 30 45 J. L. Barber, U. Berger, C. Chaemfa, S. Huber, A. Jahnke, C. Temme and K. C. Jones,
31 Analysis of per- and polyfluorinated alkyl substances in air samples from Northwest
32 Europe, *J. Environ. Monit.*, 2007, **9**, 530.
- 33 46 State of New Hampshire, *Consent Decree*, 2018.
- 34 47 F. Xiao, M. F. Simcik, T. R. Halbach and J. S. Gulliver, Perfluorooctane sulfonate (PFOS)
35 and perfluorooctanoate (PFOA) in soils and groundwater of a U.S. metropolitan area:
36 Migration and implications for human exposure, *Water Res.*, 2015, **72**, 64–74.
- 37 48 C. P. Higgins and R. G. Luthy, Sorption of perfluorinated surfactants on sediments,
38 *Environ. Sci. Technol.*, 2006, **40**, 7251–7256.
- 39 49 L. Ahrens, L. W. Y. Yeung, S. Taniyasu, P. K. S. Lam and N. Yamashita, Partitioning of
40 perfluorooctanoate (PFOA), perfluorooctane sulfonate (PFOS) and perfluorooctane
41 sulfonamide (PFOSA) between water and sediment, *Chemosphere*, 2011, **85**, 731–737.
- 42 50 J. G. Sepulvado, A. C. Blaine, L. S. Hundal and C. P. Higgins, Occurrence and fate of
43 perfluorochemicals in soil following the land application of municipal biosolids, *Environ.*
44 *Sci. Technol.*, 2011, **45**, 8106–8112.
- 45 51 J. L. Guelfo and C. P. Higgins, Subsurface Transport Potential of Perfluoroalkyl Acids at
46 Aqueous Film-Forming Foam (AFFF)-Impacted Sites, *Environ. Sci. Technol.*, 2013, **47**,
47 4164–4171.
- 48 52 J. Milinovic, S. Lacorte, M. Vidal and A. Rigol, Sorption behaviour of perfluoroalkyl
49 substances in soils, *Sci. Total Environ.*, 2015, **511**, 63–71.
- 50 53 Y. Miao, X. Guo, Dan Peng, T. Fan and C. Yang, Rates and equilibria of
51 perfluorooctanoate (PFOA) sorption on soils from different regions of China, *Ecotoxicol.*
52 *Environ. Saf.*, 2017, **139**, 102–108.
- 53 54 Q. Wang, Z. Zhao, Y. Ruan, J. Li, H. Sun and G. Zhang, Occurrence and distribution of
54

- 1
2
3 perfluorooctanoic acid (PFOA) and perfluorooctanesulfonic acid (PFOS) in natural forest
4 soils: A nationwide study in China, *Sci. Total Environ.*, 2018, **645**, 596–602.
- 5 55 Y. Li, D. P. Oliver and R. S. Kookana, A critical analysis of published data to discern the
6 role of soil and sediment properties in determining sorption of per and polyfluoroalkyl
7 substances (PFASs), *Sci. Total Environ.*, 2018, **628–629**, 110–120.
- 8 56 Z. Du, S. Deng, Y. Bei, Q. Huang, B. Wang, J. Huang and G. Yu, Adsorption behavior
9 and mechanism of perfluorinated compounds on various adsorbents—A review, *J.*
10 *Hazard. Mater.*, 2014, **274**, 443–454.
- 11 57 S. Mejia-Avenidaño, G. Munoz, S. Vo Duy, M. Desrosiers, P. Benoît, S. Sauvé and J. Liu,
12 Novel Fluoroalkylated Surfactants in Soils Following Firefighting Foam Deployment
13 During the Lac-Mégantic Railway Accident, *Environ. Sci. Technol.*, 2017, **51**, 8313–8323.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

We document soil and groundwater contamination caused by industrial airborne PFAS emission over an area of ~200 km² that has impacted over 1,200 wells.

