Organic & Biomolecular Chemistry

rsc.li/obc



ISSN 1477-0520



PAPERA. Szumna et al.The templation effect as a driving force for the self-assembly ofhydrogen-bonded peptidic capsules in competitive media

Organic & Biomolecular Chemistry

PAPER

Check for updates

Cite this: Org. Biomol. Chem., 2017, **15**, 8513

Received 2nd August 2017, Accepted 24th August 2017 DOI: 10.1039/c7ob01925d

rsc.li/obc

The templation effect as a driving force for the self-assembly of hydrogen-bonded peptidic capsules in competitive media[†]

M. Grajda, 🕩 M. J. Lewińska 🕩 and A. Szumna 🕩 *

Peptide-based cavitands (resorcin[4]arenes substituted with histidine and glutamine hydrazides) exist as monomeric species in polar solvents (DMSO and methanol). Upon complexation of fullerenes, the cavitands wrap around the hydrophobic guests forming dimeric capsular shells (as evidenced by DOSY). The self-assembly of the cavitands is based on the formation of beta-sheet-like binding motifs around the hydrophobic core. In a polar environment, these hydrogen bonded structures are kinetically stable and highly ordered as manifested by a 100-fold increase of intensity of circular dichroism bands, as well as a separate set of signals and substantial differences in chemical shifts in NMR spectra. This behavior resembles a protein folding process at the molten globule stage with non-specific hydrophobic inter-actions creating a protective and favourable local environment for the formation of secondary structures of proteins.

Our work takes inspiration from the native structures of proteins, which are exemplars of excellence in folding of ordered three-dimensional architectures in polar environments. At the initial stages of protein folding a pivotal role is attributed to hydrophobic interactions. Although these interactions are nondirectional, they lead to the formation of a hydrophobic core, a molten globule that constitutes a perfect environment for further formation of directional hydrogen bonds by lowering the local dielectric constant.^{1–3} Thus, hydrogen bonds around a hydrophobic core are stronger and contribute more significantly to the stability of the native state than hydrogen bonds exposed to the aqueous environment.

This bioinspired approach based on a combination of hydrophobic core formation with an ordering power of hydrogen bonds on the peripheries is widely used for the self-assembly of infinite fibrous structures⁴ or nanoparticles.⁵ When the number of building blocks decreases, the strength of hydrophobic clustering is reduced and the above-mentioned strategy becomes less efficient. Therefore, the formation of discrete structures made of only a few building blocks is challenging. Gibb and co-workers reported on the self-assembly of discrete capsular dimers,⁶ tetramers and hexamers⁷ in water using hydrophobic interactions only.⁸ The group of Rebek developed a series of hydrocarbon-based templates for ordering of hydrogen bonded multicomponent capsules in non-polar solvents.⁹ Fullerenes have been used for the templation of self-assembled structures based on 4-pyrimidinone derivatives (isocytosine or 2-ureido-4[1*H*]-pyrimidinone derivatives),^{10–13} amino acid functionalized naphthalenediimides,¹⁴ *S*-trityl-cysteine derivatives¹⁵ or through pyridinium–anion–pyridinium interactions.¹⁶ We have also presented resorcinarenes appended to hydrophobic peptides (Phe, Val) that form well-ordered hydrogen bonded dimers and complex fullerenes (Fig. 1).¹⁷

In all these examples, hydrogen bonded shells are formed in non-polar media (chloroform, toluene, tetrachloroethane, carbon disulfide) and they are not able to survive in polar environments (even DMSO). In this work, we demonstrate that hydrophobic interactions together with β -sheet-like binding motifs are effective (a) in inducing the self-assembly of peptidic cavitands that otherwise do not self-assemble and (b) in ordering hydrogen bonded shells around hydrophobic cores even in competitive, polar environments (Fig. 1).

In our quest towards self-assembly in polar environments we have designed resorcin[4]arene cavitands substituted with hydrophilic peptides, here containing glutamine and histidine residues (**3a** and **3b**, Table 1). The side chains of these amino acids increase solubility in polar solvents and they additionally can take part in stabilizing or competitive (destabilizing) hydrogen bonding interactions due to their dual donor/acceptor character. The final thermodynamic outcome of such a complex system of interactions is highly context-dependent. For example, polyglutamine sequences in the cores of proteins

View Article Online

Institute of Organic Chemistry, Polish Academy of Sciences, Kasprzaka 44/52, 01-22 Warsaw, Poland, E-mail: agnieszka.szumna@icho.edu.pl

⁰¹⁻²² warsaw, Polana. E-mail: agnieszka.szumna@icno.eau.pi

[†]Electronic supplementary information (ESI) available: Experimental procedures and spectra. See DOI: 10.1039/c7ob01925d



Fig. 1 Comparison of self-assembly processes for peptidic capsules observed in a non-polar environment (previous work) and the discussed process in a polar environment (this work).

stabilize β -sheets and β -barrels through hydrogen bonding interactions between side chains.^{18–21} On the other hand glutamine present in the external shells of other proteins, *e.g.* insulin does not form stabilizing interactions.

We obtained cavitands 3a and 3b in the reaction of respective amino acid hydrazides 2a or 2b and tetraformylresorcin[4]arene 1 (Table 1). Hydrazones 3a and 3b are soluble in polar solvents: DMSO and methanol but not in water. ¹H NMR spectra in DMSO-d₆ reveal C_4 -symmetric structures for both products. Analysis of ROESY NMR spectra indicates the presence of only trivial contacts (originating from covalent proximity). The intramolecular f_{-j} correlation signals that are hallmarks for dimerization via backbone interactions are not present. There are also no indications of specific interactions involving sidechains' hydrogen bonding groups. These results suggest that hydrazones 3a and 3b do not self-assemble in DMSO and exist as monomeric semi-open cavitands (Fig. 1). This conclusion is also supported by DOSY measurements that reveal a diffusion coefficient of 0.95 \times 10^{-10} m^2 s^{-1} for 3a and 0.95 \times 10^{-10} m^2 s^{-1} for **3b** (in DMSO-d₆ at 298 K), corresponding to an average diameter of molecule of ~2.2 nm (predicted value for the cavitand is 2.3 nm, the model is based on the previously reported X-ray structure of the valine derivative).¹⁷ This self-assembly behaviour and solubility characteristics of cavitands containing polar side chains are distinctly different from those of the previously reported hydrophobic derivatives that were soluble only in non-polar solvents and formed dimeric capsules in these media.17

Table 1 Synthesis of cavitands **3a** and **3b**, C_{60} complexation strategies and condition optimization of $(3a)_2 \supset C_{60}$ synthesis



					Yield ^{<i>a</i>} of:	
Entry	Path	Substrates	Equiv.	Conditions	$(\mathbf{3a})_2 \supset \mathrm{C}_{60}$	3a
1	А	3a + C ₆₀	2:1	DMSO (24 h, rt)	8%	92%
2	Α	$3a + C_{60}$	2:1	DMSO	14%	86%
				(24 h, 80 °C)		
3	А	$3a + C_{60}$	2:1	Ball-mill (2 h)	18%	82%
4	А	$3a + C_{60}$	2:10	DMSO (24 h, rt)	12%	88%
5	А	$3a + C_{60}$	2:10	DMSO	15%	85%
				(24 h, 80 °C)		
6	А	$3a + C_{60}$	2:10	Ball-mill (2 h)	19%	81%
7	В	$1 + 2a + C_{60}$	2:8:1	DMSO	24%	86%
				(24 h, 80 °C)		
8	В	$1 + 2a + C_{60}$	2:8:1	Ball-mill (2 h)	21%	79%
9	В	$1 + 2a + C_{60}$	2:8:10	DMSO	42%	58%
				(24 h, 80 °C)		
10	В	$1 + \mathbf{2a} + \mathbf{C}_{60}$	2:8:10	Ball-mill (2 h)	22%	78%
" Yield	ls were	determined	by NMR.			

In subsequent experiments, we tested the possibility of templation of self-assembly by fullerene C_{60} . For cavitands **3a** and **3b** a direct complexation experiment is not feasible due to the lack of solubility of fullerene in polar solvents. Therefore alternative ways of reaction activation were employed: (1) thermal activation of samples (heating of **3a** + C₆₀ or **3b** + C₆₀ in DMSO at 70 °C for 7 days with vigorous stirring, path A); (2) complexation during the chemical reaction of cavitand synthesis (path B); and (3) mechanochemical activation²² in the solid state (Table 1).

In all cases, the resulting products contained two types of species as revealed by ¹H NMR spectra in DMSO-d₆ (Fig. 2b and e). One set of signals corresponds to the cavitands (**3a** or **3b**) and the second one is attributed to the dimeric species with encapsulated fullerenes ((**3a**)₂ \supset C₆₀ or (**3b**)₂ \supset C₆₀). This assignment is based on ¹³C NMR spectra that reveal the signal of encapsulated fullerene at 141 ppm and on DOSY

Paper



Fig. 2 ¹H NMR (600 MHz, DMSO-d₆) spectrum of **3a** (a) and a mixture of **3a** and (**3a**)₂ \supset C₆₀ (b); DOSY spectrum of **3a** and (**3a**)₂ \supset C₆₀ (c); ¹H NMR spectrum of **3b** (d) and a mixture of **3b** and (**3b**)₂ \supset C₆₀ (e); DOSY spectrum of **3b** and (**3b**)₂ \supset C₆₀ (f).

experiments that clearly indicate a larger size of the newly formed species (Fig. 2c and f). Diffusion coefficients for $(3a)_2 \supset C_{60} (0.82 \times 10^{-10} \text{ m}^2 \text{ s}^{-1})$ and $(3b)_2 \supset C_{60} (0.80 \times 10^{-10} \text{ m}^2 \text{ s}^{-1})$ at 298 K in DMSO-d₆ correspond to the diameter of 2.7 nm (predicted value for the dimeric capsule is ~2.7 nm). Additionally, in the ROESY spectra the *f*-*j* correlation signals are present for $(3a)_2 \supset C_{60}$ and $(3b)_2 \supset C_{60}$ indicative of a close contact between hemispheres in a head-to-head arrangement (Fig. 3). This allows us to postulate the capsular structures of complexes $(3a)_2 \supset C_{60}$ and $(3b)_2 \supset C_{60}$ in DMSO-d₆. The corresponding ESI-MS experiments for the products revealed the presence of 3a and $(3a)_2 \supset C_{60}$ or 3b and $(3b)_2 \supset C_{60}$, without any traces of "empty" capsules, $(3a)_2$ or $(3b)_2$ (ESI†), which proves that the presence of a hydrophobic templating agent is essential for the self-assembly of cavitands 3a and 3b in DMSO.

In order to check the stability of hydrogen bonded species in the presence of a competitive solvent that is a hydrogen bond donor and acceptor we performed experiments in methanol. Differences in solubility (Table 2) allowed us to separate a mixture of **3b** and $(\mathbf{3b})_2 \supset C_{60}$ by selective precipitation of **3b**. Spectral data for $(\mathbf{3b})_2 \supset C_{60}$ in methanol (CH₃OH : CD₃OD 9 : 1) indicate that the complex retains its integrity (Fig. 4). The signal of encapsulated fullerene is present in the ¹³C NMR spectrum. The diffusion coefficient is in agreement with a diameter of the dimeric complex (2.65×10^{-10} m² s⁻¹ in methanol



Fig. 3 Dimerization and ordering of the structures in DMSO: ROESY effects indicative of β -sheet-type interactions (a) for (**3a**)₂ \supset C₆₀; (b) for (**3b**)₂ \supset C₆₀ and (c) predicted binding motif (the 3D structure was obtained by Val \rightarrow Gln mutation of an X-ray structure of a valine derivative¹⁷).

Table 2 Solubility table of 3a, 3b, $(3a)_2 \supset C_{60},$ and $(3b)_2 \supset C_{60}$ in MeOH and DMSO

3a	$(3a)_2 \supset C_{60}$	3b	$(\mathbf{3b})_2 \supset C_{60}$
Sol. Sol.	Insol. Sol.	Insol. Sol.	Sol. Sol.
	3a Sol. Sol.	3a $(3a)_2 \supset C_{60}$ Sol.Insol.Sol.Sol.	3a $(3a)_2 \supset C_{60}$ 3bSol.Insol.Insol.Sol.Sol.Sol.

Sol. - well-soluble; insol. - insoluble.

at 298 K corresponding to diameter 3.0 nm) and the correlation signal f-j is present in the ROESY spectrum, indicative of a close contact between hemispheres in a head-to-head arrangement. For glutamine derivatives $(3a)_2 \supset C_{60}$ and 3a the solubilities in methanol are also different, but we were not able to separate the mixture, due to co-crystallization of the species. However, from a supersaturated methanolic solution we obtained enriched crystals containing up to 60% of a $(3a)_2 \supset C_{60}$ (ESI†). After dissolution of the crystals in DMSO intact complex $(3a)_2 \supset C_{60}$ was present indicating that also glutamine-based self-assembled species are stable in an environment of methanol.

It is important to note that fullerene-templated self-assembly results in the formation of a well-defined hydrogen bonding system with a β -sheet-like binding motif and an ordered inherently chiral conformation. In the ¹H NMR spectrum, all signals for hydrogens capable of hydrogen bonding are considerably downfield shifted in $(3a)_2 \supset C_{60}$ as compared



Fig. 4 Spectra of $(3b)_2 \supset C_{60}$: (a) ¹H NMR spectrum (600 MHz, CH₃OH: CD₃OD, 9:1, v:v), (b) ¹H NMR spectrum with solvent suppression, (c) DOSY spectrum, (d) comparison of the ECD spectra of **3b** (green) and $(3b)_2 \supset C_{60}$ (orange) in MeOH (intensities of ECD spectra), (e) ECD spectra.

with cavitand 3a. For example one of the OH protons is shifted from 10.8 ppm to 13.7 ppm, and this proton is postulated to form an intramolecular hydrogen bond. Amide proton *i* is also upfield shifted in agreement with the formation of an intermolecular hydrogen bond connecting the hemispheres. The formation of a β -sheet-like binding motif is also supported by the increase of three-bond coupling constants ${}^{3}/(\text{NH-CH}_{\alpha})$ from 7.2 Hz for 3a to 8.5 Hz for $(3a)_2 \supset C_{60}$ and from 8.2 Hz for 3b to 8.9 Hz (DMSO) or 9.6 Hz (MeOH) for $(3b)_2 \supset C_{60}$. These coupling constants are directly related to the backbone dihedral φ angles as described by the Karplus relationship for peptides. Higher values are hallmarks of extended β-strand conformations ($\varphi \approx -139^\circ$) over a helix ($\varphi \approx -57^\circ$) or a random coil.^{23,24} An ordered conformation is also reflected in circular dichroism spectra that exhibit almost a 100-fold enhancement in the intensity of the lowest energy bands for $(3b)_2 \supset C_{60}$ as compared with 3b (Fig. 4d). These bands are the most sensitive to conformational inherent chirality and symmetry of these molecules, as they come from the HOMO-LUMO transition involving an inherently chiral chromophore.²⁵ These results indicate that the capsular dimers indeed retain a hydrogenbonded ordered structure in competitive medium. It is also important to note that there are no traces of dynamic exchange between species at the NMR timescale, indicating that the complexes are also kinetically persistent in this solvent, which is quite surprising considering that semi-open cavitands are also present in the solution.

In summary, we have demonstrated that wrapping of hydrogen-bonded shells around a hydrophobic template is effective in a highly disintegrative environment. DMSO is a strong competitor as a hydrogen bond acceptor and due to its hydrophobic character, while methanol is a strong hydrogen bond donor and acceptor. Therefore, both environments are competitive for aggregates based on hydrogen bonding and hydrophobic interactions. In fact, DMSO or methanol is commonly used for unfolding of natural proteins or disintegration of hydrogen bonded artificial aggregates. Here we demonstrated that the optimal combination of both interactions enables effective self-assembly, although none of the interactions is effective alone. The current system is not water-soluble, but we predict that this strategy can also be extended to aqueous solutions due to even stronger hydrophobic interactions.

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgements

This work was supported by the National Science Center (2013/09/B/ST5/01026) and the Wroclaw Centre for Networking and Supercomputing (grant no. 299). We would like to acknowledge Dr Aleksandra Butkiewicz for CD measurements.

Notes and references

- 1 D. Chandler, Nature, 2005, 437, 640-647.
- 2 H. J. Dyson, P. E. Wright and H. A. Scheraga, *Proc. Natl.* Acad. Sci. U. S. A., 2006, **103**, 13057–13061.
- 3 K. A. Dill, S. Bromberg, K. Yue, H. S. Chan, K. M. Ftebig, D. P. Yee and P. D. Thomas, *Protein Sci.*, 1995, 4, 561–602.
- 4 J. D. Hartgerink, E. Beniash and S. I. Stupp, *Science*, 2001, **294**, 1684–1688.
- 5 I. Leõn, J. Millán, E. J. Cocinero, A. Lesarri and J. A. Fernández, Angew. Chem., Int. Ed., 2013, 52, 7772– 7775.
- 6 C. L. D. Gibb and B. C. Gibb, J. Am. Chem. Soc., 2004, 126, 11408–11409.
- 7 H. Gan and B. C. Gibb, Chem. Commun., 2013, 49, 1395– 1397.
- 8 J. H. Jordan and B. C. Gibb, *Chem. Soc. Rev.*, 2015, 44, 547–585.
- 9 D. Ajami, L. Liu and J. Rebek Jr., *Chem. Soc. Rev.*, 2015, 44, 490–499.
- 10 E. Huerta, G. A. Metselaar, A. Fragoso, E. Santos, C. Bo and J. de Mendoza, *Angew. Chem., Int. Ed.*, 2007, 46, 202– 205.
- 11 E. Huerta, S. A. Serapian, E. Santos, E. Cequier, C. Bo and J. de Mendoza, *Chem. Eur. J.*, 2016, **22**, 13496–13505.
- 12 Q. Shi, K. E. Bergquist, R. Huo, J. Li, M. Lund, R. Vácha, A. Sundin, E. Butkus, E. Orentas and K. Wärnmark, *J. Am. Chem. Soc.*, 2013, **135**, 15263–15268.

- 13 D. Račkauskaite, K. E. Bergquist, Q. Shi, A. Sundin,
 E. Butkus, K. Wärnmark and E. Orentas, *J. Am. Chem. Soc.*, 2015, 137, 10536–10546.
- 14 G. D. Pantoş, J.-L. Wietor and J. K. M. Sanders, Angew. Chem., Int. Ed., 2007, 46, 2238–2240.
- 15 G. Markiewicz, A. Jenczak, M. Kołodziejski, J. J. Holstein, J. K. M. Sanders and A. R. Stefankiewicz, *Nat. Commun.*, 2017, 8, 15109.
- 16 S. J. Park, O.-H. Kwon, K.-S. Lee, K. Yamaguchi, D.-J. Jang and J.-I. Hong, *Chem. – Eur. J.*, 2008, **14**, 5353–5359.
- 17 M. Szymański, M. Wierzbicki, M. Gilski, H. Jędrzejewska, M. Sztylko, P. Cmoch, A. Shkurenko, M. Jaskólski and A. Szumna, *Chem. – Eur. J.*, 2016, 22, 3148–3155.
- 18 M. F. Perutz, T. Johnson, M. Suzuki and J. T. Finch, Proc. Natl. Acad. Sci. U. S. A., 1994, 91, 5355–5358.

- 19 M. F. Perutz, R. Staden, L. Moens and I. De Baere, *Curr. Biol.*, 1993, 3, 249–253.
- 20 M. F. Perutz, Trends Biochem. Sci., 1999, 24, 58-63.
- 21 M. R. Sawaya, S. Sambashivan, R. Nelson, M. I. Ivanova, S. A. Sievers, M. I. Apostol, M. J. Thompson, M. Balbirnie, J. J. W. Wiltzius, H. T. McFarlane, A. Ø. Madsen, C. Riekel and D. Eisenberg, *Nature*, 2007, 447, 453–457.
- 22 M. P. Szymański, H. Jędrzejewska, M. Wierzbicki and A. Szumna, *Phys. Chem. Chem. Phys.*, 2017, **19**, 15676–15680.
- 23 A. C. Wang and A. Bax, J. Am. Chem. Soc., 1996, 118, 2483– 2494.
- 24 F. Li, J. H. Lee, A. Grishaev, J. Ying and A. Bax, *ChemPhysChem*, 2015, **16**, 572–578.
- 25 H. Jędrzejewska, M. Kwit and A. Szumna, *Chem. Commun.*, 2015, **51**, 13799–13801.