Organic & Biomolecular Chemistry



View Article Online

CORRECTION

Check for updates

Cite this: Org. Biomol. Chem., 2019, **17**, 2288

Correction: Transfection efficiencies of α-tocopherylated cationic gemini lipids with hydroxyethyl bearing headgroups under high serum conditions†

Bappa Maiti,^{a,b} Mohini Kamra,^a Anjali A. Karande^c and Santanu Bhattacharya*^{a,b}

DOI: 10.1039/c9ob90009h

rsc li/obc

Correction for 'Transfection efficiencies of α -tocopherylated cationic gemini lipids with hydroxyethyl bearing headgroups under high serum conditions' by Bappa Maiti *et al.*, *Org. Biomol. Chem.*, 2018, **16**, 1983–1993.

There was an error in the ESI^{\dagger} uploaded at revision. The image for cellular internalization of fluorescein-labeled pGL3 with DOPE-T8T co-liposomes in one of the panels was included by mistake in the panel for DOPE-TH8S co-liposomes in the presence of m- β -CD (ESI, Fig. S7^{\dagger}). A revised supplementary file has been published alongside the original file.^{\dagger} This error has no impact whatsoever on the conclusions of this work.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^bDirector's Research Unit, Indian Association for the Cultivation of Science, Jadavpur, Kolkata 700032, India

^cDepartment of Biochemistry, Indian Institute of Science, Bangalore 560012, India

[†]Electronic supplementary information (ESI) available: Synthetic procedures and characterization details of gemini lipids, ¹H NMR spectra of gemini lipids, TE of neat tocopheryl lipids, optimization of the DOPE content in the co-liposomal formulations, hydrodynamic diameters of the lipoplexes, small angle X-ray diffraction of the optimized co-liposomes and lipoplexes, gel electrophoresis data and cellular internalization of lipoplexes with fluorescein labeled pGL3. See DOI: 10.1039/ c7ob02835k