


 Cite this: *RSC Adv.*, 2023, **13**, 8803

A review on arsenic in the environment: contamination, mobility, sources, and exposure

 Khageshwar Singh Patel,^{*a} Piyush Kant Pandey,^b Pablo Martín-Ramos,^{ID c} Warren T. Corns,^d Simge Varol,^{*e} Prosun Bhattacharya^f and Yanbei Zhu^{ID *g}

Arsenic is one of the regulated hazard materials in the environment and a persistent pollutant creating environmental, agricultural and health issues and posing a serious risk to humans. In the present review, sources and mobility of As in various compartments of the environment (air, water, soil and sediment) around the World are comprehensively investigated, along with measures of health hazards. Multiple atomic spectrometric approaches have been applied for total and speciation analysis of As chemical species. The LoD values are basically under $1 \mu\text{g L}^{-1}$, which is sufficient for the analysis of As or its chemical species in environmental samples. Both natural and anthropogenic sources contributed to As in air, while fine particulate matter tends to have higher concentrations of arsenic and results in high concentrations of As up to a maximum of 1660 ng m^{-3} in urban areas. Sources for As in natural waters (as dissolved or in particulate form) can be attributed to natural deposits, agricultural and industrial effluents, for which the maximum concentration of $2000 \mu\text{g L}^{-1}$ was found in groundwater. Sources for As in soil can be the initial contents, fossil fuel burning products, industrial effluents, pesticides, and so on, with a maximum reported concentration up to 4600 mg kg^{-1} . Sources for As in sediments can be attributed to their reservoirs, with a maximum reported

Received 6th February 2023

Accepted 9th March 2023

DOI: 10.1039/d3ra00789h

rsc.li/rsc-advances

^aDepartment of Applied Sciences, Amity University, Manth (Kharora), State Highway 9, Raipur-493225, CG, India. E-mail: patelkhageshwarsingh@gmail.com; kspatel@rpr.amity.edu

^bAmity University, Manth (Kharora), State Highway 9, Raipur-493225, CG, India. E-mail: pkpandey@rpr.amity.edu

^cDepartment of Agricultural and Environmental Sciences, EPS, Instituto de Investigación en Ciencias Ambientales de Aragón (IUCA), University of Zaragoza, Carretera de Cuarte, s/n, 22071 Huesca, Spain. E-mail: pmr@unizar.es

^dPS Analytical Ltd, Arthur House, Unit 11 Cray fields Industrial Estate, Orpington, Kent, BR5 3HP, UK. E-mail: wtc@psanalytical.com

^eDepartment of Geological Engineering, Faculty of Engineering, Suleyman Demirel University, Çünür, Isparta-32260, Turkey. E-mail: simgevarol@sdu.edu.tr

^fKTH-International Groundwater Arsenic Research Group, Department of Sustainable Development, Environmental Science and Engineering, KTH Royal Institute of Technology, Teknikringen 10B, SE-100 44 Stockholm, Sweden. E-mail: prosun@kth.se

^gEnvironmental Standards Research Group, Research Institute for Material and Chemical Measurement, National Metrology Institute of Japan (NMIJ), National Institute of Advanced Industrial Science and Technology (AIST), 1-1-1 Umezono, Tsukuba, Ibaraki 305-8563, Japan. E-mail: yb-zhu@aist.go.jp



Dr Khageshwar Singh Patel is currently, research Professor in Department of Applied Sciences, Amity University, Baloda Bazar Road, Raipur-493225, CG, India. He is an Alexander von Humboldt (Germany), Asia Foundation (San Francisco) and BSF, UGC, New Delhi awardee fellow, and run several national and international projects. He published >150 research papers in journals of

international repute and supervised 34 PhD students in his credit. His recent research interest is phytochemistry (identification of bioactive compounds), biofuel production and characterization as well as source appointment of toxic and strategic elements in environment.



Dr Simge Varol She was born in 1978 in Istanbul/Turkey and graduated from Geological Engineering Department of Süleyman Demirel University (SDU, Isparta, Turkey) in 2000. She completed her master's and doctorate degrees in 2006 and 2011, respectively, in Geological Engineering department of SDU. Her areas of expertise include topics such as hydrogeology, hydrology, hydrogeochemistry,

medical geology, ground and surface water quality, and water chemistry. She has 80 national and international scientific articles, 35 scientific papers, and 4 scientific book chapters related to her field of expertise. She has also been the Deputy Director of Water Institute in SDU since 2018. She is married and has 2 sons.



concentration up to 2500 mg kg⁻¹. It is notable that some reported concentrations of As in the environment are several times higher than permissible limits. However, many aspects of arsenic environmental chemistry including contamination of the environment, quantification, mobility, removal and health hazards are still unclear.

Introduction

Arsenic is a toxic element (metalloid) linked with a broad variety of neurologic, cardiovascular, dermatologic, and carcinogenic effects, including peripheral neuropathy, diabetes, ischemic heart disease, melanosis, keratosis, and impairment of liver function.¹ Chronic arsenic pollution is now recognized as a worldwide problem, with 21 countries experiencing arsenic contamination of the environment.^{2–9} Arsenic susceptibility plays an important role in manifestation of arsenicosis depending on methylation capacity, variation in host genome and individual epigenetic pattern.¹⁰

Arsenic is present in the environment in various inorganic and organic chemical forms: arsenite (As(III)), arsenate (As(V)), monomethylarsonic acid (MMA), dimethylarsinic acid (DMA), trimethyl arsine oxide (TMAO), arsenobetaine (AsB), *etc.*¹¹ The toxicity, mobility and solubility differ among species, in such a way that inorganic As(III) is more toxic than As(V), and in turn, organic As is less toxic.^{12,13}

There are several routes of human exposure to arsenic from both natural and anthropogenic sources.^{14,15} Regarding geogenic sources, the Earth's crust is an abundant natural source of arsenic,¹⁶ with an average concentration of As of $\approx 5 \text{ mg kg}^{-1}$.¹⁷ It is present in more than 200 different minerals, the most common of which is arsenopyrite.¹⁸ In relation to anthropogenic sources, mining, metal smelting and burning of fossil fuels are the major industrial processes that contribute to arsenic contamination of air, water, and soil.¹⁹ Much of the arsenic in the atmosphere comes from high-temperature processes such as coal-fired power plants^{20,21} and burning vegetation, but also from volcanic activity.²²

Prolonged intake of toxic iAs (inorganic arsenic) *via* air, water and food cause arsenicosis. Analytical techniques for As quantification at low levels in specific environmental samples are

required. Many methodologies for detection and quantification of As were reported.^{23–26}

Air is a potential source for As exposure in industrial areas mainly due to emission with airborne particulate matter from smelting of ores and coal combustion. The potential pathways of arsenic exposure and their reduction were reviewed.^{27,28} Arsenic chemistry, and factors controlling the sorption/desorption, mobility, uptake of As by plants and reduction of translocation in plant tissues and release of As from sediments into groundwater have been reviewed.^{29–31}

The aim of the review is to provide up-to-date information on arsenic contamination of the environment, quantification, sources, mobility and health hazards. The important scientific knowledge gaps and critical areas for future research are discussed.

Results and discussion

Methodology for As quantification

Arsenic contamination of environment is an urgent global issue, and therefore, a reliable and rapid method with good sensitivity and selectivity, portability and robustness to detect As at trace levels both in field and laboratory samples in view of reduced environmental and health risks are required. Several methods, *e.g.*, colorimetric, electrochemical, biological, electrophoretic, surface sensing and spectroscopic methods, are employed for As detection but most of them are suffering with poor sensitivity and selectivity. Many field kits for rapid detection of As at trace levels are used but their reliabilities are not sufficient.³² However, differentiation of the species of As is a quite complex analytical task. Numerous speciation procedures have been studied that include electrochemical, chromatographic, spectrometric and hyphenated techniques (Table 1). It is notable that different concentration units are used in Table 1, *i.e.*, $\mu\text{g L}^{-1}$, ng g^{-1} , and ng mL^{-1} , such units can be treated approximately equivalent to one another. Regardless the relatively close value to one another, the LoD values depend on the chemical species of As even measured by with an identical method. The LoD values are basically under $1 \mu\text{g L}^{-1}$, which are sufficient for the analysis of As or its chemical species as discussed in the following text. The repeatability with a relative standard deviation under 10% and a recovery value over 80% are sufficient for quantitative discussion.

Commonly used instrumental methods based on atomic absorption, atomic emission and mass spectroscopy methodologies *i.e.*, hydride generation atomic absorption spectrometry (HG-AAS), inductively coupled plasma-mass spectrometry (ICP-MS) and graphite furnace atomic absorption spectrometry (GF-AAS) were reported for monitoring of arsenic at the 0.05 mg L^{-1} ($50 \mu\text{g L}^{-1}$) MCL (maximum contaminant level). Among them, ICP-MS is a most sensitive multielement technique with selectivity depending on the version of the instrument whereas HG-AAS and GF-AAS are single element selective technique with relatively lower sensitivity.^{32,91} The ICP-MS with tandem



Yanbei Zhu, Senior researcher, National Metrology Institute of Japan (NMIJ), National Institute of Advanced Industrial Science and Technology (AIST), Japan. Yanbei received his PhD in March of 2005 from Nagoya University. After two years of postdoc research in Nagoya University, he joined NMIJ/AIST in April of 2007 and started the research on development of certified reference materials (CRMs) and related

techniques for elemental analysis in food and environmental samples. Yanbei is focusing on quantitative elemental analysis based on ICP-MS related techniques, as well as development of devices and instruments fascinating the sample pretreatment process and on-site analysis.





Table 1 The sample preparation methods, separation and detection for arsenic species in different matrixes (LC-ESI-MS/MS – liquid chromatography–electrospray ionization tandem mass spectrometry; DRC – dynamic reaction cell; CEUV – capillary electrophoresis–ultraviolet detector)^a

Matrix	Species	Sample pretreatment	Separation/ detection technique	Analytical features	Ref.
Whole blood and urine	As(III), As(V), MMA, DMA, AB	Dilution with HgCl ₂ and ultrafiltration	LC-ICP-MS	LoD/ $\mu\text{g L}^{-1}$: <0.3	32
Urine	As(III), As(V), MMA and DMA	Dilution with deionized water and filtration	IC- ICP-MS	LoD/ $\mu\text{g L}^{-1}$: 0.11 As(III), 0.25 As(V), 0.18 MMA, 0.17 DMA, 0.75 AB; repeatability/%: 1.9 As(III) 2.7 As(V), 2.1 MMA, 1.9 DMA, 2.8 AB Recovery: 85% to 100%	34
Urine	As(III), As(V), MMA, DMA and AB	Dilution with deionized water and filtration	HPLC-ICP-MS		33
Urine	As(III), As(V), MMA, DMA and TMAO	Dilution with mobile phase and filtration	HPLC-HG-AAS		35
Urine	As(III), As(V), MMA and DMA	Dilution HCl and L-cysteine	HPLC-ET-AAS		36
Urine	As(III), As(V), MMA and DMA	Dilution with deionized water and filtration	HPLC-HG-ICP-MS	LoD/ $\mu\text{g L}^{-1}$: 0.37 As(III), 0.22 As(V), 0.18 MMA, 0.17 DMA; precision/%: 4.1 As(III), 5.4 As(V), 6.0 MMA, 6.6 DMA	37 and 38
Whole blood	MMA, DMA	Centrifuge	CE-ICP-MS	LoD/ $\mu\text{g L}^{-1}$: 1; LoQ/ $\mu\text{g L}^{-1}$: 0.8 As(III), 1.0 As(V), 1.0 MMA, 0.9 DMA	39 and 40
Fish and oyster tissues	As(III), As(V), MMA and DMA	Lyophilization/microwave digestion	CE-ICP-MS	—	41–44
Fish	As(III), As(V), MMA and DMA	Ultrasonic extraction and 4 different experimental conditions	HG-AFS	LoD/ $\mu\text{g kg}^{-1}$: 0.62 As(III), 2.1 As(V), 1.8 MMA, 5.4 DMA; RSD: 6.8% As(III), 10.3% As(V), 8.5% MMA, 7.4% DMA; recovery: >93%	45–49
Fish sauce	AB, AC, TMAO	Extraction with water/methanol (1 + 1, v/v)/shaking/centrifugation	HPLC-ICP-MS	LoD/ ng g^{-1} : 3.07	50
Beverages (soft drink, lemon juice, beer)	As(III), As(V), MMA and DMA	Sample were passed through a C ₁₈ sep-pack and filtered	HPLC-ICP-MS	LoD/ ng L^{-1} : 0.2, 0.2, 0.3 and 0.5 for As(III), DMA, MMA and As(V), respectively; RSD of As(III), DMA, MMA and As(V) were 1.2, 2.1, 2.5 and 3.0%, respectively	51
Cereals	As(III), As(V), MMA and DMA	Ultrasonic extraction with H ₃ PO ₄ and Triton XT-114	HG-AFS	LoD/ ng g^{-1} : 1.3, 0.9, 1.5 and 0.6 for As(III), As(V), DMA and MMA, respectively; recoveries: >90%; repeatability: 3% for As(III), 5% for DMA and 6% for As(V) and MMA	52–55
Wines	As(III), As(V), MMA and DMA	Treatment with cysteine in HCl for total As; dilution with citrate buffer or acetic acid for As species	HG-AFS	LoD/ $\mu\text{g L}^{-1}$: 0.12, 0.27, 0.15 and 0.13 (as As); RSD: 2–6%, 5–9%, 3–7% and 2–5% for As(III), As(V), MMA and DMA, respectively	56 and 57
Vegetables	As(III), As(V), MMA and DMA	Ultrasonic extraction with H ₃ PO ₄ and Triton XT-114	HG-AFS	LoD/ ng g^{-1} : 3.1 As(III), 3.0 As(V), 1.5 DMA and 1.9 MMA; recovery: >91%	58–61
Milk	As(III), As(V)	Ultrasonic extraction with/without KI	HG-AFS	LoD/ ng g^{-1} : 8.1 and 10.3 for As(III), As(V); RSD: 5.7 and 5.5% for As(III) and As(V)	45, 62 and 63
Water	As(III), As(V), MMA	Treatment with cysteine, KI, urea or acids	HG-AAS, CE-UV, LC-ICP-MS	LoD/ $\mu\text{g L}^{-1}$: 0.10 As(III), As(V), 0.19 (DMA) for HG-AAS, 100 (As ^{III} , DMA) to 500 (As ^V) for CE-UV and 0.1 (DMA, MMA) to 0.2 (As ^{III} , As ^V) for LC-ICP-MS; precision: <5% RSD, recovery: 80–110% except CE-UV where only 50%	64

Table 1 (Contd.)

Matrix	Species	Sample pretreatment	Separation/ detection technique	Analytical features	Ref.
Water and urine sample	Inorganic, organic As	Treatment with/without cysteine	HG-ICP-MS	LoD/ng L ⁻¹ ; 6	65 and 66
Water and reference materials	As(III), As(V)	Treatment with HCl and NaBH ₄	HG-AAS	LoD/μg L ⁻¹ ; 0.1 for As(III) and 0.06 for total As. Precision (RSD) level is 2.9% for As(III) and 3.1% for total As	67 and 68
Water	As(III), As(V)	None	HG-AAS	LoD/μg L ⁻¹ ; 0.019 total As, 0.031 As(III)	69
Water	As(III), As(V)	Reaction with cysteine, NaBH ₄	HG-AAS	LoD/μg L ⁻¹ ; 0.1	70
Water	As(III), As(V)	Reaction with cysteine	HG-ETAAS	—	71
Water	As(III), As(V)	None	HG-AFS	—	72
Water	As(III), As(V)	pH adjustment	HG-AAS	LoD/μg L ⁻¹ ; 0.07–0.4 As(V) and 0.1–0.5 As(III + V); recovery: 90–102%	69
Water	As(III), As(V)	Treatment with KMnO ₄	ICP-AES	LoD/μg L ⁻¹ ; 0.1–0.6	72 and 73
Water	As(III), As(V), MMA and DMA	None	HPIC-ICP-MS	LoD/μg L ⁻¹ ; 0.22 As(III), 0.69 As(V); LoQ/μg L ⁻¹ ; 50	74
Water	Inorganic As	Pre-reduction of As(V) with cysteine	HG-AAS	LoD/μg L ⁻¹ ; 0.15; LoQ/μg L ⁻¹ ; 0.5; RSD (<i>n</i> = 10) <8%	75 and 76
Seawater	As(III)	Complexation with PDC	ETAAS	LoD/μg L ⁻¹ ; 0.008; RSD (<i>n</i> = 11): 4.5%	76 and 77
Water	As(III), As(V), MMA and DMA	None	HG-ICP-AES	Recovery: As(V) 97.6%, As(III) 100%, MMA 99.8%, and DMA 99.9%	76 and 78
Water	As(III), As(V)	Treatment with KMnO ₄	ETAAS	LoD/μg L ⁻¹ ; 0.35; recovery: 93.5–106.4%; RSD: 3–7%	76 and 79
Water	As(III), As(V), MMA, DMA and AB	None	HPIC-ICP-MS	LoD/μg L ⁻¹ ; 0.017 As(III), 0.026 As(V), 0.026 MMA, 0.023 DMA, 0.024 AB; LoQ/μg L ⁻¹ ; 0.056 As(III), 0.085 As(V), 0.088 MMA, 0.076 DMA, 0.080 AB	76 and 80
Groundwater	As(III), As(V), MMA, TMAO, PAA and PAO	None	HPIC-ICP-MS	LoQ/μg L ⁻¹ ; 0.2–0.8	81
Hot spring water	As(III), As(V), MMA, DMA, TMAO, TMA, AC and AB	None	HPIC-ICP-MS	LoD (μg L ⁻¹); 0.2; RSD (<i>n</i> = 6); <2%	82 and 83
Human hair	As(III), As(V)	Reduction of As(V) to As(III)	HG-AAS	LoD/μg L ⁻¹ ; 0.2 As(III), 0.5 As(V); RSD: 2.1% As(III) and 2.5% As(V)	76 and 84
Sediment and fly ash	Water soluble and phosphate-exchangeable As(III) and As(V)	Extraction with water and phosphate buffer	HG-AAS	LoD/μg L ⁻¹ ; 0.06–0.10; LoQ/μg L ⁻¹ ; 0.20–0.31. Repeatability expressed as RSD: <1%	76 and 85
Landfill leachate	As(III), As(V), MMA, DMA, TMAO and AB	Filtration	HPIC-ICP-MS	LoD vary between 11 ng L ⁻¹ for DMA to 27 ng L ⁻¹ for As(V). LoQ ranged between 36 and 90 ng L ⁻¹	81, 86 and 87
Municipal landfill leachates	As(III), As(V), MMA and DMA	Filtration	LC-ESI-MS/MS and HPIC-DR-ICP-MS	Recovery: 68–94%	81 and 88
Soil	Total As	Extraction with HNO ₃ , acetic acid, EDTA and Mehlich II	HG-AAS	—	89 and 90

^a As(III): arsenous acid, arsenite, As(V): arsenic acid, arsenate, MMA: monomethylarsenic acid, DMA: dimethylarsenic acid, TMAO: trimethylarsine oxide, TMAP: trimethylarsoniopropionate, TETRA, TMA: tetramethylarsonium ion, AB: arsenobetaine, AC: arsenocholine, PAO: phenylarsonic acid, PAA: phenylarsonic acid, TMAO: trimethylarsonic acid, DPAA: diphenylarsonic acid.





Fig. 1 Schematic diagram for quantification of As species in environmental samples.

quadrupole mass spectrometers (ICP-QQQ) is useful to overcome isobaric interferences with reduced background and improved selectivity. The tandem inorganic-organic mass spectrometry, *i.e.*, HPLC-ICP-MS + ESI-MS, is applicable to detect unknown arsenic compounds.⁹² However, they need extraction/separation of the analyte prior to analysis, depending on nature of the arsenic species (water- or lipid-soluble) and the matrix (biotic or abiotic), except water sample analysis.

For speciation of As in environmental samples, various approaches *i.e.* liquid-liquid, liquid-solid and solid phase extraction, sonication, pressurized liquid extraction, microwave-assisted extraction, supercritical fluid extraction, enzymatic hydrolysis, conventional sorbent, functional nanomaterials extractant, multi-sorbent based procedure, and derivatization of total arsenic were used.⁹³ Methanol or acetonitrile-water and methanol-chloroform or hexane are generally employed for extraction of polar arsenic and nonpolar species. A 1 mol L⁻¹ phosphoric acid is adequate for extraction of arsenic species from soil and sediment samples. A preconcentration step using freeze-drying or evaporation may be necessary to detect low abundance of arsenic species. Enzymes, *e.g.* pepsin, trypsin, pronase E, and lipase can be used to digest arsenic associated to lipids, proteins, peptide bonds, or cell walls, to increase extraction efficiencies.

Generally, chromatography and capillary electrophoresis are two major techniques used for separation of As species from complex matrices. Chromatography *i.e.* anion exchange, cation exchange, reversed-phase, ion pair, and hydrophilic interaction liquid chromatography (HILIC) are proposed for separation of various arsenic species such as As(III), As(V), MMA, DMA, AsB, arsenocholine (AsC), oxo-arsenosugars (oxoAsS), thio-arsenosugars (thioAsS), phenylarsenicals; AsB, AsC, TMAO, and

tetramethylarsonium ion.⁹⁴ The schematic diagram for quantification of As in environmental samples is shown in Fig. 1.

Sample collection, preservation, and pretreatment

The cleaned and sterilized FEP (fluorinated ethylene propylene) container should be used for collection of environmental samples and refrigerated at $-20\text{ }^{\circ}\text{C}$ in a dark place to control microbial activities. Water samples collected for As speciation analysis are filtered in the field using a $0.45\text{ }\mu\text{m}$ filter into opaque polyethylene bottles.⁹⁵ The sample storage duration should be minimized to avoid change in species stability. To prevent loss of arsenic species during sampling, samples (soil, plant, or water) should be collected in a sealed polyethylene bottle/bag. All samples should be stored in the freezer immediately after collection until the sample is prepared for analysis.

The preparation steps before analysis of each type of sample differ. Soil samples are air-dried, crushed lightly and sieved through a 2 mm sieve and used for analysis. The plant sample is placed in an oven dryer at $40\text{ }^{\circ}\text{C}$ to constant weight, the sample is ground, sieved and stored in a desiccator in brown glass bottles to avoid exposure to light and moisture until needed for analysis. Sample preparation for solid samples usually includes procedures such as mincing, freeze-drying, grinding, homogenization, sieving followed by extraction. After sampling of the fresh plant sample, it should be kept in the freezer ($-80\text{ }^{\circ}\text{C}$) to prevent species change. In addition, a dry and ground plant and soil sample can be stored at $-20\text{ }^{\circ}\text{C}$ for up to one year.⁹³

Statistical significance of the data is determined using analysis of variance (ANOVA) at $p < 0.05$ confidence level. Mobile phase concentrations and pH were optimized to get maximum peak separation. The limit of detection (LoD) is determined using the spike and blank based procedures. The standard





Fig. 2 The different mechanism controlling the mobility of arsenic in environmental component.

reference materials (SRMs) and certified reference materials (CRMs) are used to test and validate the accuracy of a method. SRM 1640 (NIST) and various CRM samples are widely employed to check a calibration curve for trace elements in

water, while CRMs used depend on the sample matrix and the arsenic species to be analyzed.⁹⁴

Distribution, mobility and sources of As in the environment

Arsenic exists in the form of various chemical species differing by their physicochemical behavior, in toxicity, bioavailability and biotransformation. The determination of arsenic species is an important issue for environmental, clinical and food chemistry.^{94,96}

In this review, the distribution, mobility, and sources of As in air, water, soil, sediment, plants, foods and marine organism; remediation of As from water, soil and sediment; disposal of contaminated samples, and health hazard assessment are discussed. The overall cycles of As in different environmental compartments are illustrated in Fig. 2. These cycles involve various chemical species such as As(III, V), MMA(III, V), and DMA(III, V), as well as chemical and/or biological reactions resulting reduction, oxidation, and methylation of As. Transportations of As species among soil, water, plant, animal, air, and sediment should be considered to understand the impact of As in the environment.

Table 2 Arsenic contents reported in ambient air/particulate matter (PM) over the world^a

Site	Particulate (<i>n</i>)	Concentration range (mean)	Source type	Ref
Berlin Museum, Germany	Ambient air/dust (30)	Max, 48 ng m ⁻³ /3507 mg kg ⁻¹	Anthropogenic	104
Cornwall, UK	Household dust (99)	3–1079 (84) mg kg ⁻¹	Anthropogenic, natural	114
Los Angeles, USA	PM _{2.5} (16)	As(III)/As(V) = <1.2–44.0 (7.4 ± 10.4)/<0.99–18.7 (5.2 ± 4.6) ng m ⁻³	Anthropogenic	117
Huelva, Spain	PM ₁₀	As(V)/As(III) = (1.2–2.1)/(6.5–7.8) ng m ⁻³	Anthropogenic	118
Canada	PM ₁₀	0.3 μg m ⁻³	Anthropogenic	120
Foshan, Hangzhou, Harbin, Yinchuan and Zhengzhou, China	PM ₁₀	96.9–185.2 ng m ⁻³	Anthropogenic	105
Tainan, Taiwan	PM _{2.5} /PM _{10–2.5} /SPM (180)	1.09–9.51/18–4.14/0.99–2.85 (1.94) ng m ⁻³ 22.7–155.8 (96.7) ng g ⁻¹	Anthropogenic, natural	119 and 106
Beijing, China	PM _{2.5} /PM ₁₀	8.1–128 (43.5 ± 28.0)/10–73.4 (61.8 ± 49.7) ng m ⁻³	Anthropogenic	115
AlaShan, China	PM _{2.5} /PM ₁₀	1.0–9.8 (3.7 ± 2.2)/1.1–12.4(5.4 ± 3.0) ng m ⁻³	Anthropogenic	115
Seoul, South Korea	PM _{2.5} /PM ₁₀	1.9–27.0 (9.0 ± 6.4)/2.8–216.4(34.4 ± 58.9) ng m ⁻³	Anthropogenic, natural	115
Gosan, South Korea	PM _{2.5} /PM ₁₀	0–8.9(1.7 ± 1.9)/0.2–6.8 (2.2 ± 2.0) ng m ⁻³	Anthropogenic, natural	115
Wuhan, China	PM _{2.5} (579)	6.81–10.7 ng m ⁻³	Anthropogenic	113
Hyderabad, India	PM ₁₀ and PM _{2.5} (72)	0.866 and 0.0022 μg m ⁻³	Anthropogenic	107
Raipur, India	PM ₁₀ (24)	5–47 (7 ± 1) ng m ⁻³	Anthropogenic	108
Ambargarh, Rajnandgaon, India	Road dust (13)	165–329 (238 ± 29) mg kg ⁻¹	Anthropogenic	116
Baoji, China	Road dust	9.0–42.8 (19.8) mg kg ⁻¹	Anthropogenic	112
China	SPM (918)	8.67–20.5 (11.7) ng m ⁻³	Anthropogenic	121
Dhaka, Bangladesh	SPM (8)	1.50–4.20 (3.06) ng m ⁻³	Anthropogenic	109
Lahore, Pakistan	SPM (20)	1140–1600 ng m ⁻³	Anthropogenic	110
Kolkata, India	Street dust (9)	<2–18.71 (7.96 ± 0.95) mg kg ⁻¹	Anthropogenic	111

^a SPM = suspended particulate matter, PM₁₀ = coarse particulate matter, PM_{2.5} = fine particulate matter.



Air. Arsenic is stable in the air and is emitted by both natural and anthropogenic sources. The natural sources include volcanic eruption, emission from soils or sediments by microbial reduction, dispersion of As containing particles by wind, evaporation from arsenic compounds and marine organism, *etc.*⁹⁷ Anthropogenic sources are high-temperature processes such as smelting of non-ferrous metals, burning of fossil fuels, vegetation and wastes, *etc.* They are emitted into the atmosphere primarily as As_2O_3 and arsines, and which are further transformed into arsenite, arsenate and organoarsenic compounds. Ultimately, they are transported by winds and settled into earth crusts by dry and wet depositions.⁹⁸

Arsenic in air is present mainly in particulate forms as inorganic As and released into the atmosphere primarily as oxides that adsorb on particulate matter (PM), which are dispersed by the wind to remote areas.⁹⁹ The particulates then settle down both by dry and wet deposition. Approximately two thirds of the atmospheric flux of As is of anthropogenic origin.^{100,101}

US OSHA regulations regarding iAs are found in 29 Code of Federal Regulations (CFR) 1910.1018. The permissible exposure level (PEL) for iAs is $10 \mu\text{g m}^{-3}$ of air, averaged over an 8 hour period without regarding to the use of a respirator. The action level is $5 \mu\text{g m}^{-3}$ of air. A medical surveillance program must be established for all employees exposed at or above the action level, for at least 30 days per year without regarding to the use of respirators.^{102,103} Typical background levels for arsenic in the rural, urban, and industrial areas are in the ranges of 0.2–1.5, 0.5–3, and $<50 \text{ ng m}^{-3}$, respectively.¹⁰¹ Arsenic content in ambient air/associated to $\text{PM}^{104-121}$ in different locations

worldwide is summarized in Table 2. Concentration range of As in particulate matter around the world is further illustrated in Fig. 3 based on the data given in Table 2. Inorganic As(III) and As(V) both occurred predominantly in ambient PM and preferably associated to the fine PM in accumulation mode with particle size of 0.2–2.0 μm .^{118,119,121} Concentrations (associated to ambient SPM, PM_{10} and $\text{PM}_{2.5}$) in various Chinese, Korean, Mongolian, Taiwanese, American and Canadian cities have been reported over the 2.4–185.2 ng m^{-3} range, with higher values in fine PM.^{105,106,113,115,119-121} Similar studies have been carried out for different South Asian cities. In Raipur, total As in the PM_{10} and $\text{PM}_{2.5}$ ranged from 37.0–501 and 27.0–293 $\mu\text{g m}^{-3}$, respectively.¹⁰⁸ In Dhaka city, As concentration in the SPM varied from 0.0015 to 0.0042 $\mu\text{g m}^{-3}$.¹⁰⁹ In Hyderabad, total As in the PM_{10} and $\text{PM}_{2.5}$ were in the ranges of 0.685–1.132 and 0.0010–0.0030 $\mu\text{g m}^{-3}$ with an annual average of 0.866 and 0.0022 $\mu\text{g m}^{-3}$, respectively.¹⁰⁷ The As in the SPM was detected in the range of 1140–1600 ng m^{-3} in different parts of the city of Lahore.¹¹⁰ The main reason for these high concentrations of As in Lahore city is attributed to anthropogenic pollution. Moreover, in the street, road and indoor dust, several folds higher As content was reported.^{104,111,112,114,116}

Water. Water receives As from natural deposits and from both agricultural and industrial effluents.¹²¹ It is found in the atmosphere, soils and rocks, natural waters, and other organisms. As is mobilized in the environment through a combination of natural processes and anthropogenic activities, *e.g.*, atmospheric dry and wet depositions, weathering reactions, biological activity and volcanic emissions, mining activity, combustion of fossil fuels, use of arsenical pesticides, herbicides, livestock feed additive, wood preservatives, *etc.*¹⁵

Arsenic in the air gradually precipitates on the soil as arsenic oxides and arsines. Arsenic accumulated on the soil surface is then slowly mixed into the groundwater with the effect of surface waters and rain.¹²³ In groundwater, As is released by rock weathering, long flow paths, use of phosphate fertilizers, irrigation of As enriched soils, *etc.* and influenced by physical geochemical characteristics of aquifers, water pumping rates, *etc.*^{28,124-127}

Arsenic in natural waters is mostly found in either dissolved or in particulate form. The most common forms of arsenic in natural waters are arsenite and arsenate.¹²⁸⁻¹³⁰ In well-oxygenated water and sediments, nearly all arsenic is present in the arsenate form.¹³¹⁻¹³⁴ The MMA and DMA (dimethyl arsenic acid) are also present in some water.^{15,28,135-137}

The solubility and mobility of As in the environment increase with increasing alkalinity and salinity. Its movement is controlled by adsorption/desorption and precipitation/dissolution reactions.¹³⁸ Three major modes of arsenic biotransformation occurred in the environment: redox transformation between As(III) and As(V), the reduction and methylation of As, and the bioproduction of organoarsenic compounds.^{8,139-141} Arsenic contamination with sources in water¹⁴²⁻¹⁶⁹ of some locations over the world are summarized in Table 3. Concentration range of As in water samples around the world is illustrated in Fig. 4 based on the data given in Table 3.

The allowable limit values of As for drinking and irrigation purposes recommended are 10 and 100 $\mu\text{g L}^{-1}$,



Fig. 3 Concentration range of As reported in particulate matter around the world (concentrations are given for total As, except for those specified).



Table 3 Distribution of As in water^a, $\mu\text{g L}^{-1}$

Location	Sample type (<i>n</i>)	Concentration	Sources	Source	Ref.
Eastern plains of Zimbabwe	SW(20)	1–96	Abandoned mine dumps	Natural	150
Tarkwa, Ghana	GW(40)	<5.2–69.4	Acid mine drainage	Natural	144
Kemerovo region, southwestern Siberia, Russia	POW	1700	Arsenopyrite leaching	Natural	153
Okavango Delta, Botswana	SW(9)	1.1–3.1	Arsenopyrite weathering	Natural	149
	GW (25)	1.8–116.6			
Ogun state, Nigeria	GW (20)	40–160	Arsenopyrite weathering	Natural	147
Prestea, Ghana	SW(13)	150–8250 (1568 ± 278)	Arsenopyrites	Natural	166
Paraiba do Sul delta, Brazil	GW	>50–163	Authigenic sulfides	Anthropogenic	156
Korba, India	PW (26)	8.0–30 (17.2 ± 2.1)	Coal burning	Anthropogenic, natural	167
Nile Delta, Egypt	SW	1.2–18.2	Fertilizers, detergents, herbicides	Anthropogenic	142
USA	GW(7000)	10–50	Igneous rocks	Natural	168
Hungary, Croatia, Serbia, Romania	GW	225/610/2000/13–200	Igneous rocks	Natural	151
Raipur, India	SEW (7)	26–51 (40 ± 7)	Industrial activities	Anthropogenic	158
Raipur, India	IWW (34)	600–7800 (2900 ± 1400)	Industrial activities	Anthropogenic	169
Lomé coastal region, Togo	SW (23)	3000–6460	Industrial and sea waste	Anthropogenic	164
Greece, Europe	GW	30–4500	Mineral leaching	Natural	151
Koekemoerspruit, South Africa	SW (40)	12.3–119	Minerals	Natural	145
Upper Moulouya, Morocco	SW (14)	50.08–199.6	Minerals	Natural	146
Tarkwa, Ghana	SW (12)	0.5–73	Minerals	Natural	143
South Africa	GW (10)	0.1–172.53 (32.21)	Minerals	Natural	157
Tarkwa, Ghana	GW (16)	<1–1760	Mining activities	Natural	152
African continent	SW/GW	10 000/1760	Mining activities	Natural	178 and 179
River water, Peru	SW (151)	0.1–93.1 (54.5)	Mining and agriculture activities	Anthropogenic, natural	148
Rift Valley, Ethiopia	SW(53)	<0.1–405	Rhyolitic rock	Natural	161
Rift Valley, Ethiopia	SW(11)	2.39–566 (165 ± 215)	Rhyolitic rock	Natural	162
	GW (54)	0.60–190 (22.4 ± 33.5)			
Rift Valley, Ethiopia	SW (05)/GW(25)/SPW(14)	0.02–96/<0.1–278/<0.1–156	Rock weathering	Natural	163
Uruguay	GW (46)	1.72–120.5	Rock weathering	Natural	154
Ambagrh Tehsil, India	FW (20)	29–98 (58 ± 7)	Sulfide minerals	Natural	159
Ambagrh Tehsil, India	PW (24)	10–53 (22 ± 5)	Sulfide minerals	Natural	165
Ambagrh Tehsil, India	GW (20)	148–985 (506 ± 118)	Sulfide minerals	Natural	160
Okavango Delta Botswana	SW	<1 to 188	Sulfide minerals	Natural	155
Yatenga and Zondoma provinces, Burkina Faso	GW(45)	<0.5–1630	Sulfide minerals	Natural	124

^a FW = field water, GW = groundwater, IWW = industrial wastewater, POW = pore water, PW = pond water, SEW = sewage water, SPW = spring water, SW = surface water.

respectively.^{170,171} Millions of people, especially from developing countries, use groundwater containing As for drinking purposes. Arsenic in the tube-well, well and surface water of several Asian countries (Bangladesh, Cambodia, China, India, Iran, Nepal, Pakistan, and Vietnam; Bangladesh, China, Cambodia, Iran, Pakistan, and Vietnam; China, Iran, and Pakistan) have been reported due to agricultural and industrial activities.⁴ In the particular case of China, Bangladesh and India, values of As content in field, surface, pond, ground, sewage and industrial waste water have been reported, and ascribed to mineralization of arsenopyrite.^{158–160,165,167,169,172–177}

In African countries, both surface and groundwater are also seriously polluted with arsenic due to industrial activities.¹⁷⁸ Examples for Botswana, Ethiopia, Ghana, Morocco, Nigeria,

South Africa and Tanzania^{42,145,146,150,155} are presented in Table 3. In some cases, this pollution has a natural origin (rock weathering), such as the As contents of up to 566 $\mu\text{g L}^{-1}$ in surface water reported in Rift Valley, Ethiopia,^{162,163} or groundwater pollution, with contents of up to 6150 $\mu\text{g L}^{-1}$, reported in Johannesburg, South Africa.^{177,178} Nonetheless, in other cases this contamination is a consequence of mining operations, which, for instance, have led to As contents of up to 8250 $\mu\text{g L}^{-1}$ in surface waters of Ghana and Togo.^{164,166}

Likewise, in Latin America, the population of various countries, including Argentina, Bolivia, Brazil, Chile, Colombia, Cuba, Ecuador, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Peru, and Uruguay^{15,143,144,147–149,152,154,156,157,163,180} are facing problems with As contaminated water too (see Table 3).






Fig. 4 Concentration range of As reported in water samples around the world.

Arsenic pollution is mainly due to natural resources,¹⁸¹ and is particularly serious in Argentina, Chile, and Mexico.¹⁸² In North America, contents of up to 2600 µg L⁻¹ have been reported.^{148,154,156} Western part of USA¹⁶⁸ and many countries of Europe,¹⁸³ such as Greece, Hungary, Romania, Croatia, Siberia, Turkey, and Spain are affected by elevated As levels, ranging from 30 to 4500 µg L⁻¹.^{151,153,184-186}

Soil. The background levels of As depends on soil types, ranging from 5–10 mg kg⁻¹.¹⁸⁷ The soil environment is contaminated with As by natural and anthropogenic sources. Arsenic in soils results from anthropogenic activities, and the main sources in the soil are naturally occurring minerals, such as arsenopyrite, gelignite, realgar, orpiment, *etc.*,¹⁷ and the mineralization of FeAsS and As₂S₃-like species in soil results in its pollution with As.¹⁸⁸

The major inputs of As in soil are associated with burning of fossil fuels and biomass, mining activities, industrial effluents, use of pesticides (*e.g.* monosodium methyl arsenate, disodium methyl arsenate and cacodylic acid) in agriculture, lumber preservatives (*e.g.* ammoniacal copper zinc arsenate and chromated copper arsenate), manufacturing of glass and ceramics (arsenic trioxide), alloys (brass and bronze), optics and electronic materials (light emitting or laser diodes, gallium arsenide microchips and circuit fiber optic crystals), leatherwork-leather preservative, pigments and paints.^{8,102}

Arsenic fate is a complex due to existence in many chemical forms naturally. In minerals, arsenicals and surface soils occur in arsenate forms, changing valence under different redox conditions, and adsorb to many soil clays and Fe, Mn and Al oxides.¹⁸⁹ Arsenic soil mobility and bioavailability is of great interest for human health assessment, and it depends on several physical and chemical factors: *e.g.* pH, redox potential, organic matter content, clay mineralogy, texture, Ca, Al, Mn, Fe and PO₄³⁻ content, organic ligands – fulvic and humic acids, and microbial activity of the soil and/or irrigation water.¹⁹⁰ The solubility and speciation of As in a contaminated soil is controlled by pH values of the soil extract, in such a way that the mobilization of As is increased in alkaline medium, resulting in an enhanced phytoavailability.¹⁹¹ Various elements (*e.g.*, Al, Ca, Fe, and Mn) enact a remarkable effect in As phyto availability in soil, and its translocation in different parts of the plants; root microbe-induced transformation system can also impact the fate of As in the rhizosphere; and phosphate and organic matter in the root system tend to reduce As phytotoxicity.^{122,192,193}

The As soil contamination and sources at various sites of the world are summarized in Table 4. The concentration of total As in 17 locations of surface, agricultural, garden and rhizospheric varied over 126–1600 mg kg⁻¹.^{159,194,195} The maximum value was observed in rhizospheric soil, probably due to plant and microorganism interactions. Although high concentrations have been reported all over India (in Assam, Bihar, Uttar Pradesh and West Bengal states), with values of up to 41.2 mg kg⁻¹,¹⁹⁶⁻¹⁹⁸ remarkably high values have been observed in Chhattisgarh state: with As concentrations reaching 164 mg kg⁻¹ in the surface soil of Korba basin¹⁹⁹ and of up to 4600 mg kg⁻¹ in Ambagarh Tehsil.^{159,194} Concerning other countries, contents over the 7–46, 3–266, 6.1–16.7, 9.7–110, 6–20, 7–1500 and 18.3 mg kg⁻¹ intervals have been reported in Bangladesh, Pakistan, Nepal, China, Vietnam, Iran, and South Africa, respectively.^{195,200-213}

Sediment. As is also present in the sediment bound to clay minerals and Fe, and Mn oxides, As is a redox sensitive element, present in the sediments of surface water reservoirs, formed by continuous weathering and erosion,²¹⁴ and its bioavailability for biota is dependent on both physical parameters, *e.g.*, pH, redox (200–500 mV), TOC (total organic carbon) and temperature.²¹⁵⁻²¹⁷ In general, when water conditions become oxidizing, at higher pH and organic values and low temperature, mobilized metals (As, Fe, *etc.*) is typically removed from solution as a solid precipitate, reducing the bioavailability of As for biota. The dissolution of oxides of Fe and Mn is found to be an effective mechanism for As mobility from sediment to groundwater.^{218,219}

Examples of pollution with As of the sediments of various surface water bodies (ponds, lagoons, river estuaries and sea) are summarized in Table 4. Concentration range of As in sediment and soil is illustrated in Fig. 5 based on the data given in Table 4. Over all total As concentration in 17 different types of sediments originated from pond, river, lake, lagoon, sea and estuarine was observed in interval between 2.5–2500 mg kg⁻¹ with maximum value in estuarine sediments due to industrial activities. The highest reported values are associated with



Table 4 Distribution of As in soil and sediment, mg kg⁻¹

Location	Sample type (<i>n</i>)	Concentration	Source	Source type	Ref
Kinghorn Loch, UK	Sediment	86–185	Aluminum industrial waste	Anthropogenic	226
Sindh, Pakistan	Soil	8.7–46.2	Arsenopyrite	Natural	201
Sindh, Pakistan	Lake sediment	11.3–55.8	Arsenopyrite	Natural	201
China	Agricultural soil (1648)	0.4–175.8 (10.40)	Atmospheric dusts	Anthropogenic	204
China	Soil	110	Chemical weapons	Anthropogenic	212
Pakistan	Soil	110–266	Coal burning	Anthropogenic	202
Korba, India	Surface soil (30)	49–164 (107 ± 32)	Coal mining and burning	Anthropogenic	199
Korba, India	Pond sediment (26)	36–154 (95 ± 12)	Coal mining and burning	Anthropogenic	167
China	Agricultural soil	9.7	Fertilizers	Anthropogenic	213
Bangladesh	Soil (12)	7–27.5 (14.5)	Groundwater	Natural	200
Florida, USA	Lake sediment (6)	39.0–53.5 (47.3)	Herbicides	Anthropogenic	225
USA	Buffalo river sediment(111)	2.6–417.0 (14.03 ± 39.21)	Industrial activities	Anthropogenic	224
Assam, India	Brahmaputra sediment (8)	Up to 37	Industrial activities	Anthropogenic	223
China	Coastal sediment	9.75	Industrial activities	Anthropogenic	229
Bothnian Bay/Bothnian Sea/Gulf of Finland/Gulf of Bothnia	Sediment	109–239/35–63/14–16/15–23	Industrial activities	Anthropogenic	228
North Egyptian lake	Lagoon sediment (21)	10–44 (25)	Industrial and agricultural activities	Anthropogenic	227
Bihar, middle Gangetic plain, India	Agricultural soil (19)	3.528–14.690	Irrigation water	Natural	196
Ballia, UP, India	Soil (30)	5.40–15.43 (11.12)	Irrigation water	Natural	198
Nepal	Soil	6.1–16.7	Irrigation water	Anthropogenic	203
Bangladesh	Paddy soils	46	Irrigation water	Anthropogenic	209
Vietnam	Agricultural soil	6–20	Irrigation water	Anthropogenic	206
Pakistan	Soil	3.0–3.81	Minerals	Natural	210
Iran	Soil (17)	7–795	Minerals	Natural	207
Iran	Soil (18)	105.4–1500 (1017)	Minerals	Natural	211
Bihar, India	Ganga sediment (19)	9.119–20.056	Minerals	Natural	196
Bihar middle Gangetic plain, India	Sediment (24)	6.9–14.2 (11.8)	Minerals	Natural	222
Colombia	Suratá River sediment	Up to 484	Mining	Natural	220
Brazil	Patos Lagoon sediment (15)	Up to 50 (2.5)	Mining	Natural	230
U.K. estuarine	Sediments	2–2500	Mining activities	Natural	221
USA	Fruit tree soil	>200	Pesticides	Anthropogenic	205
Ambagrh Tehsil, India	Surface soil(20)	58–302 (192 ± 28)	Sulfide minerals	Natural	195
Ambagrh Tehsil, India	Paddy soil (20)	44–270 (126 ± 28)	Sulfide minerals	Natural	159
Ambagrh Tehsil, India	Rhizospheric soil (16)	220–4600 (1600 ± 700)	Sulfide minerals	Natural	194
Ambagrh Tehsil, India	Pond sediment (24)	10–256 (53 ± 26)	Sulfide minerals	Natural	165
South Africa	Garden soil (4)	0.1–65.3 (18.3 ± 11.7)	Urban setting	Anthropogenic	208
West Bengal, India	Agricultural soil	19.40 ± 0.38–41.24 ± 0.48	Wet precipitation	Anthropogenic, natural	197

metalliferous mining activities: a concentration of 484 mg kg⁻¹ was registered in the stream sediments from Suratá river (southwestern area of Santurbán paramo, Colombia) near a gold mining area,²²⁰ and 2500 mg kg⁻¹ was recorded in the sediments of estuaries in southwest England.²²¹ Nonetheless, natural pollution can also lead to dangerous values: for instance, Ganga river – originated from Himalayas – carries sediments contaminated with As at concentrations in the 4.8–19.7 mg kg⁻¹ range,^{196,222} and contents of up to 37 mg kg⁻¹ have

been reported for Brahmaputra river – which flows from Southern Tibet.²²³

Remarkably high values of As in sediment were reported for two Indian locations, *i.e.* Ambagrh tehsil and Korba city, with values up to 256 and 154 mg kg⁻¹, respectively. Pollution of As in these locations were attributed to arsenopyrite leaching¹⁶⁵ and coal burning and alumina roasting,¹⁶⁷ respectively.

As content in the Buffalo bottom sediments (0–90 cm depth) were identified over range of 14.03 to 26.78 mg kg⁻¹ due to industrial activities.²²⁴ In the Florida lake sediment (30–45 cm





Fig. 5 Concentration range of As reported in sediment and soil around the world.

depth), high concentration of As at 147.5 mg kg^{-1} was accumulated due to use of arsenical herbicide ($\text{Na}(\text{CH}_3)_2\text{AsO}_4$) to the golf course and lawns throughout the state.²²⁵ Similarly, significantly high concentration of As at 160 mg kg^{-1} was observed in the red mud sediment (by-product of alumina production sludge) of Kinghorn Loch.²⁶ The metalliferous mining activities concentrated remarkable high concentration of As at 2500 mg kg^{-1} in the estuary's sediments of estuaries of southwest England.²²¹ Industrial activities also released As in the Brazilian lagoon, China coastal area, Egyptian lake, and Northern marine sediments over range of 2.5 to 239 mg kg^{-1} , respectively.²²⁷⁻²³⁰

Arsenic health hazards

Arsenic is known to be a human carcinogen involving in the generation of reactive oxygen species (ROS), genetic change and signal transduction pathways. Humans are more sensitive than other animals due to their lower efficiency of arsenic methylation.²³¹ It has been estimated that approximately 94 to 220 million people in the world are potentially exposed to ground-water containing excessive levels of As, most of whom are in Asia (94%) with some in Africa and South America.¹⁸³

Human exposure to As in contaminated areas occurred through oral, respiratory, or dermal routes.²³² Non-occupational human exposure to As is primarily associated with the ingestion of food and water. Food is generally the principal contributor to the daily intake of total arsenic, while drinking water can also be significant source of exposure in some areas.²³³ The daily intake of total As from food and beverages is generally between 20 and $300 \mu\text{g}$ per day.²³⁴ Pulmonary exposure may contribute up to

approximately $10 \mu\text{g}$ per day in a smoker and about $1 \mu\text{g}$ per day in a non-smoker, which values can be higher in polluted areas.

The concentration of metabolites of inorganic As in urine (inorganic arsenic, MMA and DMA) was used as a biomarker for the exposure and is generally in the range from 5 to $20 \mu\text{g As per L}$.^{235,241} In workplaces with up-to-date occupational hygiene practices, exposure concentrations generally do not exceed $100 \mu\text{g m}^{-3}$.²³⁶ Arsenic in hair and nails can be used as indicators of past As exposure, which requires to prevent external arsenic contamination of the samples.

The short-term effects of As exposure may include vomiting, diarrhea having blood, abdominal pain, dizziness, loss of sensation in limbs, skin problems, irritation, hair fall, muscle cramps, etc. Meanwhile, the long term As exposure can cause skin disorders and increased risk of diabetes, high blood pressure, and several types of cancer.²³⁷

Arsenic is a protoplasmic poisonous element mainly deactivate the sulfhydryl group of respiration, enzymes and mitosis cells. The toxic iAs (inorganic arsenic compounds) in human is methylated into non-toxic MMA and DMA, and excreted out through urine. However, the MMA is an intermediate species to form other arsenicals. Intake of iAs by air, water and/or food cause toxicity known as arsenicosis. Low exposure can lead adverse physiological effects (*i.e.*, vomiting and nausea, damaging of blood vessels, reduction in production of erythrocytes and leukocytes, change in heartbeat and body sensation, etc.). However, the prolonged exposure can develop lesions in skin, cancers, diabetes mellitus, hypertension, and neurological, pulmonary, vascular, cardiovascular diseases.²³⁸ The health effect risk (*i.e.* composition and size, type, magnitude, frequency, route, and duration of exposure) associated with intake of iAs is assessed by multiple parameters, *e.g.* average total dose (ATD), estimated daily intake (EDI), hazard quotient (HQ), hazard index (HI), cancer risk (CR) and CRlim methods. Dose-response assessment quantitatively evaluates a relationship between the amount of exposure to a contaminant and the possibility of adverse health effects. The following equations (i)–(vi) are used for evaluation the health hazard parameters.²³⁹⁻²⁴¹

$$\text{ATD} = \text{Asw} \times \text{IR} \quad (\text{i})$$

$$\text{EDI} = C_m \times \text{DI}/\text{BW} \quad (\text{ii})$$

$$\text{HQ} = \text{EDI}/\text{Rfd} \quad (\text{iii})$$

$$\text{HI} = \sum \text{HQ}_i \quad (\text{iv})$$

$$\text{CR}_{\text{lim}} = \text{Rfd} \times \text{BW}/C_m \quad (\text{v})$$

$$\text{Cancer risk} = \text{CDI} \times (\text{PF}) \quad (\text{vi})$$

where, scripts Asw, IR, C_m , DI, BW, Rfd, CDI, HI, HQ_i , CR_{lim} and PF represent arsenic contamination of water (mg L^{-1}), water ingestion rate (L per day), mean concentration of As in food, amount of food consumed per day (g per day), mean body weight of a person (kg), reference dose, chronic daily intake, hazard index, summation of HQ of noncarcinogens, the



maximum allowable food consumption rate (kg d^{-1}); and potency factor (mg per kg per day), respectively.

The permissible exposure limit for As in air, water and food recommended are $10 \mu\text{g m}^{-3}$ (over for 8 hour period), $10 \mu\text{g L}^{-1}$, and 0.5 mg kg^{-1} ,^{91,103} respectively. The recommended food dose (g per day) PF (oral route) for arsenic is $1.5 \text{ mg per kg per day}$, supposing that the cancer risk is acceptable when the value of cancer risk (CR) is lie in interval: 10^{-4} and 10^{-6} . Dietary exposures to total arsenic were highly variable, with a mean of $50.6 \mu\text{g per day}$ (range of $1.01\text{--}1081 \mu\text{g per day}$) for females and $58.5 \mu\text{g per day}$ (range of $0.21\text{--}1276 \mu\text{g per day}$) for males. U.S. dietary intake of inorganic arsenic has been estimated to range from 1 to $20 \mu\text{g per day}$, with grains and produce expected to be significant contributors to dietary inorganic arsenic intake.^{100,242–245} The permissible arsenic concentrations in water samples are $10 \mu\text{g L}^{-1}$ in the World Health Organization²⁴⁶ and United State Environmental Protection Agency (USEPA)²⁴⁰ guidelines for drinking water. In addition, guide levels for the protection of the aquatic biota and drinking water proposed by local authorities are $15 \mu\text{g L}^{-1}$ and $10 \mu\text{g L}^{-1}$ respectively.²⁴⁷ Accordingly, arsenic can cause adverse effects not only for humans but also for plant and animal species living in aquatic environments. Especially in aquatic ecosystem, arsenic concentration above the permissible limit affects various physiological systems such as growth, reproduction, ion regulation, mortification, gene expression, immune function, enzyme activities and histopathology of fish. Therefore, harmful effects may be seen for people who form the last link of the food chain and for human health.

However, in health index evaluation method factors *i.e.*, the life style, age and diet quality are not considered. For example, young children, elderly and people with long-term illnesses are at greater risk. In addition, interactions of other toxicants, and bioavailability, bioaccumulation, background concentration level of the toxicant are not considered. Toxicity data are often unavailable, and evaluated data is not sufficient to verify dose-response curve and action mode.²⁴⁸

Conclusions

In this review, data on natural and anthropogenic sources and mobility of As in the various compartments of the environment (air, water, soil and sediment, and biota) in different parts of the World has been presented, together with their main biogeochemical relations. It's apparent that arsenic contamination is an alarming problem on a global scale for animal and human health. The smelters, coal based thermal power plants and biomass burnings are potential sources for arsenic emissions, and their uses should be minimized. The contaminated water, food and soil dusts poses the greatest threat to public health from arsenic. New insights on As contamination, exposure sources, mobility and toxicity mechanisms at molecular and gene levels for understanding of its adverse effect spectrum are required.

The scientific understanding of As is still evolving even with the foundation of knowledge already established by years of research on its origin, toxicities, mobility, distribution patterns, quantification, and exposure. Further study on its interactions should help

in the development of methods of safe clean-up and exposure prevention all the way down to the trophic level of ecosystem.

Ethics approval

No ethical issue is declared in this review.

Data availability

All data are available in this review paper, and further information could be obtained from the corresponding author (<https://docs.google.com/document/d/1kz2tTt-zfVyxRTJbbxKtPA-S7LSxPosA/edit?usp=sharing&oid=102850718200296775047&rtpof=true&sd=true>).

Author contributions

Khageshwar Singh Patel: conceptualization, data curation, investigation, methodology, validation, visualization, writing – original draft. Piyush Kant Pandey and Warren T. Corns: investigation, visualization, validation, editing. Pablo Martín-Ramos: data curation, graphic drafting, editing original draft. Simge Varol: software, graphics drafting, editing – original draft. Prosun Bhattacharya: review & editing. Yanbei Zhu: preparing figures, reviewing, and editing.

Conflicts of interest

The authors declare that they have no known competing financial interests or personal relationships that could influenced the work reported in this paper.

Acknowledgements

The University Grants Commission (UGC), New Delhi is gratefully acknowledged for providing financial support through BSR grant no. F.18-1/2011(BSR)2016.

References

- 1 R. N. Ratnaike, *Postgrad. Med. J.*, 2003, **79**, 391–396, <https://pmj.bmj.com/content/postgradmedj/79/933/391.full.pdf>.
- 2 P. Bhattacharyya and M. M. Alam, Arsenic-contaminated soil toxicity and its mitigation through monocot crops, in *Contaminants in Agriculture*, ed. M. Naeem, A. Ansari and S. Gill, Springer, 2020.
- 3 B. K. Mandal and K. T. Suzuki, *Talanta*, 2002, **58**, 201–235, DOI: [10.1016/S0039-9140\(02\)00268-0](https://doi.org/10.1016/S0039-9140(02)00268-0).
- 4 M. Shahid, M. Imran, S. Khalid, B. Murtaza, N. K. Niazi, Y. Zhang, I. Hussain, Arsenic environmental contamination status in South Asia, *Arsenic in Drinking Water and Food*, 2020, pp. 13–39.
- 5 D. K. Nordstrom, *Science*, 2002, **296**, 2143–2145, DOI: [10.1126/science.1072375](https://doi.org/10.1126/science.1072375).
- 6 P. K. Pandey, S. Yadav, S. Nair and A. Bhui, *Environ. Int.*, 2002, **28**, 235–245, DOI: [10.1016/s0160-4120\(02\)00022-3](https://doi.org/10.1016/s0160-4120(02)00022-3).



- 7 M. M. Rahman, R. Naidu and P. Bhattacharya, *Environ. Geochem. Health*, 2009, **S1**, 9–21, DOI: [10.1007/s10653-008-9233-2](https://doi.org/10.1007/s10653-008-9233-2).
- 8 S. Shankar and U. Shanker, *Sci. World J.*, 2014, 304524, DOI: [10.1155/2014/304524](https://doi.org/10.1155/2014/304524).
- 9 A. Shrivastava, A. Barla, S. Singh, S. Mandraha and S. Bose, *J. Hazard. Mater.*, 2017, **324**, 526–534, DOI: [10.1016/j.jhazmat.2016.11.022](https://doi.org/10.1016/j.jhazmat.2016.11.022).
- 10 T. Sanyal, P. Bhattacharjee, S. Paul and P. Bhattacharjee, *Front. Public Health*, 2020, **8**, 464, DOI: [10.3389/fpubh.2020.00464](https://doi.org/10.3389/fpubh.2020.00464).
- 11 K. F. Akter, G. Owens, D. E. Davey and R. Naidu, *Rev. Environ. Contam. Toxicol.*, 2005, **184**, 97–149, DOI: [10.1007/0-387-27565-7_3](https://doi.org/10.1007/0-387-27565-7_3).
- 12 S. Alka, S. Shahir, N. Ibrahim, M. J. Ndejiko, D. V. Vo and F. A. Manan, *J. Cleaner Prod.*, 2021, **278**, 123805, DOI: [10.1016/j.jclepro.2020.123805](https://doi.org/10.1016/j.jclepro.2020.123805).
- 13 V. K. Sharma and M. Sohn, *Environ. Int.*, 2009, **35**, 743–759, DOI: [10.1016/j.envint.2009.01.005](https://doi.org/10.1016/j.envint.2009.01.005).
- 14 B. A. Baker, V. A. Cassano and C. Murray, *J. Occup. Environ. Med.*, 2018, **60**, e634–e639, DOI: [10.1097/JOM.0000000000001485](https://doi.org/10.1097/JOM.0000000000001485).
- 15 P. L. Smedley and D. G. Kinniburgh, *Appl. Geochem.*, 2002, **17**, 517–568, DOI: [10.1016/S0883-2927\(02\)00018-5](https://doi.org/10.1016/S0883-2927(02)00018-5).
- 16 R. Bhadauria, *Horticult. Int. J.*, 2019, **3**, 20–22, DOI: [10.15406/hij.2019.03.00106](https://doi.org/10.15406/hij.2019.03.00106).
- 17 H. Masuda, *Prog. Earth Planet. Sci.*, 2018, **5**, 68, DOI: [10.1186/s40645-018-0224-3](https://doi.org/10.1186/s40645-018-0224-3).
- 18 IARC (International Agency for Research on Cancer), 2012, <https://www.ncbi.nlm.nih.gov/books/NBK304375/>.
- 19 M. K. Upadhyay, A. Shukla, P. Yadav and S. Srivastava, *Food Chem.*, 2019, **276**, 608–618, DOI: [10.1016/j.foodchem.2018.10.069](https://doi.org/10.1016/j.foodchem.2018.10.069).
- 20 C. Wang, H. Liu, Y. Zhang, C. Zou and E. J. Anthony, *Prog. Energy Combust. Sci.*, 2018, **68**, 1–28, DOI: [10.1016/j.peccs.2018.04](https://doi.org/10.1016/j.peccs.2018.04).
- 21 Y. E. Yudovich and M. P. Ketris, Arsenic in coal: a review, *Int. J. Coal Geol.*, 2005, **61**, 141–196, DOI: [10.1016/j.coal.2004.09.003](https://doi.org/10.1016/j.coal.2004.09.003).
- 22 ENVIS (Environmental Information System), publisher (Ministry of Environment & Forest, Govt. of India), 2020, Arsenic -<https://nbrienvs.nic.in>.
- 23 M. D'Amato, G. Forte and S. Caroli, Identification and quantification of major species of arsenic in rice, *J. AOAC Int.*, 2004, **87**, 238–243, DOI: [10.1093/jaoac/87.1.238](https://doi.org/10.1093/jaoac/87.1.238).
- 24 W. Davis, R. Zeisler, J. Sieber and L. Yu, *Anal. Bioanal. Chem.*, 2010, **396**, 3041–3050, DOI: [10.1007/s00216-010-3541-y](https://doi.org/10.1007/s00216-010-3541-y).
- 25 F. Mehoul and S. W. Fowler, *Environ. Sci. Pollut. Res.*, 2022, **29**, 3288–3301, DOI: [10.1007/s11356-021-17130-0](https://doi.org/10.1007/s11356-021-17130-0).
- 26 M. S. Rahman, A. Kumar, R. Kumar, M. Ali, A. K. Ghosh and S. K. Singh, *Indian J. Occup. Environ. Med.*, 2019, **23**, 126–132, DOI: [10.4103/ijocem.IJOEM_240_18](https://doi.org/10.4103/ijocem.IJOEM_240_18).
- 27 J.-Y. Chung, S.-D. Yu and Y.-S. Hong, *J. Prev. Med. Public Health*, 2014, **47**, 253–257, DOI: [10.3961/jpmph.14.036](https://doi.org/10.3961/jpmph.14.036).
- 28 N. J. Raju, *Environ. Res.*, 2022, **203**, 111782, DOI: [10.1016/j.envres.2021.111782](https://doi.org/10.1016/j.envres.2021.111782).
- 29 A. Aftabtalab, J. Rinklebe, S. M. Shaheen, N. K. Niazi, E. Moreno-Jiménez, J. Schaller and K. H. Knorr, *Chemosphere*, 2022, **286**, 131790, DOI: [10.1016/j.chemosphere.2021.131790](https://doi.org/10.1016/j.chemosphere.2021.131790).
- 30 A. Dadwal and V. Mishra, *Clean: Soil, Air, Water*, 2017, **45**, 1600364, DOI: [10.1002/clean.201600364](https://doi.org/10.1002/clean.201600364).
- 31 M. Pigna, A. Caporale, L. Cavalca, A. Sommella and A. Violante, *Environ. Eng. Sci.*, 2015, **32**, 150505074828006, DOI: [10.1089/ees.2015.0018](https://doi.org/10.1089/ees.2015.0018).
- 32 N. Yogarajah and S. S. H. Tsai, *Environ. Sci.: Water Res. Technol.*, 2015, **1**, 426–447, DOI: [10.1039/C5EW00099H](https://doi.org/10.1039/C5EW00099H).
- 33 C. F. Harrington, R. Clough, L. R. Drennan-Harris, S. J. Hilld and J. F. Tyson, *J. Anal. At. Spectrom.*, 2011, **26**, 1561–1595.
- 34 R. Xie, W. Johnson, S. Spayd, G. S. Hall and B. Buckley, *Anal. Chim. Acta*, 2006, **578**, 186–194, DOI: [10.1016/j.aca.2006.06.076](https://doi.org/10.1016/j.aca.2006.06.076).
- 35 R. Sur and L. Dunemann, *J. Chromatogr. B: Anal. Technol. Biomed. Life Sci.*, 2004, **807**, 169–176, DOI: [10.1016/j.jchromb.2004.03.051](https://doi.org/10.1016/j.jchromb.2004.03.051).
- 36 Y. Shi, *Speciation of arsenic by chemical separations and neutron activation analysis*, 2001, <http://hdl.handle.net/10222/55802>.
- 37 K. A. Francesconi and D. Kuehnelt, *Analyst*, 2004, **129**, 373–395, DOI: [10.1039/b401321m](https://doi.org/10.1039/b401321m).
- 38 B. Radke, L. Jewell and J. Namieśnik, *Crit. Rev. Anal. Chem.*, 2012, **42**, 162–183, DOI: [10.1080/10408347.2011.634637](https://doi.org/10.1080/10408347.2011.634637).
- 39 V. L. Dressler, F. G. Antes, C. M. Moreira, D. Pozebon and F. A. Duarte, *Int. J. Mass Spectrom.*, 2011, **307**, 149–162, DOI: [10.1016/j.ijms.2011.01.026](https://doi.org/10.1016/j.ijms.2011.01.026).
- 40 M. F. Mesko, C. A. Hartwig, C. A. Bizzi, J. S. Pereira, P. A. Mello and E. M. Flores, *Int. J. Mass Spectrom.*, 2011, **307**, 123–136, DOI: [10.1016/j.ijms.2011.03.002](https://doi.org/10.1016/j.ijms.2011.03.002).
- 41 C. G. Yuan and G. B. Jiang, *J. Anal. At. Spectrom.*, 2005, **20**, 103–110.
- 42 S. Simon, H. Tran, F. Pannier and M. Potin-Gautier, *J. Chromatogr. A*, 2004, **1024**, 105–113, DOI: [10.1016/j.chroma.2003.09.068](https://doi.org/10.1016/j.chroma.2003.09.068).
- 43 K. L. Ackley, C. B'Hymer, K. L. Sutton and J. A. Caruso, *J. Anal. At. Spectrom.*, 1999, **14**, 845–850, DOI: [10.1039/A807466F](https://doi.org/10.1039/A807466F).
- 44 J. L. Gomez-Ariza, D. Sanchez-Rodas, I. Giraldez and E. Morales, *Analyst*, 2000, **125**, 401–407, DOI: [10.1039/A908884](https://doi.org/10.1039/A908884).
- 45 P. Cava-Montesinos, K. Nilles, M. L. Cervera and M. de la Guardia, *Talanta*, 2005, **66**, 895–901, DOI: [10.1016/j.talanta.2004.12.042](https://doi.org/10.1016/j.talanta.2004.12.042).
- 46 J. L. Gómez-Ariza, D. Sánchez-Rodas, R. Beltrán and I. Giraldez, *Int. J. Environ. Anal. Chem.*, 1999, **74**, 203–213, DOI: [10.1080/03067319908031426](https://doi.org/10.1080/03067319908031426).
- 47 C. G. Yuan and G. B. Jiang, *J. Anal. At. Spectrom.*, 2005, **20**, 103–110, DOI: [10.1039/B416102E](https://doi.org/10.1039/B416102E).
- 48 Y. Bohari and A. Astruc, *J. Anal. At. Spectrom.*, 2001, **16**, 774–778, DOI: [10.1039/b101591p](https://doi.org/10.1039/b101591p).
- 49 T. Johannes and V. Elterena, *Water Res.*, 2000, **36**, 2967, DOI: [10.1016/S0043-1354\(01\)00527-9](https://doi.org/10.1016/S0043-1354(01)00527-9).



- 50 O. Muñoz, D. Vélez and R. Montoro, *Analyst*, 1999, **124**, 601–607, DOI: [10.1039/A809426H](https://doi.org/10.1039/A809426H).
- 51 N. M. M. Coelho, L. M. Coelho, E. S. De Lima, A. Pastor and M. De La Guardia, *Talanta*, 2005, **66**, 818–822, DOI: [10.1016/j.talanta.2004.11.037](https://doi.org/10.1016/j.talanta.2004.11.037).
- 52 M. M. Reyes, M. L. Cervera, R. C. Campos and M. De la Guardia, *Spectrochim. Acta, Part B*, 2007, **62**, 1078–1082, DOI: [10.1016/j.sab.2007.07.011](https://doi.org/10.1016/j.sab.2007.07.011).
- 53 D. T. Heitkemper, N. P. Vela, K. R. Stewart and C. S. Westphal, *J. Anal. At. Spectrom.*, 2001, **16**, 299–306, DOI: [10.1039/B007241I](https://doi.org/10.1039/B007241I).
- 54 N. P. Vela and D. T. Heitkemper, *J. AOAC Int.*, 2004, **87**, 244–252, PMID: 15084107.
- 55 I. Pizarro, M. Gómez, M. A. Palacios and C. Cámara, *Anal. Bioanal. Chem.*, 2003, **376**, 102–109, DOI: [10.1007/s00216-003-1870-9](https://doi.org/10.1007/s00216-003-1870-9).
- 56 H. C. Rezende, I. L. Almeida, L. M. Coelho, N. M. Coelho and T. L. Marques, *Sample Prep.*, 2014, **2**, 31–48, DOI: [10.2478/sample-2014-0004](https://doi.org/10.2478/sample-2014-0004).
- 57 M. A. Vieira, P. Grinberg, C. R. Bobeda, M. N. Reyes and R. C. Campos, *Spectrochim. Acta, Part B*, 2009, **64**, 459–476, DOI: [10.1016/j.sab.2009.04.010](https://doi.org/10.1016/j.sab.2009.04.010).
- 58 M. N. M. Reyes, M. L. Cervera, R. C. Campos and M. de la Guardia, *Talanta*, 2008, **75**, 811–816, DOI: [10.1016/j.sab.2009.04.010](https://doi.org/10.1016/j.sab.2009.04.010).
- 59 N. P. Vela, D. T. Heitkemper and K. R. Stewart, *Analyst*, 2001, **126**, 1011–1017, DOI: [10.1039/B102420P](https://doi.org/10.1039/B102420P).
- 60 Y. Bohari and A. Astruc, *J. Anal. At. Spectrom.*, 2001, **16**, 774–778, DOI: [10.1039/B101591P](https://doi.org/10.1039/B101591P).
- 61 J. A. Caruso, D. T. Heitkemper and C. B'Hymer, *Analyst*, 2001, **126**, 136–140, DOI: [10.1039/b009825f](https://doi.org/10.1039/b009825f).
- 62 X. Ai, L. Wu, M. Zhang, X. Hou, L. Yang and C. J. Zheng, *Agric. Food Chem.*, 2014, **62**, 8586–8593, DOI: [10.1021/jf501638k](https://doi.org/10.1021/jf501638k).
- 63 H. Yu, C. Li, Y. Tian and X. Jiang, *Microchem. J.*, 2020, **152**, 104312, DOI: [10.1016/j.microc.2019.104312](https://doi.org/10.1016/j.microc.2019.104312).
- 64 R. Clough, C. F. Harrington, S. J. Hill, Y. Madrid and J. F. Tyson, *J. Anal. At. Spectrom.*, 2013, **28**, 1153–1195, DOI: [10.1039/C3JA90039H](https://doi.org/10.1039/C3JA90039H).
- 65 A. I. Mihaltan, T. Frentiu, M. Ponta, D. Petreus, M. Frentiu, E. Darvasi and C. Marutoiu, *Talanta*, 2013, **109**, 84–90, DOI: [10.1016/j.talanta.2013.01.056](https://doi.org/10.1016/j.talanta.2013.01.056).
- 66 M. I. Leybourne, K. H. Johannesson and A. Asfaw, *Rev. Mineral. Geochem.*, 2014, **79**, 371–390, DOI: [10.2138/rmg.2014.79.6](https://doi.org/10.2138/rmg.2014.79.6).
- 67 G. Samanta, T. R. Chowdhury, B. K. Mandal, B. K. Biswas, U. K. Chowdhury, G. K. Basu, C. R. Chanda, D. Lodh and D. Chakraborti, *Microchem. J.*, 1999, **62**, 174–191, DOI: [10.1006/mchj.1999.1713](https://doi.org/10.1006/mchj.1999.1713).
- 68 A. M. Abdel-Lateef, R. A. Mohamed and H. H. Mahmoud, *Adv. Chem. Sci.*, 2013, **2**, 110–113.
- 69 K. F. Akter, G. Owens, D. E. Davey and R. Naidu, *Rev. Environ. Contam. Toxicol.*, 2005, **184**, 97–149, DOI: [10.1007/0-387-27565-7_3](https://doi.org/10.1007/0-387-27565-7_3).
- 70 V. Bühl, C. Álvarez, K. Kordas, M. Pistón and N. Mañay, *J. Environ. Pollut. Hum. Health*, 2015, **3**, 46–51, DOI: [10.12691/jephh-3-2-4](https://doi.org/10.12691/jephh-3-2-4).
- 71 E. Pitzalis, D. Ajala, M. Onor, R. Zamboni and A. D'Ulivo, *Anal. Chem.*, 2007, **79**, 6324–6333, DOI: [10.1021/ac070513p](https://doi.org/10.1021/ac070513p).
- 72 D. Sutherland, P. M. Swash, A. C. Macqueen, L. E. McWilliam, D. J. Ross and S. C. Wood, *Environ. Technol.*, 2002, **23**, 385–403, DOI: [10.1080/09593332508618444](https://doi.org/10.1080/09593332508618444).
- 73 T. Rupasinghe, T. J. Cardwell, R. W. Catrall, I. D. Potter and S. D. Kolev, *Anal. Chim. Acta*, 2004, **510**, 225–230, DOI: [10.1016/j.aca.2004.01.006](https://doi.org/10.1016/j.aca.2004.01.006).
- 74 I. Komorowicz and D. Barańkiewicz, *Rapid Commun. Mass Spectrom.*, 2014, **28**, 159–168, DOI: [10.1002/rcm.6774](https://doi.org/10.1002/rcm.6774).
- 75 G. G. Bortoleto and S. Cadore, *Talanta*, 2005, **67**, 169–174, DOI: [10.1016/j.talanta.2005.02.012](https://doi.org/10.1016/j.talanta.2005.02.012).
- 76 A. Gonzalez, M. L. Cervera, S. Armenta and M. De la Guardia, *Anal. Chim. Acta*, 2009, **636**, 129–157, DOI: [10.1016/j.aca.2009.01.065](https://doi.org/10.1016/j.aca.2009.01.065).
- 77 P. Herbello-Hermelo, M. C. Barciela-Alonso, A. Bermejo-Barrera and P. Bermejo-Barrera, *J. Anal. At. Spectrom.*, 2005, **20**, 662–664, DOI: [10.1039/B416772D](https://doi.org/10.1039/B416772D).
- 78 J. Narvaez, P. Richter and M. I. Toral, *Anal. Bioanal. Chem.*, 2005, **381**, 1483–1487, DOI: [10.1007/s00216-005-3122-7](https://doi.org/10.1007/s00216-005-3122-7).
- 79 L. M. Coelho, N. M. Coelho, M. A. Arruda and M. de la Guardia, *Talanta*, 2007, **71**, 353–358, DOI: [10.1016/j.talanta.2006.04.007](https://doi.org/10.1016/j.talanta.2006.04.007).
- 80 S. N. Ronkart, V. Laurent, P. Carbonnelle, N. Mabon, A. Copin and J. P. Barthélemy, *Chemosphere*, 2007, **66**, 738–745, DOI: [10.1016/j.chemosphere.2006.07.056](https://doi.org/10.1016/j.chemosphere.2006.07.056).
- 81 I. Komorowicz and D. Barańkiewicz, *Talanta*, 2011, **84**, 247–261, DOI: [10.1016/j.talanta.2010.10.065](https://doi.org/10.1016/j.talanta.2010.10.065).
- 82 Y. Morita, T. Kobayashi, T. Kuroiwa and T. Narukawa, *Talanta*, 2007, **73**, 81–86, DOI: [10.1016/j.talanta.2007.03.005](https://doi.org/10.1016/j.talanta.2007.03.005).
- 83 M. Popp, S. Hann and G. Koellensperger, *Anal. Chim. Acta*, 2010, **668**, 114–129, DOI: [10.1016/j.aca.2010.04.036](https://doi.org/10.1016/j.aca.2010.04.036).
- 84 X. Li, J. Jia and Z. Wang, *Anal. Chim. Acta*, 2006, **560**, 153–158, DOI: [10.1016/j.aca.2005.12.054](https://doi.org/10.1016/j.aca.2005.12.054).
- 85 J. C. González, I. Lavilla and C. Bendicho, *Talanta*, 2003, **59**, 525–534, DOI: [10.1016/S0039-9140\(02\)00541-6](https://doi.org/10.1016/S0039-9140(02)00541-6).
- 86 E. V. Hoek, J. T. Elteren and R. N. J. Comansi, *Int. J. Environ. Anal. Chem.*, 1996, **63**, 67–79, DOI: [10.1080/03067319608039811](https://doi.org/10.1080/03067319608039811).
- 87 M. Ponthieu, P. Pinel-Raffaitin, I. Hecho, L. Mazeas, D. Amouroux, O. F. X. Donard and M. Potin-Gautier, *Water Res.*, 2007, **41**, 3177–3185, DOI: [10.1016/j.watres.2007.04.026](https://doi.org/10.1016/j.watres.2007.04.026).
- 88 Y. Li, G. K. Low, J. A. Scott and R. Amal, *Chemosphere*, 2010, **79**, 794–801, DOI: [10.1016/j.chemosphere.2010.03.010](https://doi.org/10.1016/j.chemosphere.2010.03.010).
- 89 R. Leschber, R. D. Davis and P. L'Hermite, *Chemical methods for assessing bio-available metals in sludges and soils*, 1985.
- 90 J. Száková, P. Tlustoš, W. Goessler, Z. Frková and J. Najmanová, *J. Hazard. Mater.*, 2009, **172**, 1244–1251, DOI: [10.1016/j.jhazmat.2009.07.143](https://doi.org/10.1016/j.jhazmat.2009.07.143).
- 91 USEPA (US Environmental Protection Agency), *The Arsenic Rule*, 2015, <https://www.epa.gov/sites/production/files/2015-09/documents/train1-background.pdf>.



- 92 F. Ardini, G. Dan and M. Grotti, *J. Anal. At. Spectrom.*, 2020, **35**, 215–237, DOI: [10.1039/C9JA00333A](https://doi.org/10.1039/C9JA00333A).
- 93 M. Z. U. Kamal, M. Y. Miah, Arsenic Speciation Techniques in Soil Water and Plant: An Overview, *Arsenic Monitoring, Removal and Remediation*, Intech Open, 2021, DOI: [10.5772/intechopen.99273](https://doi.org/10.5772/intechopen.99273).
- 94 M. S. Reid, K. S. Hoy, J. R. M. Schofield, J. S. Uppal, Y. Lin, X. Lu, H. Peng and X. C. Le, *TrAC, Trends Anal. Chem.*, 2020, **123**, 115770, DOI: [10.1016/j.trac.2019.115770](https://doi.org/10.1016/j.trac.2019.115770).
- 95 S. J. Stetson, C. Lawrence, S. Whitcomb and C. Kanagy, *MethodsX*, 2021, **8**, 101183, DOI: [10.1016/j.mex.2020.101183](https://doi.org/10.1016/j.mex.2020.101183).
- 96 K. A. Francesconi and D. Kuehnelt, *Analyst*, 2004, **129**, 373–395, DOI: [10.1039/b401321m](https://doi.org/10.1039/b401321m).
- 97 IPCS (International Programme on Chemical Safety), *Arsenic and arsenic compounds*, WHO, 2001.
- 98 W. T. Frankenberger, *Environmental Chemistry of Arsenic*, Marcel Dekker, Inc, New York, 2002.
- 99 J. Matschullat, *Sci. Total Environ.*, 2000, **249**, 297–312, DOI: [10.1016/S0048-9697\(99\)00524-0](https://doi.org/10.1016/S0048-9697(99)00524-0).
- 100 ATSDR (Agency for Toxic Substances and Disease Registry), *Toxicological profile for arsenic, Update*, U.S. Department of Health & Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, 2007, <http://www.atsdr.cdc.gov/toxprofiles/tp2.html>.
- 101 WHO (World Health Organization), *Arsenic and arsenic compounds*, World Health Organization, International Programme on Chemical Safety, Geneva, 2nd edn, 2001, Environmental Health Criteria 224.
- 102 B. A. Baker, V. A. Cassano and C. Murray, *J. Occup. Environ. Med.*, 2018, **60**, e634–e639, DOI: [10.1097/JOM.0000000000001485](https://doi.org/10.1097/JOM.0000000000001485).
- 103 USOSHA (US Occupational Safety and Health Administration), *29 CFR 1910.1018. OSHA Standard for inorganic arsenic*, 2001, https://www.osha.gov/pls/oshaweb/owadisp.show_document?p_table=STANDARDS&p_id=10023, accessed August 8, 2022.
- 104 K. Deering, E. Spiegel, C. Quaisser, D. Nowak, R. Schierl, S. B. Reilly and M. Garí, *Environ. Monit. Assess.*, 2019, **191**, 375, DOI: [10.1007/s10661-019-7495-z](https://doi.org/10.1007/s10661-019-7495-z).
- 105 J. Duan and J. Tan, *Atmos. Environ.*, 2013, **74**, 93–101, DOI: [10.1016/j.atmosenv.2013.03.031](https://doi.org/10.1016/j.atmosenv.2013.03.031).
- 106 G.-C. Fang, C.-C. Lin, J.-H. Huang and Y.-L. Huang, *Aerosol Air Qual. Res.*, 2011, **11**, 218–229, DOI: [10.4209/aaqr.2010.09.0075](https://doi.org/10.4209/aaqr.2010.09.0075).
- 107 S. Gummeneni, Y. B. Yusup, M. Chavali and S. Z. Samadi, *Atmos. Res.*, 2011, **101**, 752–764, DOI: [10.1016/j.atmosres.2011.05.002](https://doi.org/10.1016/j.atmosres.2011.05.002).
- 108 S. Gupta, Studies on trace element composition and sources of aerosols, PhD thesis submitted to Pt. Ravishankar Shukla University, Raipur, India, 2010.
- 109 M. F. Islam, S. S. Majumder, A. Al Mamu, Md. B. Khan, M. A. Rahman and A. Salam, *Open J. Air Pollut.*, 2015, **4**, 86–98, DOI: [10.4236/ojap.2015.42009](https://doi.org/10.4236/ojap.2015.42009).
- 110 M. I. Jalees and Z. Asim, *Earth Sci.*, 2016, **75**, 842, DOI: [10.1007/s12665-016-5604-7](https://doi.org/10.1007/s12665-016-5604-7).
- 111 S. Kar, B. Nath, A. C. Samal and S. C. Santra, *Curr. Sci.*, 2006, **90**, 158–160.
- 112 X. Lu, L. Y. Li, L. Wang, K. Lei, J. Huang and Y. Zhai, *Atmos. Environ.*, 2009, **43**, 2489–2496, DOI: [10.1016/j.atmosenv.2009.01.048](https://doi.org/10.1016/j.atmosenv.2009.01.048).
- 113 X. Mao, X. Hu, Y. Wang, W. Xia, S. Zhao and Y. Wan, *Environ. Sci. Pollut. Res.*, 2020, **17**, 21654–21665, DOI: [10.1007/s11356-020-08626-2](https://doi.org/10.1007/s11356-020-08626-2).
- 114 D. R. S. Middleton, M. J. Watts, D. J. Beriro, E. M. Hamilito, G. S. Leonardi, T. Fletcher, R. M. Close and D. A. Polya, *Environ. Sci.: Processes Impacts*, 2017, **19**, 517–527, DOI: [10.1039/C6EM00690F](https://doi.org/10.1039/C6EM00690F).
- 115 E. Park, D. Kim, C. Park, S. Song, B. Lee, Y. Hon, X. Pan, J. Wang, Y. Zhang and K. Park, *Korean J. Environ. Health Sci.*, 2010, **36**, 60–372, DOI: [10.5668/JEHS.2010.36.5.360](https://doi.org/10.5668/JEHS.2010.36.5.360).
- 116 K. S. Patel, A. Yadav, Y. K. Sahu, L. Lata, H. Milosh, W. T. Corns and P. Martín-Ramos, *J. Hazard., Toxic Radioact. Waste*, 2020, **24**, 05019006., DOI: [10.1061/\(ASCE\)HZ.2153-5515.0000475](https://doi.org/10.1061/(ASCE)HZ.2153-5515.0000475).
- 117 E. S. Rabano, N. T. Castillo, K. J. Torre and P. A. Solomon, *J. Air Waste Manage.*, 1989, **39**, 76–80, DOI: [10.1080/08940630.1989.10466511](https://doi.org/10.1080/08940630.1989.10466511).
- 118 D. Sánchez-Rodas, M. Ana, S. Campa, J. D. Rosa, V. Oliveira, J. L. Gómez-Ariza, X. Querol and A. Alastuey, *Chemosphere*, 2007, **66**, 1485–1493, DOI: [10.1016/j.chemosphere.2006.08.043](https://doi.org/10.1016/j.chemosphere.2006.08.043).
- 119 Y. Tsai, S. Kuo and Y. Lin, *Atmos. Environ.*, 2003, **37**, 3401–3411, DOI: [10.1016/S1352-2310\(03\)00358-3](https://doi.org/10.1016/S1352-2310(03)00358-3).
- 120 S. Wang and C. N. Mulligan, *Sci. Total Environ.*, 2006, **366**, 701–721, DOI: [10.1016/j.scitotenv.2005.09.005](https://doi.org/10.1016/j.scitotenv.2005.09.005).
- 121 J. Wang, Y. Wan, L. Cheng, W. Xia, Y. Li and S. Xu, *Atmos. Pollut. Res.*, 2020, **11**, 785–792, DOI: [10.1016/j.apr.2020.01.006](https://doi.org/10.1016/j.apr.2020.01.006).
- 122 A. D. Roble, P. Polizzi, M. B. Romero, L. N. C. Boudet, S. Medici, A. Costas and M. Gerpe, *Environ. Earth Sci.*, 2016, **75**, 1479, DOI: [10.1007/s12665-016-6273-2](https://doi.org/10.1007/s12665-016-6273-2).
- 123 J. O. Nriagu, P. Bhattacharya, A. B. Mukherjee, J. Bundschuh, R. Zevenhoven and R. H. Loeppert, *Trace Met. Other Contam. Environ.*, 2007, **9**, 3–60, DOI: [10.1016/S1875-1121\(06\)09001-8](https://doi.org/10.1016/S1875-1121(06)09001-8).
- 124 P. L. Smedley, J. Knudsen and D. Maiga, *Appl. Geochem.*, 2007, **22**, 1074–1092, DOI: [10.1016/j.apgeochem.2007.01.001](https://doi.org/10.1016/j.apgeochem.2007.01.001).
- 125 M. Chakraborty, A. Mukherjee and K. M. Ahmed, *Curr. Pollut. Rep.*, 2015, **1**, 220–247, DOI: [10.1007/s40726-015-0022-0](https://doi.org/10.1007/s40726-015-0022-0).
- 126 N. Akhtar, M. I. Syakir Ishak, S. A. Bhawani and K. Umar, *Water*, 2021, **13**, 2660, DOI: [10.3390/w13192660](https://doi.org/10.3390/w13192660).
- 127 B. Pradhan, S. Chand, S. Chand, P. R. Rou and S. K. Naik, *Groundwater Sustain. Develop.*, 2023, **20**, 100868, DOI: [10.1016/j.gsd.2022.100868](https://doi.org/10.1016/j.gsd.2022.100868).
- 128 Y. Y. Wang, L. Y. Chai and W. C. Yang, in *Arsenic Pollution Control in Nonferrous Metallurgy*, ed. L.Y. Chai, Springer, Singapore, 2019, pp. 1–15, DOI: [10.1007/978-981-13-6721-2_1](https://doi.org/10.1007/978-981-13-6721-2_1).



- 129 A. Penezić, M. L. Tercier-Waeber, M. Abdou, C. Bossy, L. Dutruch, E. Bakker and J. Schäfer, *Mar. Chem.*, 2020, **223**, 103804, DOI: [10.1016/j.marchem.2020.103804](https://doi.org/10.1016/j.marchem.2020.103804).
- 130 F. U. R. Qaiser, F. Zhang, R. R. Pant, C. Zeng, N. G. Khan and G. Wang, *Sci. Total Environ.*, 2023, **857**, 159408, DOI: [10.1016/j.scitotenv.2022.159408](https://doi.org/10.1016/j.scitotenv.2022.159408).
- 131 Y. W. Chen, X. Yu and N. Belzile, *J. Geochem. Explor.*, 2019, **205**, 106349, DOI: [10.1016/j.gexplo.2019.106349](https://doi.org/10.1016/j.gexplo.2019.106349).
- 132 Q. Zheng, S. Tu, J. Hou, C. Ni, M. Wang, L. Ren, M. Wang, M. Cao, S. Xiong and W. Tan, *Water Res.*, 2021, **203**, 117558, DOI: [10.1016/j.watres.2021.117558](https://doi.org/10.1016/j.watres.2021.117558).
- 133 B. C. Astles, J. Chételat, M. J. Palmer and J. C. Vermaire, *PLoS One*, 2022, **17**, e0279412, DOI: [10.1371/journal.pone.0279412](https://doi.org/10.1371/journal.pone.0279412).
- 134 S. R. Fung, E. A. Hull, K. Burkart, J. E. Gawel, A. R. Horner-Devine and R. B. Neumann, *Water Resour. Res.*, 2022, **58**, e2022WR032564, DOI: [10.1029/2022WR032564](https://doi.org/10.1029/2022WR032564).
- 135 G. Raber, K. A. Francesconi, K. J. Irgolic and W. Goessler, *Fresenius' J. Anal. Chem.*, 2000, **367**, 181–188, DOI: [10.1007/s002160051621](https://doi.org/10.1007/s002160051621).
- 136 S. J. Flora, *Arsenic: chemistry, occurrence, and exposure*, in *Handbook of arsenic toxicology*, Academic Press, 2015, pp. 1–49.
- 137 Y. C. Jang, Y. Somanna and H. J. I. J. Kim, *Int. J. Appl. Environ. Sci.*, 2016, **11**, 559–581.
- 138 M. L. Brusseau and J. Chorover, Chapter 8-Chemical Processes Affecting Contaminant Transport and Fate, *Environ. Pollut. Sci.*, 2020, 113–130, DOI: [10.1016/B978-0-12-814719-1.00008-2](https://doi.org/10.1016/B978-0-12-814719-1.00008-2).
- 139 D. L. Alonso, R. Pérez, C. K. Y. A. Okio and E. Castillo, *J. Environ. Manage.*, 2020, **264**, 110478, DOI: [10.1016/j.jenvman.2020.110478](https://doi.org/10.1016/j.jenvman.2020.110478).
- 140 M. M. Hussain, J. Wang, I. Bibi, M. Shahid, N. K. Niazi, J. Iqbal, I. A. Mian, S. M. Shaheen, S. Bashir, N. S. Shah and J. Rinklebe, *J. Hazard. Mater.*, 2021, **403**, 124027, DOI: [10.1016/j.jhazmat.2020.124027](https://doi.org/10.1016/j.jhazmat.2020.124027).
- 141 X. Meng, R. R. Dupont, D. L. Sorensen, A. R. Jacobson and J. E. McLean, *Appl. Geochem.*, 2016, **66**, 250–263, DOI: [10.1016/j.apgeochem.2016.01.004](https://doi.org/10.1016/j.apgeochem.2016.01.004).
- 142 A. R. Abdel-Moati, *Water, Air, Soil Pollut.*, 1990, **51**, 117–132, DOI: [10.1007/BF00211509](https://doi.org/10.1007/BF00211509).
- 143 K. A. Asante, T. Agusa, A. Subramanian, O. D. Ansa-Asare, C. A. Biney and S. Tanabe, *Chemosphere*, 2007, **66**, 1513–1522, DOI: [10.1016/j.chemosphere.2006.08.022](https://doi.org/10.1016/j.chemosphere.2006.08.022).
- 144 P. Bhattacharya, O. Sracek, B. Eldvall, R. Asklund, G. Barmen, G. Jacks, J. Koku, J.-E. Gustafsson, N. Singh and B. B. Balfors, *J. Afr. Earth Sci.*, 2012, **66–67**, 72–84, DOI: [10.1016/j.jafrearsci.2012.03.005](https://doi.org/10.1016/j.jafrearsci.2012.03.005).
- 145 B. M. Dzoma, R. A. Moralo, L. E. Motsei, R. V. Ndou and F. R. Bakunzi, *J. Anim. Vet. Adv.*, 2010, **9**, 3026–3033, DOI: [10.3923/javaa.2010.3026.3033](https://doi.org/10.3923/javaa.2010.3026.3033).
- 146 M. L. El Hachimi, L. El Founti, A. Bouabdli, N. Saidi, M. Fekhoui and N. Tassé, *Rev. Sci. Eau*, 2007, **20**, 1–13, DOI: [10.7202/014903ar](https://doi.org/10.7202/014903ar).
- 147 A. M. Gbadebo, *32nd International Geological Congress*, Florence, Italy, CRC Press, 2005, pp. 85–92, DOI: [10.1201/9780203970829](https://doi.org/10.1201/9780203970829).
- 148 G. George and B. M. Gqaza, *J. Adv. Agric. Technol.*, 2015, **2**, 29–35, DOI: [10.12720/joaat.2.1.29-33](https://doi.org/10.12720/joaat.2.1.29-33).
- 149 P. Huntsman-Mapila, T. Mapila, M. Letshwenyo, P. Wolski and C. Hemond, *Appl. Geochem.*, 2006, **21**, 1376–1391, DOI: [10.1016/j.apgeochem.2006.05.003](https://doi.org/10.1016/j.apgeochem.2006.05.003).
- 150 S. B. Jonnalagadda and G. Nenzou, *J. Environ. Sci. Health, Part A: Environ. Sci. Eng. Toxic Hazard. Subst. Control*, 1996, **11**, 2547–2555, DOI: [10.1080/10934529609376509](https://doi.org/10.1080/10934529609376509).
- 151 I. A. Katsoyiannis, M. Mitrakas and A. I. Zouboulis, *Desalin. Water Treat.*, 2015, **54**, 2100–2107, DOI: [10.1080/19443994.2014.933630](https://doi.org/10.1080/19443994.2014.933630).
- 152 J. M. Kusimi and B. A. Kusimi, *J. Geochem. Explor.*, 2012, **112**, 252–261, DOI: [10.1016/j.gexplo.2011.09.003](https://doi.org/10.1016/j.gexplo.2011.09.003).
- 153 E. V. Lazareva, O. V. Shuvaeva and V. G. Tsimbalist, *Geochem.: Explor., Environ., Anal.*, 2002, **2**, 263–268, DOI: [10.1144/1467-787302-030](https://doi.org/10.1144/1467-787302-030).
- 154 I. Machado, L. Falchi, V. Bühl and N. Mañay, *Sci. Total Environ.*, 2020, **721**, 137787, DOI: [10.1016/j.scitotenv.2020.137787](https://doi.org/10.1016/j.scitotenv.2020.137787).
- 155 N. Mladenov, P. Wolski, G. M. Hettiarachchi, M. Murray-Hudson, H. Enriquez, S. Damaraju and W. Masamba, *J. Hydrol.*, 2014, **518**, 326–341, DOI: [10.1016/j.jhydrol.2013.09.026](https://doi.org/10.1016/j.jhydrol.2013.09.026).
- 156 N. Mirlean, V. E. Andrus, P. Baisch, G. Griep and M. R. Casartelli, *Mar. Pollut. Bull.*, 2003, **46**, 1480–1484, DOI: [10.1016/S0025-326X\(03\)00257-1](https://doi.org/10.1016/S0025-326X(03)00257-1).
- 157 R. Mudzielwana, M. W. Gitari, S. A. Akinyemi, A. O. Talabi and P. Ndungu, *Groundwater Sustain. Develop.*, 2020, **10**, 100336, DOI: [10.1016/j.gsd.2020.100336](https://doi.org/10.1016/j.gsd.2020.100336).
- 158 K. S. Patel, N. S. Dahariya, S. Chakradhari, P. K. Sahu, K. P. Rajhans, S. Ramteke, L. Lata and H. Milos, *Am. J. Anal. Chem.*, 2015, **6**, 787–796, DOI: [10.4236/ajac.2015.610075](https://doi.org/10.4236/ajac.2015.610075).
- 159 K. S. Patel, B. L. Sahu, S. Ramteke and E. Bontempi, *J. Environ. Prot.*, 2016, **7**, 689–698, DOI: [10.4236/jep.2016.75061](https://doi.org/10.4236/jep.2016.75061).
- 160 K. S. Patel, B. L. Sahu, N. S. Dahariya, A. Bhatia, R. K. Patel, L. Matini, O. Sracek and P. Bhattacharya, *Appl. Water Sci.*, 2017, **7**, 1817–1826, DOI: [10.1007/s13201-015-0355-2](https://doi.org/10.1007/s13201-015-0355-2).
- 161 T. Rango, G. Bianchini, L. Beccaluva and R. Tassinari, *J. Afr. Earth Sci.*, 2010, **57**, 479–491, DOI: [10.1016/j.jafrearsci.2009.12.005](https://doi.org/10.1016/j.jafrearsci.2009.12.005).
- 162 T. Rango, A. Vengosh, G. Dwyer and G. Bianchini, *Water Res.*, 2013, **47**, 5801–5818, DOI: [10.1016/j.watres.2013.07.002](https://doi.org/10.1016/j.watres.2013.07.002).
- 163 C. Reimann, K. Bjorvatn, B. Frengstad, Z. Melaku, R. Tekle-Haimanot and U. Siewers, *Sci. Total Environ.*, 2003, **311**, 65–80, DOI: [10.1016/S0048-9697\(03\)00137-2](https://doi.org/10.1016/S0048-9697(03)00137-2).
- 164 M. H. Rezaie-Boroon, K. Ghandi and K.-M. Folly, *Fresenius Environ. Bull.*, 2011, **20**, 1853–1865. <http://www.psp-parlar.de>.
- 165 B. L. Sahu, S. Ramteke, K. P. Rajhans, K. S. Patel, I. Wysocka and I. Jaron, *Water Resour.*, 2018, **45**, 992–1001, DOI: [10.1134/S0097807818060131](https://doi.org/10.1134/S0097807818060131).



- 166 Y. Serfor-Armah, B. J. B. Nyarko, S. B. Dampare and D. Adomako, *Water, Air, Soil Pollut.*, 2006, **175**, 181–192, DOI: [10.1007/s11270-006-9127-9](https://doi.org/10.1007/s11270-006-9127-9).
- 167 R. Sharma, K. S. Patel, L. Lata and H. Milosh, *J. Environ. Prot.*, 2017, **8**, 358–379, DOI: [10.4236/jep.2017.83027](https://doi.org/10.4236/jep.2017.83027).
- 168 A. H. Welch, M. S. Lico and J. L. Hughes, *Groundwater*, 1988, **26**, 333–347, DOI: [10.1111/j.1745-6584.1988.tb00397.x](https://doi.org/10.1111/j.1745-6584.1988.tb00397.x).
- 169 A. Yadav, P. Sahu, S. Chakradhari, K. P. Rajhans, S. Ramteke, N. Dahariya, K. S. Patel and G. Agnihotri, *J. Environ. Prot.*, 2016, **7**, 52–59, DOI: [10.4236/jep.2016.71005](https://doi.org/10.4236/jep.2016.71005).
- 170 FAO (Food and Agricultural Organization), *Water quality guidelines for Maximum Crop Production*, Food and Agricultural Organization/UN, Rome, Italy, 1985.
- 171 WHO (World Health Organization), *Arsenic in Drinking-water*, World Health Organization, Geneva, Switzerland, 2011.
- 172 S. A. Ahmad, M. H. Khan and M. Haque, *Risk Manag. Healthc. Pol.*, 2018, **11**, 251–261, DOI: [10.2147/RMHP.S153188](https://doi.org/10.2147/RMHP.S153188).
- 173 X. Cao, Y. Lu, C. Wang, M. Zhang, J. Yuan, A. Zhang, S. Song, Y. Baninla, K. Khan and Y. Wang, *Ecotoxicol. Environ. Saf.*, 2019, **186**, 109628, DOI: [10.1016/j.ecoenv.2019.109628](https://doi.org/10.1016/j.ecoenv.2019.109628).
- 174 R. Saha, N. C. Dey, M. Rahman, P. Bhattacharya and G. H. Rabbani, *Front. Environ. Sci.*, 2019, **7**, 57, DOI: [10.3389/fenvs.2019.00057](https://doi.org/10.3389/fenvs.2019.00057).
- 175 M. Shahid, N. K. Niazi, C. Dumat, R. Naidu, S. Khalid, M. M. Rahman and I. Bibi, *Environ. Pollut.*, 2018, **242(A)**, 307–319, DOI: [10.1016/j.envpol.2018.06.083](https://doi.org/10.1016/j.envpol.2018.06.083).
- 176 A. K. Singh, *Proceedings of 11th National Symposium on Hydrology with focal theme on Water Quality*, Roorkee, 2004.
- 177 A. K. Thakur, V. Gupta, P. Bhattacharya, M. Jakariya and M. T. Islam, *Groundwater Sustain. Develop.*, 2021, **12**, 100504, DOI: [10.1016/j.gsd.2020.100504](https://doi.org/10.1016/j.gsd.2020.100504).
- 178 R. Irunde, J. Ijumulana, F. Ligate, J. Maity, A. Ahmad, J. Mtamba, F. Mtalo and P. Bhattacharya, *Groundwater Sustain. Develop.*, 2022, **18**, 100746, DOI: [10.1016/j.gsd.2022.100746](https://doi.org/10.1016/j.gsd.2022.100746).
- 179 T. A. Abiye and P. Bhattacharya, Arsenic concentration in groundwater: Archetypal study from South Africa, *Groundwater Sustain. Develop.*, 2019, **9**, 100246, DOI: [10.1016/j.gsd.2019.100246](https://doi.org/10.1016/j.gsd.2019.100246).
- 180 I. Quino Lima, M. Ormachea Muñoz, O. E. Ramos Ramos, P. Bhattacharya, R. Quispe Choqu, J. Quintanilla Aguirre and O. Sracek, *Groundwater Sustain. Develop.*, 2019, **8**, 281–293, DOI: [10.1016/j.gsd.2018.11.013](https://doi.org/10.1016/j.gsd.2018.11.013).
- 181 M. I. Litter, M. A. Armienta, R. E. Villanueva-Estrada, E. C. V. Lepori and V. Olmos, *Sci. Rev.*, 2019, **1**, 54–73. <https://www.researchgate.net/publication/337714294>.
- 182 M. I. Litter, A. M. Ingallinella, V. Olmos, M. Savio, G. Difeo, L. Botto, M. Farfan, S. Taylor, S. Frangie, J. Herkovits, I. Schalamuk, M. J. González, E. Berardozzi, F. S. G. Einschlag, P. Bhattacharya and A. Ahmad, *Sci. Total Environ.*, 2019, **676**, 756–766, DOI: [10.1016/j.scitotenv.2019.04.262](https://doi.org/10.1016/j.scitotenv.2019.04.262).
- 183 J. Podgorski and M. Berg, *Science*, 2020, **368**, 845–850, DOI: [10.1126/science.aba1510](https://doi.org/10.1126/science.aba1510).
- 184 H. Rowland, E. Omoregie, R. Millot, C. Jimenez, J. Mertens, C. Baciú, S. Hug and M. Berg, *Appl. Geochem.*, 2011, **26**, 1–17, DOI: [10.1016/j.apgeochem.2010.10.006](https://doi.org/10.1016/j.apgeochem.2010.10.006).
- 185 A. Molinari, L. Guadagnini, M. Marcaccio and A. Guadagnini, *Water Res.*, 2019, **149**, 522–532, DOI: [10.1016/j.watres.2018.09.049](https://doi.org/10.1016/j.watres.2018.09.049).
- 186 F. A. P. Reyes, G. B. Crosta, P. Frattini, S. Basiricò and D. R. Pergola, *J. Hydrol.*, 2015, **529**, 1530–1549, DOI: [10.1016/j.jhydrol.2015.08.029](https://doi.org/10.1016/j.jhydrol.2015.08.029).
- 187 G. Abbas, B. Murtaza, I. Bibi, M. Shahid, N. K. Niazi, M. I. Khan, M. Amjad, M. Hussain and T. Natasha, *Int. J. Environ. Res. Public Health*, 2018, **15**, 59, DOI: [10.3390/ijerph15010059](https://doi.org/10.3390/ijerph15010059).
- 188 T. Itabashi, J. Li, Y. Hashimoto, M. Ueshima, H. Sakanakura, T. Yasutaka, Y. Imoto and M. Hosomi, *Environ. Sci. Technol.*, 2019, **53**, 14186–14193, DOI: [10.1021/acs.est.9b03864](https://doi.org/10.1021/acs.est.9b03864).
- 189 ITRC (Technology and Regulatory Council), *Arsenic: Fate and transport*, Washington, DC, 2017, <https://bcsc-1.itrcweb.org/7-1-fate-and-transport/>.
- 190 A. Violante, in *Advances in agronomy*, ed. D. L. Sparks, Elsevier Inc., Newark (DE), 2013, pp. 111–176, DOI: [10.1016/B978-0-12-405942-9.00003-7](https://doi.org/10.1016/B978-0-12-405942-9.00003-7).
- 191 P. H. Masscheleyn, R. D. Delaune and W. H. Patrick, Effect of redox potential and pH on arsenic speciation and solubility in a contaminated soil, *Environ. Sci. Technol.*, 1991, **25**, 1414–1419, DOI: [10.1021/es00020a008](https://doi.org/10.1021/es00020a008).
- 192 S. M. G. G. Azam, T. C. Sarker and S. Naz, *Toxicol. Mech. Methods*, 2016, **26**, 565–579, DOI: [10.1080/15376516.2016.1230165](https://doi.org/10.1080/15376516.2016.1230165).
- 193 S. B. Singh and P. K. Srivastava, *SN Appl. Sci.*, 2020, **2**, 153, DOI: [10.1007/s42452-019-1932-z](https://doi.org/10.1007/s42452-019-1932-z).
- 194 K. S. Patel, B. L. Sahu, S. Ramteke, N. K. Jaiswal, L. Borgese, A. Gianoncelli and E. Bontempi, *Am. J. Anal. Chem.*, 2015, **6**, 822–829, DOI: [10.4236/ajac.2015.610078](https://doi.org/10.4236/ajac.2015.610078).
- 195 A. Yadav, P. K. Sahu, K. S. Patel, L. Lata, H. Milosh, W. T. Corns, J. Allen and P. Martín-Ramos, *J. Hazard., Toxic Radioact. Waste*, 2020, **24**, DOI: [10.1061/\(ASCE\)HZ.2153-5515.0000478](https://doi.org/10.1061/(ASCE)HZ.2153-5515.0000478).
- 196 M. Kumar, A. L. Ramanathan, M. M. Rahman and R. Naidu, *Sci. Total Environ.*, 2016, **573**, 1103–1114, DOI: [10.1016/j.scitotenv.2016.08.109](https://doi.org/10.1016/j.scitotenv.2016.08.109).
- 197 A. Shrivastava, A. Barla, S. Singh, S. Mandraha and S. Bose, *J. Hazard. Mater.*, 2017, **324**, 526–534, DOI: [10.1016/j.jhazmat.2016.11.022](https://doi.org/10.1016/j.jhazmat.2016.11.022).
- 198 S. Srivastava and Y. K. Sharma, *Environ. Monit. Assess.*, 2013, **185**, 4995–5002, DOI: [10.1007/s10661-012-2920-6](https://doi.org/10.1007/s10661-012-2920-6).
- 199 R. Sharma, A. Yadav, S. Ramteke, K. S. Patel, L. Lata, H. Milosh, W. T. Corns and P. Martín-Ramos, *J. Hazard., Toxic Radioact. Waste*, 2019, **23**, DOI: [10.1061/\(ASCE\)HZ.2153-5515.0000460](https://doi.org/10.1061/(ASCE)HZ.2153-5515.0000460).
- 200 M. A. Ali, A. B. M. Badruzzaman, M. A. Jalil, M. D. Hossain, M. F. Ahmed, A. A. Masud, Md. Kamruzzaman and M. A. Rahman, *Arsenic in Plant-Soil Environment in Bangladesh*, 2003, <https://www.researchgate.net/>



- [publication/237455644_Arsenic_in_Plant_Soil_Environment_in_Bangladesh.](https://pubs.rsc.org/doi/10.1039/C3AY01334A)
- 201 M. B. Arain, T. G. Kazi, J. A. Baig, M. K. Jamali, H. I. Afridi, A. Q. Shah, N. Jalbani and R. A. Sarfraz, *Food Chem. Toxicol.*, 2009, **47**, 242–248, DOI: [10.1016/j.fct.2008.11.009](https://doi.org/10.1016/j.fct.2008.11.009).
- 202 K. D. Brahman, T. G. Kazi, J. A. Baig, H. I. Afridi, A. Khan, S. S. Arain and M. B. Arain, *Chemosphere*, 2014, **110**, 182–189, DOI: [10.1016/j.chemosphere.2013.11.035](https://doi.org/10.1016/j.chemosphere.2013.11.035).
- 203 A. M. Dahal, M. Fuerhacker, A. Mentler, K. B. Karki, R. R. Shrestha and W. E. H. Blum, *Environ. Pollut.*, 2008, **155**, 157–163, DOI: [10.1016/j.envpol.2007.10.024](https://doi.org/10.1016/j.envpol.2007.10.024).
- 204 Y. Gong, Y. Qu, S. Yang, S. Tao, T. Shi, Q. Liu, Y. Chen, Y. Wu and J. Ma, *Environ. Res.*, 2020, **186**, 109525, DOI: [10.1016/j.envres.2020.109525](https://doi.org/10.1016/j.envres.2020.109525).
- 205 M. A. Higgins, M. J. Metcalf and G. A. Robbins, *J. Environ. Qual.*, 2022, **51**, 66–77, DOI: [10.1002/jeq2.20304](https://doi.org/10.1002/jeq2.20304).
- 206 Y. Huang, K. Miyauchi, G. Endo, D. Le Don, N. C. Manh and C. Inoue, *Earth Sci.*, 2016, **75**, 757, DOI: [10.1007/s12665-016-5535-3](https://doi.org/10.1007/s12665-016-5535-3).
- 207 N. Karimi, S. M. Ghaderian and H. Schat, *Int. J. Environ. Sci. Technol.*, 2013, **10**, 743–752, DOI: [10.1007/s13762-013-0227-y](https://doi.org/10.1007/s13762-013-0227-y).
- 208 A. Mathee, T. Kootbodien, T. Kapwata and N. Naicker, *Environ. Res.*, 2018, **167**, 524–527, DOI: [10.1016/j.envres.2018.08.012](https://doi.org/10.1016/j.envres.2018.08.012).
- 209 A. A. Meharg and Md. M. Rahman, *Environ. Sci. Technol.*, 2003, **37**, 229–234, DOI: [10.1021/es0259842](https://doi.org/10.1021/es0259842).
- 210 Z. U. Rehman, S. Khan, K. Qin, M. L. Brusseau, M. T. Shah and I. Din, *Sci. Total Environ.*, 2016, **550**, 321–329, DOI: [10.1016/j.scitotenv.2016.01.094](https://doi.org/10.1016/j.scitotenv.2016.01.094).
- 211 S. Zandsalimi, N. Karimi and A. Kohandel, *Int. J. Environ. Sci. Technol.*, 2011, **8**, 331–338, DOI: [10.1007/BF03326220](https://doi.org/10.1007/BF03326220).
- 212 Y. Zhang, Y. Zhu, S. Zhao, D. Li, H. Xi and Y. Wang, *Environ. Sci. Pollut. Res. Int.*, 2022, **29**, 28957–28972, DOI: [10.1007/s11356-021-18482-3](https://doi.org/10.1007/s11356-021-18482-3).
- 213 Y. Zhou, L. Niu, K. Liu, S. Yin and W. Liu, *Sci. Total Environ.*, 2018, **616–617**, 56–163, DOI: [10.1016/j.scitotenv.2017.10.232](https://doi.org/10.1016/j.scitotenv.2017.10.232).
- 214 A. Dargahi, Reservoir Sedimentation, in *Encyclopedia of Lakes and Reservoirs. Encyclopedia of Earth Sciences Series*, ed. L. Bengtsson, R. W. Herschy and R. W. Fairbridge, Springer, Dordrecht, 2012, DOI: [10.1007/978-1-4020-4410-6](https://doi.org/10.1007/978-1-4020-4410-6).
- 215 A. John and J. S. Leventhal, *Bioavailability of metals. Preliminary Compilation of Descriptive Geoenvironmental Mineral Deposit Models*, U.S. Geological Survey Open-File Report, 1995, vol. 95–831, pp. 10–19, <https://pubs.usgs.gov/of/1995/0831/report.pdf>.
- 216 B. Kominkova, J. Nabelkova and D. Starmanova, in *Urban Environment. Alliance for Global Sustainability Bookseries (Science and Technology: Tools for Sustainable Development)*, ed. S. Rauch and G. Morrison, Springer, Dordrecht, 2012, vol. 19, DOI: [10.1007/978-94-007-2540-9_28](https://doi.org/10.1007/978-94-007-2540-9_28).
- 217 P. O. Lamie, *Proceedings of the annual international conference on soils, sediments, water and energy*, 2012, vol. 17, article 3, <https://scholarworks.umass.edu/soilsproceedings/vol17/iss1/3>.
- 218 S. Hossain, C. T. Oguchi, S. Hachinohe, T. Ishiyama and H. Hamamoto, *Geophys. Res.*, 2014, **16**, 4807.
- 219 N. P. Nikolaidis, G. M. Dobbs, J. Chen and J. A. Lackovic, *Environ. Pollut.*, 2004, **129**, 479–487, DOI: [10.1016/j.envpol.2003.11.005](https://doi.org/10.1016/j.envpol.2003.11.005), PMID: 15016468.
- 220 A. L. Alonso, R. Pérez, C. K. Y. A. Okio and E. Castillo, *J. Environ. Manage.*, 2020, **264**, 110478, DOI: [10.1016/j.jenvman.2020.110478](https://doi.org/10.1016/j.jenvman.2020.110478).
- 221 W. J. Langston, *J. Mar. Biol. Assoc.*, 1980, **60**, 869–881, DOI: [10.1017/S0025315400041953](https://doi.org/10.1017/S0025315400041953).
- 222 M. Kumar and A. L. Ramanathan, *Int. J. Environ. Res. Public Health*, 2018, **15**, 183, DOI: [10.3390/ijerph15020183](https://doi.org/10.3390/ijerph15020183).
- 223 R. Borah, K. Taki, A. Gogoi, P. Das and M. Kumar, *Ecotoxicol. Environ. Saf.*, 2018, **161**, 769–776, DOI: [10.1016/j.ecoenv.2018.06.038](https://doi.org/10.1016/j.ecoenv.2018.06.038).
- 224 A. Gawedzki and K. W. Forsythe, *Int. J. Ecol.*, 2012, 496740, DOI: [10.1155/2012/496740](https://doi.org/10.1155/2012/496740).
- 225 T. J. Whitmore, M. A. Riedinger-Whitmore, J. M. Smoak, K. V. Kolasa, E. A. Goddard and R. Bindler, *J. Paleolimnol.*, 2008, **40**, 869–884, DOI: [10.1007/s10933-008-9204-8](https://doi.org/10.1007/s10933-008-9204-8).
- 226 J. P. Olszewska, A. A. Meharg, K. V. Heal, M. Carey, I. D. M. Gunn, K. R. Searle, I. J. Winfield and B. M. Spears, *Environ. Sci. Technol.*, 2016, **50**, 9044–9052, DOI: [10.1021/acs.est.6b00942](https://doi.org/10.1021/acs.est.6b00942).
- 227 A. E.-M. A. El-Badry and M. M. Khalifa, *Ecologia*, 2017, **7**, 20–28, DOI: [10.3923/ecologia.2017.20.28](https://doi.org/10.3923/ecologia.2017.20.28).
- 228 M. Leivuori and H. Vallius, in: *Arsenic in marine sediments*, ed. K. Loukola-Ruskeeniemi and P. Lahermo, *Arsenic in Finland: Distribution, Environmental Impacts and Risks*, Geological Survey of Finland, 2004, pp. 89–96. (in Finnish with synopsis in English).
- 229 X. Liu, B. Zeng and G. Lin, *Mar. Pollut. Bull.*, 2022, **175**, 113350, DOI: [10.1016/j.marpolbul.2022.113350](https://doi.org/10.1016/j.marpolbul.2022.113350).
- 230 N. Mirlean, V. E. Andrus, P. Baisch, G. Griep and M. R. Casartelli, *Mar. Pollut. Bull.*, 2003, **46**, 1480–1484, DOI: [10.1016/S0025-326X\(03\)00257-1](https://doi.org/10.1016/S0025-326X(03)00257-1).
- 231 Q. Zhou and S. Xi, *Regul. Toxicol. Pharmacol.*, 2018, **99**, 78–88, DOI: [10.1016/j.yrtph.2018.09.010](https://doi.org/10.1016/j.yrtph.2018.09.010).
- 232 Y.-S. Hong, K.-H. Song and J.-Y. Chung, *J. Prev. Med. Public Health*, 2014, **47**, 245–252, DOI: [10.3961/jpmph.14.035](https://doi.org/10.3961/jpmph.14.035).
- 233 K. Khosravi-Darani, Y. Rehman, I. A. Katsoyiannis, E. Kokkinos and A. I. Zouboulis, *Water*, 2022, **14**, 1884, DOI: [10.3390/w14121884](https://doi.org/10.3390/w14121884).
- 234 J. Ng, A. Gomez-Caminero, P. Howe, M. Hughes, E. M. Kenyon, D. R. Lewis, M. R. Moore, A. Aitio and G. Becking, *Environmental Health Criteria 224: Arsenic and Arsenic Compounds Beds*. WHO, ILO & UNEP, Geneva, 2001.
- 235 M. F. Hughes, *Environ. Health Perspect.*, 2006, **114**, 1790–1796, DOI: [10.1289/ehp.9058](https://doi.org/10.1289/ehp.9058).
- 236 K. Welch, I. Higgins, M. S. Oh and C. Burchfiel, *Arch. Environ. Health*, 2013, **37**, 325–335, DOI: [10.1080/00039896.1982.10667586](https://doi.org/10.1080/00039896.1982.10667586).
- 237 K. S. M. Abdul, S. S. Jayasinghe, E. P. Chandana, C. De Silva and P. M. C. Jayasumana, *Environ. Toxicol. Pharmacol.*, 2015, **40**, 828–846, DOI: [10.1016/j.etap.2015.09.016](https://doi.org/10.1016/j.etap.2015.09.016).



- 238 M. Jaishankar, T. Tseten, N. Anbalagan, B. B. Mathew and K. N. Beeregowda, *Interdiscip. Toxicol.*, 2014, 7, 60–72, DOI: [10.2478/intox-2014-0009](https://doi.org/10.2478/intox-2014-0009).
- 239 S. Bhatia, G. Balamurugan and A. Baranwal, *Front. Environ. Sci.*, 2014, 2, 49, DOI: [10.3389/fenvs.2014.00049](https://doi.org/10.3389/fenvs.2014.00049).
- 240 USEPA (US Environmental Protection Agency), *Columbia river basin fish contaminant survey; 1996-1998*, USEPA, Seattle, WA, USA, 2002.
- 241 S. Varol, *J. Water Health*, 2021, 19, 97–107, DOI: [10.2166/wh.2020.107](https://doi.org/10.2166/wh.2020.107).
- 242 R. A. Schoof, J. Eickhoff, L. J. Yost, E. A. Creclius, D. W. Cragin, D. M. Meacher and D. B. Menzel, *Dietary exposure to inorganic arsenic*, in *Arsenic exposure and health effects III*, Elsevier Science Ltd, 1999, pp. 81–88.
- 243 S. S. H. Tao and P. Michael Bolger, *Food Addit. Contam.*, 1999, 16, 465–472, DOI: [10.1080/026520399283759](https://doi.org/10.1080/026520399283759).
- 244 L. M. Del Razo, G. G. Garcia-Vargas, J. Garcia-Salcedo, M. F. Sanmiguel, M. Rivera, M. C. Hernandez and M. E. Cebrian, *Food Chem. Toxicol.*, 2002, 40, 1423–1431, DOI: [10.1016/s0278-6915\(02\)00074-1](https://doi.org/10.1016/s0278-6915(02)00074-1).
- 245 O. Muñoz, D. Vélez and R. Montoro, *Analyst*, 1999, 124, 601–607, DOI: [10.1039/A809426H](https://doi.org/10.1039/A809426H).
- 246 WHO (World Health Organization), *Chemical fact sheets: Arsenic*, World Health Organization, Geneva, 3rd edition incorporating 1st and 2nd addenda, 2008, vol. 1, Recommendations, pp. 306–308b, http://www.who.int/water_sanitation_health/dwq/GDW12rev1and2.pdf.
- 247 SRH (Subsecretaria de Recursos Hídricos de Argentina), *Niveles Gui á de Calidad de Aguas*, 2006, http://hidricos.obraspublicas.gov.ar/calidad_del_agua_actividades.htm, accessed 11 July 2021.
- 248 T. R. Sterner, P. J. Robinson, D. R. Mattie and G. A. Burton, *The Toxicology of chemical mixtures risk assessment for human and ecological receptors*, 2005, p. 101, DOI: [10.21236/ada458544](https://doi.org/10.21236/ada458544).

