



Cite this: RSC Adv., 2018, 8, 20894

2,3-Dihydroquinazolin-4(1H)-one as a privileged scaffold in drug design

Mariateresa Badolato,^{ab} Francesca Aiello^a and Nouri Neamati^b

2,3-Dihydroquinazolin-4-one (DHQ) belongs to the class of nitrogen-containing heterocyclic compounds representing a core structural component in various biologically active compounds. In the past decades, several methodologies have been developed for the synthesis of the DHQ framework, especially the 2-substituted derivatives. Unfortunately, multistep syntheses, harsh reaction conditions, and the use of toxic reagents and solvents have limited their full potential as a versatile fragment. Recently, use of green chemistry and alternative strategies are being explored to prepare diverse DHQ derivatives. This fragment is used as a synthon for the preparation of biologically active quinazolinones and as a functional substrate for the synthesis of modified DHQ derivatives exhibiting different biological properties. In this review, we provide a comprehensive assessment of the synthesis and biological evaluations of DHQ derivatives.

Received 2nd April 2018

Accepted 29th May 2018

DOI: 10.1039/c8ra02827c

rsc.li/rsc-advances

Introduction

Nitrogen-containing heterocyclic scaffolds are quite common fragments in drugs and biologically active compounds.^{1,2} The 2,3-dihydroquinazolin-4(1H)-one (DHQ) is an important

nitrogen-containing motif consisting of a phenyl ring condensed with a six-membered ring with two nitrogen atoms on positions 1 and 3, and a keto group on carbon 4 (Fig. 1). Most of the DHQ derivatives are substituted on the carbon 2 chiral center. Due to their attractive properties, 2-substituted DHQs are becoming a prominent synthetic intermediate for organic chemists and various methodologies are reported in the literature for their preparation as racemic mixtures. Although some asymmetric strategies have been attempted, the aminal chiral center is sensitive to racemization, making it difficult to

^aDepartment of Pharmacy, Health and Nutritional Sciences, University of Calabria, Ed. Polifunzionale, 87036, Arcavacata di Rende, CS, Italy

^bDepartment of Medicinal Chemistry, College of Pharmacy, University of Michigan, North Campus Research Complex, 1600 Huron Parkway, Ann Arbor, MI 48109, USA

Mariateresa Badolato is currently pursuing her PhD degree in Translational Medicine under the supervision of Dr Francesca Aiello and Professor Antonio Garofalo at the University of Calabria, Italy. She received her Master's degree in Pharmaceutical Technology and Chemistry at the University of Calabria in 2013. She worked for one year as a visiting scholar in Professor Neamati's laboratory at the Department of Medicinal Chemistry, College of Pharmacy, University of Michigan. With the project "RYGoldZip" guided by Dr Francesca Aiello, as a member of the team she won the Clinic Center Life Sciences award. Among various projects, she designed and synthesized several ligands for TRPV1 receptors. Her PhD dissertation focuses on the preclinical and mechanistic studies of small-molecule drugs targeting STAT3 and other signaling pathways for the treatment of various cancers.

Dr Francesca Aiello is an Assistant Professor of Medicinal Chemistry at the Department of Pharmacy, Health and Nutritional Sciences, University of Calabria, Italy. She obtained her PhD degree in Pharmaceutical and Technologies Chemistry from the University of Siena in 1998. In 1999, she obtained her qualification as a Pharmacist. She completed a Postdoctoral Research Fellowship under the supervision of Professor Neamati at the Department of Pharmacology and Pharmaceutical Sciences, School of Pharmacy, University of Southern California, Los Angeles, CA. She was a lecturer in Qualitative Analytical Medicinal Chemistry at the University of Calabria. During the past decade she received numerous grants from the University of Salerno, University of Calabria, and University of Siena to develop new anticancer chemotherapeutics, HIV-1 integrase inhibitors, and to synthesize new D3-receptor ligands. Dr Aiello has published 38 peer-reviewed manuscripts, 4 book chapters, and is an inventor in two patents in the area of Medicinal Chemistry. As leader of the project "RYGoldZip", she discovered and formulated a topical product with wound healing properties for diabetic ulcers and she won the Clinic Center Life Sciences award.






Fig. 1 2,3-Dihydroquinazolin-4(1H)-one framework.

synthesize pure enantiomers.³ The aim of the present review is to discuss reported methods for the synthesis of DHQ derivatives, highlighting their evolution towards alternative approaches and enantioselective strategies, and to summarize their use as synthons in organic chemistry and their important biological activities.

2,3-Dihydroquinazolin-4(1H)-one: a privileged scaffold

In 1988, Evans *et al.* introduced for the first time the concept of “privileged structures”. They are useful tools in the field of drug discovery since they represent suitable lead compounds for diverse receptors and the rational optimization of such structures could provide new receptor modulators and potential drugs.⁴ Medicinal chemists exploit the “privileged structures” to synthesize new libraries of compounds based on a central scaffold and screen them against various receptors implicated in different pathways, in some cases yielding biologically active compounds. In this regard, the DHQ core is emerging as

a “privileged scaffold” and a variety of its derivatives, having diverse mechanism of action, are currently used for the treatment of various diseases.^{5–11} A panel of marketed drugs with the DHQ core is shown in Fig. 2. Besides these marketed drugs, a number of new DHQ derivatives have been designed that exhibit a wide range of pharmacological properties. Because of their importance, the synthesis of substituted DHQ derivatives has attracted much attention and different synthetic strategies have been developed. Since the classical protocols involved the use of toxic reagents and solvents in harsh reaction conditions, the evolution towards simple, clean, environmentally benign and high-yielding methods is gaining momentum (Fig. 3).

Conventional procedure: cyclocondensation of anthranilamide and an aldehyde

Although numerous strategies have been developed for the construction of the DHQ core, the most common and simple synthetic route for the preparation of DHQs is the direct cyclocondensation of anthranilamide and an aldehyde (Scheme 1). In the past decades, different catalysts and organic solvents have been used to speed up and improve the general yield of the reaction (Table 1). Regardless of the used catalyst and/or solvent, the most presumed mechanism of cyclocondensation is shown in Scheme 2.¹² The first step involves the nucleophilic attack of the nitrogen of the amino group of the anthranilamide on the carbonyl carbon of the aldehyde, promoted by the catalyst, resulting in the formation of hydroxyl intermediate **I**. Next, the catalyst promotes the formation of the Schiff base (**II**) from **I** through the removal of a water molecule. Finally, the imine undergoes intramolecular cyclization by nucleophilic attack of the nitrogen of the amide group on the imine carbon, to furnish the corresponding DHQ derivative.

A variation of the traditional cyclocondensation of anthranilamide and an aldehyde is represented by the intramolecular cyclization of a Schiff base. As shown in Scheme 3, the nucleophilic attack of the nitrogen of the amide group on the imine carbon leads to the cyclic DHQ derivative.


Basic catalysis

The base-catalyzed cyclocondensation of anthranilamide and an aldehyde was the first proposed strategy for the synthesis of DHQ derivatives. In 1967, Yale *et al.* prepared a new class of DHQs in the presence of 20% aqueous NaOH in refluxing EtOH. These compounds were tested for their inhibition of cell proliferation.¹³ Later, Bonola *et al.* used NaOH or NaOEt in absolute EtOH to prepare DHQs with antibacterial and antifungal activities.¹⁴ Ericsson *et al.* patented DHQ derivatives as anti-fertility agents; these compounds were prepared from anthranilamide and 1-naphthaldehyde in the presence of NaOH in refluxing absolute EtOH.¹⁵ The NaOH-catalyzed reactions gave moderate to high yields (60–85%) in 1 h compared to NaOEt, which took 3–4 h to complete the formation of DHQs (Table 1). Depending on the available aldehyde, the use of a strong base as catalyst was not always successful. Because of this limitation, different catalysts were commonly used for the synthesis of DHQs starting from anthranilamide and an aldehyde.

Dr Nouri Neamati is the John G. Searle Professor of Medicinal Chemistry at the College of Pharmacy, University of Michigan and a member of the UM Translational Oncology Program. He obtained his PhD degree in Biomedical Sciences from the University of Texas Graduate School of Biomedical Sciences and MD Anderson Cancer Center, Houston, Texas in 1995. From 1995 to 2000, he was a postdoctoral and a research fellow at the National Institutes of Health (NIH) in Bethesda, MD. In September 2000, he joined the faculty at the School of Pharmacy, University of Southern California, with a joint appointment at the Norris Comprehensive Cancer Center. Professor Neamati is the recipient of numerous awards, including the NIH Technology Transfer, STOP CANCER, GlaxoSmithKline Drug Discovery Award, several Awards from the US Department of Defense, LUNgevity Discovery Award from the American Lung Association, and the Littlefield-AACR award in Metastatic Colon Cancer Research. He has published over 230 peer-reviewed manuscripts, 20 book chapters, and is an inventor in over 30 patents in the area of drug design and discovery. He was the founding Editor-in-Chief of Current Molecular Pharmacology, and is currently an Associate Editor of the Journal of Medicinal Chemistry as well as Current Anticancer Drug Targets. He has served as an Editorial Advisory Board member of several journals, including Expert Opinion on Drug Discovery, Expert Opinion on Investigational Drugs, and Hormones & Cancer.







Scheme 1 Direct cyclocondensation of anthranilamide and aldehyde under conventional conditions.

Acid catalysis

As shown in Table 1, more frequently, the cyclocondensation of anthranilamide and an aldehyde is performed in acid catalysis. HCl in EtOH¹⁴ or in association with concentrated HNO₃ (ref. 16) was used as catalyst under reflux conditions. As reported, DHQ derivatives were often synthesized in the presence of a catalytic amount of H₂SO₄ (ref. 17) or by using catalysts bearing sulfonic acid functionality. Among them, *p*-toluenesulfonic acid was used in different solvents, including boiling chlorobenzene,¹³ benzene,¹⁸ *N,N*-dimethylacetamide^{19,20} and EtOH.²¹ 2-(*N*-Morpholino)ethanesulfonic acid, commonly used as a buffering agent in biology, also stood out as a mild acid

catalyst for the cyclocondensation of anthranilamide and an aldehyde.²² The use of sulfanilic acid,²³ NaHSO₄,^{24,25} and SOCl₂,²⁶ as a catalyst in EtOH has also been described. Moreover, formic acid¹² as catalyst and solvent, as well as malonic acid²⁷ in aqueous EtOH (50%), were used for the synthesis of DHQs. Furthermore, boric acid (H₃BO₃),²⁸ and propylphosphonic anhydride,²⁹ usually employed in the Fisher indole synthesis and the Pictet–Spengler reaction, efficiently catalyzed the cyclocondensation of anthranilamide and aldehydes.


Iodine and metal salts as catalysts

Molecular iodine (I₂) emerged as a versatile, inexpensive and non-toxic catalyst, which serves as a Lewis acid. It is suitable in the cyclocondensation of anthranilamide and aldehydes, in ionic liquids,^{30,31} or in EtOAc.³² A synthesis in aqueous medium was also attempted. Due to poor solubility of I₂ in water, 1 mol% of I₂ as Lugol's solution (I₂/KI) was used to prepare DHQs.³³ In the presence of I₂ as catalyst, DHQ derivatives were obtained with 66–95% yield, but the reactions required up to 15 h to complete.


Table 1 Conventional conditions for the cyclocondensation of anthranilamide and an aldehyde

Nature catalyst	Catalyst	Conditions	Time	Yield	Ref.
Strong base	NaOH	EtOH, reflux	1 h	60–86%	13–15
	NaOEt	EtOH, reflux	3–4 h	20–85%	14
Strong Brønsted acid	HCl	EtOH, reflux	4 h	10%	14
	Conc. HNO ₃ /HCl	Reflux → rt	5 → 30 min	>98%	16
	H ₂ SO ₄	Solvent-free, MWI	Few min	68–78%	17
Sulfonic acid functionality	PTSA	Chlorobenzene, reflux	1 h	74%	13
		Benzene, reflux	4 h	70%	18
		DMAC, rt/reflux	1–2 h	75–95%	19 and 20
		EtOH, reflux	1 h	70–90%	21
	MES	Aq. EtOH (50%), MWI	5–20 min	83–96%	22
	Sulfanilic acid	Aq. EtOH (50%), 70 °C			23
	NaHSO ₄	EtOH, rt	0.5–5.5 h	91–97%	24
		H ₂ O, grinding rt → 60 °C	0.5–7 h	54–97%	25
	SOCl ₂	EtOH, rt	30–35 min	93–95%	26
	Formic acid	20 °C			12
	Malonic acid	Aq. EtOH (50%), rt	5–37 min	81–98%	27
	H ₃ BO ₃	Solvent-free, 120 °C	5 min	82–90%	28
Organic	T3P®	AcCN, rt	10–15 min	85–94%	29
Lewis acid	I ₂	ILs, 50 °C/80 °C	0.5–10 h	76–99%	30 and 31
		EtOAc, <i>hν</i>	1–15 h	66–93%	32
Lugol's solution	I ₂ /KI	H ₂ O, rt	2–12 h	47–95%	33
Organic Lewis acid	C ₃ Cl ₃ N ₃	AcCN, rt	8–20 min	60–96%	40
	Mn(CH ₃ COO) ₂	EtOH, reflux	5 h		34
	ZrCl ₂	EtOH, rt	9–60 min	80–97%	35
	HgCl ₂	EtOH, 60 °C	1–2 h	88–94%	36
	Cp ₂ TiCl ₂	EtOH, rt	7–9 min	95–98%	37
	InBr ₃	AcCN, rt	10–60 min	75–98%	38
	BiBr ₃	AcCN, rt	30 min	80–95%	39
	Sc(OTf) ₃	EtOH, 70 °C	20–40 min	85–92%	42
		Dry DCM, rt	4–7 h	85–94%	43
		PEG-400, 80 °C	2 h	78–90%	44
	Yb(OTf) ₃	EtOH, 80 °C	2–6 h	>95%	45
		IL, rt	6–8 h	85–96%	46
	Y(OTf) ₃	EtOH, rt	1.5 h	88–99%	47
	NH ₄ Cl	EtOH, rt/reflux	5–120 min	78–98%	48 and 49
	CAN	H ₂ O, rt → 60 °C	1–8 h	62–97%	50
	TBAHS	MeOH, reflux	2 h	64–90%	51





Scheme 2 Presumed mechanism of the cyclocondensation of anthranilamide and an aldehyde.



Scheme 3 Intramolecular cyclization of a Schiff base

Table 2 Reaction conditions for the intramolecular cyclization of a Schiff base

Substrate	Conditions	Time	% Yield	Ref.
Schiff base	NaH, THF, 0 °C → rt	16 h	42–91	52
	N ₂ , EtOH, reflux	6 h	83	53
	Fe ₃ O ₄ NPs, EtOH, reflux	1–3.5 h	94–98	54
	AcOH, reflux	1.5 h	80–92	55 and 56

Transition metal salts are excellent catalysts due to their kinetic stability, low toxicity and intrinsic metallic Lewis acidity. They were used in various organic transformations, including the cyclocondensation of anthranilamide and aldehydes.^{34,40} In particular, the use of transition metal salts reduced reaction time from hours to minutes without decreasing the yield. Compared to conventional Lewis acids, metal triflates are better catalysts in organic synthesis because of their chemical and physical properties, such as moisture and air-stability, recoverability, operational simplicity, and a strong tolerance to oxygen, nitrogen, phosphorus, and sulfur-containing reaction substrates and functional groups.⁴¹ Among them, scandium,^{42–44} ytterbium,^{45,46} and yttrium⁴⁷ triflates were extensively used in various conditions to catalyze the cyclocondensation of anthranilamide and aldehydes, leading to DHQs with high yields (78–99%) in 0.3–8 h.

As shown in Table 1, even ammonium salts stood out as good catalysts for the preparation of DHQ derivatives.^{48–51} The reactions required short time and yields were up to 60%.


Intramolecular cyclization of a Schiff base

The synthesis of DHQs through intramolecular cyclization of a Schiff base was performed under different conditions, providing DHQs with 42–99% yield in 1–16 h (Table 2). While strong basic conditions gave the final product in long time,⁵² the use of molecular nitrogen shortened the completion of the reaction.⁵³ A rapid synthesis was obtained using metal oxide nanoparticles,⁵⁴ and under catalyst-free conditions.^{55,56}

An atom-efficient method: one-pot three-component synthesis


Although the cyclocondensation of anthranilamide and aldehydes is a facile and simple approach to obtain the DHQ core, more advantageous strategies to synthesize DHQ derivatives remain a desired goal in organic chemistry. The first evolution towards a more convenient approach was moving from a two-component reaction (anthranilamide and an aldehyde) to a one-pot three-component reaction (isatoic anhydride, ammonium acetate and an aldehyde) (Scheme 4). Initially, Staiger *et al.* suggested this strategy about half a century ago,⁵⁷ and recently the advantages of multi-component reactions have been highlighted. Compared to conventional synthesis, the one-pot three-component reaction represents an attractive and atom-efficient method to efficiently prepare the DHQ framework.

Multi-component reactions are characterized by (i) atom economy, incorporating all materials used in the process into the final product; (ii) high levels of diversity achieved simply by varying the reaction components; (iii) time-efficiency, since products are formed in a single step, allowing a fast probe of a chemical hypothesis; and (iv) simple experimental procedures, ideally there is no need to isolate the intermediates. Obviously, the adoption of such a strategy minimizes both waste production and cost.⁵⁸ The plausible mechanism of the one-pot three-component synthesis of DHQs is shown in Scheme 5. Initially, the catalyst facilitates the nucleophilic attack of NH₄OAc on the carbonyl carbon of the isatoic anhydride. Nucleophilic addition of ammonium leads to



Scheme 4 Preparation of DHQ through one-pot three-component reaction.





Scheme 5 Plausible mechanism of the one-pot three-component synthesis of DHQ.

intermediate **I** while the following decarboxylation produces anthranilamide (**II**). Then, the reaction proceeds similarly to the cyclocondensation. The catalyst promotes the nucleophilic attack of the amino group of **II** on the carbonyl carbon of the aldehyde, resulting in the formation of the Schiff base (**III**) by removing a water molecule. Finally, the imine intramolecularly cyclizes by nucleophilic attack of the nitrogen of the amide group on the imine carbon, to furnish the corresponding DHQ derivative. Examples of various catalysts and organic solvents used in the reaction of isatoic anhydride, NH_4OAc and an aldehyde are listed in Table 3.

Conventional reaction conditions

In general, the one-pot three-component syntheses of DHQs were performed under acid-catalyzed conditions (Table 3). Brønsted acid catalysts, such as ethylenediamine diacetate⁵⁹ and pentafluorophenylammonium triflate⁶⁰ were used under reflux conditions, giving DHQ derivatives with 86–95% yield in about 10 h or less. Catalysts bearing sulfonic acid functionalities were also widely used in the multi-component

reactions.^{61–66} They significantly reduced the reaction time, generating DHQs with 50–98% yield. Recently, the use of *N*-halo sulfonamides as catalysts, broadly reported in organic synthesis, has been attempted in the one-pot three-component reaction.⁶⁷ Metal-catalyzed multi-component synthesis of DHQ derivatives was also reported.^{68–73} Generally, the use of metals or Lewis acids as catalysts did not lead to great yields (<90%), except when I_2 is used under solvent-free conditions at 115 °C⁶⁹ (Table 3).

Recyclable catalysts

To improve the one-pot three-component reaction using green chemistry, the use of recyclable catalysts was introduced to further minimize waste production (Table 3). Some Lewis acid catalysts had the advantage of being easily recovered after the reaction and recycled several times without considerable loss of reactivity.^{63,64,66,71}

β -Cyclodextrin (β -CD) is a cyclic oligosaccharide, composed of seven glucose units connected “head-to-tail” by 1,4-links. This cyclic heptamer has a truncated cone shape, with



Nature Catalyst	Catalyst	Conditions	Time	% Yield	Ref.
Brønsted acid	EDDA	H ₂ O, reflux	7–10 h	86–93	59
	PFPAT	Toluene, reflux	3 h	88–95	60
Sulfonic acid functionality	PTSA	EtOH or H ₂ O, reflux	3–12 h	50–71	61
	DBSA	H ₂ O, USI	1–2 h	80–91	62
	Co(<i>m</i> -NBS) ₂	EtOH/H ₂ O, reflux	0.5–4 h	82–98	65
	TBBDa or PBBS	EtOH/H ₂ O, reflux	1–3.6 h	60–95	67
	Ce(SO ₄) ₂ ·4H ₂ O	Solvent-free, 120 °C	30–50 min	85–97	63
Sulfonic acid functionality/ recyclable	Al(MS) ₃ ·4H ₂ O	EtOH/H ₂ O, reflux	0.5–6.5 h	60–96	64
	Cu[C ₆ H ₅ SO ₃] ₂ ·6H ₂ O	EtOH/H ₂ O, reflux	0.5–6 h	71–95	66
Lewis acid	I ₂	EtOH or H ₂ O, reflux	0.5–10 h	56–95	68
		Solvent-free, 115 °C	4–25 min	94–98	69
Lewis acid/recyclable	SrCl ₂ ·6H ₂ O	EtOH/H ₂ O, reflux	0.5–6 h	42–94	70
	Cu(OTf) ₂	Toluene, reflux	12–18 h	50–85	72
	Yb(OTf) ₃	DMSO, 90 °C	15 h	36–41	73
	Ga(OTf) ₃	EtOH, 70 °C	35–60 min	71–91	71
	β-CD	H ₂ O, 60 °C/reflux	1.5–5 h	78–92	76–78
Recyclable	β-CD-SO ₃ H	H ₂ O, rt → 50 °C	25 min	80–97	79
	Starch solution	EtOH, 70 °C	4–8 h	73–94	80
Heterogeneous/recyclable	Starch sulfate	Solvent-free, 100 °C	5–55 min	75–96	81
	KAl(SO ₄) ₂ ·12H ₂ O	EtOH, reflux	4–6 h	70–83	82
	Citric acid	H ₂ O, 80 °C	1–7 h	50–94	83
	VB ₁	EtOH, reflux	2–6 h	75–94	84
	Amberlyst-15	Solvent-free, MWI	3–7 min	69–87	85
	Fe ₃ O ₄ NPs	H ₂ O, reflux	1.5–6 h	51–88	88
	Al/Al ₂ O ₃ NPs	Solvent-free, 115 °C	8–30 min	65–98	89
	CuO NPs	EtOH/H ₂ O, reflux or USI	10–30 min	73–95	90
	In ₂ O ₃ NPs	H ₂ O, 80 °C	4 h	78–88	91
	AlN NPs	Drop of H ₂ O, 130 °C	3–7 h	62–73	92
	HAP NPs	H ₂ O, 110 °C	0.7–2 h	80–90	93
	SPNP	H ₂ O, reflux	1–6 h	79–97	95 and 96
	SiO ₂ -FeCl ₃	Solvent-free, 80 °C	9 min to 2 h	45–91	97
	Fe ₃ O ₄ -SBA-15	EtOH, reflux	1.5–4 h	65–78	98
	Solid acidic Catalyst	Titanium-SiO ₂	H ₂ O, 100 °C	2–8 h	86–95
SSA		EtOH, reflux	3–7 h	73–92	100
		H ₂ O, 80 °C or solvent-free	3–6 h	70–86	101
SBSSA		EtOH, 80 °C	0.5–4 h	75–90	102
SBNPSA		EtOH, reflux	2.5–3 h	81–88	103
LPCAHS-SiO ₂		H ₂ O, 80 °C	1.5–4 h	72–95	104
MCM-41-SO ₃ H		Solvent-free, 115 °C	4–20 min	75–98	105
SPC		Solvent-free, 70 °C	2.5–3.5 h	78–86	106
H ₃ BO ₃ -MCM-41		Solvent-free, 80 °C	0.3–1 h	76–94	107
Cellulose-H ₃ BO ₃		Solvent-free, rt	3–40 min	79–92	108
Al(H ₂ PO ₄) ₃		Solvent-free, 100 °C	9–17 min	80–93	109
H ₃ PO ₄ -Al ₂ O ₃		Solvent-free, 100 °C	4min to 3 h	70–90	110
PTA-DEAEC		EtOH, reflux	5–7 h	60–91	111
Copolymer-PTSA		EtOH, reflux	5–7 h	70–94	112
Heterogeneous/recyclable		Montmorillonite K-10	EtOH, reflux	0.5–7 h	70–95
	Cu-CNTs	Solvent-free, MWI	5–23 min	87–99	116
	Co-CNTs	Solvent-free	5–20 min	85–96	117
	Co-MWCNTs	EtOH, USI	6–20 min	76–97	118
	Pt-MWCNTs	EtOH, USI	8–20 min	88–96	119
Acid-surfactant-Combined	732-resin	EtOH/H ₂ O, 90 °C	0.5–5 h	85–95	122
	La ³⁺ /4 Å	AcCN, reflux	24 h	45–95	123
	Zn(PFO) ₂	EtOH/H ₂ O, reflux	6–7 h	77–86	120
	TA-SDS	EtOH/H ₂ O, grinding, rt	1 min to 24 h	87–93	121

as a catalyst for a variety of organic reactions, providing a microenvironment whereby it catalyzes reactions through the formation of non-covalent interactions.⁷⁵ β -CD is a suitable catalyst in the one-pot three-component synthesis of various DHQs producing 78–92% yield in aqueous media.^{76–78} DHQs

were also obtained in less than half an hour in the presence of an inexpensive, safe and recyclable sulfonic acid-functionalized β -CD (β -CD-SO₃H) as an efficient catalyst in green media.⁷⁹

Starch is a renewable, biodegradable, and relatively inexpensive material. A starch aqueous solution in EtOH was employed as safe, non-toxic and reusable catalyst for the preparation of DHQ derivatives with 73–94% yields.⁸⁰ Starch sulfate also reduced the reaction time to less than 1 hour.⁸¹

Furthermore, alum (KAl(SO₄)₂·12H₂O),⁸² citric acid,⁸³ and thiamine hydrochloride (VB₁),⁸⁴ catalyzed the one-pot three-component reaction in aqueous media. The protocols were longer (up to 7 h) but reported to be environmentally safe.

Amberlyst-15 is a cationic exchange resin. These types of resins, especially the macroporous ones, are recyclable catalysts for various organic syntheses, including the preparation of DHQs. Amberlyst-15 presents several advantages over conventional catalyst with respect to corrosion, product recovery, and selectivity.⁸⁵

Heterogeneous and reusable catalysts

Recently, the development of heterogeneous organic reactions has been gaining popularity. They are characterized by ease of handling, separation, recycling, and environmentally safe disposal.⁸⁶ In the last decade, the field of nanoscience and nanotechnology has had tremendous growth. Nanoparticles (NPs) are defined as materials having 1–50 nm diameter, a size range where metals can show size-dependent properties. Because of their interesting structures and high catalytic activities, due to the wide surface/volume ratio that provides many active sites per unit area, NPs and particularly magnetic particles have emerged as useful heterogeneous catalysts in terms of selectivity, reactivity, and improved yields of products. In addition, the magnetic properties of NPs make complete recovery of the catalyst possible by means of an external magnetic field.⁸⁷ As reported in Table 3, some metal and metal oxide NPs, exhibiting high surface/volume ratio, quantum size and quantum tunnel effects, efficiently catalyzed the one-pot three-component synthesis of DHQs, in 0.5–6 h.^{88–91} Nano-sized aluminum nitride, a non-toxic, low-cost and highly pure powder, was used as a solid source of ammonia in the multi-component synthesis of DHQs.⁹² Hydroxyapatite (HAP) NPs show ion-exchange ability, adsorption capacity, and acid–base properties. Because of their higher surface areas and lower particle size, HAP NPs provide greater catalytic activity in the synthesis of DHQs.⁹³ Although NPs have advantages such as simplified isolation of the product, easy recyclability and recovery of the catalyst, the naked NPs could aggregate into large clusters, limiting their use. This problem could be solved by immobilizing the NPs on mesoporous substrates characterized by large surface area, high chemical and thermal stability and good compatibility, such as silica, polymers and carbon.⁹⁴ The integration of mesoporous silica with magnetic NPs is certainly of great interest for practical applications. As shown in Table 3, different silica-supported NPs were used as catalysts for the one-pot three-component reaction, leading to 45–97% yield in 1–8 h.^{95–99} The use of solid acids, as non-toxic, low-cost and

reusable catalysts, has also emerged in the synthesis of DHQs (Table 3). Among them, several acids bearing sulfonic moiety supported on silica catalyzed the multi-component synthesis under different conditions, providing DHQs with >70% yield in 0.5–7 h.^{100–104} MCM-41 is another type of ordered mesoporous silica material. Various acid-functionalized MCM-41,^{105–107} cellulose,¹⁰⁸ and alumina,^{109,110} were used as solid recyclable acidic catalysts for the one-pot three-component synthesis of DHQs, under solvent-free conditions, affording 70–98% yield in less than 1 hour. On the other hand, solid acids used in refluxing EtOH lengthened the reaction time to 5–7 hours.^{111,112} Montmorillonite K-10 is one of the most important smectites, a phyllosilicate mineral species, used as a catalyst in organic synthesis. It is a clay with both Brønsted and Lewis acid sites, with a high cation-exchange capacity. Montmorillonite K-10 is considered a solid acid that acts as heterogeneous catalyst for diverse syntheses, including the one-pot three-component cyclocondensation, and can be easily removed from the reaction mixture.^{113,114} Other interesting solid supports used in heterogeneous catalysis are carbon nanotubes (CNTs), due to the porosity, inertness, and low interactions between catalyst and support and good mechanical strength.¹¹⁵ The deposition of metal NPs on the external surface of CNTs and multi-walled CNTs (MWCNTs) are attractive for catalysis, since they increase the reactive surface area. As reported in Table 3, CNTs supporting transition metals efficiently catalyzed the one-pot three-component synthesis of DHQs, obtained with 76–99% yield in less than half an hour.^{116–119} About the same yield was obtained using acid-surfactant-combined catalysts.^{120,121} A strong acidic cation-exchange resin (732-resin),¹²² and a 4 Å molecular sieve modified with lanthanum(III) (La³⁺/4 Å),¹²³ were employed as heterogeneous and recyclable catalysts for the multi-component synthesis of DHQs.

Greener and convenient approaches to obtain DHQ derivatives

In the new century, there has been an increasing demand for the development of sustainable chemistry. In 1998, Anastas and Warner published the “Twelve Principles of Green Chemistry”,¹²⁴ whose main purpose was the pollution prevention. In this regard, green chemistry was committed to (i) decrease pollution-generating chemicals; (ii) limit the use of dangerous chemicals and exhaustible feedstock materials and scarce resources; and (iii) reduce the harmful effects of final products.^{125–127} The development of cleaner and safer chemical processes started with the use of heterogeneous and recyclable catalysts or alternative solvents, which are not volatile, flammable or toxic. Then, it moved to performing reactions under catalyst- and/or solvent-free conditions. Finally, it incorporated the replacement of conventional thermal equipment by non-conventional sources, such as microwave or ultrasound irradiations.

Heterogeneous/recyclable catalysts

As shown in Table 4, recyclable catalysts, such as β -CD,⁷⁷ amberlyst-15,¹²⁸ and cerous methanesulfonate [Ce(MS)₃],¹²⁹



Table 4 Catalysts for green chemistry approaches to synthesize various DHQs

Nature catalyst	Catalyst	Conditions	Time	% Yield	Ref.
Recyclable	β -CD	H ₂ O, 60 °C	1.5 h	78–92	77
	Amberlyst-15	CH ₃ CN, 80 °C	10–30 min	96–98	128
	Ce(MS) ₃	H ₂ O, grinding technique	0.2–3 h	84–94	129
Solid acid	Sulfamic acid	H ₂ O, 60 °C or MeOH, rt	15–180 min	57–95	130
	H ₂ SO ₄ -SiO ₂	rt	0.2–5 h	93–97	131
	PPA-SiO ₂	Solvent-free, rt	1.5–4 h	89–93	132 and 133
	Amberlyst-15 and SiO ₂ -HClO ₄	CH ₃ CN, 80 °C	45–120 min	55–90	134
	SiO ₂ -ZnCl ₂	Solvent-free, 100 °C	6–80 min	51–95	135
	CAN-SiO ₂	CH ₃ CN, rt	10–50 min	78–94	136
	Boehmite-SSA	EtOH, 80 °C	35–130 min	85–96	137
	Boehmite-Si-DSA	EtOH, reflux	30–190 min	92–98	138
	Montmorillonite KSF	CH ₃ CN, rt	20–50 min	90–99	139
	HCNC-4	CH ₃ CN, rt	15–30 min	82–99	140
	Cellulose-SO ₃ H	CH ₃ CN, rt	40–60 min	77–92	141
	Wang-OSO ₃ H	H ₂ O, 100 °C	0.4–1.1 h	78–88	142
	Fe ₃ O ₄ -SA-PPCA	EtOH, 80 °C	30–80 min	91–95	143
	MNPs-PSA	H ₂ O, 70 °C	25–170 min	71–97	144
	SuSA	H ₂ O, 70 °C	48–60 min	86–95	145
	SO ₄ ²⁻ /ZrO ₂	EtOH, reflux	7–160 min	84–96	146
	Zr(DS) ₄	H ₂ O, rt	8–45 min	83–97	147
	<i>p</i> -SAC	H ₂ O, rt	18–90 min	64–94	148
	H ₃ PW ₁₂ O ₄₀	H ₂ O, rt	8–10 min	79–97	149
	H ₃ PW ₁₂ O ₄₀	EtOH/H ₂ O, 80 °C	12–18 h	71–94	140
	SiO ₂ -H ₃ PW ₁₂ O ₄₀	EtOH, reflux	5–40 min	88–98	151
	Poly(VPyPS)-PW	EtOH, USI	6–16 min	74–96	152
	α -Chymotrypsin	EtOH, 60 °C	30–60 min	90–98	153
Organocatalyst heterogeneous/ recyclable	Fe ₃ O ₄ -GO	EtOH, reflux	2.5–5 h	70–80	154
	GO nanosheets	H ₂ O, rt	10–30 min	85–97	155
	Co-CNTs	Solvent-free, MWI	10–35 min	75–98	156
	Ag-CNTs	EtOH, USI	5–21 min	86–97	157
	CuCl ₂ /Fe ₃ O ₄ -TEDETA	EtOH, 80 °C	25–100 min	94–98	158
	MCM-41-dtz-Ni	PEG-400, rt	10–35 min	90–98	159
	Fe ₃ O ₄ /TiCl ₂ /cellulose	EtOH, rt	6–15 min	79–96	160
	Sc(OTf) ₃	PEG-400, 80 °C	2 h	78–90	161

have been used for the cyclocondensation of anthranilamide and aldehydes. In general, the use of heterogeneous and reusable catalysts afforded high yields, up to 99%, in very short times. A number of solid acid catalysts, on various supports, such as silica,^{130–138} and clay,^{139,140} as well as different catalysts bearing sulfonic acid groups and anchored on cellulose,¹⁴¹ Wang resin,¹⁴² NPs,^{143,144} and other supports,^{145,146} were employed. Reusable acid-surfactant-combined catalysts,^{147,148} and heteropolyacids,^{149–152} have also emerged as good catalysts, due to their unique properties like high thermal stability, low cost, ease of preparation and recovery. They efficiently catalyze the synthesis of DHQs in aqueous media in 5–90 min.

Enzymes have received great attention as sustainable and biodegradable catalysts for the synthesis of biologically active compounds. Among them, α -chymotrypsin rapidly catalyzes the cyclocondensation of anthranilamide and aldehydes with 90–98% yield.¹⁵³ Transition metal-based heterogeneous systems were efficiently used as recyclable catalysts for the cyclocondensation of anthranilamide and aldehydes, yielding > 70% DHQs.^{154–161}

Alternative solvents

The choice of the solvent for a desired chemical process can have profound economic and environmental consequences. For this reason, there has been significant interest in using alternative “clean” solvents, mostly aqueous media and ionic liquids. Water is readily available, cheap, non-toxic, non-flammable and is very attractive from both an economical and environmental point of view.¹⁶² The use of ionic liquids as reaction media and catalysts also gives a solution to solvent emission and catalyst-recycling problems. Ionic liquids present many important features, such as negligible vapor pressure, non-inflammability, immiscibility with non-polar solvents, reasonable thermal and chemical stability and recyclability.¹⁶³

Various catalysts were used in aqueous media to synthesize DHQ derivatives, both from the cyclocondensation of anthranilamide and aldehydes, and the one-pot three-component reaction. A hydrotropic solution¹⁶⁴ and a deep eutectic solvent¹⁶⁵ were used under catalyst-free conditions (Table 5). Ionic liquids were also widely employed for the synthesis of DHQ derivatives, in different reaction conditions (Table 6). Imidazolium-based ionic liquids^{30,31,46,166–171} afforded DHQ derivatives with 70–99% yield, in 0.5–10 h unless used under



Table 5 Alternative solvents: aqueous media for greener approaches to synthesis of DHQs

Nature solvent	Solvent	Conditions	Time	% Yield	Ref.
Aqueous medium	Aq. EtOH (50%)	MES, MWI	5–20 min	83–96	22
		Sulfanilic acid, 70 °C			23
		Malonic acid, rt	5–37 min	81–98	27
	EtOH/H ₂ O	Al(MS) ₃ ·4H ₂ O, reflux	0.5–6.5 h	60–96	64
		Co(<i>m</i> -NBS) ₂ , reflux	0.5–4 h	82–98	65
		Cu[C ₆ H ₅ SO ₃] ₂ ·6H ₂ O, reflux	0.5–6 h	71–95	66
		TBBDA or PBBS, reflux	1–3.6 h	60–95	67
		SrCl ₂ ·6H ₂ O, reflux	0.5–6 h	42–94	70
		CuO NPs, reflux or USI	10–30 min	73–95	90
		Zn(PFO) ₂ , reflux	6–7 h	77–86	120
		TA-SDS, grinding, rt	1 min–24 h	87–93	121
		732-resin, 90 °C	0.5–5 h	85–95	122
		H ₃ PW ₁₂ O ₄₀ , 80 °C	12–18 h	71–94	150
	EtOH or H ₂ O	PTSA, reflux	3–12 h	50–71	61
		I ₂ , reflux	0.5–10 h	56–95	68
	H ₂ O	NaHSO ₄ , grinding	0.5–7 h	54–97	25
		I ₂ /KI, rt	2–12 h	47–95	33
		CAN, rt → 60 °C	1–8 h	62–97	50
		EDDA, reflux	7–10 h	86–93	59
		DBSA, USI	1–2 h	80–91	62
		β-CD, 60 °C/reflux	1.5–5 h	78–92	76–78
		β-CD-SO ₃ H, rt → 50 °C	25 min	80–97	79
		CA, 80 °C	1–7 h	50–94	83
		Magnetic Fe ₃ O ₄ NPs, reflux	1.5–6 h	51–88	88
		In ₂ O ₃ NPs, 80 °C	4 h	78–88	91
		HAP NPs, 110 °C	0.7–2 h	80–90	93
		SPNP, reflux	1–6 h	79–97	95 and 96
		Titanium-SiO ₂ , 100 °C	2–8 h	86–95	99
		SSA, 80 °C	3–6 h	70–86	101
		LPCAHS-SiO ₂ , 80 °C	1.5–4 h	72–95	104
		Ce(MS) ₃ , grinding	0.2–3 h	84–94	129
		Sulfamic acid, 60 °C or MeOH, rt	15–180 min	57–95	130
		Wang-OSO ₃ H, 100 °C	0.4–1.1 h	78–88	142
		MNPs-PSA, 70 °C	25–170 min	71–97	144
		SuSA, 70 °C	48–60 min	86–95	145
		Zr(DS) ₄ , rt	8–45 min	83–97	147
		<i>p</i> -SAC, rt	18–90 min	64–94	148
		H ₃ PW ₁₂ O ₄₀ , rt	8–10 min	79–97	149
		GO nanosheets, rt	10–30 min	85–97	155
Hydrotropic solution	NaPTS (50%)	Catalyst-free, 60 °C	50–95 min	78–95	164
Deep eutectic solvent	L-(+)-TA-DMU	Catalyst-free, 90 °C	4 h	79	165

microwave irradiation (MWI).¹⁷² The triazolium-based reactions take less than half an hour to complete the one-pot three-component reaction.¹⁷³ Ionic liquids bearing sulfonic acid functionality were used under solvent-free conditions and rapidly catalyzed the synthesis of DHQs.^{174,175} Although basic ionic liquids^{171–178} and a glycerol based ionic liquid with a boron core¹⁷⁹ were also used under solvent-free conditions, they catalyzed the reaction in a longer time (10–90 min).

Catalyst- and solvent-free reactions

The challenge for a sustainable environment requires the development of greener and cleaner chemical processes that can avoid the use of harmful solvents and catalysts. In this sense, new strategies have been developed to synthesize DHQ derivatives in solvent- and/or catalyst-free conditions (Table 7). The cyclocondensation of anthranilamide and aldehydes was

performed in the presence of sodium dihydrogen phosphate (NaH₂PO₄),¹⁸⁰ acidic catalysts,^{181–183} and various heterogeneous systems, under solvent-free conditions, leading to various DHQs with 50–98% yield in a few minutes. On the contrary, a simple and environmentally benign procedure, without any catalyst, lengthened the reaction time to hours, although producing similar yield.^{184–192} The one-pot three-component reaction was also efficiently performed under solvent-free conditions in the presence of heterogeneous catalysts using conventional heating or MWI. The absence of catalyst again afforded 74–97% yield but in longer reaction times (1–6 h).^{193–197} Urea or thiourea was used in the multi-component synthesis of DHQs as the ammonia surrogate under catalyst-free conditions.¹⁹⁸ A very efficient and fast synthesis of DHQs was attempted under both catalyst- and solvent-free conditions (>90% yield, in 3–10 min).^{199,200}



Table 6 Alternative solvents: ionic liquids for green chemistry approaches to synthesize various DHQs

Ionic liquid	Structure	Conditions	Time	% Yield	Ref.
[bmim ⁺][BF ₄ [−]]		I ₂ , 50 °C I ₂ , 80 °C Catalyst-free, 70 °C	4–10 h 0.5–1 h 1.5–2.5 h	76–98 90–99 78–94	30 31 166
[bmim ⁺][Br [−]]		I ₂ , 80 °C Yb(OTf) ₃ , rt Catalyst-free, 80 °C	7–10 h 6–8 h	82–91 85–96	167 46 168
[bmim ⁺][PF ₆ [−]]		Solvent-free, 75 °C	35–75 min	77–94	169
[bmim ⁺][HSO ₄ [−]]		H ₂ O, reflux	3–4 h	70–85	170
[msim ⁺][HSO ₄ [−]]		EtOH/H ₂ O, reflux	25–45 min	80–95	171
[bdbim ⁺][Br [−]]		MWI, 100 °C	3–7 min	89–97	172
IPTT		EtOH, 30 °C	1–24 min	10–96	173
[PY(CH ₂) ₄ SO ₃ H][HSO ₄ [−]]		A300SiO ₂ solvent-free, 110 °C	10–14 min	81–90	174
[tpps ⁺][TS [−]]		Solvent-free, 80 °C	4–30 min	85–96	175
TBAB		Solvent-free, 100 °C Solvent-free, 105 °C or H ₂ O, 70 °C	45–90 min 25–75 min	72–84 67–91	176 177
Basic ionic liquid		rt	3–5 h	80–90	178
H[Gly ₂ B]		Solvent-free, 60 °C	10–55 min	83–92	179

New energy sources: microwave and ultrasound irradiations

MWI in organic synthesis is commonly used because it facilitates heat transfer better in chemical reactions. The efficiency of MWI heating results in a dramatic reduction in reaction times to minutes as compared to conventional heating methods taking several hours. From an economic and environmental viewpoint, the use of MWI provides unique chemical processes, characterized by enhanced reaction rates, sometimes higher yields, greater selectivity, and ease of manipulation.^{201,202} Previously, efforts were made to synthesize DHQs under MWI

(Table 8). Among them, the cyclocondensation of anthranilamide, under acidic catalysis,²⁰³ as well as the one-pot three-component reaction, in the presence of L-proline in water.²⁰⁴ The ultrasound irradiation (USI)-assisted reactions have also become increasingly popular in organic synthesis. Due to faster reactions, MWI and USI allow the elimination or minimization of side products formation. They are frequently used in the pharmaceutical industry and may pave the way towards a greener and more sustainable approach to chemical synthesis.²⁰⁵ A variety of organic reactions were carried out



Table 7 Catalyst- and/or solvent-free synthesis of DHQs

Reaction	Catalyst	Conditions	Time	% Yield	Ref.
Cyclocondensation	H ₂ SO ₄	Solvent-free, MWI	Few min	68–78	17
	H ₃ BO ₃	Solvent-free, 120 °C	5 min	82–90	28
	H ₃ BO ₃ -MCM-41	Solvent-free, 80 °C	20–65 min	76–94	107
	Cellulose-H ₃ BO ₃	Solvent-free, rt	3–40 min	79–92	108
	Co-CNTs	Solvent-free, rt	5–20 min	85–96	117
	PPA-SiO ₂	Solvent-free, rt	1.5–4 h	89–93	132
	Co-CNTs	Solvent-free, MWI	10–35 min	75–98	156
	H ₃ BO ₃ or NaH ₂ PO ₄	Solvent-free, 120 °C	3–15 min	50–92	180
	Citric acid	Solvent-free, grinding	10–20 min	65–98	181
	Lemon juice	Solvent-free, rt	7–10 min	82–85	182
	Lactic acid	Solvent-free, 70 °C	0.3–6 h	80–92	183
	Catalyst-free	L-(+)-TA-DMU, 90 °C	4 h	79	165
		AcOH, rt	4 h	35–49	184
		Dry MeOH, reflux	5 h	55–73	185
		TFE, reflux	0.4–56 h	40–98	186
		DCM, reflux	2–3 d	98	187
		PEG-400, 100–110 °C	4–10 h	78–92	188
		Glycerol, 80 °C	1–4 min	87–95	189
		H ₂ O, 90 °C	1–6 h	67–94	190 and 191
		H ₂ O, reflux	0.5–27 h	73–99	192
One-pot Three-component	Ce(SO ₄) ₂ ·4H ₂ O	Solvent-free, 120 °C	30–50 min	85–97	63
	I ₂	Solvent-free, 115 °C	4–25 min	94–98	69
	Starch sulfate	Solvent-free, 100 °C	5–55 min	75–96	81
	Amberlyst-15	Solvent-free, MWI	3–7 min	69–87	85
	Al/Al ₂ O ₃ NPs	Solvent-free, 115 °C	8–30 min	65–98	89
	SiO ₂ -FeCl ₃	Solvent-free, 80 °C	9–120 min	45–91	97
	SSA	Solvent-free, rt	3–6 h	70–86	101
	MCM-41-SO ₃ H	Solvent-free, 115 °C	4–20 min	75–98	105
	SPC	Solvent-free, 70 °C	2.5–3.5 h	78–86	106
	Cellulose-H ₃ BO ₃	Solvent-free, rt	3–40 min	79–92	108
	Al(H ₂ PO ₄) ₃	Solvent-free, 100 °C	9–17 min	80–93	109
	H ₃ PO ₄ -Al ₂ O ₃	Solvent-free, 100 °C	4–180 min	70–90	110
	Cu-CNTs	Solvent-free, MWI	5–23 min	87–99	116
	SiO ₂ -ZnCl ₂	Solvent-free, 100 °C	6–80 min	51–95	135
	Catalyst-free	NaPTS (50%), 60 °C	3–5 h	80–90	178
		AcOH, reflux	1–2.5 h	79–97	193
		TFE, reflux	3 h	80–97	194
		[bmim ⁺][PF ₆ ⁻], 75 °C	15–55 min	74–93	195
		PEG-400, 120–125 °C	1–6 h	80–97	196
		Glycerol, 80 °C	2 h	88–90	197
		EtOH, reflux	6 h	80–92	198
		Solvent-free, 70 °C	10 min	87–96	199
		Solvent-free, 120 °C or MWI	3 min	90–97	200

within short times under USI, including the synthesis of DHQs, in the presence of amberlyst-15 (ref. 206) and in other different conditions (Table 8).

Alternative synthetic strategies

Although the cyclocondensation of anthranilamide and the one-pot three-component reaction of isatoic anhydrides, ammonium acetates and aldehydes are the main ways to synthesize DHQ derivatives, other synthetic strategies have also been developed (Table 9).

Cyclocondensation of anthranilamide and different substrates

Anthranilamide was the most common starting material for the preparation of DHQs. Other than aldehydes, other substrates

were used for the cyclocondensation with anthranilamide in different conditions to give DHQ derivatives (Scheme 6).

First, the cyclocondensation of anthranilamide and oxo-compounds, such as benzil,²⁰⁷ 2-oxo(alkyl)acetates,²⁰⁸ and 4'-bromoacetophenone,²⁰⁹ were attempted to give DHQs in different conditions. The obtained 2,2-disubstituted DHQ was then cleaved in ethanolic/methanolic hydroxide to give the respective derivative. Alcohols were used in a ruthenium-catalyzed cyclocondensation with anthranilamide.²¹⁰ Cyclocondensations of anthranilamide and *gem*-dibromomethylarenes, as aldehyde equivalents, were performed in the presence of potassium *tert*-butoxide (*t*-BuOK), in anhydrous pyridine and *N,N*-dimethylformamide (DMF).²¹¹ Anthranilamide also reacts with dicyanoepoxide to give DHQs in refluxing CH₃CN.²¹²




Table 8 New energy sources for greener approaches to synthesize DHQs

Reaction	Energy source	Catalyst	Conditions	Time	% Yield	Ref.
Cyclocondensation	MWI	H ₂ SO ₄	Solvent-free	Few min	68–78	17
		MES	Aq. EtOH (50%), 600W	5–20 min	83–96	22
		Cu-CNTs	Solvent-free, 300W	5–23 min	87–99	116
		Catalyst-free	Solvent-free, 300W	3 min	90–97	200
		PTSA	AcOH, 300W	5–20 min	66–95	203
	USI	Poly(VPyPS)-PW	EtOH, rt	6–16 min	74–96	152
		Ag-CNTs	EtOH, 75 °C	5–21 min	86–97	157
		Amberlyst-15	CH ₃ CN, rt	1–5 min	95–98	206
		One-pot three-component	MWI	Amberlyst-15	Solvent-free, 360W	3–7 min
Co-CNTs	Solvent-free, > 500W			10–35 min	75–98	156
Catalyst-free	[bdbim ⁺][Br [−]], 100 °C			45–90 min	72–84	172
Catalyst-free	Solvent-free, 300W			1–6 h	80–97	200
L-Proline	H ₂ O, 100 °C, 250W			7–8 min	84–95	204
USI	DBSA		H ₂ O, 40–42 °C	1–2 h	80–91	62
	CuO NPs		EtOH/H ₂ O, reflux	10–30 min	73–95	90
	Co-MWCNTs		EtOH, 35 kHz, 40 °C	6–20 min	76–97	118
	Pt-MWCNTs		EtOH, 60 °C	8–20 min	88–96	119


Table 9 Reaction conditions for the alternative synthetic strategies

Starting material	Substrate	Conditions	Time	% Yield	Ref.
Anthranilamide	Benzyl	i: ZnCl ₂ , AcOH, reflux	3 h	76	207
		ii: NaOH, EtOH, rt	1 h	88	
	2-Oxo(alkyl)acetate	i: PTSA, toluene, reflux	4–7 h	65–80	208
		ii: KOH, MeOH, rt	24 h	94	
	4'-Bromo acetophenone alcohol	I ₂ , THF, 50 °C	6 h	86	209
		5mol% (PPh ₃) ₃ Ru(CO)H ₂ , 5mol% xantphos, 2.5eq. crononitrile, 20mol% NH ₄ Cl, toluene, N ₂ , reflux	14 h	69–78	210
		<i>t</i> -BuOK, Py/DMF, 80 °C	4–4.5 h	60–90	211
	Gem-dibromomethylarene	CH ₃ CN, reflux	20 h	45–82	212
	Dicyanoepoxide	5mol% Ph ₃ PauNTf ₂ , toluene, 100 °C	12 h	60–97	213
	Terminal alkynes	5mol% PtBr ₂ or Au(PPh ₃)Cl, MeOH, 80 °C	24 h	70–98	214
2-Amino-benzonitrile	Alkyne	5mol% PtBr ₂ or Au(PPh ₃)Cl, MeOH, 80 °C	24 h	70–98	214
		ZnCl ₂ , DMF, reflux	1–24 h	47–88	215–217
	Aldehyde	CHOH, H ₂ O, 80 °C	0.5–2 h	82–96	218
		K ₃ PO ₄ , H ₂ O, 100 °C	8 h	28–80	219
		1,3-Dipropylimidazole, solvent-free, rt	1–2 h	18–98	220
		i: 20% KOH, reflux	7 h	39–49	221
		ii: AcOH, rt	4 h		
		Amberlyst A26 OH, EtOH/H ₂ O, 50–60 °C	2.5–4 h	75–93	222
		TiCl ₄ /Zn, THF, reflux	2 h	79–91	223
		TiCl ₄ /Sm, THF, reflux	2 h	71–92	224
<i>o</i> -Nitrobenzamide	Aldehyde	Fe, AcOH, 115 °C	30 min	73–94	225
		SnCl ₂ ·H ₂ O, EtOH, rt	2–4 h	82–86	226
		SmI ₂ , THF, rt	3–4 h	60–85	227
		SmI ₂ , MeOH, reflux	20 h	69–89	228
	Aldehyde	SmI ₂ , THF/MeOH, rt → reflux	2–4 h	69–88	229
		i: SiCl ₄ , Py, rt	6–24 h	36–81	230
<i>o</i> -Azidobenzamide	Amine	ii: SMEAH, toluene, reflux	24 h		
		iii: ClCO ₂ Et, Py, rt	48 h	46–88	
		i: Triphosgene, dry THF, rt	30 min	72	231
	Aldehyde	ii: 28% aq. NH ₄ OH, THF, rt	2 h	94	
		iii: PTSA, MeOH, rt	2 h	84	
		CuBr, L-proline, DMSO, 100 °C	5 h	85	232
		CuBr ₂ , K ₂ CO ₃ , DMF, 130 °C	4 h	67	233
		NaBH ₄ , diglyme, 85 °C	1 h	50	234
		NaBH ₄ , AcOH, 50 °C	48 h	50–52	235
		NaBH ₄ CN, AcOH, rt	24 h	33	236
2-Halobenzamide	Aldehyde	NiCl ₂ , NaBH ₄ , dry MeOH, rt	0.5–24 h	79–92	237
	Aniline	NH ₄ OAc, AcOH, 50 °C → 90 °C	90–150 min	76–81	238
	Benzyaniline	O ₂ , AcOH, rt	24 h	62–83	239





Scheme 6 Alternative synthetic strategies of cyclocondensation of anthranilamide and different substrates.



Scheme 7 Cyclocondensation of 2-aminobenzonitrile and an aldehyde for the preparation of DHQs.

The direct hydroamination/hydroarylation and double hydroamination of alkynes, followed by the cyclocondensation with anthranilamide was also exploited for the synthesis of DHQ derivatives.^{213,214} These alternative strategies afforded DHQs with 45–97% yield in 1–24 h (Table 9).

Reductive cyclocondensation of 2-aminobenzonitrile/*o*-nitrobenzamide/*o*-azidobenzamide and aldehydes

An alternative method starting from 2-aminobenzonitrile and aldehydes was developed to synthesize DHQs (Scheme 7) and two plausible mechanisms have been proposed. The first was the formation of a Schiff base and subsequent cyclocondensation in the presence of various catalysts (Scheme 8a), yielding 18–98% DHQs in 0.5–8 h (Table 9).^{215–220} The second proposed mechanism was the hydration of the 2-aminobenzonitrile to anthranilamide in different conditions (Scheme 8b) by means of potassium hydroxide (KOH),²²¹ or amberlyst A26 OH.²²²

The reductive cyclization of *o*-nitrobenzamide^{223–227} and/or *o*-azidobenzamide^{228,229} and aldehydes yield DHQ derivatives in the presence of metallic catalysts with 60–94% yield in short reaction times (Scheme 9; Table 9).

Other strategies


Other strategies have been developed to synthesize DHQ derivatives starting from various substrates (Scheme 10) giving 39–94% yield in 0.5–48 h (Table 9). One such method employed

2-aminobenzoic acid as a starting material to prepare DHQ derivatives. The condensation of the 2-aminobenzoic acid with amines in the presence of SiCl_4 led to the corresponding substituted anthranilamides, subsequently reduced with sodium bis(2-methoxyethoxy)-aluminum hydride to *o*-aminobenzylamines. Then, the latter compounds were cyclized into DHQs by means of ethylchloroformate/pyridine.²³⁰ Another strategy converted the 2-aminobenzoic acid in isatoic anhydride by means of triphosgene in dry THF, then converted to anthranilamide using 28% NH_4OH solution. The cyclocondensation with the aldehyde in the presence of *p*-toluenesulfonic acid, in refluxing MeOH, gave the respective DHQ.²³¹ The copper-catalyzed cyclocondensation of 2-halobenzamide and aldehyde in the presence of aqueous ammonia,²³² and 2-halobenzamide with anilines,²³³ also gave DHQs derivatives. DHQs could also be prepared by reduction from 4(3*H*)-quinazolinones (QZs) with NaBH_4 ,^{234,235} or NaBH_4CN .²³⁶ DHQs were also obtained by reductive desulfurization of 2-thioxo-4(3*H*)-quinazolinones with nickel boride, using nickel(II) chloride (NiCl_2) and NaBH_4 .²³⁷ The intramolecular cyclization of 2-phenyl-ethyl anthranilate in the presence of NH_4OAc led to the respective DHQs.²³⁸ The oxidation of benzylamines to the corresponding *N*-benzylbenzaldimines was also investigated and used for the synthesis of DHQ derivatives.²³⁹

Enantioselective synthesis of DHQ derivatives


Most of the reported methods allow the synthesis of DHQ derivatives as racemic mixtures. In certain examples, the (*S*)-enantiomer of DHQs had better antiproliferative activity, compared to the (*R*)-enantiomer, even though the racemic mixture showed a similar potency to the pure *S*-enantiomer.³ Chinigo *et al.* first proved that a (*S*)-enantiomer of DHQ binds to tubulin, showing antiproliferative activity in different cancer cell lines. They obtained the pure (*S*)-enantiomer using trifluoroacetic acid in CH_3CN with 34–86% yield and 79–91% enantiomeric excess percentage (ee%) in 1.5 h.³ Although the





Scheme 8 (a) Plausible mechanism of the cyclocondensation of 2-aminobenzonitrile and an aldehyde through formation of Schiff base. (b) Plausible mechanism of the cyclocondensation of 2-amino-benzonitrile and an aldehyde through hydration to anthranilamide.

enantioselective synthesis of DHQs is difficult due to an unstable aминаl stereo-center that is sensitive to racemization, some asymmetric strategies have been developed to obtain pure



Scheme 9 Reductive cyclocondensation of *o*-nitrobenzamide/*o*-azidobenzamide and an aldehyde.

(*S*)-enantiomers of DHQs (Table 10). Mainly, they consist of the cyclocondensation of anthranilamide and aldehydes in the presence of chiral catalysts that promote the formation of pure enantiomers, or the intramolecular amidation of *N*-Boc imines and anthranilamide (Scheme 11). The use of chiral phosphoric acidic catalysts is common in the enantioselective synthesis of DHQs, although they need a longer time (15–48 h) to complete the reaction. Various chiral phosphoric acids were employed to catalyze the cyclocondensation of anthranilamide and aldehydes,^{240–243} affording 67–99% yield and 26–99 ee%. An amidation of *N*-Boc imines and anthranilamide,^{244,245} gave pure (*S*)-enantiomer in 10–96 ee%. Scandium(III)-catalytic systems were also effectively employed for the enantioselective cyclocondensation of anthranilamide and an aldehyde (Table 10).^{246,247}

DHQ as an intermediate in organic chemistry

In addition to their many significant pharmacological activities, DHQs also play a central role as intermediates in organic synthesis. In particular, they can be easily oxidized to the biologically active QZs. The quinazolinone ring (Fig. 4) is frequently encountered in organic chemistry as well as in medicinal chemistry.^{248–254} The QZ core is present in the structure of numerous natural products, especially alkaloids,^{255,256} and in some drugs,²⁵⁷ exhibiting various pharmacological properties. QZs also represent a privileged scaffold and many protocols are reported in literature for the synthesis of this important synthon. Among them, the dehydrogenation of DHQ derivatives (Scheme 12) has emerged as an easy and fast strategy to prepare QZs under different oxidant conditions (Table 11). Initially, QZs were obtained by dehydrogenation of DHQs by means of ZnCl₂ in the presence of air, with 42% yield and in 10 h.²⁰² A catalyst-free reaction in an open flask in refluxing EtOH was completed in few hours.²⁵⁸ The oxidation of DHQs by the addition of 2,3-dichloro-5,6-dicyano-1,4-benzoquinone also gave the corresponding oxidized derivatives with 83% yield.²³¹ Various oxidizing agents were used for the dehydrogenation of DHQs to efficiently generate QZs, with 26–92% yield.^{259–262} Metal-catalyzed dehydrogenation was also attempted, affording the oxidized derivatives in moderate to good yields (18–85%) but in 16–24 h.^{263–265} Biocatalysis using laccase/*N*-hydroxybenzotriazole gave QZs in 62–87% yield.²⁶⁶

DHQ as a versatile fragment in drug design

The purpose of drug discovery has always been the design and development of “magic bullets” targeting a single key biomolecule in a central pathway of a specific disease. This led to the dominant paradigm “one target, one drug”, which might be inadequate to achieve a therapeutic effect for complex diseases.^{267,268} For this reason, the polypharmacology research and the design of multitarget compounds is considerably emerging,^{269,270} contributing to overcome some of the limitations of classical approach, in term of risks and costs.²⁷¹ DHQ is a versatile fragment that can be easily functionalized at different positions. The introduction of specific moieties in the DHQ nucleus leads to the ability to interact with multiple





2-Aryl DHQ derivatives


Quinolin-5-yl DHQ induced cytochrome *c* mediated apoptosis and autophagy in human leukemia MOLT-4 cells as demonstrated by flow cytometry, microscopy, LC3 immunofluorescence, and western blot analysis.²⁷⁵ Other pathways have been suggested to clarify the antitumor activity of DHQ derivatives. DHQs bearing a phenyl substituent and piperidine/piperazine moiety on C5 and C7 respectively were identified as selective inhibitors of p38 MAP kinase. They efficiently repressed the production of TNF- α in monocyte, THP-1 cells and LPS-stimulated whole blood (IC₅₀ values in the nanomolar range). These analogs had good clearance but low oral bioavailability in rats. However, the introduction of a bulky *t*-butyl substituent on the piperidine nitrogen significantly increased the oral exposure in rats.²⁷⁶ Other DHQ derivatives were discovered by a high-throughput screening as inhibitors of the Hedgehog pathway, involved in embryonic development and oncogenesis. The biochemical mechanism of action of these DHQs was the inhibition of the AAA+ ATPase motor cytoplasmic dynein that converts chemical energy into mechanical force and regulates many cellular processes, including ciliary trafficking, formation of mitotic spindle and organelle transport. These AAA+ ATPase inhibitors could be useful to study cellular processes that require microtubule motor.²⁷⁷ Different targets were proposed to cause the antitumor activity of 2-aryl DHQs. Among them are cathepsin B and H, known to facilitate invasion, angiogenesis and metastasis through degradation of extracellular matrices and are associated with cancer progression.

Some 2-aryl DHQ derivatives were identified, through molecular docking studies, as reversible inhibitors of cathepsin B and H with effective inhibitory constant (K_i) values in the

Table 10 Catalysts for the enantioselective synthesis of DHQs

Catalyst	Structure	Conditions	Time	% Yield	ee%	Ref.
TFA		CH ₃ CN, 45 °C	1.5 h	34–85	79–91	3
C ₈ -TRIP		Toluene, 5 Å MS, –45 °C	24 h	67–94	26–98	240
9-Anthracenyl-TRIP		CHCl ₃ , rt	24–48 h	73–99	90–99	241
		CHCl ₃ , rt	24 h	30–94	10–96	244
(R,R)-PhDAP		1,2-DFB, rt	15 h	99–100	80–86	242
9-Anthracenyl-SPINOL		CHCl ₃ , 3 Å MS, rt	24 h	88–99	59–98	243
BINOL-derived phosphoric acid		CHCl ₃ , –15 °C	24–36 h	60–85	80–96	245
Sc(OTf) ₃ /Pybox		DCM, 4 Å MS, rt	6–48 h	80–94	86–98	246
Sc(OTf) ₃ /L6		DCM, 4 Å MS, rt	48 h	81–94	87–98	247





Scheme 11 Enantioselective synthesis of DHQs.





Fig. 4 Quinazolin-4(3H)-one framework.



Scheme 12 Dehydrogenation of DHQs as intermediate in organic synthesis.

nanomolar range.²⁷⁸ Through virtual and *in vitro* screening, 2-aryl DHQs were found to be potent tankyrase 2 inhibitors (IC_{50} ranging from low micromolar to nanomolar values). Crystal structures of tankyrase 2 with inhibitors showed that they interact with the nicotinamide-binding site of the catalytic domain of the enzyme. These compounds also inhibited Wnt/ β -catenin pathway in a cell-based assay.²⁷⁹ Development of hybrid compounds that incorporate two pharmacophores in a single molecule with improved potency is gaining momentum. A new

class of hybrids was designed and synthesized from DHQ and 3,5-diaryl isoxazoline/isoxazoles, linked through alkane spacers, and their antitumor activity was evaluated. Derivatives with 3C chain spacers showed excellent potency against lung cancer, exhibiting nanomolar IC_{50} values. Flow cytometric analysis revealed G2/M cell cycle arrest, usually associated with the inhibition of tubulin polymerization. Furthermore, these compounds disrupt microtubules, inhibit cyclin B1, CDK1, and induce apoptosis in cancer cells.²⁸⁰ *N*-indolylmethyl substituted spiroindoline-3,2'-DHQ were hypothesized as potential sirtuin inhibitors. Sirtuins, whose family consists of seven members (SIRT1-7), are important targets for cancer therapy, being up-regulated in several types of cancer. In particular, SIRT1 has several substrates, including p53 and NF- κ B, and its inhibition leads to the re-expression of silenced tumor suppressor genes and the subsequent decrease of cancer cell growth. These new DHQ derivatives were obtained through Pd/C–Cu-mediated coupling cyclization and tested *in vitro* using a yeast homologue of mammalian SIRT1, Sir 2 protein, showing dose-dependent inhibition. Molecular docking analysis showed that the benzene ring of the DHQ occupied the deep hydrophobic pocket of the protein, while the NH and the sulfonyl groups form H-bonding interaction with select amino acid residues (Val412 and Gly415).²⁸¹ Over half a century ago, DHQ derivatives bearing a sulfonamide moiety on C7 showed diuretic activity, causing natriuresis and chloruresis and a slight increase of potassium excretion after oral administration.²⁸² Among them, fenquizone is a FDA-approved drug for the treatment of edema and hypertension.^{283–286} The substitution of C8 with N in the benzene ring of DHQ and the introduction of

Table 11 Reaction conditions for the dehydrogenation of DHQs

Catalyst	Conditions	Time	% Yield	Ref.
ZnCl ₂	AcOH, air, reflux	10 h	42	202
Catalyst-free	EtOH, air, reflux	3 h	65	258
DDQ	MeOH, reflux	2 h	83.5	231
MnO ₂	DCM, rt	2 h	39–79	221
MnO ₂	CHCl ₃ , rt	5–20 h	26–75	259
KMnO ₄	DMF, reflux	2–3 h	85–90	191
KMnO ₄	Acetone, rt	1 h		202
KMnO ₄	DMAC, MWI, 210 W	3–5 min	60–92	260
O ₂	AcOH, 150 °C	24 h	46–72	239
SO ₂	DMF/H ₂ O, air or N ₂ , 90 °C	5 h	65–92	261
K ₂ S ₂ O ₈	CH ₃ CN, 90 °C	3–16 h	55–90	262
CuBr	K ₂ CO ₃ , DMSO, air, 130 °C	24 h	18–85	263
FeCl ₃	K ₂ CO ₃ , toluene, 120 °C	16 h	45–85	264
Ph ₃ PAuNTf ₂	Toluene, 100 °C	24 h		265
Laccase/HBT	O ₂ , citrate buffer (pH 4.5), 45 °C	20–30 h	62–87	266






Fig. 5 Pharmacological activities of DHQ derivatives.


a pyridyl moiety on C2 led to 1,2-dihydro-2-(3-pyridyl)-3H-pyrido [2,3-*d*]pyrimidin-4-one derivatives. They also showed diuretic activity.²⁸⁷ Some DHQ derivatives also show broad-spectrum antimicrobial properties. 2-(5-Nitro-2-thienyl) DHQs showed antibacterial activity against *Hemophilus vaginalis* and *Escherichia coli* strains, responsible for bacterial vaginitis, exhibiting low minimal inhibitory concentration (MIC, ranging from 0.4 to 12.5 $\mu\text{g ml}^{-1}$).²⁸⁸ More recently, some spiro-oxindole DHQs have shown significant antibacterial activity against both Gram-positive and Gram-negative bacterial strains, in particular *E. coli*. In order to explain the mechanism of this antibacterial activity, DHQs were docked on *E. coli* biotin carboxylase (EcBC) enzyme. EcBC is a known target of the fatty acid biosynthetic pathway since it catalyzes the ATP-dependent carboxylation of the vitamin biotin. Molecular docking results showed that the DHQ core fit in the hydrophobic enclosure formed by Ile437 and His236, and the different substituents form H-bonds as well as hydrophobic interactions with specific amino acid residues.²⁸⁹

Tuberculosis, an infectious disease caused by *Mycobacterium tuberculosis*, is a worldwide leading cause of death. Multiple antibiotics are needed to treat tuberculosis over a long period of time, but the development of multiple drug-resistant tuberculosis limits complete recovery. Some spiro-DHQ derivatives were tested for their *in vitro* inhibitory activity against *Mycobacterium tuberculosis* H37Rv chorismate mutase, a promising target for the identification of new antitubercular drugs. These derivatives showed a moderate inhibition at relatively high doses (30 μM).²⁹⁰ 2-Aryl DHQs also show moderate to good antifungal activities, in particular against *Candida albicans* and *Aspergillus niger*.⁴³ A class of 2-thioxo DHQ derivatives showed an excellent antimicrobial activity by inhibiting myeloperoxidase. This

enzyme plays an important role in host defense and contributes to inflammation. They reversibly inhibited myeloperoxidase, exhibiting IC_{50} values in the nanomolar range.²⁹¹ 4-Alkynyl-3,4-dihydro-2(1H)-quinazolinones, bearing a second substituent at position 4, such as cycloalkyl, aryl and trifluoromethyl were identified as potent HIV-1 non-nucleoside reverse transcriptase inhibitors, inhibiting wild-type and various mutant forms of HIV-1. These compounds also showed a good oral bioavailability.^{292–294} 2-Aryl DHQ derivatives, with or without substituents on N1 and N3, were found to interact with human serum albumin (HSA), using fluorescence spectroscopy. The therapeutic effects of drugs depend on their absorption, distribution, metabolism and excretion, and can be influenced by the binding affinities of drugs with HSA. In particular, strong binding can reduce free drug concentrations in plasma whereas weak binding can decrease lifetime and/or distribution of drugs. Results of site marker competitive experiments revealed that DHQs spontaneously bind to HSA on site II, subdomain IIIA, though hydrophobic forces. Various substituents in the benzene ring of DHQ could increase the interactions with HSA, forming additional van der Waals forces and H-bonds.^{295–297} Transition metals, such as iron, zinc and copper, play important roles in the human body. Specifically, copper is a catalytic cofactor for a variety of metalloenzymes and physiological processes. Increased levels of copper in the body could be involved in the production of reactive oxygen species (ROS), causing imbalance in cellular functions and several diseases. For this reason, the development of fluorescent chemosensors for biologically active transition metal ions are becoming attractive. In this regard, 2-aryl DHQs were efficiently used as fluorescent probes to selectively detect Cu^{2+} ions.²⁹⁸



Table 12 Mono- and di-substituted DHQs



R	X	Y	R ₁	Activity/mechanism	Ref.
Alkyl, cycloalkyl	Phenyl, benzyl	H	H, alkyl, alkoxy, amino, nitro	Anti-inflammatory	298 and 299
Dimethyl	Alkyl, cycloalkyl, aryl	H	Heterocycles	PKC θ inhibitors	301
H	Phenyl	H	Amide, alkylamino, sulfonamide	p38 MAPK inhibitors	302
H, phenyl, furyl, phenylamino	H	Alkylamino, cycloalkylamino, hydroxy, alkoxy	H	Antibacterial	311 and 312
Phenyl	H	Alkylamino	H	Antimalarial	313
Thienyl, furyl, pyrrolidinyl	H	Phenyl	H	Antiviral	314 and 315
Thienyl, pyridinyl, indolyl	H	Phenyl	H	Antiprotozoal/shiga toxin	316
Phenyl	H	Furan-2-ylmethyl, benzyl	H, hydroxy, methoxy	TSHR inhibitors	45 and 305
Alkyl	H	Biphenyl with <i>o</i> -tetrazole	H, alkoxy	Angiotensin II receptor antagonists	234
Alky, spiro	H	Alkyl	Phenyl, benzyloxy	Antifungal/lysozyme	317
Pyridinyl-1 <i>H</i> -pyrazolyl	H	Alkyl, cycloalkyl	H, alkyl, halo	Insecticidal/calcium channels	318
Alkyl, phenyl	H	Alkyl	H, alkyl	Anticonvulsant/Na ⁺ /Ca ²⁺ exchanger inhibitors	230 and 319
H	H	Heterocycles	H, heterocycles	CDK5 inhibitors	320
H	H	Heterocycles	H	M ₁ and M ₄ receptors agonists	321
Keto	H	2-Oxoindolinyl	H	Antitumor	322
H	H	Phenyl	Amino, phenyl	Antitumor/p38 MAPK inhibitors	323
Keto, spiro	H	Alkyl, aryl	Methoxy, oxazolyl	IMPDH II inhibitors	324
H, methyl	Acyl, alkyl, phenyl	Aryl	H	Analgesic and anti-inflammatory	300 and 326
H	Acyl	Alkyl, phenyl	H	Choleretic and antibrillatory	327


A series of 2-disubstituted DHQ derivatives showed inhibitory activity against cholinesterases (ChE), involved in the lysis of choline-based esters that act as neurotransmitters. Acetylcholinesterase is present in chemical synapses and in red blood cell membranes, and butyrylcholinesterase is found in blood plasma. ChE inhibitors are the only effective therapeutic approach for the symptomatic treatment of Alzheimer's disease. These derivatives inhibited both enzymes (IC₅₀ values in micromolar range), better than or comparable to the standard drug galantamine. Molecular docking studies revealed that the benzene ring of DHQs can fit into the choline-binding site while the phenyl ring at position 2 is oriented towards the peripheral anionic site. The NH group of DHQs and Asp72 form an ion-dipole interaction, while substituents in the benzene ring are involved in H-bonding interactions.¹⁶

Mono- and di-substituted DHQ derivatives


As a privileged scaffold, the DHQ nucleus is amenable to modification. The most common functionalization was the single introduction of different chemical groups on heterocyclic N1 and mainly on N3, and substitutions on both positions. The resulting mono- and di-substituted derivatives showed specific activities depending on the substituents (Table 12).

1-Substituted DHQs can be synthesized similarly to the unsubstituted derivatives, starting from anthranilic acid²⁸⁴ and isatoic anhydride,^{299,300} bearing substituents on the amino group and the heterocyclic N respectively, or 2-fluorobenzonitrile (Scheme 13).³⁰¹ Most DHQ derivatives bearing aryl substituents on N1 showed good anti-inflammatory properties.^{298,299} Some 1-(2-phenylethyl) DHQs were found to inhibit IL-





Scheme 13 Synthetic strategies for 1-substituted DHQ derivatives.




Scheme 14 Synthetic strategies for 3-substituted DHQ derivatives.

2 production through inhibition of the protein kinase C- θ (PKC θ),³⁰¹ while 1-aryl-6-substituted DHQs inhibited p38 *in vitro* at nanomolar concentrations.³⁰²

More commonly, 3-substituted DHQs were synthesized by cyclocondensation of 2-aminobenzamides,^{303–305} or 2-nitrobenzamides,³⁰⁶ substituted on the amide groups, with different aldehydes. 3-substituted DHQs could be also obtained through the one-pot three-component reaction of isatoic anhydride with amines and aldehydes (Scheme 14).^{307–310} DHQ derivatives

bearing cyclic amino substituents on N3 showed antibacterial activity,^{311,312} while 3-alkylamino DHQs exhibited antimalarial effects.³¹³ Different heterocycles like thienyl, furyl and pyrrolidinyl in position 2 or 3-phenyl DHQs were responsible for antiviral activity,^{314,315} and for the protective effect against Shiga toxin.³¹⁶ 3-(Furan-2-ylmethyl) derivatives were reported as thyroid-stimulating hormone receptor agonists.^{45,305} 3-Biphenyl substituted DHQs with a tetrazole ring in the *ortho* position acted as angiotensin II receptor antagonists.³¹⁴ 3-Alkyl substituted DHQs bind to lysozyme showing antifungal activity.³¹⁷ Certain 3-alkyl and 3-cycloalkyl DHQ derivatives exhibit insecticidal properties by targeting calcium channels.³¹⁸ On the other hand, 3-alkyl and 3-alkylamino substituted 3,4-dihydro-2(1H)quinazolinones show anticonvulsant activity²³⁰ and inhibited the Na⁺/Ca²⁺ exchanger.³¹⁹ Other 3,4-dihydro-2(1H)quinazolinones bearing heterocyclic substituents at position 3 showed neuroprotective and antipsychotic properties, inhibiting cyclin-dependent kinase 5 (CDK5),³²⁰ and targeting M₁ and M₄ muscarinic acetylcholine receptors,³²¹ respectively. 3-(2-Oxoindolin-3-yl) DHQ derivatives showed antiproliferative activity against human tumor cell lines (IC₅₀ in the micromolar range).³²² Other 3-aryl and 3-alkyl DHQs also showed antitumor properties, inhibiting p38 MAP kinase,³²³ and inosine 5'-monophosphate dehydrogenase type II (IMPDH II).³²⁴

The less studied 1,3-disubstituted DHQs could be obtained in two different ways: through the cyclization of *N*-substituted isatoic anhydride and substituted arylidene anilines,²⁹⁹ or the reaction of 3-substituted DHQ with chloride (Scheme 15).³²⁵ 1,3-



Scheme 15 Synthetic strategies for 1,3-disubstituted DHQ derivatives.



Disubstituted DHQ derivatives mostly showed anti-analgesic and inflammatory activities.^{300,326} 1-Acyl DHQs bearing alkyl and phenyl substituents on N showed choleric and anti-fibrillatory activities.³²⁷

Conclusions

The importance of the DHQ nucleus has emerged due to its versatility as suitable substrate for functionalization and its remarkable bioactivities. Many procedures have been reported for the synthesis of DHQ derivatives as racemic mixture. Although the cyclocondensation of anthranilamide and an aldehyde and the one-pot three-component reaction of isatoic anhydride, ammonium acetate and an aldehyde represent the most common ways to obtain DHQ derivatives, several other methods have been suggested. Many approaches have been investigated, from classical to greener and more sustainable reaction conditions. Recently, enantioselective strategies have been attempted in order to obtain pure (*S*)-enantiomers. Furthermore, the DHQ scaffold is an important intermediate in organic chemistry and can easily be oxidized into QZ scaffold. On the other hand, various important bioactivities have been associated to the DHQ scaffold and reported in the literature. On this basis, the DHQ nucleus imposes itself as a privileged scaffold in drug design and an interesting fragment for drug discovery.

Conflicts of interest

There are no conflicts to declare.

References

- 1 K. Imtiaz, I. Aliya, A. Waqas and S. Aamer, *Eur. J. Med. Chem.*, 2015, **90**, 124–169.
- 2 P. D. Leeson and B. Springthorpe, *Nat. Rev. Drug Discovery*, 2007, **6**, 881–890.
- 3 G. M. Chinigo, M. Paige, S. Grindrod, E. Hamel, S. Dakshanamurthy, M. Chruszcz, W. Minor and M. L. Brown, *J. Med. Chem.*, 2008, **51**, 4620–4631.
- 4 B. E. Evans, K. E. Rittle, M. G. Bock, R. M. DiPardo, R. M. Freidinger, W. L. Whitter, G. F. Lundell, D. F. Veber, P. S. Anderson, R. S. L. Chang, V. J. Lotti, D. J. Cerino, T. B. Chen, P. J. Kling, K. A. Kunkel, J. P. Springer and J. Hirshfield, *J. Med. Chem.*, 1988, **31**, 2235–2246.
- 5 D. A. Horton, G. T. Bourne and M. L. Smythe, *Chem. Rev.*, 2003, **103**, 893–930.
- 6 <https://www.drugbank.ca>.
- 7 R. Williams, C. M. Niswender, Q. Luo, U. Le, P. J. Conn and C. W. Lindsley, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 962–966.
- 8 Y. Kobayashi, Y. Nakano, K. Hoshikuma, Y. Yokoo and T. Kamiya, *Planta Med.*, 2001, **67**, 628–633.
- 9 A. N. Hughes, I. Rafi, M. J. Griffin, A. H. Calvert, D. R. Newell, J. A. Calvete, A. Johnston, N. Clendeninn and A. V. Boddy, *Clin. Cancer Res.*, 1999, **5**, 111–118.
- 10 R. V. Coombs, R. P. Danna, M. Denzer, G. E. Hardtmann, B. Huegi, G. Koletar, J. Koletar, H. Ott, E. Juckiewicz, J. W. Perrine, E. I. Takesue and J. H. Trapold, *J. Med. Chem.*, 1973, **16**, 1237–1245.
- 11 S. Seyedmousavi, H. Rafati, M. Ilkit, A. Tolooee, M. T. Hedayati and P. Verweij, *Methods and Protocols*, 2016, **1508**, 107–139.
- 12 M. Shahraki, S. M. Habibi-khorassani and Y. Narouei, *Am. J. Chem. Appl.*, 2015, **2**, 83–90.
- 13 H. L. Yale and M. Kalkstein, *J. Med. Chem.*, 1967, **10**, 334–336.
- 14 G. Bonola and E. Sianesi, *J. Med. Chem.*, 1970, **13**, 329–332.
- 15 R. J. Ericsson and E. Reif, *US pat.* 3651061, 1972.
- 16 M. Sarfraz, N. Sultana, U. Rashid, M. S. Akram, A. Sadiq and M. I. Tariq, *Bioorg. Chem.*, 2017, **70**, 237–244.
- 17 N. Saroja, E. Laxminarayana and K. R. S. Prasad, *Indian J. Heterocycl. Chem.*, 2014, **24**, 67–70.
- 18 S. D. Sharma and V. Kaur, *Synthesis*, 1989, **1989**, 677–680.
- 19 M. Hour, L. Huang, S. Kuo, K. Bastow, Y. Nakanishi, E. Hamel and K. Lee, *J. Med. Chem.*, 2000, **43**, 4479–4487.
- 20 S. Kuo, M. Hour, L. Huang and K. Lee, *US pat* 6479499 B1, 2002.
- 21 J. J. Naleway, C. M. J. Fox, D. Robinhold, E. Terpetschnig, N. A. Olson and P. Haugland, *Tetrahedron Lett.*, 1994, **35**, 8569–8572.
- 22 V. B. Labade, P. V. Shinde and M. S. Shingare, *Tetrahedron Lett.*, 2013, **54**, 5778–5780.
- 23 A. G. Al-Sehemi, M. Pannipara and A. Kalam, *Spectrochim. Acta, Part A*, 2017, **171**, 97–103.
- 24 X. Liu, D. Hu and H. Shen, *Asian J. Chem.*, 2012, **24**, 1365–1367.
- 25 M. Wang, J. Gao, Z. Song and L. Wang, *Org. Prep. Proced. Int.*, 2012, **44**, 159–163.
- 26 R. Navudu, G. R. Mannem, T. Margani, U. M. R. Vanga and H. B. Bollikolla, *Asian J. Chem.*, 2016, **28**, 1321–1324.
- 27 S. Esfandiari, M. T. Maghsoodlou, S. M. Habibi-Khorassani, S. Kiaee and J. Aboonajmi, *Iranian Journal of Organic Chemistry*, 2012, **4**, 827–830.
- 28 M. J. Mphahlele, M. M. Maluleka and T. Khoza, *Bull. Chem. Soc. Ethiop.*, 2014, **28**, 81–90.
- 29 M. Desroses, M. Scobie and T. Helleday, *New J. Chem.*, 2013, **37**, 3595–3597.
- 30 X. Wang, K. Yang, J. Zhou and S. Tu, *J. Comb. Chem.*, 2010, **12**, 417–421.
- 31 X. Wang, K. Yang, M. Zhang and C. Yao, *Synth. Commun.*, 2010, **40**, 2633–2646.
- 32 Y. Nagasawa, Y. Matsusaki, T. Nobuta, N. Tada, T. Miura and A. Itoh, *RSC Adv.*, 2015, **5**, 63952–63954.
- 33 F. Miklós, V. Hum and F. Fülöp, *ARKIVOC*, 2014, **2014**, 25–37.
- 34 D. M. Boghaei, M. M. Najafpour and V. McKee, *Acta Crystallogr., Sect. E: Struct. Rep. Online*, 2009, **E65**, o193.
- 35 M. Abdollahi-Alibeik and E. Shabani, *Chin. Chem. Lett.*, 2011, **22**, 1163–1166.
- 36 N. Ramesh, M. G. Rao, R. Valara, V. U. Rao and B. H. Babu, *Med. Chem. Res.*, 2016, **25**, 1945–1951.
- 37 Y. Luo, Y. Wu, Y. Wang, H. Sun, Z. Xie, W. Zhang and Z. Gao, *RSC Adv.*, 2016, **6**, 66074–66077.



- 38 L. Rajaka, N. R. Penumati, K. Nagaiah, Y. Poornachandra and C. G. Kumar, *Synth. Commun.*, 2015, **45**, 1893–1901.
- 39 K. R. Gopinath, H. S. Shekar, K. J. Rajendraprasad, H. Nagabhushana and M. Krishnappa, *WJPPS*, 2016, **5**, 1272–1279.
- 40 M. Sharma, S. Pandey, K. Chauhan, D. Sharma, B. Kumar and P. M. S. Chauhan, *J. Org. Chem.*, 2012, **77**, 929–937.
- 41 S. Kobayashi, M. Sugiura, H. Kitagawa and W. W. Lam, *Chem. Rev.*, 2002, **102**, 2227–2302.
- 42 J. X. Chen, H. Y. Wu and W. K. Su, *Chin. Chem. Lett.*, 2007, **18**, 536–538.
- 43 S. Vasudhevan and R. J. Karunakaran, *Int. J. PharmTech Res.*, 2013, **5**, 2844–2853.
- 44 P. Sivaguru, K. Parameswaran and A. Lalitha, *Tetrahedron Lett.*, 2016, **57**, 2549–2553.
- 45 E. E. Englund, S. Neumann, E. Eliseeva, J. G. McCoy, S. Titus, W. Zheng, N. Southall, P. Shin, W. Leister, C. J. Thomas, J. Inglese, C. P. Austin, M. C. Gershengorn and W. Huang, *MedChemComm*, 2011, **2**, 1016–1020.
- 46 S. Wang, K. Yang and X. Wang, *Chin. J. Org. Chem.*, 2011, **8**, 1235–1239.
- 47 Y. Shang, L. Fan, X. Li and M. Liu, *Chin. Chem. Lett.*, 2015, **26**, 1355–1358.
- 48 S. Li, W. Jhang, T. Liou and D. Yang, *Dyes Pigm.*, 2015, **114**, 259–266.
- 49 A. Shaabani, A. Maleki and H. Mofakham, *Synth. Commun.*, 2008, **38**, 3751–3759.
- 50 M. Wang, J. J. Gao, Z. G. Song and L. Wang, *Chem. Heterocycl. Compd.*, 2011, **47**, 851–855.
- 51 R. Venkatesh, S. Kasaboina, H. K. Gaikwad, S. Janardhan, R. Bantu, L. Nagarapu, G. N. Sastry and S. K. Banerjee, *Eur. J. Med. Chem.*, 2015, **96**, 22–29.
- 52 R. Noel, J. Martinez, D. Buisson, L. Johannes, D. Gillet, J. Barbier and J. Cintrat, *J. Med. Chem.*, 2013, **56**, 3404–3413.
- 53 R. M. Christie and S. Moss, *J. Chem. Soc., Perkin Trans. 1*, 1985, **1**, 2779–2783.
- 54 A. Ghorbani-Choghamarani, Z. Darvishnejad and M. Norouzi, *Appl. Organomet. Chem.*, 2015, **29**, 707–711.
- 55 P. Hanumanthu, S. K. V. Seshavatharam, C. V. Ratnam and N. V. S. Rao, *Proc. Indian Acad. Sci.*, 1976, **84**, 57–63.
- 56 P. R. Kumar and S. Reddy, *Synth. Commun.*, 1992, **22**, 2499–2508.
- 57 R. P. Staiger, C. L. Moyer and G. R. Pitcher, *J. Chem. Eng. Data*, 1963, **8**, 454–456.
- 58 A. Dömling, *Chem. Rev.*, 2006, **106**, 17–86.
- 59 M. Narasimhulu and Y. R. Lee, *Tetrahedron*, 2011, **67**, 9627–9634.
- 60 S. Khaksar and M. Gholami, *Res. Chem. Intermed.*, 2015, **41**, 3709–3718.
- 61 M. Baghbanzadeh, P. Salehi, M. Dabiri and G. Kozehgary, *Synthesis*, 2006, **2**, 344–348.
- 62 B. Chen, J. Li and G. Chen, *Ultrason. Sonochem.*, 2015, **23**, 59–65.
- 63 A. Davoodnia, M. Khashi and N. Tavakoli-Hoseini, *Chin. J. Catal.*, 2014, **35**, 1054–1058.
- 64 Z. Song, L. Liu, Y. Wang and X. Sun, *Res. Chem. Intermed.*, 2012, **38**, 1091–1099.
- 65 Z. Song, X. Wan and S. Zhao, *Indian J. Chem. Technol.*, 2012, **19**, 118–123.
- 66 M. Wang, T. T. Zhang, Y. Liang and J. J. Gao, *Monatsh. Chem.*, 2012, **143**, 835–839.
- 67 R. Ghorbani-Vaghei, A. Shahriari, Y. Maghbooli and J. Mahmoudi, *Res. Chem. Intermed.*, 2017, **43**, 983–993.
- 68 L. Zeng and C. Cai, *J. Heterocycl. Chem.*, 2010, **47**, 1035–1039.
- 69 S. Rostamizadeh, A. M. Amani, R. Aryan, H. R. Ghaieni and N. Shadjou, *Synth. Commun.*, 2008, **38**, 3567–3576.
- 70 M. Wang, T. T. Zhang, Y. Liang and J. J. Gao, *Chin. Chem. Lett.*, 2011, **22**, 1423–1426.
- 71 J. Chen, D. Wu, F. He, M. Liu, H. Wu, J. Ding and W. Su, *Tetrahedron Lett.*, 2008, **49**, 3814–3818.
- 72 X. Zhu, S. R. Kang, L. Xia, J. Lee, N. Basavegowda and Y. R. Lee, *Mol. Diversity*, 2015, **19**, 67–75.
- 73 L. Sancineto, N. Iraci, S. Massari, V. Attanasio, G. Corazza, M. L. Barreca, S. Sabatini, G. Manfroni, N. R. Avanzi, V. Cecchetti, C. Pannecouque, A. Marcello and O. Tabarrini, *ChemMedChem*, 2013, 1941–1953.
- 74 J. Szejtli, *Chem. Rev.*, 1998, **98**, 1743–1753.
- 75 S. V. Bhosale and S. V. Bhosale, *Mini-Rev. Org. Chem.*, 2007, **4**, 231–242.
- 76 K. Ramesh, K. Karnakar, G. Satish, K. H. V. Reddy and Y. V. D. Nageswar, *Tetrahedron Lett.*, 2012, **53**, 6095–6099.
- 77 K. Ramesh, K. Karnakar, G. Satish, B. S. P. Anil Kumar and Y. V. D. Nageswar, *Tetrahedron Lett.*, 2012, **53**, 6936–6939.
- 78 D. R. Patil, P. G. Ingole, K. Singh and D. S. Dalal, *J. Inclusion Phenom. Macrocyclic Chem.*, 2013, **76**, 327–332.
- 79 J. Wu, X. Du, J. Ma, Y. Zhang, O. Shi, L. Luo, B. Song, S. Yang and D. Hu, *Green Chem.*, 2014, **16**, 3210–3217.
- 80 M. T. Maghsoodlou, N. Khorshidi, M. R. Mousavi, N. Hazeri and S. M. Habibi-Khorassani, *Res. Chem. Intermed.*, 2015, **41**, 7497–7508.
- 81 H. R. Shaterian and F. Rigi, *Res. Chem. Intermed.*, 2015, **41**, 721–738.
- 82 M. Dabiri, P. Salehi, S. Otokesh, M. Baghbanzadeh, G. Kozehgary and A. A. Mohammadi, *Tetrahedron Lett.*, 2005, **46**, 6123–6126.
- 83 A. Ghorbani-Choghamarani and T. Taghipour, *Lett. Org. Chem.*, 2011, **8**, 470–476.
- 84 Y. Chen, W. Shan, M. Lei and L. Hu, *Tetrahedron Lett.*, 2012, **53**, 5923–5925.
- 85 M. P. Surpur, P. R. Singh, S. B. Patil and S. D. Samant, *Synth. Commun.*, 2007, **37**, 1965–1970.
- 86 S. Ikegami and H. Hamamoto, *Chem. Rev.*, 2009, **109**, 583–593.
- 87 M. Moreno-Man and R. Pleixats, *Acc. Chem. Res.*, 2003, **36**, 638–643.
- 88 Z. Zhang, H. Lü, S. Yang and J. Gao, *J. Comb. Chem.*, 2010, **12**, 643–646.
- 89 M. Z. Kassaei, S. Rostamizadeh, N. Shadjou, E. Motamedi and M. Esmaeizadeh, *J. Heterocycl. Chem.*, 2010, **47**, 1421–1424.
- 90 J. Zhang, D. Ren, Y. Ma, W. Wang and H. Wu, *Tetrahedron*, 2014, **70**, 5274–5282.



- 91 S. Santra, M. Rahman, A. Roy, A. Majee and A. Hajra, *Catal. Commun.*, 2014, **49**, 52–57.
- 92 M. Hajjami, A. Ghorbani-Choghamarani, Z. Yousofvand and M. Norouzi, *J. Chem. Sci.*, 2015, **127**, 1221–1228.
- 93 N. Ravazi and B. Akhlaghinia, *New J. Chem.*, 2016, **40**, 447–457.
- 94 J. Lu and P. H. Toy, *Chem. Rev.*, 2009, **109**, 815–838.
- 95 A. Gharib, B. R. Hashemipour, M. Jahangir, M. Roshani and R. Safaei, *Org. Chem. Int.*, 2013, **2013**, 848237.
- 96 A. Gharib, L. Vojdanifard, N. Noroozi Pesyan, B. R. Hashemi Pou Khorasani, M. Jahangir and M. Roshani, *Bulg. Chem. Commun.*, 2014, **46**, 667–679.
- 97 G. Majid, A. Kobre, M. Hamed and S. Hamid Reza, *Chin. J. Chem.*, 2011, **29**, 1617–1623.
- 98 A. Maleki, M. Rabbani and S. Shahrokh, *Appl. Organomet. Chem.*, 2015, **29**, 809–814.
- 99 R. Mekala, M. V. Madhubabu, G. Dhanunjaya, S. Regati, K. B. Chandrasekhar and J. Sarva, *Synth. Commun.*, 2017, **47**, 121–130.
- 100 P. Salehi, M. Dabiri, M. A. Zolfigol and M. Baghbanzadeh, *Synlett*, 2005, **7**, 1155–1157.
- 101 M. Dabiri, P. Salehi, M. Baghbanzadeh, M. A. Zolfigol, M. Agheb and S. Heydari, *Catal. Commun.*, 2008, **9**, 785–788.
- 102 K. Niknam, M. Mohammadizadeh and S. Mirzaee, *Chin. J. Chem.*, 2011, **29**, 1417–1422.
- 103 K. Niknam, N. Jafarpour and E. Niknam, *Chin. Chem. Lett.*, 2011, **22**, 69–72.
- 104 A. Ghorbani-Choghamarani and P. Zamani, *J. Iran. Chem. Soc.*, 2012, **9**, 607–613.
- 105 S. Rostamizadeh, A. M. Amani, G. Mahdavinia, H. Sepehrian and S. Ebrahimi, *Synthesis*, 2010, **8**, 1356–1360.
- 106 A. Shokrolahi, A. Zali, M. A. Zare and K. Esmailpour, *Iran. J. Catal.*, 2012, **2**, 91–94.
- 107 P. Sivaguru, K. Parameswaran, M. Kiruthiga, P. Vadivel and A. Lalitha, *J. Iran. Chem. Soc.*, 2015, **12**, 95–100.
- 108 H. R. Shaterian and F. Rigi, *Res. Chem. Intermed.*, 2014, **40**, 2983–2999.
- 109 H. R. Shaterian, A. R. Oveisi and M. Honarmand, *Synth. Commun.*, 2010, **40**, 1231–1242.
- 110 H. R. Shaterian, N. Fahimi and K. Aziz, *Res. Chem. Intermed.*, 2014, **40**, 1879–1898.
- 111 S. Li, Q. Zhang and Y. Peng, *Monatsh. Chem.*, 2015, **146**, 1859–1864.
- 112 A. Saffar-Teluri and S. Bolouk, *Monatsh. Chem.*, 2010, **141**, 1113–1115.
- 113 P. Salehi, *Synth. Commun.*, 2006, **36**, 2287–2292.
- 114 S. M. Roopan, F. N. Khan, J. S. Jin and R. S. Kumar, *Res. Chem. Intermed.*, 2011, **37**, 919–927.
- 115 F. Serp, M. Corrias and P. Kalck, *Appl. Catal., A*, 2003, **253**, 337–358.
- 116 J. Safari and S. Gandomi-Ravandi, *J. Mol. Catal. A: Chem.*, 2013, **371**, 135–140.
- 117 J. Safari and S. Gandomi-Ravandi, *C. R. Chim.*, 2013, **16**, 1158–1164.
- 118 J. Safari and S. Gandomi-Ravandi, *J. Mol. Struct.*, 2014, **1072**, 173–178.
- 119 J. Safari and S. Gandomi-Ravandi, *J. Saudi Chem. Soc.*, 2017, **21**, S415–S424.
- 120 L. Wang, L. Hu, J. Shao, J. Yu and L. Zhang, *J. Fluorine Chem.*, 2008, **129**, 1139–1145.
- 121 R. Sharma, A. K. Pandey and P. M. S. Chauhan, *Synlett*, 2012, **23**, 2209–2214.
- 122 M. Wang, T. T. Zhang, J. J. Gao and Y. Liang, *Chem. Heterocycl. Compd.*, 2012, **6**, 897–902.
- 123 Á. Magyar and Z. Kell, *Catal. Lett.*, 2016, **146**, 1153–1162.
- 124 P. T. Anastas and J. C. Warner, *Green Chemistry: Theory and Practice*, Oxford University Press, Oxford, 1998.
- 125 D. L. Hjerensen, D. L. Schett and J. M. Boese, *J. Chem. Educ.*, 2000, **77**, 1543–1547.
- 126 R. Mestres, *Environ. Sci. Pollut. Res.*, 2005, **12**, 128–132.
- 127 J. L. Tucker, *Org. Process Res. Dev.*, 2006, **10**, 315–319.
- 128 P. V. N. S. Murthy, D. Rambabu, G. Rama Krishna, C. Malla Reddy, K. R. S. Prasad, M. V. Basaveswara Rao and M. Pal, *Tetrahedron Lett.*, 2012, **53**, 863–867.
- 129 M. Wang, J. Gao, Z. Song and L. Wang, *J. Heterocycl. Chem.*, 2012, **49**, 1250–1253.
- 130 A. Rostami and A. Tavakoli, *Chin. Chem. Lett.*, 2011, **22**, 1317–1320.
- 131 A. D. Roy, K. Jayalakshmi, S. Dasgupta, R. Roy and B. Mukhopadhyay, *Magn. Reson. Chem.*, 2008, **46**, 1119–1126.
- 132 H. R. Shaterian and A. R. Oveisi, *Chin. J. Chem.*, 2009, **27**, 2418–2422.
- 133 M. Singh and N. Raghav, *Bioorg. Chem.*, 2015, **59**, 12–22.
- 134 S. B. Bharate, N. Muppararu, S. Manda, J. B. Bharate, R. Mudududdla, R. R. Yadav and R. A. Vishwakarma, *ARKIVOC*, 2012, **2012**, 308–318.
- 135 M. Ghashang, *Orient. J. Chem.*, 2012, **28**, 1213–1218.
- 136 S. D. Dindulkar, J. Oh, V. M. Arole and Y. T. Jeong, *C. R. Chimie*, 2014, **17**, 971–979.
- 137 A. Ghorbani-Choghamarani and B. Tahmasbi, *New J. Chem.*, 2016, **40**, 1205–1212.
- 138 M. Hajjami, A. Ghirbani-Choghamarani, R. Ghafouri-Nejad and B. Tahmasbi, *New J. Chem.*, 2016, **40**, 3066–3074.
- 139 B. A. Dar, A. K. Sahu, P. Patidar, Y. Parial, P. Sharma, M. Sharma and B. Singh, *Am. J. Chem.*, 2012, **2**, 248–254.
- 140 B. A. Dar, A. K. Sahu, P. Patidar, P. R. Sharma, N. Mupparapu, D. Vyas, S. Maity, M. Sharma and B. Singh, *J. Ind. Eng. Chem.*, 2013, **19**, 407–412.
- 141 B. V. Subba Reddy, A. Venkateswarlu, C. Madan and A. Vinu, *Tetrahedron Lett.*, 2011, **52**, 1891–1894.
- 142 A. V. Dhanunjaya, B. P. Vyakuntswararao, T. Bhaskarkumar, N. R. Jogdand, D. Kalita, J. K. D. Lilakar, V. Siddaiah, P. D. Sanasi and A. Raghunadh, *Tetrahedron Lett.*, 2015, **56**, 4714–4717.
- 143 A. Ghorbani-Choghamarani and G. Azadi, *RSC Adv.*, 2015, **5**, 9752–9758.
- 144 A. Rostami, B. Tahmasbi, H. Gholami and H. Taymorian, *Chin. Chem. Lett.*, 2013, **24**, 211–214.
- 145 M. Ghashang, S. S. Mansoor and K. Aswin, *Res. Chem. Intermed.*, 2015, **41**, 3447–3460.



- 146 M. Abdollahi-Alibeik and E. Shabani, *J. Iran. Chem. Soc.*, 2014, **11**, 351–359.
- 147 H. R. Safaei, M. Shekouhy, S. Khademi, V. Rahmanian and M. Safaei, *J. Ind. Eng. Chem.*, 2014, **20**, 3019–3024.
- 148 M. Rahman, I. Ling, N. Abdullah, R. Hashim and A. Hajra, *RSC Adv.*, 2015, **5**, 7755–7760.
- 149 X. Zong, Y. Zhao, W. Luo, X. H. Yu, J. K. Wang and Y. Pan, *Chin. Chem. Lett.*, 2010, **21**, 778–781.
- 150 M. Tajbakhsh, R. Hosseinzadeh, P. Rezaee and M. Tajbakhsh, *Chin. J. Catal.*, 2014, **35**, 58–65.
- 151 H. Alinezhadi, E. Soleymani and M. Zare, *Res. Chem. Intermed.*, 2017, **43**, 457–466.
- 152 J. Wang, Y. Zong, R. Fu, Y. Niu, G. Yue, Z. Quan, X. Wang and Y. Pan, *Ultrason. Sonochem.*, 2014, **21**, 29–34.
- 153 S. Zhang, Z. Xie, L. Liu, M. Liang and Z. Le, *Chin. Chem. Lett.*, 2017, **28**, 101–104.
- 154 A. Maleki and S. Shahrokh, *1st International Electronic Conference on Materials*, 26 May–10 June 2014.
- 155 N. Kausar, I. Roy, D. Chattopadhyay and A. R. Das, *RSC Adv.*, 2016, **6**, 22320–22330.
- 156 J. Safari and S. Gandomi-Ravandi, *J. Mol. Catal. A: Chem.*, 2014, **390**, 1–6.
- 157 J. Safari and S. Gandomi-Ravandi, *RSC Adv.*, 2014, **4**, 11654–11660.
- 158 A. Ghorbani-Choghamarani and M. Norouzi, *J. Mol. Catal. A: Chem.*, 2014, **395**, 172–179.
- 159 F. Havasi, A. Ghorbani-Choghamarani and F. Nikpour, *Microporous Mesoporous Mater.*, 2016, **224**, 26–35.
- 160 B. B. F. Mirjalili, A. Bamoniri and S. Azad, *J. Iran. Chem. Soc.*, 2017, **14**, 47–55.
- 161 P. Sivaguru, K. Parameswaran and A. Lalitha, *Tetrahedron Lett.*, 2016, **57**, 2549–2553.
- 162 W. Wei, C. C. K. Keh, C. Li and R. S. Varma, *Clean Technol. Environ. Policy*, 2005, **7**, 62–69.
- 163 A. K. Chakraborti and S. R. Roy, *J. Am. Chem. Soc.*, 2009, **131**, 6902–6903.
- 164 A. Patil, M. Barge, G. Rashinkar and R. Salunkhe, *Mol. Diversity*, 2015, **19**, 435–445.
- 165 S. K. Ghosh and R. Nagarajan, *RSC Adv.*, 2015, **6**, 27378–27387.
- 166 M. Dabiri, P. Salehi and M. Baghbanzadeh, *Monatsh. Chem.*, 2007, **138**, 1191–1194.
- 167 M. Zhang, Y. Liu and X. Wang, *Chin. J. Org. Chem.*, 2014, **34**, 1682–1686.
- 168 M. Li and C. Feng, *Acta Crystallogr., Sect. E: Struct. Rep. Online*, 2009, **E65**, o2145.
- 169 J. Chen, W. Su, M. Liu and C. Jin, *Green Chem.*, 2007, **9**, 972–975.
- 170 N. B. Darvatkar, S. V. Bhilare, A. R. Deorukhkar, D. G. Raut and M. M. Salunkhe, *Green Chem. Lett. Rev.*, 2010, **3**, 301–306.
- 171 H. Alinezhad, M. Tajbakhsh and N. Ghobadi, *Res. Rev. Mater. Sci. Chem.*, 2014, **3**, 123–135.
- 172 M. Nikpassand, L. Zare Fekri, K. F. Sina, S. Z. Abed and O. Marvi, *Russ. J. Gen. Chem.*, 2015, **85**, 1959–1964.
- 173 S. Nagarajan, T. M. Shaikh and E. Kandasamy, *New J. Chem.*, 2015, **39**, 9693–9699.
- 174 G. Yassaghi, A. Davoodnia, S. Allameh, A. Zare-Bidaki and N. Tavakoli-Hoseini, *Bull. Korean Chem. Soc.*, 2012, **33**, 2724–2730.
- 175 F. S. Toosi and M. Khakzadi, *Res. Chem. Intermed.*, 2015, **41**, 311–317.
- 176 A. Davoodnia, S. Allameh, A. R. Fakhari and N. Tavakoli-Hoseini, *Chin. Chem. Lett.*, 2010, **21**, 550–553.
- 177 M. A. B. Fard, A. Mobinikhaledi and M. Hamidinasab, *Synth. React. Inorg. Met.-Org. Chem.*, 2014, **44**, 567–571.
- 178 O. Obaiah, N. N. Kebbahalli, R. M. Goravanahalli, P. S. Chottanahalli, R. S. Kanchugarakoppal and M. Kempegowda, *Eur. J. Chem.*, 2014, **5**, 671–675.
- 179 H. R. Safaei, M. Shekouhy, V. Shafiee and M. Davoodi, *J. Mol. Liq.*, 2013, **180**, 139–144.
- 180 Z. Karimi-Jaberi and L. Zarei, *S. Afr. J. Chem.*, 2012, **65**, 36–38.
- 181 Q. Ding, J. Zhang, J. Chen, M. Liu, J. Ding and H. Wu, *J. Heterocycl. Chem.*, 2012, **49**, 375–380.
- 182 V. B. Ningdale, U. N. Chaudhary and K. A. Shaikh, *Arch. Appl. Sci. Res.*, 2013, **5**, 82–87.
- 183 S. Zhaleh, N. Hazeri and M. T. Maghsoodlou, *Res. Chem. Intermed.*, 2016, **42**, 6381–6390.
- 184 D. Maitraie, G. Venkat Reddy, V. V. V. N. S. Rama Rao, S. Ravi Kanth, P. Shanthan Rao and B. Narsaiah, *J. Fluorine Chem.*, 2002, **118**, 73–79.
- 185 O. A. Maloshitskaya, J. Sinkkonen, V. V. Alekseyev, K. N. Zelenin and K. Pihlaja, *Tetrahedron*, 2005, **61**, 7294–7303.
- 186 R. Z. Qiao, B. L. Xu and Y. W. Wang, *Chin. Chem. Lett.*, 2007, **18**, 656–658.
- 187 Y. Zheng, M. Bian, X. Deng, S. Wang and Z. Quan, *Arch. Pharm. Chem. Life Sci.*, 2013, **346**, 119–126.
- 188 P. Yerram, R. Chowrasia, S. Seeka and S. J. Tangenda, *Eur. J. Chem.*, 2013, **4**, 462–466.
- 189 R. Ramesh, N. Nagasundaram, D. Meignasundar, P. Vadivel and A. Lalitha, *Res. Chem. Intermed.*, 2017, **43**, 1767–1782.
- 190 Z. Xie, S. Zhang, G. Jiang, D. Sun and Z. Le, *Green Chem. Lett. Rev.*, 2015, **8**, 95–98.
- 191 M. Tímea, F. Miklós, L. Lázár and F. Fülöp, *ARKIVOC*, 2016, **2016**, 247–258.
- 192 M. Wang, T. T. Zhang and Z. G. Song, *Chin. Chem. Lett.*, 2011, **22**, 427–430.
- 193 Z. Karimi-Jaberi and R. Arjmandi, *Monatsh. Chem.*, 2011, **142**, 631–635.
- 194 S. Khaksar and S. M. Talesh, *C. R. Chimie*, 2012, **15**, 779–783.
- 195 K. R. Rao, R. Mekala, A. Raghunadh, S. B. Meruva, S. P. Kumar, D. Kalita, E. Laxminarayana, B. Prasad and M. Pal, *RSC Adv.*, 2015, **5**, 61575–61579.
- 196 M. Hajjami, F. Bakhti and E. Ghiasbeygi, *Croat. Chem. Acta*, 2015, **88**, 197–205.
- 197 A. Olyaei, F. Rahbarian and M. Sadeghpour, *Chem. Heterocycl. Compd.*, 2015, **51**, 899–902.
- 198 P. P. Naidu, A. Raghunadh, K. R. Rao, R. Mekala, J. M. Babu, B. R. Rao, V. Siddaiah and M. Pal, *Synth. Commun.*, 2014, **44**, 1475–1482.



- 199 L. Gao, H. Ji, L. Rong, D. Tang, Y. Zha, Y. Shi and S. Tu, *J. Heterocycl. Chem.*, 2011, **48**, 957–960.
- 200 A. D. Gupta, S. Samanta and A. K. Mallik, *Org. Prep. Proced. Int.*, 2015, **47**, 356–360.
- 201 V. Polshettiwar and R. S. Varma, *Pure Appl. Chem.*, 2008, **80**, 777–790.
- 202 H. M. Hügel, *Molecules*, 2009, **14**, 4936–4972.
- 203 F. Li, Y. Feng, Q. Meng, W. Li, Z. Li, Q. Wang and F. Tao, *ARKIVOC*, 2007, **2007**, 40–50.
- 204 K. Kumari, D. S. Raghuvanshi and K. N. Singh, *Indian J. Chem.*, 2012, **51B**, 860–865.
- 205 R. S. Varma, *Green Chem. Lett. Rev.*, 2007, **1**, 37–45.
- 206 D. Rambabu, S. K. Kumar, B. Y. Sreenivas, S. Sandra, A. Kandale, P. Misra, M. V. B. Rao and M. Pal, *Tetrahedron Lett.*, 2013, **54**, 495–501.
- 207 J. A. Moore, G. J. Sutherland, R. Sowerby, E. G. Kelly, S. Palermo and W. Webster, *J. Org. Chem.*, 1969, **34**, 887–892.
- 208 M. Soural, P. Funk, L. Kvapil, P. Hradil, J. Hlaváč and V. Bertolasi, *ARKIVOC*, 2010, **2010**, 255–265.
- 209 M. Zhang, K. Yang and X. Wang, *Acta Crystallogr., Sect. E: Struct. Rep. Online*, 2010, **E66**, o1069.
- 210 A. J. A. Watson, A. C. Maxwell and J. M. J. William, *Org. Biomol. Chem.*, 2012, **10**, 240–243.
- 211 K. H. Narasimhamurthy, S. Chandrappa, K. S. Sharath, K. B. Harsha, H. Ananda and K. S. Rangappa, *RSC Adv.*, 2014, **4**, 34479–34486.
- 212 A. E. O. Lalami, S. Boukhris, N. Habbadi, N. B. Larb and A. Souizi, *J. Maroc. Chim. Heterocycl.*, 2002, **1**, 37–43.
- 213 N. T. Patil, R. D. Kavthe, V. S. Raut, V. S. Shinde and B. Sridhar, *J. Org. Chem.*, 2010, **75**, 1277–1280.
- 214 N. T. Patil, P. G. V. V. Lakshmi and V. Singh, *Eur. J. Org. Chem.*, 2010, 4719–4731.
- 215 J. Tang, J. Li, L. Zhang, S. Ma, D. Shi, Q. Zhang, L. Yang, X. Wang, X. Liu and C. Liu, *J. Heterocycl. Chem.*, 2012, **49**, 533–542.
- 216 J. Tang, D. Shi, L. Zhang, Q. Zhang and J. Li, *Synth. Commun.*, 2010, **40**, 632–641.
- 217 L. Zhang, X. Yang, D. Shi and J. Chen, *Acta Crystallogr., Sect. E: Struct. Rep. Online*, 2008, **E64**, o450.
- 218 P. N. Borase, P. B. Thale and G. S. Shankarling, *RSC Adv.*, 2016, **6**, 63078–63083.
- 219 X. Wy, S. Oschatz, A. Block, A. Spannenberg and P. Langer, *Org. Biomol. Chem.*, 2014, **12**, 1865–1870.
- 220 H. Chai, J. Li, L. Yang, M. Liu, D. Yang, Q. Zhang and D. Shi, *Chin. J. Chem.*, 2014, **32**, 865–870.
- 221 C. Balakumar, P. Lamba, D. P. Kishore, B. L. Narayana, K. V. Eao, K. Rajwinder, A. R. Rao, B. Shireesha and B. Narsajah, *Eur. J. Med. Chem.*, 2010, **45**, 4904–4913.
- 222 F. Tamaddon and F. Pouramini, *Synlett*, 2014, **25**, 1127–1131.
- 223 D. Shi, L. Rong, J. Wang, Q. Zhuang, X. Wang and H. Hu, *Tetrahedron Lett.*, 2003, **44**, 3199–3201.
- 224 D. Shi, J. Wang, L. Rong, Q. Zhuang, S. Tu and H. Hu, *J. Chem. Res.*, 2003, **S**, 671–673.
- 225 R. A. Bunce and B. Nammalwar, *J. Heterocycl. Chem.*, 2011, **48**, 991–997.
- 226 M. Wang, G. Dou and D. Shi, *J. Comb. Chem.*, 2010, **12**, 582–586.
- 227 G. Cai, X. Xu, Z. Li, W. P. Weber and P. Lu, *J. Heterocycl. Chem.*, 2002, **39**, 1271–1272.
- 228 W. Su and B. Yang, *Aust. J. Chem.*, 2002, **55**, 695–697.
- 229 W. Su and B. Yang, *J. Chem. Res.*, 2002, **S**, 604–605.
- 230 M. J. Kornet, *J. Heterocycl. Chem.*, 1992, **29**, 103–105.
- 231 N. Ho, R. S. Harapanhalli, B. A. Dahman, K. Chen, K. Wang, S. J. Adelstein and A. L. Kassiss, *Bioconjugate Chem.*, 2002, **13**, 357–364.
- 232 H. Asakwa and M. Matano, *Chem. Pharm. Bull.*, 1979, **27**, 1287–1298.
- 233 Z. Xu, Y. Zhang, H. Fu, H. Zhong, K. Hong and W. Zhu, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 4005–4007.
- 234 J. I. Levin, P. S. Chan, T. Bailey, A. S. Katocs and A. M. Venkatesan, The synthesis of 2,3-dihydro-4(1H)-quinazolinone angiotensin II receptor antagonists, *Bioorg. Med. Chem. Lett.*, 1994, **4**, 1141–1146.
- 235 S. Guo, Y. Li, L. Tao, W. Zhang and X. Fan, *RSC Adv.*, 2014, **4**, 59289–59296.
- 236 B. Hu, L. Wang, J. Xiang, L. Yang and Y. Tang, *Chin. Chem. Lett.*, 2015, **26**, 369–372.
- 237 J. M. Khurana and G. Kukreja, *J. Heterocycl. Chem.*, 2003, **40**, 677–679.
- 238 P. Hradil, L. Kvapil, J. Hlaváč, T. Weidlich, A. Lyčka and K. Lemr, *J. Heterocycl. Chem.*, 2000, **37**, 831–837.
- 239 T. B. Nguyen, L. Ermolenko and A. Al-Mourabit, *Green Chem.*, 2013, **15**, 2713–2717.
- 240 X. Cheng, S. Vellalath, R. Goddard and B. List, *J. Am. Chem. Soc.*, 2008, **130**, 15786–15787.
- 241 M. Rueping, A. P. Antonchick, E. Sugiono and K. Grenader, *Angew. Chem., Int. Ed.*, 2009, **48**, 908–910.
- 242 T. Honjo, R. J. Phipps, V. Rauniyar and D. Toste, *Angew. Chem., Int. Ed.*, 2012, **51**, 9684–9688.
- 243 D. Huang, X. Li, F. Xu, L. Li and X. Lin, *ACS Catal.*, 2013, **3**, 2244–2247.
- 244 D. Cheng, Y. Tian and S. Tian, *Adv. Synth. Catal.*, 2012, **354**, 995–999.
- 245 B. Zhang, L. Shi and R. Guo, *Catal. Lett.*, 2015, **145**, 1718–1723.
- 246 M. Prakash and V. Kesavan, *Org. Lett.*, 2012, **14**, 1896–1899.
- 247 T. Deng, H. Wang and C. Cai, *J. Fluorine Chem.*, 2015, **169**, 72–77.
- 248 J. I. Levin, A. M. Venkatesan, P. S. Chan, J. S. Baker, G. Francisco, T. Bailey, G. Vice, A. Katocs, F. Lai and J. Coupet, *Bioorg. Med. Chem. Lett.*, 1994, **4**, 1135–1140.
- 249 Y. Kurogi, Y. Inoue, K. Tsutsumi, S. Nakamura, K. Nagao, H. Yoshitsugu and Y. Tsuda, *J. Med. Chem.*, 1996, **39**, 1433–1437.
- 250 V. J. Ram, Farhanullah, B. K. Tripathi and A. K. Srivastava, *Bioorg. Med. Chem.*, 2003, **11**, 2439–2444.
- 251 A. R. Maarouf, E. R. El-Bendary and F. E. Goda, *Arch. Pharm. Pharm. Med. Chem.*, 2004, **337**, 527–532.
- 252 S. M. Mosaad, M. K. Mohsen, M. M. K. Emad, A. Nageh, I. A. E. Sherein and F. A. Marwa, *Eur. J. Med. Chem.*, 2010, **45**, 3311–3319.



- 253 M. N. Noolvi, H. M. Patel, V. Bhardwaj and A. Chauhan, *Eur. J. Med. Chem.*, 2011, **46**, 2327–2346.
- 254 Z. Xu, Y. Zhang, H. Fu, H. Zhong, K. Hong and W. Zhu, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 4005–4007.
- 255 S. B. Mhaske and N. P. Argade, *Tetrahedron*, 2006, **62**, 9787–9826.
- 256 U. A. Kshirsagar, *Org. Biomol. Chem.*, 2015, **13**, 9336–9352.
- 257 T. Panneer Selvam and P. Vijayaraj Kumar, *Res. Pharm.*, 2015, **1**, 1–21.
- 258 R. Pater, *J. Heterocycl. Chem.*, 1971, **8**, 699–702.
- 259 Y. Peng, Y. Zeng, G. Oiu, L. Cai and V. W. Pike, *J. Heterocycl. Chem.*, 2010, **47**, 1240–1245.
- 260 M. Bakavoli, O. Sabzevari and M. Rahimizadeh, *Chin. Chem. Lett.*, 2007, **18**, 1466–1468.
- 261 A. Romero, J. Salazar and S. E. López, *Synthesis*, 2013, **45**, 2043–2050.
- 262 J. K. Laha, K. S. S. Tummalapalli, A. Nair and N. Patel, *J. Org. Chem.*, 2015, **80**, 11351–11359.
- 263 B. Hu, L. Wang, L. Yang, J. Xiang and Y. Tang, *Eur. J. Org. Chem.*, 2015, 4504–4509.
- 264 M. Kumar, Richa, S. Sharma, V. Bhatt and N. Kumar, *Adv. Synth. Catal.*, 2015, **357**, 2862–2868.
- 265 L. Wang, B. Hu, J. Xiang, J. Cui, X. Hao, T. Liang and Y. Tang, *Tetrahedron*, 2014, **70**, 8588–8591.
- 266 M. Heidary, M. Khoobi, S. Ghasemi, Z. Habibi and M. A. Faramarzi, *Adv. Synth. Catal.*, 2014, **356**, 1789–1794.
- 267 M. L. Bolognesi, *Curr. Med. Chem.*, 2013, **20**, 1639–1645.
- 268 M. L. Bolognesi and A. Cavalli, *ChemMedChem*, 2016, **11**, 1190–1192.
- 269 R. Morphy, C. Kay and Z. Rankovic, *Drug Discovery Today*, 2004, **9**, 641–651.
- 270 J. L. Mendina-Franco, M. A. Giulianotti, G. S. Welmarker and R. A. Houghten, *Drug Discovery Today*, 2013, **18**, 495–501.
- 271 R. Morphy and Z. Rankovic, *J. Med. Chem.*, 2005, **48**, 6523–6543.
- 272 N. M. Abdel Gawad, H. H. Georgey, R. M. Youssef and N. A. El-Sayed, *Eur. J. Med. Chem.*, 2010, **45**, 6058–6067.
- 273 E. Hamel, C. M. Lin, J. Plowman, H. Wang, K. Lee and K. D. Paull, *Biochem. Pharmacol.*, 1996, **51**, 53–59.
- 274 S. H. Hwang, A. Rait, K. F. Pirollo, Q. Zhou, V. M. Yenugonda, G. M. Chinigo, M. L. Brown and E. H. Chang, *Mol. Cancer Ther.*, 2008, **7**, 559–568.
- 275 S. Kumar, S. K. Guru, A. S. Pathania, N. Mupparapu, A. Kumar, F. Malik, S. B. Bharate, Q. N. Ahmed, R. A. Vishwakarma and S. Bhushan, *Toxicol. Rep.*, 2014, **1**, 1013–1025.
- 276 J. E. Stelmach, L. Liu, S. B. Patel, J. V. Pivnichny, G. Scapin, S. Singh, C. E. C. A. Hop, Z. Wang, J. R. Strauss, P. M. Cameron, E. A. Nichols, S. J. O'Keefe, E. A. O'Neil, D. M. Schmatz, C. D. Schmatz, C. M. Thompson, D. M. Zaller and J. B. Doherty, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 277–280.
- 277 A. J. Firestone, J. S. Weinger, M. Maldonado, K. Barlan, L. D. Langston, M. O'Donnell, V. I. Gelfand, T. M. Kapoor and J. K. Chen, *Nature*, 2012, **484**, 125–129.
- 278 M. Singh and N. Raghav, *Bioorg. Chem.*, 2015, **59**, 12–22.
- 279 J. Nkizinkiko, B. V. S. S. Kumar, V. U. Jeankumar, T. Haikarainen, J. Koivunen, C. Madhuri, P. Yogeeswari, H. Venkannagari, E. Obaji, T. Pihlajaniemi, D. Sriram and L. Lehtiö, *Bioorg. Med. Chem.*, 2015, **23**, 4139–4149.
- 280 A. Kamal, E. V. Bharathi, J. S. Reddy, M. J. Ramaiah, D. Dastagiri, M. K. Reddy, A. Viswanath, T. L. Reddy, T. B. Shaik, S. N. C. V. L. Pushpavalli and M. P. Bhadra, *Eur. J. Med. Chem.*, 2011, **46**, 691–703.
- 281 D. Rambabu, G. Raja, B. Y. Sreenivas, G. P. K. Seerapu, K. L. Kumar, G. S. Deora, D. Haldar, M. V. B. Rao and M. Pal, *Bioorg. Med. Chem. Lett.*, 2013, 1351–1357.
- 282 E. Cohen, B. Klarberg and J. R. Vaughan, Organic Chemical Research Section, *Lederle Laboratories Division*, American Cyanamid Company, Pearl River, New York, 1959, vol. 81, pp. 5508–5509.
- 283 C. Ferrando, J. M. Foy, F. W. Pratt and I. R. Purvis, *J. Pharm. Pharmacol.*, 1981, **33**, 219–222.
- 284 A. Marzo, G. Quadro and E. Treffner, *J. Chromatogr. B: Biomed. Sci. Appl.*, 1983, **272**, 95–102.
- 285 A. Cupisti, F. Ciardella, E. Morelli, L. Dani, S. Lupetti, S. Luchi, M. Meola and G. Barsotti, *Contrib. Nephrol.*, 1987, **58**, 184–186.
- 286 F. V. Costa, C. Borghi, S. Boschi, A. Mussi and E. Ambrosini, *J. Clin. Pharmacol.*, 1990, **30**, 254–261.
- 287 H. A. Parish and R. D. Gilliom, *J. Med. Chem.*, 1982, **25**, 98–102.
- 288 R. J. Alaimo and H. E. Russell, *J. Med. Chem.*, 1972, **15**, 335–336.
- 289 J. Zhang, J. Zhao, L. Wang, J. Liu, D. Ren and Y. Ma, *Tetrahedron*, 2016, **72**, 936–943.
- 290 D. Rambabu, S. K. Kumar, B. Y. Sreenivas, S. Sandra, A. Kandale, P. Misra, M. V. B. Rao and M. Pal, *Tetrahedron Lett.*, 2013, **54**, 495–501.
- 291 Y. Li, T. Ganesh, B. A. Diebold, Y. Zhu, J. W. McCoy, S. M. E. Smith, A. Sun and D. Lambeth, *ACS Med. Chem. Lett.*, 2015, **6**, 1047–1052.
- 292 T. J. Tucker, T. A. Lyle, C. M. Wiscount, S. F. Britcher, S. D. Young, W. M. Sanders, W. C. Lumma, M. E. Goldman, J. A. O'Brien, R. G. Ball, C. F. Homnick, W. A. Schleif, E. A. Emini, J. R. Huff and P. S. Anderson, *J. Med. Chem.*, 1994, **37**, 2437–2444.
- 293 J. W. Corbett, S. S. Ko, J. D. Rodgers, S. Jeffrey, L. T. Bacheler, R. M. Klabe, S. Diamond, C. Lai, S. R. Rabel, J. A. Saye, S. P. Adams, G. L. Trainor, P. S. Anderson and S. K. Erickson-Viitanen, *Antimicrob. Agents Chemother.*, 1999, **43**, 2893–2897.
- 294 J. W. Corbett, S. S. Ko, J. D. Rodgers, L. A. Gearhart, N. A. Magnus, L. T. Bacheler, S. Diamond, S. Jeffrey, R. M. Klabe, B. C. Cordova, S. Garber, K. Logue, G. L. Trainor, P. S. Anderson and S. K. Erickson-Viitanen, *J. Med. Chem.*, 2000, **43**, 2019–2030.
- 295 F. Liu, Y. Wang, L. Wang, J. Ou, M. Wang and S. Liu, *Molecules*, 2012, **17**, 2000–2014.
- 296 Y. Wang, M. Zhu, F. Liu, X. Wu, D. Pan, J. Liu, S. Fan, Z. Wang, J. Tang, R. Na, Q. X. Li, R. Hua and S. Liu, *Molecules*, 2016, **21**, 1373.



- 297 Y. Wang, M. Zhu, J. Liu, R. Na, F. Liu, X. Wu, S. Fan, Z. Wang, D. Pan, J. Tang, Q. X. Li, R. Hua and S. Liu, *Appl. Sci.*, 2017, **7**, 200.
- 298 P. N. Borase, P. B. Thale and G. S. Shankarling, *Dyes Pigm.*, 2016, **134**, 276–284.
- 299 K. Ozaki, Y. Yamada, T. Oine, T. Ishizuka and Y. Iwasawat, *J. Med. Chem.*, 1985, **28**, 568–576.
- 300 Y. S. Sadanandam, K. R. M. Reddy and A. B. Rao, *Eur. J. Med. Chem.*, 1987, **22**, 169–173.
- 301 O. I. El-Sabbagh, S. M. Ibrahim, M. M. Baraka and H. Kothayer, *Arch. Pharm. Chem. Life Sci.*, 2010, **343**, 274–281.
- 302 T. Katoh, T. Takai, T. Yukawa, T. Tsukamoto, E. Watanabe, H. Mototani, T. Arita, H. Hayashi, H. Nakagawa, M. G. Klein, H. Zou, H. Sang, G. Snell and Y. Nakada, *Bioorg. Med. Chem.*, 2016, **24**, 2466–2475.
- 303 A. Schlapbach, R. Heng and F. Di Padova, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 357–360.
- 304 H. L. Yale, *J. Heterocycl. Chem.*, 1977, **14**, 1357–1359.
- 305 S. Neumann, W. Huang, E. Eliseeva, S. Titus, C. J. Thomas and M. C. Gershengorn, *Endocrinology*, 2010, **151**, 3454–3459.
- 306 X. Zhu, S. R. Kang, L. Xia, J. Lee, N. Basavegowda and Y. R. Lee, *Mol. Diversity*, 2015, **19**, 67–75.
- 307 C. L. Yoo, J. C. Fetters and M. J. Kurth, *J. Org. Chem.*, 2005, **70**, 6941–6943.
- 308 L. Wang, L. Hu, J. Shao, J. Yu and L. Zhang, *J. Fluorine Chem.*, 2008, **129**, 1139–1145.
- 309 J. M. Khurana and S. Kumar, *Green Chem. Lett. Rev.*, 2011, **4**, 321–325.
- 310 A. Bharathi, S. M. Roopan, A. Kajbafvala, R. D. Padmaja, M. S. Darsana and G. N. Kumari, *Chin. Chem. Lett.*, 2014, **25**, 324–326.
- 311 A. Takács, A. Fodor, J. Németh and Z. Hell, *Synth. Commun.*, 2014, **44**, 2269–2275.
- 312 M. J. Namkung and T. L. Fletcher, *J. Med. Chem.*, 1970, **13**, 329–332.
- 313 A. A. Mohammadi, H. Rohi and A. A. Soorki, *J. Heterocycl. Chem.*, 2013, **50**, 1129–1133.
- 314 E. R. Derbyshire, J. Min, W. A. Guiguemde, J. A. Clark, M. C. Connelly, A. D. Magalhães, R. K. Guy and J. Clardy, *Antimicrob. Agents Chemother.*, 2014, **58**, 1516–1522.
- 315 C. D. S. Nelson, D. W. Carney, A. Derdowsky, A. Lipovsky, G. V. Gee, B. O'Hara, P. Williard, D. DiMaio, J. K. Sello and W. J. Atwood, *mBio*, 2013, **4**, e00729.
- 316 D. W. Carney, C. D. Nelson, B. D. Ferris, J. P. Stevens, A. Lipovsky, T. Kazakov, D. DiMaio, W. J. Atwood and J. K. Sello, *Bioorg. Med. Chem.*, 2014, **22**, 4836–4847.
- 317 R. Noel, N. Gupta, V. Pons, A. Goudet, M. D. Garcia-Castillo, A. Michau, J. Martinez, D. Buisson, L. Johannes, D. Gillet, J. Barbier and J. Cintrat, *J. Med. Chem.*, 2013, **56**, 3404–3413.
- 318 K. Hemalatha, G. Mdhumitha, L. Ravi, V. G. Khanna, N. A. Al-Dhabi and M. V. Arasu, *J. Photochem. Photobiol., B*, 2016, **161**, 71–79.
- 319 Y. Zhou, Q. Feng, F. Di, Q. Liu, D. Wang, Y. Chen, L. Xiong, H. Song, Y. Li and Z. Li, *Bioorg. Med. Chem.*, 2013, **21**, 4968–4975.
- 320 H. Hasegawa, M. Muraoka, K. Matsui and A. Kojima, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 3471–3475.
- 321 R. M. Rzaia, M. R. Kaller, G. Liu, E. Magal, T. T. Nguyen, T. D. Osslund, D. Powers, V. J. Santora, V. N. Viswanadhan, H. Wang, X. Xiong, W. Zhong and M. H. Norman, *Bioorg. Med. Chem.*, 2007, **15**, 6574–6595.
- 322 Y. Uruno, Y. Konishi, A. Suwa, K. Takai, K. Tojo, T. Nakako, M. Sakai, T. Enomoto, H. Matsuda, A. Kitamura and T. Sumiyoshi, *Bioorg. Med. Chem.*, 2015, **25**, 5357–5361.
- 323 Y. Wang, X. Tang, Z. Shao, J. Ren, D. Liu, P. Proksch and W. Lin, *J. Antibiot.*, 2014, **67**, 395–399.
- 324 J. A. Hunt, F. Kallashi, R. D. Ruzek, P. J. Sinclair, I. Ita, S. X. McCormick, J. V. Pivnichny, C. E. C. A. Hop, S. Kumar, Z. Wang, S. J. O'Keefe, E. A. O'Neill, G. Porter, J. E. Thompson, A. Woods, D. M. Zaller and J. B. Doherty, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 467–470.
- 325 H. L. Birch, G. M. Buckely, N. Davies, H. J. Dyke, E. J. Frost, P. J. Glibert, D. R. Hannah, A. F. Haughan, M. J. Madigan, T. Morgan, W. R. Pitt, A. J. Ratcliffe, N. C. Ray, M. D. Richard, A. Sharpe, A. J. Taylor, J. M. Whitworth and S. C. Williams, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 5335–5339.
- 326 K. Okumura, T. Oine, Y. Yamada, G. Hayashi and M. Nakama, *J. Med. Chem.*, 1968, **11**, 348–352.
- 327 G. Bonola, P. Da Re, M. J. Magistretti, E. Massarani and I. Setnikar, *J. Med. Chem.*, 1968, **11**, 1136–1139.

