



Cite this: *Chem. Commun.*, 2020, 56, 1529

Received 20th November 2019,
Accepted 3rd January 2020

DOI: 10.1039/c9cc09050a

rsc.li/chemcomm

Total synthesis and biological evaluation of simplified aplyronine analogues as synthetically tractable anticancer agents†

Talia R. Pettigrew,^{‡a} Rachel J. Porter,^{‡a} Stephen J. Walsh,^{IDab}
Michael P. Housden,^a Nelson Y. S. Lam,^{IDa} Jason S. Carroll,^b
Jeremy S. Parker,^{IDc} David R. Spring,^{IDa} and Ian Paterson^{ID*}

The aplyronines are a family of highly cytotoxic marine natural products with potential application in targeted cancer chemotherapy. To address the severe supply issue, function-oriented molecular editing of their macrolactone scaffold led to the design of a series of simplified aplyronine analogues. Enabled by a highly convergent aldol-based route, the total synthesis of four analogues was achieved, with a significant improvement in step economy *versus* previous compounds, and their cancer cell growth inhibition in the HeLa cell line was determined. The modular strategy presented offers a means for significantly shortening their chemical synthesis to facilitate the continued development of this promising class of anticancer agent.

Isolated in 1993 by Yamada and co-workers from the sea hare *Aplysia kurodai*,¹ the aplyronines are a family of complex polyketides comprised of a 24-membered macrolactone and an elaborate side chain terminating in an *N*-vinyl-*N*-formamide moiety.^{1–3} Aplyronines A (1, Fig. 1) and C (2) bear a C29 *N,N*-dimethylalanine (DMAIa) residue, while aplyronine D (3) contains a *N,N*-dimethylglycine (DMGIy). Aplyronines A and D further feature a *N,N,O*-trimethylserine (TMSer) residue at C7. Aplyronine A was found to display sub-nanomolar growth inhibitory activity (mean GI_{50} = 0.2 nM) in the NCI 60 cell line panel, prompting investigations into its mode of action in disrupting cytoskeletal dynamics.⁴ The 11-carbon side chain of the aplyronines is conserved across several related anticancer macrolides (reidispogiolide A (4), scytophycin C (5) and rhizopodin (6)), which all bind to actin as their protein target.⁵ Additionally, aplyronine A was found to bind to tubulin through the formation of an unprecedented actin–aplyronine–tubulin heterotrimeric complex.⁶ Interestingly, actin binding was shown




Fig. 1 The aplyronine family of cytotoxic marine macrolides, highlighting the structural differences between the congeners from C7 and C29 acyl substitution. Key regions implicated in protein binding are denoted, as well as their structural homology to related actin-binding macrolides.

to be a necessary but insufficient marker for their cytotoxicity, where aplyronine A is roughly 50 times more cytotoxic than aplyronine C, despite registering similar actin depolymerising ability. This structure–activity relationship (SAR) highlighted the synergistic role of the macrolactone, specifically the C1–C13 region, with the C7 TMSer important in forming the crucial interaction with tubulin in aplyronines A and D.⁷

Their extraordinary potency and novel mode of action mark the aplyronines out as promising candidates as novel payloads for antibody–drug conjugates.^{8,9} While recent total syntheses of the aplyronines have been achieved in 29–38 steps LLS (and up to 80 total steps),^{9,10} their structural complexity and natural scarcity (2.6 mg of aplyronine D was isolated from 300 kg of *Aplysia kurodai*)^{1b} adversely impact their further development. To address these challenges, we turned towards a function-oriented molecular editing of the aplyronine scaffold to help

^a University Chemical Laboratory, Lensfield Road, Cambridge, CB2 1EW, UK

^b Cancer Research UK Cambridge Institute, University of Cambridge, Cambridge, CB2 0RE, UK

^c Early Chemical Development, Pharmaceutical Sciences, R&D, AstraZeneca, Macclesfield, UK

† Electronic supplementary information (ESI) available: Full experimental procedures and discussion. See DOI: 10.1039/c9cc09050a

‡ Authors contributed to this work equally.



improve the step economy while retaining the biological activity.¹¹ Herein, we report the concise total synthesis of a series of simplified aplyronine analogues in 24–26 steps LLS (37–39 total steps) and their preliminary biological evaluation.


Inspection of the aplyronine A–actin complex crystal structure,⁵ and taking previous SAR studies³ into account, indicated that the C16–C22 region of the macrolactone did not appear to form any significant interaction with the protein binding site (see the ESI† for discussion). As outlined in Scheme 1, we proposed that this region was primarily acting as a conformational constraint for the macrocycle and that its replacement by a simple (CH₂)₇ unit should preserve the 3D structure and critical C1–C13 binding region, and hence maintain its biological activity. Additionally, our proposed


aplyrologues involved replacing the C25 hydroxyl and C31 acetoxy groups with methyl ethers, which simplifies the protecting group strategy, as well as removing the propensity for the acetate to eliminate (Scheme 1).⁹ Moreover, these latter modifications mirror the functionalities present in the equivalent side chains (Fig. 1) present in reidispongiolide A (**4**), scytopycin C (**5**) and rhizopodin (**6**). We envisaged that these designed aplyrologues¹² would preserve the core pharmacophore, while significantly improving the synthetic tractability for this promising class of anticancer agent.

Building on our recent total synthesis of aplyronines A and D,⁹ we planned a late-stage diversification to construct aplyrologues bearing the aplyronine D side chain, as in **7** (Scheme 1), as well as the scytopycin–aplyronine A/D (bearing the C7 TMSer: **8**) and C (**9**) analogues. This analysis leads back to the common scaffold **10**, which would be forged through an aldol coupling sequence with the C28–C34 ketone *E-11* and a macrocyclic aldehyde. The macrocycle itself can then be constructed by a Horner–Wadsworth–Emmons (HWE) olefination with the simplified C15–C27 aldehyde **12** and the previously reported C1–C14 phosphonate **13**.^{9a}


Construction of the C15–C27 aldehyde **12** (Scheme 2) commenced with a Ti(OiPr)₃-mediated aldol reaction¹³ between the Roche ester-derived ketone *S-13*¹⁴ and the aldehyde **14** to give the 1,4-*syn*, 3,4-*syn* adduct **15**. An Evans–Tishchenko reduction¹⁵ then delivered the 1,3-*anti* configured alcohol **16**, which was methylated, desilylated and oxidised to afford the aldehyde **12** in five steps from **13** and **14**. Notably, this expedient sequence represented more than a three-fold reduction in the step count relative to the analogous fragment required for the aplyronines.⁹

Synthesis of the modified C28–C34 side chain (Scheme 3) commenced with an analogous titanium aldol reaction between the ketone *R-13* and acetaldehyde to give **17**. Adduct **17** was reduced to give alcohol **18** followed by methylation, PMB ether cleavage and ester reduction to afford the diol **19**. At this point, a double oxidation under Swern conditions gave ketoaldehyde **20**, which was directly subjected to a Wittig olefination¹⁶ with phosphonium salt **21** to generate the enamides *E/Z-11*. Final completion of the configurationally pure C28–C34 ketone *E-11* was accomplished by I₂-mediated isomerisation.





With this first series of aplyrologues in hand, attention now turned towards determining their *in vitro* cytotoxicity. It was considered prudent to initially examine the activity of our previously synthesised samples of aplyronines A (1), C (2) and D (3), with discodermolide included as a control (Table 1).



While the expected sub-nanomolar potencies were found for aplyronines A and D, we did not find any significant difference between these congeners in the same HeLa cell line (Table 1, entries 1–3). However, our synthetic aplyrologues did not register the same exquisite potency compared with the native aplyronines. Nevertheless, the compounds *R*- and *S*-8 (entries 5 and 6), bearing the scytophycin side chain, as well as 7 (entry 4) still gave inhibitory activities in the nM range. Interestingly, both the configuration of the TMSer residue and the nature of the C29 functionality as the ketone or DMGly ester did not appear to greatly influence the biological activity, suggesting that the deleterious effect may be attributable to an oversimplification of the southern hemisphere region of the macrolactone.

Table 1 Antiproliferative activity of synthetic aplyronines and aplyrologues against the HeLa cell line (discodermolide was included as a control)

Entry	Compound	IC ₅₀ ^a (nM)
1	Aplyronine A (1)	0.31
2	Aplyronine C (2)	3.67
3	Aplyronine D (3)	0.26
4	7 (aplyrologue D)	124
5	S-8 (aplyronine A, D/scytophycin C hybrid)	164
6	R-8 (aplyronine A, D/scytophycin C hybrid)	192
7	9 (aplyronine C/scytophycin C hybrid)	>1000
8	Discodermolide	3.40

^a *n* = 3. See the ESI for cytotoxicity curves and further experimental details.

In conclusion, the preliminary biological results obtained for this first series of aplyrologues suggests that the southern hemisphere of the macrocycle may play a subtle but vital role in reinforcing the proposed actin–tubulin interaction that the aplyronines uniquely mediate. Additionally, the biological data obtained for our synthetic aplyronines does not replicate the greater cytotoxicity previously reported for a natural sample of aplyronine D relative to aplyronine A,^{1b} signifying the scope for structural diversification of the side chain. In future work, further judicious molecular editing of the southern hemisphere region can be flexibly adapted towards the incorporation of more elaborate macrocyclic tethers. We envisage that our modular synthetic platform offers a means for enhancing the synthetic tractability of the aplyronines to facilitate the continued development of this promising class of anticancer agent.²²

We thank Trinity College (TRP), AstraZeneca (RJP and SJW), EPSRC (RJP and SJW), Downing College (MPH) and the Woolf Fisher Trust (NYSL) for financial support, Dr Nika Anžicek and Dr Simon Williams for helpful discussions, and the National Mass Spectrometry Centre (Swansea) for mass spectra.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- (a) K. Yamada, M. Ojika, T. Ishigaki, Y. Yoshida, H. Ekimoto and M. Arakawa, *J. Am. Chem. Soc.*, 1993, **115**, 11020; (b) M. Ojika,

- H. Kigoshi, K. Suenaga, Y. Imamura, K. Yoshikawa, T. Ishigaki, A. Sakakura, T. Mutou and K. Yamada, *Tetrahedron*, 2012, **68**, 982.
- (a) M. Ojika, H. Kigoshi, T. Ishigaki, I. Tsukada, T. Tsuboi, T. Ogawa and K. Yamada, *J. Am. Chem. Soc.*, 1994, **116**, 7441; (b) H. Kigoshi, M. Ojika, T. Ishigaki, K. Suenaga, T. Mutou, A. Sakakura, T. Ogawa and K. Yamada, *J. Am. Chem. Soc.*, 1994, **116**, 7443.
- (a) H. Kigoshi, K. Suenaga, M. Takagi, A. Akao, K. Kanematsu, N. Kamei, Y. Okugawa and K. Yamada, *Tetrahedron*, 2002, **58**, 1075; (b) M. Ojika, H. Kigoshi, Y. Yoshida, T. Ishigaki, M. Nisiwaki, I. Tsukada, M. Arakawa, H. Ekimoto and K. Yamada, *Tetrahedron*, 2007, **63**, 3138.
- K. Hirata, S. Muraoka, K. Suenaga, T. Kuroda, K. Kato, H. Tanaka, M. Yamamoto, M. Takata, K. Yamada and H. Kigoshi, *J. Mol. Biol.*, 2006, **356**, 945.
- (a) K.-S. Yeung and I. Paterson, *Angew. Chem., Int. Ed.*, 2002, **41**, 4632; (b) J. S. Allingham, V. A. Klenchin and I. Rayment, *Cell. Mol. Life Sci.*, 2006, **63**, 2119.
- M. Kita, Y. Hirayama, K. Yoneda, K. Yamagishi, T. Chinen, T. Usui, E. Sumiya, M. Uesugi and H. Kigoshi, *J. Am. Chem. Soc.*, 2013, **135**, 18089.
- M. Kita and H. Kigoshi, *Nat. Prod. Rep.*, 2015, **32**, 534.
- (a) R. V. J. Chari, M. L. Miller and W. C. Widdison, *Angew. Chem., Int. Ed.*, 2014, **53**, 3796; (b) K. C. Nicolaou and S. Rigol, *Angew. Chem., Int. Ed.*, 2019, **58**, 11206; (c) D. J. Newman, *Mar. Drugs*, 2019, **17**, 324.
- (a) N. Anžicek, S. Williams, M. P. Housden and I. Paterson, *Org. Biomol. Chem.*, 2018, **16**, 1343; (b) I. Paterson, S. J. Fink, L. Y. W. Lee, S. J. Atkinson and S. B. Blakey, *Org. Lett.*, 2013, **15**, 3118.
- I. Hayakawa, K. Saito, S. Matsumoto, S. Kobayashi, A. Taniguchi, K. Kobayashi, Y. Fujii, T. Kaneko and H. Kigoshi, *Org. Biomol. Chem.*, 2017, **15**, 124.
- P. A. Wender, R. V. Quiroz and M. C. Stevens, *Acc. Chem. Res.*, 2015, **48**, 752.
- (a) K. Futaki, M. Takahashi, K. Tanabe, A. Fujiedo, H. Kigoshi and M. Kita, *ACS Omega*, 2019, **4**, 8598; (b) T. Ohyoshi, A. Takano, M. Namiki, T. Ogura, Y. Miyazaki, Y. Ebihara, K. Takeno, I. Hawakawa and H. Kigoshi, *Chem. Commun.*, 2018, **54**, 9537.
- J. G. Solsona, J. Nebot, P. Romea and F. Urpí, *J. Org. Chem.*, 2005, **70**, 6533.
- I. Paterson, G. J. Florence, K. Gerlach, J. P. Scott and N. Sereinig, *J. Am. Chem. Soc.*, 2001, **123**, 9535.
- D. A. Evans and A. H. Hoveyda, *J. Am. Chem. Soc.*, 1990, **112**, 6447.
- I. Paterson, C. J. Cowden, V. S. Rahn and M. D. Woodrow, *Synlett*, 1998, 915.
- I. Paterson, K.-S. Yeung and J. B. Smaill, *Synlett*, 1993, 774.
- E. J. Corey, R. K. Bakshi, S. Shibata, C. P. Chen and V. K. Singh, *J. Am. Chem. Soc.*, 1987, **109**, 7925.
- I. Paterson, C. J. Cowden, V. S. Rahn and M. D. Woodrow, *Synlett*, 1998, 915.
- J. Inanaga, K. Hirata, H. Saeki, T. Katsuki and M. Yamaguchi, *Bull. Chem. Soc. Jpn.*, 1979, **52**, 1989.
- E. P. Boden and G. E. Keck, *J. Org. Chem.*, 1985, **50**, 2394.
- For other recent work from our group, see: (a) I. Paterson and N. Y. S. Lam, *J. Antibiot.*, 2017, **71**, 215; (b) N. Y. S. Lam and I. Paterson, *Eur. J. Org. Chem.*, DOI: 10.1002/ejoc.201901243; (c) N. Y. S. Lam, G. Muir, V. R. Challa, R. Britton and I. Paterson, *Chem. Commun.*, 2019, **55**, 9717; (d) M. Anketell, T. Sharrock and I. Paterson, *Angew. Chem., Int. Ed.*, DOI: 10.1002/anie.201914042.