

Targeting cancer cells with folic acid–iminoboronate fluorescent conjugates†

Pedro M. S. D. Cal,^a Raquel F. M. Frade,^a Vijay Chudasama,^b Carlos Cordeiro,^c Stephen Caddick^b and Pedro M. P. Gois^{*a}

Cite this: *Chem. Commun.*, 2014, 50, 5261


Received 1st October 2013,
Accepted 4th November 2013

DOI: 10.1039/c3cc47534d

www.rsc.org/chemcomm

Herein we present the synthesis of fluorescent 2-acetylbenzeneboronic acids that undergo B–N promoted conjugation with lysozyme and *N*-(2-aminoethyl) folic acid (EDA-FA), generating conjugates that are selectively recognized and internalized by cancer cells that over-express folic acid receptors.

Boronic acids are well-defined neutral, planar trivalent Lewis acids that are readily converted into negatively charged tetravalent borates by forming reversible complexes with Lewis base donors such as amines.¹ Appreciation of the unique B–N bond properties triggered a burgeoning interest for this bonding motif.² B–N bonds have been extensively exploited to construct self-assembled molecularly defined nanostructures, polymeric materials and molecular sensors for the detection of carbohydrates.³ Recently, the isosterism between B–N and C–C bonds was also recognized as a powerful tool to tune the stereo electronic properties of organic molecules.⁴ In this context, we used the B–N bond to prepare natural product-like structures and heterocycles with inhibitory activity against human neutrophil elastase.⁵ More recently, we found that this bonding motif could be exploited to efficiently modify proteins at their N-terminal or at lysine's ϵ -amino group, *via* the formation of alkylic iminoboronates in aqueous media (Scheme 1).⁶ Despite their stability, these modifications were also shown to be reversible in the presence of fructose, dopamine and glutathione, as they presumably induce hydrolysis of the iminoboronate by disruption of the B–N bond.² Based on this, we became very interested in understanding if the B–N bond could also be used to assemble constructs that are able to internalize into cells. If so, this simple, selective and reversible methodology to functionalize proteins may



Scheme 1 Protein modification *via* the formation of iminoboronates. Proposed structure of N–B based cancer cell targeting fluorescent conjugates.

develop into an innovative strategy to design conjugates that may selectively target and deliver cargo to cancer cells.⁷

In this communication, we address for the first time, the synthesis and evaluation of cancer cell targeting fluorescent conjugates, in which the recognition biomolecule is connected to the fluorescent motif *via* a B–N linkage (Scheme 1). In our previous study, 2-acetylbenzeneboronic acid afforded the most stable iminoboronate in buffer solutions (pH range 6–9).⁶ Consequently, this scaffold was selected to integrate the fluorescent motifs as outlined in Scheme 1.

We initiated this study with the synthesis of fluorescent derivatives of 2-acetylbenzeneboronic acid. As shown in Scheme 2, 2,4-dihydroxyacetophenone was alkylated selectively with ethyl 4-bromobutyrate in a 5 g scale to yield compound **2** in 87% isolated yield. After this, phenol **2** was converted to a triflate, which then underwent a Miyaura borylation reaction with bis(pinacolato)diboron to afford **3** in 64% yield. Hydrolysis of the ester with TFA in aqueous media led to acid **4**, which became the core structure for connecting fluorescent probes to targeting molecules. Dansyl and nitrobenzofurazan fluorescent motifs were modified with ethylenediamine and ethanolamine, respectively, and each subsequently reacted with **4**. Removal of the pinacol group using a boronic acid resin under acidic conditions afforded the final fluorescent compounds **5** and **6** in moderate yields (Scheme 2). Once prepared, compounds **5** and **6** were immediately tested for their ability to conjugate with lysozyme. Gratifyingly, both compounds retained their ability

^a Instituto de Investigação do Medicamento (iMed.Ulisboa), Faculdade de Farmácia, Universidade de Lisboa Av. Prof. Gama Pinto, 1649-003 Lisboa, Portugal.


E-mail: pedrogois@ff.ul.pt; Fax: +351 217946470; Tel: +351 217946400

^b Department of Chemistry, University College London, 20 Gordon St, London, UK

^c Centro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências da Universidade de Lisboa, 1749-016 Lisboa, Portugal

† Electronic supplementary information (ESI) available: Experimental procedures, full DFT reference list, atomic coordinates for the optimized species, and spectral data for all compounds. See DOI: 10.1039/c3cc47534d







to functionalize the protein, yielding the expected constructs in ammonium acetate buffer (50 mM, pH 7.0) at room temperature as confirmed by ESI-FTICR-MS (Schemes 3 and 4). Encouraged by these results, we next evaluated the possibility of designing a cancer cell targeting conjugate, in which the fluorescent motif is linked to the targeting molecule by a B–N bond.

Folic acid is essential for cell functioning, however, a considerable percentage of cancer cell lines over-express receptors for this small vitamin due to their fast cell division and growth.⁸ Consequently, folic acid has been extensively used as a recognition moiety in many conjugates.⁹ Based on this, the conjugation of **5** and **6** with *N*-(2-aminoethyl) folic acid (EDA-FA) was evaluated. As expected, upon addition of 100 μ M of **5** or **6** in DMSO to EDA-FA in ammonium acetate buffer (50 mM, pH 7.0) at room temperature, their corresponding iminoboronates were formed readily, as characterized by high resolution mass spectrometry – HRMS (Schemes 3 and 4). Following this, the ability of the conjugates to differentiate between cancer and non-cancer cell lines was studied. To do this, conjugate **8** was generated prior to exposure to cells and tested against human non-small lung cancer cells (NCI-H460) and non-cancer human kidney embryonic cells (HEK 293). Interestingly, laser scanning confocal images of HEK 293 and NCI-H460 treated with **8** revealed that only the cancer cell line that over-expressed the FA receptor up took fluorescence, which suggests a selective folate-receptor mediated internalization of the B-N construct **8**.

To further elucidate this mechanism, the cancer cell line NCI-H460 was treated with **6** without EDA-FA. In agreement with a folate-receptor mediated internalization mechanism, without the recognition moiety, no fluorescence was detected inside the cells. In contrast, when the same compound **6** was allowed to form a conjugate with EDA-FA, the previously generated B-N construct **10** smoothly underwent internalization as shown in Scheme 4. Finally, the NCI-H460 cells were treated with EDA-FA



Scheme 3 Reaction of lysozyme (10 μM) with **5** (200 μM) in acetate buffer (50 mM, pH 7) at room temperature. Zoom of the (7+) charge state of the ESI-FTICR-MS spectrum. Reaction of EDA-FA (100 μM) with compound **5** (100 μM) in acetate buffer (50 mM, pH 7) to obtain **8** confirmed by HRMS. HEK (A) and NCI-H460 (B) cells were incubated with compound **8** at 20 μM for 4 h. Images show an overlay of the emission of compound **8** in the 460–580 nm region upon excitation at 458 nm (green), and the emission of the membrane marker in the 650–700 nm region upon excitation at 514 nm (red).



Scheme 4 Reaction of lysozyme (10 μ M) with **6** (200 μ M) in acetate buffer (50 mM, pH 7) at room temperature. Zoom of the (7+) charge state of the ESI-FTICR-MS spectra after 30 min and 2 h of reaction. Reaction of EDA-FA (100 μ M) with compound **6** (100 μ M) in acetate buffer (50 mM, pH 7) to obtain **10** as confirmed by HRMS. NCI-H460 (C) and NCI-H460 without (D) and with pre-treatment with EDA-FA (E) were incubated with compound **10**. The procedure is described in Scheme 3.

