

Cite this: *Chem. Sci.*, 2026, 17, 2582

All publication charges for this article have been paid for by the Royal Society of Chemistry

The dynamic chemistry of the boron–nitrogen bond

Federico Fratello,  ^{†*a} Giorgio Capocasa,  ^{†*a} Martina De Angelis,  ^{†a} Greta Sandri,^a Osvaldo Lanzalunga,  ^a Chiara Massera  ^{*b} and Stefano Di Stefano  ^{*a}

Here we report that fully reversible B←N bond formation/cleavage is a promising tool for the achievement of dynamic libraries (DLs) of rapidly interconverting compounds. The composition of a number of minimal DLs of adducts between phenylboronic acid catechol ester **1** and a series of nitrogen-based aromatic heterocycles (*NArHets*) is demonstrated to be predictable taking into account the association constants related to the formation processes of the single adducts involved. Furthermore, such composition can be controlled over time by the use of activated carboxylic acids (ACAs). Depending on the amount of added ACA, a B←N based DL can be either overturned in terms of composition, transiently overexpressing an adduct initially under-expressed, or transiently fully disassembled into its building blocks.

Received 3rd October 2025
Accepted 26th November 2025

DOI: 10.1039/d5sc07665j

rsc.li/chemical-science

Introduction

Dynamic combinatorial chemistry (DCC)^{1–8} has been an intense field of investigation for the past two decades due to its implications in different topics such as supramolecular recognition,^{9–12} catalysis,^{13–17} dissipative systems,^{18–29} and systems chemistry.^{30–36} Both covalent and supramolecular reversible bonds have been largely employed to constitute dynamic libraries (with this term, DLs, one typically refers to collections of compounds able to interconvert by exchanging building blocks under equilibrium conditions), with the latter generally having the advantage of very fast kinetics on the human timescale. Nevertheless, the presence of a catalyst may confer the same feature also to covalent dynamic bonds such as imine, acetal, and olefin bonds, among others. Here we report a systematic investigation of the exchange reactions involving the Lewis-pairs formed between phenylboronic acid catechol ester (**1**) and several nitrogen-based aromatic heterocycles (*NArHets*), specifically a series of pyridines and *N*-methylimidazole. Such reaction entails the formation/cleavage of the dynamic, covalent (dative) B←N bond. The process turns out to be fast on the human timescale and fully reversible, even in the absence of any catalyst. The B←N exchange process involving catechol boronic esters and *NArHets* has been already exploited

to build a variety of complex molecular structures.^{37–40} Several examples involve the achievement of macrocycles,^{41–45} molecular cages,^{46,47} rotaxanes,^{48,49} supramolecular adducts,^{50–55} covalent organic frameworks (COFs),^{56–61} and polymeric materials.^{62–67}

However, despite the episodic use of the B←N bond in the design of (supra)molecular architectures, we feel that a systematic investigation of its properties is still missing. In this work, we aim to build a deeper mechanistic comprehension of the properties of such bonds and to frame the B←N Lewis pair formation within the field of dynamic combinatorial chemistry as a versatile tool for the generation of DLs.

Experimental section

Materials

All the *NArHets*, amines, catechol, phenylboronic acid, and tribromoacetic acid were purchased from Fluorochem, TCI or Merck. Deuterated chloroform was purchased from Fluorochem. Non-deuterated solvents were purchased from Carlo Erba. Deuterated chloroform was passed through short plugs of Na₂SO₄ (anhydrous) and Al₂O₃ (activated, basic) before use to remove excess water and traces of acids. NMR spectra were recorded on either a BrukerDPX300 or a BrukerDPX400 spectrometer and were internally referenced to the residual proton solvent signal. See SI for further details.

Synthesis of **1**

Catechol (451 mg, 4.1 mmol) was added to a dichloromethane (14 mL) suspension of phenylboronic acid (500 mg, 4.1 mmol). Then, ethyl acetate was added dropwise under stirring until a homogeneous solution was obtained. The resulting mixture

^aDipartimento di Chimica, Università di Roma La Sapienza and ISB-CNR Sede Secondaria di Roma – Meccanismi di Reazione, P.le A. Moro 5, I-00185 Roma, Italy. E-mail: federico.fratello@uniroma1.it; giorgio.capocasa@uniroma1.it; stefano.distefano@uniroma1.it

^bDipartimento di Scienze Chimiche, della Vita e della Sostenibilità Ambientale, Università degli Studi di Parma, Parco Area delle Scienze 17/A, 43124 Parma, Italy. E-mail: chiara.massera@unipr.it

[†] These authors contributed equally.



was stirred at room temperature overnight. The solution was dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure. Recrystallization from hot hexane afforded **1** as clear needle-like crystals (650 mg, 81%). See SI for further information.

Results and discussion

First, a series of titration experiments was carried out in order to quantify the strength of the interaction between phenylboronic acid catechol ester **1**, which can be considered a benchmark boron-based substrate, and a series of *N*ArHets in CDCl_3 at 25 °C, see Fig. 1a for the relevant equilibria.

The equilibrium constants (K_{bind}) for the formation of complexes **1**·**L** with **L** = {**2**–**6**} were obtained by plotting the chemical shift of the ^1H NMR signal of proton H_{ortho} on **1** against the *N*ArHets concentration. A 1 : 1 binding model was fitted to the experimental data (see Fig. 1c for the case of the **1**·**3** adduct and SI for the other cases). For complexes **1**·**2** and **1**·**5**, the obtained K_{bind} values are in good accordance with the values previously measured in the same solvent.³⁸

Upon addition of *N*ArHets **2**–**6** to a solution of **1**, the formation of complexes **1**·(**2**–**6**) occurs immediately and smoothly with no detectable side-product accumulating in solution. These equilibria are fast on the ^1H NMR timescale, as



Fig. 1 (a) Binding processes between phenylboronic acid catechol ester **1** and *N*ArHets **2**–**6**. (b) Overall proposed transformation occurring in the presence of amines **7**–**11**. (c) ^1H -NMR titration of 5.0 mM **1** with **3** in CDCl_3 at 25 °C (see Fig. S23–S32 for the other titrations). (d) Brønsted correlation between $\log K_{\text{bind}}$ (or $\log K_{\text{obs}}$, for aliphatic amines **7**–**11**) and $\text{p}K_{\text{a}}(\text{H}^+)$ for compounds **2**–**11** (for pyridines, $\rho = 0.68$, $R^2 = 0.97$, see Table S2 for details). The linear correlation only holds for *N*ArHets **2**–**6**; the data points relative to amines **7**–**11** lay outside the line. (e) Comparison of ^1H -NMR traces of **1** (bottom, black), a mixture of **1** and **5** (centre, green), and a mixture of **1** and **8** (top, gold). Spectra recorded at 25 °C; all compounds are 10 mM in CDCl_3 . Only the aromatics portion is shown. See Fig. S51 for full spectra. The H_{ortho} protons on the phenyl moiety on **1** are rendered equivalent in the ^1H -NMR spectra by the fast rotation about the Ph–B bond.





Fig. 3 Linear correlation between the experimental stabilisation energies derived from K_{bind} measurements, and the computed stabilisation energies for the 1-NArHet adducts.

the $r^2\text{SCAN-3c}$ level of theory, using the SMD solvation model (chloroform).^{73,74} Single point energies and frequencies of the optimised structure were then computed at the $r^2\text{SCAN-3c/def2-QZVPP(SMD)}$ level of theory, following a benchmark study focused on Lewis acid–base interaction.⁷⁵

The stabilisation energies (E_{stab}) of the interaction between **1** and NArHets were calculated as the difference between the energy of the adduct (E_{adduct}) and the sum of the energies of the donor NArHet (E_{d}), and the acceptor **1** (E_{a}), i.e. $E_{\text{stab}} = E_{\text{adduct}} - (E_{\text{d}} + E_{\text{a}})$.⁷⁶ E_{stab} values were then compared with those derivable

from the experimentally measured binding constants. Fig. 3 shows a good linear correlation between experimental and theoretical data, with a systematic discrepancy, on average, of about $0.57 \text{ kcal mol}^{-1}$ (see Table S27). Moreover, the calculated B–N distances are comparable with those obtained from X-ray crystallography (see Table S28). To rationalise the similar H_{ortho} chemical shift saturation values observed for the 1-NArHets adducts, Hirshfeld partial charges were computed (see Table S28). The averaged H_{ortho} partial charges for each 1-NArHets range from 0.0313 for **1·5**, to 0.0329 for **1·2** with the average value between all the adducts being 0.0321. These values are all very similar, but they are markedly different than the one obtained for free **1**, which is 0.0479. The computed partial charges fit well with lower, very similar values of the H_{ortho} chemical shift for all the adducts with respect to **1**.

Having gained some chemical understanding of the dynamic B←N bond, we turned to building and controlling dynamic libraries of increasing complexity by exploiting its properties.

Therefore, six competitive experiments, each one involving two couples of adducts **1·L** (with $L = \text{NArHets } 2\text{--}5$) were carried out in CDCl_3 at 25°C as depicted in Fig. 4, where the equilibrium for the couples **1·2** and **1·3** is shown. In all cases, compound **1** and the two given NArHets were added in equimolar amount (12.5 mM), and it was found that the exchange reaction was fast on the ^1H NMR time scale, with the equilibrium reached immediately after the addition of the reagents (first spectrum recorded). K_{eq} for each exchange reaction can be calculated from the ratio between the corresponding K_{bind} for each adduct. For example, in the case of the equilibrium **1·2** + **3**



Fig. 4 An exchange experiment in which pyridine **2** (red) and 4-methyl pyridine **3** (blue) compete for phenylboronic acid catechol ester **1** (CDCl_3 , 25°C). All the equilibria are fast on the ^1H -NMR time scale with adduct **1·3** prevailing on adduct **1·2** under competitive conditions (trace c). Trace a 12.5 mM **2**, trace b 12.5 mM **1** and **2**, trace c 12.5 mM **1**, **2** and **3**, trace d 12.5 mM **1** and **3**, trace e 12.5 mM **3**. The black circles are relative to the signals on the **1** backbone.





Fig. 5 Top: Simplified reaction scheme for the consecutive generation of DLs containing compounds **1** (black), **3** (blue), **4** (pink), **6** (light purple), and **5** (green), in order of addition together with complexes **1·3**, **1·4**, **1·6**, and **1·5** (in order of increasing stability). Bottom: NMR spectra of **1** (trace a); a mixture of **1** and **3** (trace b); a mixture of **1**, **3**, and **4** (trace c); a mixture of **1**, **3**, **4**, and **6** (trace d); a mixture of **1**, **3**, **4**, **6**, and **5** (trace e). In each spectrum, the signal relative to each H α proton is highlighted in the same colour as the NARHet (see structures). The H_{ortho} proton is marked with a circle in the colour corresponding to the most prevalent adduct. To the right of each spectrum, a pie chart reports the calculated speciation of each DL. See Tables S17 and S18 for the ChemEQL-assisted speciation of the mixtures. All compounds are added 12.5 mM in CDCl₃, the spectra are recorded at 25 °C. Only the aromatics region is shown. See Fig. S53 for the full spectra. * = in the absence of any ligands, **1** is in equilibrium with a small amount (<5% in our conditions) of phenylboronic acid and catechol due to partial hydrolysis from traces of water in the solvent. The addition of NARHets suppresses this side-reaction.

\rightleftharpoons **1·3** + **2** which corresponds to trace c in Fig. 3, $K_{\text{eq}} = K_{\text{bind}}(\mathbf{1}\cdot\mathbf{3})/K_{\text{bind}}(\mathbf{1}\cdot\mathbf{2}) = 4.4$ with adduct **1·3** abundantly prevailing over **1·2** (free **1** 30% of **1**_{tot}, **1·3** 50% of **1**_{tot} and **1·2** 20% of **1**_{tot}, see Fig. S45 and the related caption in the SI for details on the mixture composition obtained both experimentally and theoretically). As expected, since in the case of both adducts **1·2** and **1·3** the adopted conditions do not allow for a complete binding, the presence of two molar equivalents of bases (**2** + **3** with respect to **1**) in the competitive experiment causes a further up-field shift of all signals related to the boronic ester (identified by

a black circle in Fig. 4). As for the remaining five DLs the speciation was calculated and the theoretical amount of free **1** was satisfactorily compared to that calculated from NMR data (see Tables S7–S9). Being the strongest binder for ester **1**, 4-dimethylaminopyridine **5** shifts the equilibrium toward adduct **1·5** whenever it is involved (see Fig. S47, S49, and S50). These experiments clearly demonstrate the ease of achieving DLs of B←N adducts able to rapidly interconvert on the human timescale, with a predictable composition.





Fig. 6 A minimal DL composed of 1·5, 1·3, and related free building blocks driven by ACA 12-CO₂H (see text). All compounds (1, 3, 5 and 12-CO₂H) are added in solution at 15.0 mM concentration (CDCl₃, 25 °C). On each spectrum, the relevant signals for 3 are marked in blue, while those of 5 are coloured in green. Spectra (a–e) are recorded over the course of the reaction. Spectrum (f) shows the signals of 5H⁺, obtained by adding in solution 5 and trifluoroacetic acid at 15.0 mM concentration.

Next, we performed consecutive competition experiments, building DLs with an increased number of components (Fig. 5 shows the reaction scheme, the NMR spectra, and the computed speciation for each DL). Thus, 1.0 mol equiv. of 3 was added to a 12.5 mM CDCl₃ solution of 1 (1.0 mol equiv., black trace). Complex 1·3 partially formed and, as expected, the signal relative to the H_{ortho} proton on 1 was shifted upfield, while those belonging to 3 moved downfield (blue trace). Addition of 1.0 mol equiv. of 4 to this mixture partially displaced 3, as evidenced by the upfield shift of the protons on the latter (pink trace). The H_{ortho} signal also moved upfield, indicating increased binding of 1, which was distributed between its free form and the bound ones (1·3 and 1·4). Then, 1.0 mol equiv. of 6 was added to the solution, causing a further upfield shift of the signals belonging to 3 and 4, as well as of the H_{ortho} proton (light purple trace). Finally, 1.0 mol equiv. of 5 were added to the mixture. This resulted in an upfield shift of the signals belonging to all the previously added ligands and brought the H_{ortho} proton to its saturation value of 7.50 ppm, indicating complete binding.

Thus, at the end of the experiment, the DL contained compounds 1, 3, 4, 6, and 5 (in order of addition) together with complexes 1·3, 1·4, 1·6, and 1·5 (in order of increasing stability). After each addition, the computed amount of free 1 was satisfactorily compared to that estimated from NMR data (see Fig. S53, Tables S17 and S18).

Next, we investigated the chance to control over time a minimal DL composed of two B–N adducts and related NAr-Hets using an activated carboxylic acid (ACA) as a stimulus. ACAs are used for the operation of dissipative chemical systems,

which possess one or more Brønsted basic sites.^{77–89} For instance, the DL 1·3 + 5 ⇌ 1·5 + 3 was rapidly obtained by adding in CDCl₃ 15.0 mM equimolar 1, 3 and 5 (Fig. 6, ¹H NMR, trace a). As expected from the *K*_{bind} values reported in Fig. 1a, adduct 1·5 largely prevails on 1·3. Addition of 15.0 mM ACA 12-CO₂H (tribromoacetic acid) causes the extensive protonation of 5, the strongest base present in solution, which is subtracted from the equilibrium. The latter is now shifted to the left with 1·3 prevailing on 1·5.

Comparison between traces a and b of Fig. 6 shows that all signals are shifted down-field soon after the addition of 12-CO₂H (trace b). Such shifts are due to protonation of 5 to 5H⁺, whose signals are found down-field shifted with respect to the corresponding ones in adduct 1·5 and to partial association of 3 to 1, which causes the down-field shift of the signals related to both moieties 3 and 1.

However, the new state (Fig. 6, trace b) is not an equilibrium one since the ACA conjugate base 12-CO₂[−] slowly decarboxylates and the corresponding, just formed carbanion 12[−] retakes the proton from 5H⁺ to give free pyridine 5 and bromoform 12 (traces b to e).

Consequently, the initial equilibrium with 1·5 prevailing on 1·3 is restored (trace e). Thus, when the decarboxylation of ACA 12-CO₂H is complete, the DL goes back to the initial equilibrium state. The above interpretation of the experimental results is strictly consistent with the higher sensitivity of the affinity of the basic nitrogen atoms for the proton than for the boron atom of 1 (for log *K*_{bind} vs. p*K*_aH⁺ ρ = 0.68, see Fig. 1d and S33 and the related caption for more details). Furthermore, a computer assisted calculation of the DL speciation based on the p*K*_aH⁺ of





Fig. 7 Behaviour of a DL based on the B←N bond under the action of excess ACA 12-CO₂H. Traces (a) 10 mM **1**; (b) adduct **1·5** obtained by mixing 10 mM **1** with 10 mM **5**; (c) adduct **1·3** obtained by mixing 10 mM **1** and 10 mM **3**; (d) equilibrated solution of 10 mM **1**, **3** and **5** before addition of 30 mM ACA 12-CO₂H ($t = 0$); (e–i) monitoring over time of solution in trace d after addition of ACA 12-CO₂H. CDCl₃, 25 °C. Dashed vertical lines are guides for the eye. On each spectrum, the relevant signals for **1** are marked in pink, those of **3** in blue, while those of **5** are coloured in green.

3 and **5** and pK_{bind} of **1·3** and **1·5** confirms such reading (see Tables S21 and S22). However, another corroborating experiment that further shows the fidelity of an ACA driven DL based on the B←N bond was carried out in the presence of excess acid 12-CO₂H.

Compounds **1**, **3** and **5**, 10.0 mM each, were added in CDCl₃ at 25 °C. The thermodynamic equilibrium with adducts **1·5** prevailing on **1·3** was immediately reached as expected (see state D and trace d in Fig. 7 where the ¹H NMR spectra of free **1**, **1·3** and **1·5** are also reported as traces a, b and c, respectively, for the sake of comparison). Addition of 30.0 mM 12-CO₂H causes protonation of both pyridines **3** and **5**, which are liberated in solution affording free **1** (see trace e in Fig. 7 recorded at $t = 9$ min from addition of the ACA). The new state E, which can be defined as a dissipative state, persists as long as the excess acid is present (traces e and f, recorded at time $t = 9$ min and 7 h, are both consistent with the presence of excess acid). When the excess is over (from state F onward), the less basic **3** is first deprotonated forming adduct **1·3**, as shown by trace g ($t = 36$ h), which roughly corresponds to state G of Fig. 7. At this point the strongest base **5** is still mostly protonated and cannot effectively engage in bonding with **1**. Then, ACA 12-CO₂H is further consumed and free base **5** starts to be available for the formation of **1·5** at the expense of **1·3**. Eventually, when ACA 12-CO₂H is exhausted, the system goes back to the initial equilibrium state with **1·5** again prevailing on **1·3** (see trace i in Fig. 7, $t = 5$ days, which corresponds again to state D). A corollary of this experiment is that ACA 12-CO₂H can also be used to temporally drive the simple binding equilibrium between ester **1** and bases **2–6**. And in fact, Fig. S40 shows that addition of ACA 12-CO₂H to complex **1·5** causes the liberation of **1** in solution due to the transient protonation of **5**. When the decarboxylation

of 12-CO₂H is over, adduct **1·5** is reversibly and completely restored. The process turned out to be fully reversible as demonstrated by an experiment in which three subsequent cycles were triggered by three successive additions of 12-CO₂H. At the end of each cycle (see Fig. S44), adduct **1·5** was found completely reassembled, also proving that bromoform is not detrimental to the B←N bond chemistry to any extent. Thus, similarly to what was observed in the case of the transimination reaction,^{27,29,89} the use of ACA also allows the B←N bond exchange to drive dissipative²⁷ dynamic libraries (DDLs) over time in a predictable fashion.²⁸

Conclusions

In this report we show that the B←N bond between phenylboronic acid catechol ester and a series of *N*-based aromatic heterocycles is fully reversible and can be conveniently used to obtain dynamic libraries of interconverting compounds. In chloroform, such B←N exchange reaction turns out to be rapid on the ¹H NMR time scale and its thermodynamic fate easily predictable from the formation equilibrium constants (K_{bind}) of the exchanging adducts. Moreover, it is shown that the exchange reactions among B←N adducts as well as their formation can be finely controlled over time in a dissipative fashion using activated carboxylic acids (ACAs). It is expected that such reversible interaction will be exploited in the near future for a number of applications such as (i) the generation of dynamic libraries composed of more complex chemical structures whose composition can be controlled in a dissipative fashion, (ii) the design of boron-based receptors for NArHet anchoring groups with the chance to control the related binding process over time, and (iii) the achievement of stimuli



responsive materials and dynamic polymer networks based on the exchange of the *N*-donor species.

Author contributions

Conceptualization: F. F, G. C, and S. D. S. Data curation: F. F, G. C, M. D. A, and C. M. Formal analysis: G. C, F. F, and C. M. Funding acquisition: S. D. S and C. M. Investigation: all authors. Methodology: F. F, G. C, O. L, and M. D. A. Supervision: F. F, C. M, and S. D. S. Writing: F. F, G. C, M. D. A, C. M, and S. D. S.

Conflicts of interest

There are no conflicts to declare.

Data availability

CCDC 2413767, 2413768 and 2492086 contain the supplementary crystallographic data for this paper.^{90a-c}

The data supporting this article have been included as part of the supplementary information (SI). Supplementary information: details on the synthesis of **1**, titration of **1** with *NAR*Hets **2–6**, crystallographic data, NMR spectra, ORCA input and output files, and speciation of the DDLs obtained. See DOI: <https://doi.org/10.1039/d5sc07665j>.

Acknowledgements

We are grateful for the support by Ateneo 2022 Sapienza (RG1221815C85AF91). Furthermore, we thank the European Union-NextGenerationEU under the Italian Ministry of University and Research (MUR), for the PRIN project “Chemically-Driven Autonomous Molecular Machines and Other Dissipative Systems” (No. 2022X779KE). CM acknowledges the support from the COMP-HUB and COMP-R Initiatives, funded by the ‘Departments of Excellence’ program of the Italian Ministry for University and Research (MIUR, 2018–2022 and MUR, 2023–2027).

Notes and references

‡ The log K_{bind} values also nicely correlate with Mayr’s nucleophilicity parameter; the points related to the log K_{obs} of aliphatic amines behave differently (see Fig. S61 and Table S29).

- S. J. Rowan, S. J. Cantrill, G. R. L. Cousins, J. K. M. Sanders and J. F. Stoddart, *Angew. Chem., Int. Ed.*, 2002, **41**, 898–952.
- J. Reek and S. Otto, *Dynamic Combinatorial Chemistry*, Wiley-VCH Verlag GmbH & Co, Weinheim, 2010.
- J. Li, P. Nowak and S. Otto, *J. Am. Chem. Soc.*, 2013, **135**, 9222–9239.
- A. Hermann, *Chem. Soc. Rev.*, 2014, **43**, 1899–1933.
- Y. Jin, Q. Wang, P. Taynton and W. Zhang, *Acc. Chem. Res.*, 2014, **47**, 1575–1586.
- M. Mondal and A. K. H. Hirsch, *Chem. Soc. Rev.*, 2015, **44**, 2455–2488.
- M. He and J.-M. Lehn, *J. Am. Chem. Soc.*, 2019, **141**, 18560–18569.
- M. Ciaccia and S. Di Stefano, *Org. Biomol. Chem.*, 2015, **13**, 646–654.
- P. T. Corbett, S. Otto and J. K. M. Sanders, *Chem.–Eur. J.*, 2004, **10**, 3139–3143.
- M. Rauschenberg, S. Bomke, U. Karst and B. J. Ravoo, *Angew. Chem., Int. Ed.*, 2010, **49**, 7340–7345.
- P. Nowak, M. Colomb-Delsuc, S. Otto and J. Li, *J. Am. Chem. Soc.*, 2015, **137**, 10965–10969.
- A. Canal-Martín and R. Pérez-Fernández, *Nat. Commun.*, 2021, **12**, 163.
- B. Brisig, J. K. M. Sanders and S. Otto, *Angew. Chem., Int. Ed.*, 2003, **42**, 1270–1273.
- J. W. Sadownik and D. Philp, *Angew. Chem., Int. Ed.*, 2008, **47**, 9965–9970.
- G. Gasparini, M. Dal Molin and L. J. Prins, *Eur. J. Org. Chem.*, 2010, 2429–2440.
- P.-A. R. Breuil and J. N. H. Reek in *Dynamic Combinatorial Chemistry*, ed. J. Reek and S. Otto, Wiley-VCH Verlag GmbH & Co, Weinheim, 2010, p. 91.
- F. Schaufelberger and O. Ramström, *J. Am. Chem. Soc.*, 2016, **138**, 7836–7839.
- M. Tena-Solsona, C. Wanzke, B. Riess, A. R. Bausch and J. Boekhoven, *Nat. Commun.*, 2018, **9**, 2044.
- C. M. E. Kriebisch, A. M. Bergmann and J. Boekhoven, *J. Am. Chem. Soc.*, 2021, **143**, 7719–7725.
- P. S. Schwarz, L. Tebcharani, J. E. Heger, P. Müller-Buschbaum and J. Boekhoven, *Chem. Sci.*, 2021, **12**, 9969–9976.
- M. A. Wurbs, P. S. Schwarz, J. Heckel, A. M. Bergmann, A. Walther and J. Boekhoven, *ChemSystemsChem*, 2021, **3**, e2100015.
- A. H. J. Engwerda, J. Southworth, M. A. Lebedeva, R. J. H. Scanes, P. Kukura and S. P. Fletcher, *Angew. Chem., Int. Ed.*, 2020, **59**, 20361–20366.
- M. G. Howlett, R. J. H. Scanes and S. P. Fletcher, *JACS Au*, 2021, **1**, 1355–1361.
- S. Yang, G. Schaeffer, E. Mattia, O. Markovitch, K. Liu, A. S. Hussain, J. Ottelé, A. Sood and S. Otto, *Angew. Chem., Int. Ed.*, 2021, **60**, 11344–11349.
- B. Liu, M. A. Beatty, C. G. Pappas, K. Liu, J. Ottelé and S. Otto, *Angew. Chem., Int. Ed.*, 2021, **60**, 13569–13573.
- P. S. Schwarz, M. Tena-Solsona, K. Dai and J. Boekhoven, *Chem. Commun.*, 2022, **58**, 1284–1297.
- D. Del Giudice, M. Valentini, G. Melchiorre, E. Spatola and S. Di Stefano, *Chem.–Eur. J.*, 2022, **28**, e202200685.
- D. Del Giudice, E. Spatola, M. Valentini, G. Ercolani and S. Di Stefano, *ChemSystemsChem*, 2022, **4**, e202200023.
- M. Valentini, G. Ercolani and S. Di Stefano, *Chem. Eur. J.*, 2024, **30**, e202401104.
- J. M. A. Carnall, C. A. Waudby, A. Belenguer, M. C. A. Stuart, J. J.-P. Peyralans and S. Otto, *Science*, 2010, **327**, 1502.
- J. Li, P. Nowak and S. Otto, *J. Am. Chem. Soc.*, 2013, **135**, 9222–9239.
- G. Ashkenasy, T. M. Hermans, S. Otto and A. F. Taylor, *Chem. Soc. Rev.*, 2017, **46**, 2543–2554.



- 33 A. G. Orrillo, A. La-Venia, A. M. Escalante and R. L. E. Furlan, *Chem.–Eur. J.*, 2018, **24**, 3141–3146.
- 34 D. Larsen and S. R. Beeren, *Chem. Sci.*, 2019, **10**, 9981–9987.
- 35 D. Carbajo, Y. Pérez, J. Bujons and I. Alfonso, *Angew. Chem., Int. Ed.*, 2020, **59**, 17202–17206.
- 36 S. Otto, *Acc. Chem. Res.*, 2022, **55**, 145–155.
- 37 P. M. Mitrasinovic, *Curr. Org. Synth.*, 2012, **9**, 233–246.
- 38 Z.-H. Zhao, C.-H. Li and J.-L. Zuo, *SmartMat*, 2023, **4**, e1187.
- 39 C. He, J. Dong, C. Xu and X. Pan, *ACS Polym. Au*, 2023, **3**, 5–27.
- 40 B. Chen and F. Jäkle, *Angew Chem. Int. Ed. Engl.*, 2024, **63**, e202313379.
- 41 N. Farfán, H. Höpfl, V. Barba, M. Eugenia Ochoa, R. Santillan, E. Gómez and A. Gutiérrez, *J. Organomet. Chem.*, 1999, **581**, 70–81.
- 42 N. Christinat, R. Scopelliti and K. Severin, *Chem. Commun.*, 2004, **10**, 1158–1159.
- 43 N. Christinat, R. Scopelliti and K. Severin, *J. Org. Chem.*, 2007, **72**, 2192–2200.
- 44 D. Salazar-Mendoza, J. Cruz-Huerta, H. Höpfl, I. F. Hernández-Ahuactzi and M. Sanchez, *Cryst. Growth Des.*, 2013, **13**, 2441–2454.
- 45 C. J. Hartwick, S. P. Yelgaonkar, E. W. Reinheimer, G. Campillo-Alvarado and L. R. MacGillivray, *Cryst. Growth Des.*, 2021, **21**, 4482–4487.
- 46 B. Icli, E. Sheepwash, T. Riis-Johannessen, K. Schenk, Y. Filinchuk, R. Scopelliti and K. Severin, *Chem. Sci.*, 2011, **2**, 1719–1721.
- 47 A. Dhara and F. Beuerle, *Chem.–Eur. J.*, 2015, **21**, 17391–17396.
- 48 N. Christinat, R. Scopelliti and K. Severin, *Chem. Commun.*, 2008, **31**, 3660–3662.
- 49 X. Xiao, D. Xiao, G. Sheng, T. Shan, J. Wang, X. Miao, Y. Liu, G. Li, Y. Zhu, J. L. Sessler and F. Huang, *Sci. Adv.*, 2023, **9**, eadi1169.
- 50 G. Campillo-Alvarado, E. C. Vargas-Olvera, H. Höpfl, A. D. Herrera-España, O. Sánchez-Guadarrama, H. Morales-Rojas, L. R. MacGillivray, B. Rodríguez-Molina and N. Farfán, *Cryst. Growth Des.*, 2018, **18**, 2726–2743.
- 51 G. Campillo-Alvarado, K. P. D'mello, D. C. Swenson, S. V. Santhana Mariappan, H. Höpfl, H. Morales-Rojas and L. R. MacGillivray, *Angew Chem. Int. Ed. Engl.*, 2019, **58**, 5413–5416.
- 52 G. Campillo-Alvarado, C. Li, Z. Feng, K. M. Hutchins, D. C. Swenson, H. Höpfl, H. Morales-Rojas and L. R. MacGillivray, *Organometallics*, 2020, **39**, 2197–2201.
- 53 A. D. Herrera-España, H. Höpfl and H. Morales-Rojas, *ChemPlusChem*, 2020, **85**, 548–560.
- 54 K. K. Ray, G. Campillo-Alvarado, H. Morales-Rojas, H. Höpfl, L. R. MacGillivray and A. V. Tivanski, *Cryst. Growth Des.*, 2020, **20**, 3–8.
- 55 J. D. Loya, S. A. Agarwal, N. Lutz, E. W. Reinheimer and G. Campillo-Alvarado, *CrystEngComm*, 2024, **26**, 4017–4021.
- 56 E. Sheepwash, V. Krampl, R. Scopelliti, O. Sereda, A. Neels and K. Severin, *Angew. Chem., Int. Ed.*, 2011, **50**, 3034–3037.
- 57 J. Cruz-Huerta, D. Salazar-Mendoza, J. Hernández-Paredes, I. F. Hernández Ahuactzi and H. Höpfl, *Chem. Commun.*, 2012, **48**, 4241–4243.
- 58 A. J. Stephens, R. Scopelliti, F. F. Tirani, E. Solari and K. Severin, *ACS Mater. Lett.*, 2019, **1**, 3–7.
- 59 F. Li, X. Xiao and Q. Xu, *Chem*, 2023, **9**, 13–15.
- 60 H. Zhang, Y. Li, L. Chen, Y. Yang, H. Lin, S. Xiang, B. Chen and Z. Zhang, *Chem*, 2023, **9**, 242–252.
- 61 C.-H. Liu, L. Chen, H. Zhang, Y. Li, H. Lin, L. Li, J. Wu, C. Liu, Z.-M. Ye, S. Xiang, B. Chen and Z. Zhang, *Chem*, 2023, **9**, 3532–3543.
- 62 N. Christinat, E. Croisier, R. Scopelliti, M. Cascella, U. Röthlisberger and K. Severin, *Eur. J. Inorg. Chem.*, 2007, 5177–5181.
- 63 E. Sheepwash, N. Luisier, M. R. Krause, S. Noé, S. Kubik and K. Severin, *Chem. Commun.*, 2012, **48**, 7808–7810.
- 64 N. Luisier, R. Scopelliti and K. Severin, *Soft Matter*, 2016, **12**, 588–593.
- 65 J. Cruz-Huerta, G. Campillo-Alvarado, H. Höpfl, P. Rodríguez-Cuamatzi, V. Reyes-Márquez, J. Guerrero-Álvarez, D. Salazar-Mendoza and N. Farfán-García, *Eur. J. Inorg. Chem.*, 2016, 355–365.
- 66 F. Vidal, J. Gomezcoello, R. A. Lalancette and F. Jäkle, *J. Am. Chem. Soc.*, 2019, **141**, 15963–15971.
- 67 X. Wang, L. Zhang, X. Wang, T. Cheng, M. Xue, Q. Dong, Y. Peng and Q. Zhang, *Adv. Funct. Mater.*, 2024, 202401362.
- 68 X. Sun, B. M. Chapin, P. Metola, B. Collins, B. Wang, T. D. James and E. V. Anslyn, *Nat. Chem.*, 2019, **11**, 768–778.
- 69 C. R. Groom, I. J. Bruno, M. P. Lightfoot and S. C. Ward, *Acta Crystallogr., Sect. B: Struct. Sci., Cryst. Eng. Mater.*, 2016, **72**, 171–179.
- 70 W. Clegg, A. J. Scott, F. E. S. Souza and T. B. Marder, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.*, 1999, **55**, 1885–1888.
- 71 F. Neese, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2012, **2**, 73–78.
- 72 F. Neese, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2025, **15**, e70019.
- 73 S. Grimme, A. Hansen, S. Ehlert and J.-M. Mewes, *J. Chem. Phys.*, 2021, **154**, 064103.
- 74 A. V. Marenich, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. B*, 2009, **113**, 6378–6396.
- 75 P. Erdmann, L. M. Sigmund, M. Schmitt, T. Hähnel, L. B. Dittmer and L. Greb, *ChemPhysChem*, 2024, **25**, e202400761.
- 76 R. Lo, D. Manna, M. Lamanec, M. Dračinský, P. Bouř, T. Wu, G. Bastien, J. Kaleta, V. M. Miriyala, V. Špirko, A. Mašínová, D. Nachtigallová and P. Hobza, *Nat. Commun.*, 2022, **13**, 2107.
- 77 J. A. Berrocal, C. Biagini, L. Mandolini and S. Di Stefano, *Angew. Chem., Int. Ed.*, 2016, **55**, 6997–7001.
- 78 C. Biagini, F. Di Pietri, L. Mandolini, O. Lanzalunga and S. Di Stefano, *Chem.–Eur. J.*, 2018, **24**, 10122–10127.
- 79 C. Biagini, G. Capocasa, V. Cataldi, D. Del Giudice, L. Mandolini and S. Di Stefano, *Chem.–Eur. J.*, 2019, **25**, 15205–15211.



- 80 C. Biagini and S. Di Stefano, *Angew. Chem., Int. Ed.*, 2020, **59**, 8344–8354.
- 81 D. Del Giudice, F. Tavani, M. Di Berto Mancini, F. Fratelloreto, M. Busato, D. Oliveira De Souza, F. Cenesi, O. Lanzalunga, S. Di Stefano and P. D'Angelo, *Chem.–Eur. J.*, 2022, **28**, e202103825.
- 82 F. Fratelloreto, F. Tavani, M. Di Berto Mancini, D. Del Giudice, G. Capocasa, I. Kieffer, O. Lanzalunga, S. Di Stefano and P. D'Angelo, *J. Phys. Chem. Lett.*, 2022, **13**, 5522–5529.
- 83 D. Del Giudice, F. Fratelloreto, C. Sappino and S. Di Stefano, *Eur. J. Org Chem.*, 2022, e202200407.
- 84 M. Valentini, F. Fratelloreto, M. Conti, R. Cacciapaglia, D. Del Giudice and S. Di Stefano, *Chem.–Eur. J.*, 2023, **29**, e202301835.
- 85 D. Del Giudice and S. Di Stefano, *Acc. Chem. Res.*, 2023, **56**, 889–899.
- 86 G. Capocasa, F. Fratelloreto, S. Correale Cavallari, M. Valentini, O. Lanzalunga and S. Di Stefano, *Chem.–Eur. J.*, 2024, **30**, e202303897.
- 87 F. Fratelloreto, G. Capocasa, A. D'Arrigo, M. De Angelis, O. Lanzalunga and S. Di Stefano, *ChemSystemsChem*, 2025, **7**, e202500008.
- 88 M. De Angelis, G. Capocasa, F. Ranieri, G. Mazzocanti, F. Fratelloreto, S. Manetto, A. Fagnano, C. Massera, G. Olivo, F. Ceccacci, A. Ciogli and S. Di Stefano, *Angew. Chem., Int. Ed.*, 2025, e202513917.
- 89 G. Melchiorre, L. Visieri, M. Valentini, R. Cacciapaglia, A. Casnati, L. Baldini, J. A. Berrocal and S. Di Stefano, *J. Am. Chem. Soc.*, 2025, **147**, 11327–11335.
- 90 (a) CCDC 2413767: Experimental Crystal Structure Determination, 2025, DOI: [10.5517/ccdc.csd.cc2m0qgv](https://doi.org/10.5517/ccdc.csd.cc2m0qgv); (b) CCDC 2413768: Experimental Crystal Structure Determination, 2025, DOI: [10.5517/ccdc.csd.cc2m0qhw](https://doi.org/10.5517/ccdc.csd.cc2m0qhw); (c) CCDC 2492086: Experimental Crystal Structure Determination, 2025, DOI: [10.5517/ccdc.csd.cc2pn6wg](https://doi.org/10.5517/ccdc.csd.cc2pn6wg).

