

Cite this: *Chem. Commun.*, 2014, 50, 5871

Received 1st December 2013,  
Accepted 14th April 2014

DOI: 10.1039/c3cc49143a

[www.rsc.org/chemcomm](http://www.rsc.org/chemcomm)


**A series of five-coordinate  $M^{II}$ -semiquinonate ( $M = Fe, Mn, Co$ ) complexes were synthesized and characterized, including the first example of a mononuclear  $Fe^{II}$ -semiquinonate. Intermediates were observed in the reactions of  $M^{II}$ -phenSQ ( $M = Fe, Co$ ) with  $O_2$ . Evidence for the relevance of these intermediates to the intradiol catechol dioxygenases was obtained by characterization of the oxidized semiquinone-derived product, muconic anhydride, resulting from the reaction of  $[PhTt^{tBu}]Co^{II}(3,5-DBSQ)$  with  $O_2$ .**

Intradiol catechol dioxygenases are non-heme iron enzymes that catalyze the oxidative cleavage of the C1–C2 bond of catechols.<sup>1</sup> The state of the enzyme that reacts with  $O_2$  contains a five-coordinate metal site. The activity of the enzyme and its synthetic analogs has been attributed to the partial  $Fe^{II}$ -semiquinonate (SQ) character within the  $Fe^{III}$ -catecholate species,<sup>2</sup> which results in the formation of a  $Fe^{III}$ -alkylperoxy intermediate upon addition of  $O_2$ . The extradiol catechol dioxygenases contain iron or manganese active sites and catalyze the oxidative cleavage of C2–C3 bond of catechols.<sup>1,3</sup> During catalytic turnover, superoxo- $Fe^{II}$ -semiquinonate and  $Fe^{II}$ -alkylperoxy intermediates have been detected.<sup>4</sup> Interestingly, comparable extradiol-cleaving activities were obtained by substituting the native iron with manganese or cobalt suggesting the intermediacy of superoxo- $M^{II}$ -semiquinonate species ( $M = Fe, Mn, Co$ ) in the catalytic cycles.<sup>5</sup> Moreover, reactions of redox-active ligand complexes with dioxygen have received recent attention due to their potential utility in stoichiometric and catalytic transformations.<sup>6</sup>

In spite of the implications of the  $Fe^{II}$ -semiquinonate species in both intradiol and extradiol dioxygenases, to the best of our knowledge, well characterized mononuclear  $Fe^{II}$ -semiquinonate complexes are unknown. Related complexes were reported recently

by Fiedler and co-workers, including a mononuclear  $Fe^{II}$ -(imino)-semiquinonate complex<sup>7</sup> and a semiquinonate-bridged diiron(II) complex.<sup>8</sup> Herein, we report the first well characterized mononuclear  $Fe^{II}$ -semiquinonate complex and its  $Mn^{II}$  and  $Co^{II}$  analogues –  $[PhTt^{tBu}]M(phenSQ)$  ( $M = Fe, Mn, Co$ ,  $PhTt^{tBu}$  = phenyltris-((*tert*-butylthio)methyl)borate, phenSQ = 9,10-phenanthrene-semiquinonate). The suitability of these complexes in modelling catalytic intermediates of the intradiol dioxygenases was evaluated by  $O_2$  reactivity studies. A related complex,  $[PhTt^{tBu}]Co(3,5-DBSQ)$  ( $3,5-DBSQ$  = 3,5-di-*tert*-butyl-1,2-semiquinonate) exhibited the intradiol reactivity, suggesting the relevance of the observed intermediates to the intradiol catechol dioxygenases.

$[PhTt^{tBu}]M(phenSQ)$  were synthesized using two complementary preparative routes, Scheme 1. Metathesis of  $[PhTt^{tBu}]MI$  ( $M = Fe, Mn, Co$ ) with  $Tl(phenSQ)$  yielded  $[PhTt^{tBu}]M(phenSQ)$  in excellent yields (89–95%). A similar method was applied to the synthesis of  $[PhTt^{tBu}]Co(3,5-DBSQ)$  by replacing  $Tl(phenSQ)$  with  $Tl(3,5-DBSQ)$ . Alternatively, the oxidative addition of phenQ to monovalent metal precursors,  $[PhTt^{tBu}]M(PMe_3)$  ( $M = Fe, Co$ )<sup>9</sup> afforded  $[PhTt^{tBu}]M(phenSQ)$  ( $M = Fe, Co$ ) in good yields (65–70%). High resolution mass spectroscopy (HRMS) data combined with  $^1H$  NMR spectral analyses confirmed the composition and purity of the  $[PhTt^{tBu}]M^{II}-(SQ)$  complexes (Fig. S4–S7, S11–S14, ESI<sup>†</sup>).



**Scheme 1** Synthetic routes to  $[PhTt^{tBu}]M(SQ)$  complexes ( $M = Fe, Mn, Co$ ) and  $O_2$  reactivity of  $[PhTt^{tBu}]Co(3,5-DBSQ)$ .

Department of Chemistry & Biochemistry, University of Delaware, Newark, DE 19716, USA. E-mail: [riordan@udel.edu](mailto:riordan@udel.edu); Tel: +1 302 831 1073

<sup>†</sup> Electronic supplementary information (ESI) available: Experimental details, new compound characterization data and crystallographic data, etc. CCDC 969713 ( $[PhTt^{tBu}]Fe(phenSQ)$ ), 969714 ( $[PhTt^{tBu}]Co(phenSQ)$ ), 969715 ( $[PhTt^{tBu}]CoI$ ), 969716 ( $[PhTt^{tBu}]MnI$ ), 969717 ( $[PhTt^{tBu}]Mn(phenSQ)$ ) and 992704 ( $[PhTt^{tBu}]Co(3,5-DBSQ)$ ). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c3cc49143a







Fig. 1 Electronic spectra of the  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{SQ})$  complexes. The insert highlights ligand field transitions.

The electronic spectra of the  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{SQ})$  complexes are contained in Fig. 1.  $[\text{PhTt}^{t\text{Bu}}]\text{Fe}(\text{phenSQ})$  shows two features of low intensity at 600 nm ( $\epsilon = 868 \text{ M}^{-1} \text{ cm}^{-1}$ ) and 935 nm ( $\epsilon = 539 \text{ M}^{-1} \text{ cm}^{-1}$ ), the latter being consistent with a typical ligand field transition for five-coordinate, high-spin  $\text{Fe}^{\text{II}}$  complexes.<sup>10</sup> No apparent ligand field transition was observed for  $[\text{PhTt}^{t\text{Bu}}]\text{Mn}(\text{phenSQ})$ , indicating a high-spin  $\text{Mn}^{\text{II}}$  center.  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(\text{phenSQ})$  exhibits two features at 683 ( $\epsilon = 1310 \text{ M}^{-1} \text{ cm}^{-1}$ ) and 803 ( $\epsilon = 1290 \text{ M}^{-1} \text{ cm}^{-1}$ ), both of which agree well with the ligand field transitions of five-coordinate, high-spin  $\text{Co}^{\text{II}}$ .<sup>11</sup> Unlike the previously reported  $[\text{Tp}^{\text{Cum},\text{Me}}]\text{Co}(3,5\text{-DBSQ})$ ,<sup>12</sup> the ligand field transitions of  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$  were not observed in the spectrum presumably due to the overlap with the intense band at 784 nm ( $\epsilon = 5470 \text{ M}^{-1} \text{ cm}^{-1}$ ).<sup>13</sup>

The description of the metal complex electronic structures as  $\text{M}^{\text{II}}\text{-SQ}$  is supported by their infrared spectra. In contrast to the IR spectra of the respective  $[\text{PhTt}^{t\text{Bu}}]\text{M}$  complexes,  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{phenSQ})$  exhibit intense bands between  $1500 \text{ cm}^{-1}$  and  $1600 \text{ cm}^{-1}$ , which are consistent with the  $\text{C}=\text{O}$  stretches of phenSQ (Fig. S15–S17, ESI†).<sup>14</sup> Although  $[\text{PhTt}^{t\text{Bu}}]\text{CoI}$  also shows two bands in the  $1400\text{--}1450 \text{ cm}^{-1}$  range, the much more intense band at  $1462 \text{ cm}^{-1}$  of  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$  is tentatively assigned to the  $\nu(\text{C}=\text{O})$  of 3,5-DBSQ (Fig. S18, ESI†).<sup>15</sup>

All the complexes are five-coordinate as deduced by X-ray diffraction analyses (Fig. S20, ESI†). The coordination geometry of the  $[\text{PhTt}^{t\text{Bu}}]\text{Mn}(\text{phenSQ})$  lies between trigonal-bipyramidal and square pyramidal ( $\tau_5 = 0.58$ ),<sup>16</sup> whereas the geometries of  $[\text{PhTt}^{t\text{Bu}}]\text{Fe}(\text{phenSQ})$  ( $\tau_5 = 0.14$ ),  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(\text{phenSQ})$  ( $\tau_5 = 0.01$ ) and  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$  ( $\tau_5 = 0.17$ ) are best described as distorted square pyramids. Key bond lengths support the redox state assignment of the bidentate ligand as semiquinonate, Table 1 and Fig. S20 (ESI†). For the  $\text{M}^{\text{II}}$ -phenSQ complexes, the C–O distances are in the range of  $1.28\text{--}1.30 \text{ \AA}$  and C–C distances are in the range of  $1.41\text{--}1.44 \text{ \AA}$ . These bond distances are characteristic of a bound phenSQ ligand.<sup>14a,17</sup> For  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$ , the average C–O distance is  $1.314(2) \text{ \AA}$ . This distance is certainly among the longest C–O distances for 3,5-DBSQ, but is not unprecedented.<sup>18</sup> Furthermore, the “four long/two short” quinoid distortion in the semiquinonate ring further supports its electronic structure

Table 1 Molecular structure of  $[\text{PhTt}^{t\text{Bu}}]\text{Fe}(\text{phenSQ})$  and selected metric parameters of the  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{phenSQ})$  complexes. Hydrogen atoms are omitted for clarity. (Structures of the other new complexes are available in the ESI)



	Fe(phenSQ)	Mn(phenSQ)	Co(phenSQ)
M1–O1	2.064(2)	2.109(2)	1.922(1)
M1–O2	2.015(2)	2.080(2)	1.912(1)
O1–C22	1.285(2)	1.282(4)	1.301(2)
O2–C35	1.285(2)	1.280(4)	1.297(2)
C35–C22	1.435(3)	1.436(5)	1.411(2)
$\tau$ -value	0.14	0.58	0.01

description. All complexes are air-sensitive in solution as demonstrated by  $\text{O}_2$  reactivity studies, *vide infra*, which can be rationalized by the  $\text{M}^{\text{II}}\text{-SQ}$  charge distribution.<sup>2a–c</sup> Space filling models (Fig. S21, ESI†) indicate  $\text{O}_2$  accessibility to both the metal centers and the semiquinonate ligands.

The  $\text{M}^{\text{II}}\text{-SQ}$  complexes showed paramagnetic  $^1\text{H}$  NMR spectra. Their effective magnetic moments measured in solution by the Evans method are  $[\text{PhTt}^{t\text{Bu}}]\text{Mn}(\text{phenSQ})$  and  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$   $\mu_{\text{eff}} = 5.01(6) \mu_{\text{B}}$  and  $2.91(2) \mu_{\text{B}}$ , respectively. These values are very close to the spin-only values for  $S = 2$  and  $S = 1$  systems, indicating strong antiferromagnetic coupling between high-spin divalent metal centers and the SQ radicals.  $[\text{PhTt}^{t\text{Bu}}]\text{Fe}(\text{phenSQ})$  and  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(\text{phenSQ})$  display  $\mu_{\text{eff}} = 4.65(2) \mu_{\text{B}}$  and  $3.43(3) \mu_{\text{B}}$ , respectively. These values are higher than the spin-only values for  $S = 3/2$  and  $S = 1$  systems, but lower than expected for non-spin coupled systems,<sup>12</sup> suggesting either weaker antiferromagnetic coupling or more likely, non-negligible spin–orbit coupling.

The cyclic voltammograms (CV) were measured in THF to evaluate the redox characteristics of the  $\text{M}^{\text{II}}\text{-SQ}$  complexes (Fig. S22, ESI†).  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(\text{phenSQ})$  and  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$  exhibit reversible reduction events at  $-0.97 \text{ V}$  and  $-0.82 \text{ V}$  (*vs.*  $\text{Fc}^{+/\text{0}}$ ) which are assigned as ligand-centered reductions, phenSQ/phenCat and 3,5-DBSQ/DBCat, respectively. These redox potentials match very well with the values reported for  $\text{Me}_4\text{cyclam}$  supported  $\text{Co}^{\text{II}}$ -semiquinonate complexes ( $-1.00 \text{ V}$  for phenSQ/phenCat and  $-0.85 \text{ V}$  for 3,5-DBSQ/DBCat).<sup>19</sup> The redox events for  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{phenSQ})$  ( $\text{M} = \text{Mn, Fe}$ ) are irreversible on the electrochemical timescale, exhibiting  $E_{\text{c}}$  values of  $-1.17 \text{ V}$  for  $[\text{PhTt}^{t\text{Bu}}]\text{Fe}(\text{phenSQ})$  and  $-1.11 \text{ V}$  for  $[\text{PhTt}^{t\text{Bu}}]\text{Mn}(\text{phenSQ})$ , assigned as phenSQ/phenCat reductions. The trend in reduction potentials among the  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{phenSQ})$  species,  $\text{Fe} < \text{Mn} < \text{Co}$ , indicates the Co complex is most readily reduced. The oxidations of  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(\text{phenSQ})$  and  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$  at  $0.26 \text{ V}$  and  $0.47 \text{ V}$ , respectively are also irreversible.

To evaluate the utility of five-coordinate  $\text{M}^{\text{II}}\text{-SQ}$  complexes to model putative intermediates in catechol dioxygenase catalysis,






Fig. 2 Electronic spectral changes during the oxygenation of  $[\text{PhTt}^{t\text{Bu}}]\text{Fe}(\text{phenSQ})$  at  $-90\text{ }^\circ\text{C}$  in toluene. (a) Spectral changes after exposing  $[\text{PhTt}^{t\text{Bu}}]\text{Fe}(\text{phenSQ})$  to  $\text{O}_2$ . Intermediate growth indicated by red arrow. Spectra were recorded at 2 min intervals. (b) Spectra of intermediate and product collected at  $-90\text{ }^\circ\text{C}$ . The spectrum of the product was obtained by warming the sample to  $0\text{ }^\circ\text{C}$  for 5 min to allow the decay of the intermediate and cooling back to  $-90\text{ }^\circ\text{C}$  for 15 min.

electronic spectroscopy was employed to monitor the reaction of  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{phenSQ})$  with  $\text{O}_2$ .<sup>20</sup> Even at very low temperature ( $-90\text{ }^\circ\text{C}$ ) rapid spectroscopic changes were observed upon addition of  $\text{O}_2$  to  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{phenSQ})$  ( $\text{M} = \text{Fe, Co}$ ), producing intermediates which decay to the thermodynamic products in 5–10 minutes upon warming to higher temperatures (Fig. 2 and Fig. S28, ESI†).<sup>21</sup> Indirect evidence for the relevance of these synthetic intermediates to the intradiol dioxygenases was obtained using  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$ . Upon reaction with  $\text{O}_2$  in THF for 16 hours,  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$  produced the intradiol cleavage product, muconic anhydride in 16% yield, Scheme 1 (see ESI† for details). This result, together with a previous discovery that the five coordinate complex  $[\text{Tp}^{i\text{Pr}_2}]\text{Mn}(3,5\text{-DBSQ})$  also produces the intradiol product,<sup>22</sup> suggests that the semiquinonate character of the ligand may contribute to the intradiol cleaving reactivity, even for different metals. To the best of our knowledge, this is the first example of intradiol reactivity of a  $\text{Co}^{II}(3,5\text{-DBSQ})$  complex.<sup>23</sup>

In summary, a series of five-coordinate  $\text{M}^{II}$ -semiquinonate ( $\text{M} = \text{Fe, Mn, Co}$ ) complexes supported by the tris(thioether) ligand  $[\text{PhTt}^{t\text{Bu}}]$  were synthesized and characterized, including the first example of a mononuclear  $\text{Fe}^{II}$ -semiquinonate complex. While  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$  was found to be a reactivity model for the intradiol catechol dioxygenases,  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{phenSQ})$  ( $\text{M} = \text{Fe, Co}$ ) serve as potential precursors to model the putative intermediates in intradiol dioxygenase catalysis. Our current efforts are focused on additional spectroscopic and structural characterization of the intermediates produced from the low temperature reactions of  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{phenSQ})$  with  $\text{O}_2$ . Also under investigation are the reactions of  $[\text{PhTt}^{t\text{Bu}}]\text{M}(\text{phenSQ})$  with superoxide to model intermediate(s) of relevance in extradiol dioxygenase catalysis.

The US National Science Foundation supported this research program *via* CHE-1112035 to CGR. The X-ray diffractometer (CHE-1048367) and LIFDI mass spectrometer (CHE-1229234) acquisitions were supported by NSF.

## Notes and references

- (a) T. D. H. Bugg and S. Ramaswamy, *Curr. Opin. Chem. Biol.*, 2008, **12**, 134–140; (b) M. Costas, M. P. Mehn, M. P. Jensen and L. Que, Jr., *Chem. Rev.*, 2004, **104**, 939–986.
- For experimental work, see: (a) D. D. Cox and L. Que, Jr., *J. Am. Chem. Soc.*, 1988, **110**, 8085–8092; (b) H. G. Jang, D. D. Cox and L. Que, Jr., *J. Am. Chem. Soc.*, 1991, **113**, 9200–9204; (c) T. Funabiki, A. Fukui, Y. Hitomi, M. Higuchia, T. Yamamoto, T. Tanaka, F. Tani and Y. Naruta, *J. Inorg. Biochem.*, 2002, **91**, 151–158. For theoretical work, see: (d) T. Funabiki and T. Yamazaki, *J. Mol. Catal. A: Chem.*, 1999, **150**, 37–47; (e) N. Nakatani, Y. Hitomi and S. Sakaki, *J. Phys. Chem. B*, 2011, **115**, 4781–4789.
- J. D. Lipscomb, *Curr. Opin. Struct. Biol.*, 2008, **18**, 644–649.
- E. G. Kovaleva and J. D. Lipscomb, *Science*, 2007, **316**, 453–457.
- (a) J. P. Emerson, E. G. Kovaleva, E. R. Farquhar, J. D. Lipscomb and L. Que, Jr., *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 7347–7352; (b) A. J. Fielding, E. G. Kovaleva, E. R. Farquhar, J. D. Lipscomb and L. Que, Jr., *JBIC, J. Biol. Inorg. Chem.*, 2011, **16**, 341–355.
- (a) S. Mukherjee, T. Weyhermuller, E. Bothe, K. Wieghardt and P. Chaudhuri, *Dalton Trans.*, 2004, 3842–3853; (b) C. A. Lippert, S. A. Arnstein, C. D. Sherrill and J. D. Soper, *J. Am. Chem. Soc.*, 2010, **132**, 3879–3892; (c) C. J. Rolle, III, K. I. Hardcastle and J. D. Soper, *Inorg. Chem.*, 2008, **47**, 1892–1894.
- M. M. Bittner, S. V. Lindeman and A. T. Fiedler, *J. Am. Chem. Soc.*, 2012, **134**, 5460–5463.
- A. E. Baum, S. V. Lindeman and A. T. Fiedler, *Chem. Commun.*, 2013, **49**, 6531–6533.
- (a) M. T. Mock, C. V. Popescu, G. P. A. Yap, W. G. Dougherty and C. G. Riordan, *Inorg. Chem.*, 2008, **47**, 1889–1891; (b) J. A. Dupont, G. P. A. Yap and C. G. Riordan, *Inorg. Chem.*, 2008, **47**, 10700–10707.
- E. I. Solomon, T. C. Brunold, M. I. Davis, J. N. Kemsley, S. K. Lee, N. Lehnert, F. Neese, A. J. Skulan, Y. S. Yang and J. Zhou, *Chem. Rev.*, 2000, **100**, 235–349.
- (a) M. Ciampolini, N. Nardi and G. P. Speroni, *Coord. Chem. Rev.*, 1966, **1**, 222–233; (b) Z. Dori and H. B. Gray, *Inorg. Chem.*, 1968, **7**, 889–892.
- M. Ruf, B. C. Noll, M. D. Groner, G. T. Yee and C. G. Pierpont, *Inorg. Chem.*, 1997, **36**, 4860–4865.
- Analogous to  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(3,5\text{-DBSQ})$ , an intense band at 818 nm was observed for  $[\text{PhTt}^{t\text{Bu}}]\text{Co}(\text{phenSQ})$  at  $-90\text{ }^\circ\text{C}$ .
- (a) A. Caneschi, A. Dei, F. Fabrizi de Biani, P. Gutlich, V. Ksenofontov, G. Levchenko, A. Hoefer and F. Renz, *Chem. – Eur. J.*, 2001, **7**, 3926–3930; (b) S. Roy, B. Sarkar, D. Bubrin, M. Niemeyer, S. Zalis, G. K. Lahiri and W. Kaim, *J. Am. Chem. Soc.*, 2008, **130**, 15230–15231.
- (a) F. Hartl and A. Vlcek, Jr., *Inorg. Chem.*, 1996, **35**, 1257–1265; (b) P. A. Wicklund, L. S. Beckmann and D. G. Brown, *Inorg. Chem.*, 1976, **15**, 1996–1997.
- A. W. Addison, T. N. Rao, J. Reedijk, J. Vanrijn and G. C. Verschoor, *J. Chem. Soc., Dalton Trans.*, 1984, 1349–1356.
- C. G. Pierpont and R. M. Buchanan, *Coord. Chem. Rev.*, 1981, **38**, 45–87.
- M. Graf, G. Wolmershäuser, H. Kelm, S. Demeschko, F. Meyer and H.-J. Krüger, *Angew. Chem., Int. Ed.*, 2010, **49**, 950–953.
- A. Caneschi, A. Dei, D. Gatteschi and V. Tangoulis, *Inorg. Chem.*, 2002, **41**, 3508–3512.
- A (triphos)Ir( $\text{O}_2$ )(phenSQ) complex modeling the intradiol dioxygenase intermediate was formed from the reaction of  $[(\text{triphos})\text{Ir}(\text{phenCat})]\text{BPh}_4$  with  $\text{O}_2$ . See: P. Barbaro, C. Bianchini, C. Mealli and A. Meli, *J. Am. Chem. Soc.*, 1991, **113**, 3181–3183.
- No intermediate was observed upon reaction of  $[\text{PhTt}^{t\text{Bu}}]\text{Mn}(\text{phenSQ})$  with  $\text{O}_2$  at  $-90\text{ }^\circ\text{C}$ .
- H. Komatsuzaki, A. Shiota, S. Hazawa, M. Itoh, N. Miyamura, N. Miki, Y. Takano, J. Nakazawa, A. Inagaki, M. Akita and S. Hikichi, *Chem. – Asian J.*, 2013, **8**, 1115–1119.
- A tetramine- $\text{Co}^{II}(3,5\text{-DBSQ})$  complex reacted with  $\text{O}_2$ , leading to the ring cleavage of the catechol. However, the organic products were not unambiguously characterized as intradiol or extradiol products. See: S. Nakashima, H. Ohya-Nishiguchi, N. Hirota, S. Tsuboyama and T. Chijimatsu, *Bull. Chem. Soc. Jpn.*, 1992, **65**, 1225–1232.

