



Cite this: *Org. Biomol. Chem.*, 2019, **17**, 7874

Received 28th June 2019,  
Accepted 6th August 2019

DOI: 10.1039/c9ob01460h

rsc.li/obc

## Microcalorimetry and fluorescence show stable peptide nucleic acid (PNA) duplexes in high organic content solvent mixtures†

Samuel Núñez-Pertíñez,<sup>id</sup> Thomas R. Wilks<sup>id</sup>\* and Rachel K. O'Reilly<sup>id</sup>\*

The selectivity of nucleic acid hybridisation can be exploited to template chemical reactions, enabling materials discovery by chemical evolution. However, to date the range of reactions that can be used has been limited to those that are compatible with aqueous media, since the addition of organic co-solvents can have a large impact on the stability of nucleic acid duplexes. Peptide nucleic acids (PNAs) are promising in this regard because previous studies have suggested they may be stable as duplexes in high organic content solvent mixtures. Here, we use micro-differential scanning calorimetry (micro-DSC) to confirm for the first time that double-stranded PNA (dsPNA) is stable in *N,N*-dimethylformamide (DMF)/water mixtures up to 95 vol% DMF. Using fluorescence, we corroborate these results and show that the isothermal annealing of PNA in high DMF content solution is also rapid. These findings suggest that PNA could enable the use of a range of water-sensitive chemistries in nucleic acid templating applications.

Nucleic acids have the remarkable ability to recognise complementary strands. This selective hybridisation has been exploited to template the synthesis of oligomers<sup>1,2</sup> and for the production of combinatorial libraries<sup>3,4</sup> in an analogous manner to the genetically controlled biosynthesis of proteins.<sup>5</sup> Interestingly, this approach to organic synthesis has the potential to perform directed chemical evolution: whole libraries of DNA-tagged molecules can be screened in one pot, allowing the selection and identification of those that perform a particular function, such as catalysis or enzyme inhibition.<sup>6–10</sup>

One major limitation of this nucleic acid-templated synthesis (NATS) is the incompatibility of many chemical reactions with aqueous solution. Aqueous solution is required because nucleic acid duplexes are greatly destabilised by the addition of organic cosolvents.<sup>11</sup> Overcoming this limitation

would allow the use of a wider variety of chemical transformations, increasing the scope of NATS technology.

Peptide nucleic acids (PNAs) are DNA analogues in which the phosphate and sugar backbone is replaced by *N*-(2-aminoethyl)glycine linkages.<sup>12</sup> PNAs have been used as molecular tags to encode the identity of molecules in combinatorial chemical libraries,<sup>13</sup> and as templates in several examples of templated synthesis.<sup>14–16</sup> PNAs are capable of selective recognition through hybridisation and, in contrast to native DNA and RNA, are soluble in a range of organic solvents such as 1,4-dioxane, dimethylformamide (DMF), *N*-methylpyrrolidone (NMP) and dimethyl sulfoxide (DMSO).<sup>15</sup> Previous studies have also shown that double-stranded PNA (dsPNA) is stable in up to 70 vol% DMF/water mixtures.<sup>17</sup> They are therefore promising candidates to allow extension of NATS into high organic content mixtures, or even pure organic solvents.

The duplex melting temperature ( $T_m$ ) is the most commonly used proxy of the thermal stability of dsPNA, and is defined as the temperature at which half the possible duplexes have formed. There are three main techniques available to measure the  $T_m$  of nucleic acid duplexes (see Table 1): ultraviolet (UV) absorbance, fluorescence and calorimetry. The formation of dsPNA in hydrated organic solvent was studied by Sen and Nielsen through UV absorbance, and confirmed duplex stability up to 50<sup>18</sup> and 70 vol%<sup>17</sup> DMF. However, study of higher organic content mixtures was not possible because of DMF's interference with the UV absorbance measurements. This limitation also holds true for a wide range of organic solvents with high UV cut-offs. This problem can be avoided by

**Table 1** Main features of the three available techniques to determine the thermal stability of nucleic acid duplexes

	UV absorbance	Fluorescence	Calorimetry
Sensitivity	Moderate	<b>High</b>	Low
Chemical modification	<b>None required</b>	Required	<b>None required</b>
Solvent range	Narrow	<b>Wide</b>	<b>Wide</b>

School of Chemistry, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK. E-mail: T.R.Wilks@bham.ac.uk, R.O'Reilly@bham.ac.uk

† Electronic supplementary information (ESI) available: Experimental methods, HPLC and LC-MS characterisation, UV absorbance, micro-DSC and fluorescence data. See DOI: 10.1039/c9ob01460h







**Fig. 2** Evolution of the melting temperature ( $T_m$ ) of dsPNA as a function of the volumetric DMF content. 15-mer dsPNA micro-DSC measurements (open dashed red circle), 10-mer dsPNA micro-DSC measurements (solid blue dot), 10-mer dsPNA UV absorbance reported data (open blue circle) and 10-mer PNA-DNA hybrid duplex micro-DSC measurements (black triangle). No transition was observed in 100 vol% DMF solutions (data not shown). (Sequences: [10]-DNA/PNA1 5'/N – TCA CTA GAT G – C/3' [10]-DNA/PNA2 – 5'/N – CAT CTA GTG A – C/3' [15]-PNA1 N – CGC CGT CAC TAG ATG – C [15]-PNA2 N – CAT CTA GTG ACG GCG – C.) For further details and standard error evaluation see section 4 ESI†.

pure organic solvent. However, our results demonstrate that this is not the case. A representation of the  $T_m$  as a function of the molar percentage (mol%) of DMF rather than vol% illustrates why: because DMF and water have similar densities but different molecular weights, previous studies had only explored a relatively small range of DMF:water molar ratios, making extrapolation of the trend to high DMF content prone to inaccuracy (see Fig. S6 and S7 ESI†).

The stability of PNA-DNA hybrid duplexes was also studied by micro-DSC in different DMF/water mixtures. It was found that addition of DMF had a much greater impact on the stability of DNA-PNA duplexes than on dsPNA.


Our results led us to speculate that a longer PNA duplex might remain stable at even higher DMF vol%, because increasing the length of a nucleic acid duplex generally results in a greater  $T_m$ , at least for short sequences.<sup>22</sup> In pure aqueous solutions, extending the length of unmodified PNA is not an option because strong inter-strand interactions render it insoluble,<sup>23</sup> but we speculated that the addition of DMF would allow longer PNA strands to dissolve and hybridise in a reversible and well-controlled manner. These hypotheses were tested with a 15-mer dsPNA (see Fig. 2). As expected, the addition of five extra nucleobases increased the  $T_m$  in 90 vol% DMF from  $33 \pm 1.1$  °C to  $67.7 \pm 0.3$  °C. Furthermore, while the 10-mer dsPNA duplex was not observed in solutions with a DMF content higher than 90 vol%, it was found that the 15-mer sequence formed stable dsPNA in up to 95 vol% DMF. A previous study has demonstrated that reducing the concentration of water to 5 vol% enables the use of water-sensitive transformations such as the pyrrolidine catalysed aldol reaction to afford nucleic acid-tem-

plated products in high yield (up to 88%), illustrating the impact of our findings on future NATS applications.<sup>24</sup>

Nucleic acid duplexes are stabilised by a combination of hydrophobic, hydrogen bonding,  $\pi$ - $\pi$  stacking, van der Waals and electrostatic interactions.<sup>25</sup> The reduction in the  $T_m$  that we observed at lower molar water content indicated that the hydrophobic effect accounted for a considerable fraction of this stabilisation for dsPNA. It was previously proposed by Sen and Nielsen that as DMF content increases, the major contributor to duplex stability shifts towards hydrogen bonding.<sup>17</sup> As DMF has the ability to accept hydrogen bonds, this suggests that alternative, non-hydrogen bonding solvents hold potential for further stabilisation of the duplex and could be identified in the future by micro-DSC.

Finally, to assess the potential of dsPNA for use in more complex multi-step templated syntheses,<sup>2</sup> we investigated the isothermal annealing of PNA duplexes in 90 vol% DMF and aqueous solution (Fig. 3).

One 10-mer PNA strand was labelled with a tryptophan group at the N' end and the fluorescence in solution recorded for several minutes to ensure a stable baseline. The fluorescence cuvette was then removed, the complementary strand labelled with a 5-(dimethylamino)naphthalene-sulfonamide (dansyl) group at the C' was added, and the cuvette shaken to



**Fig. 3** Isothermal annealing of a 10-mer dsPNA in 90 vol% DMF: (a) plot of the emission of the tryptophan-labelled PNA strand over time. The addition of the complementary quencher-labelled PNA strand is labelled with a red arrow. (b) Thermal denaturation of the isothermally annealed sample, and subsequent re-annealing upon cooling. (The donor moiety in [10]-PNA1-F was Tryptophan, the acceptor moiety in [10]-PNA2-Q was Dansyl. Tryptophan was excited at 290 nm and the emission was recorded at 340 nm. Dansyl emission was recorded at 550 nm.) For further details see section 5 ESI†.



ensure thorough mixing, before being replaced in the instrument and measurements continued.

In 90 vol% DMF, annealing was found to be rapid (Fig. 3a), with full quenching observed by the time the cuvette had been replaced in the instrument following addition of the complementary strand (a delay of around 12 s). It was also found that the freshly dissolved PNA strands required a heating/cooling cycle in the desired solvent prior to isothermal annealing in order to display these fast kinetics, otherwise annealing took approximately 1.5 h to complete (see Fig. S14 ESI†). This was interpreted as the effect of kinetically trapped states present in the dry pellet that provided an additional barrier to dsPNA formation. The isothermally annealed PNA was subsequently thermally denatured and was observed to re-anneal upon cooling, demonstrating the full reversibility of the process (Fig. 3b). This supported the micro-DSC evidence of dsPNA formation in 90 vol% DMF and suggests that multistep PNA templated synthesis in organic solvents may be possible in the future.

In conclusion, it has been demonstrated that microcalorimetry allows reliable determination of the thermal stability of dsPNA in solvent mixtures where the widely employed UV absorbance method cannot be used as a result of the high UV cut-off of the solvent. This technique was subsequently used to study the stability of 10-mer and 15-mer dsPNA duplexes in 0 to 100 vol% DMF showing the formation of dsPNA in up to 95 vol% DMF. A remarkable stability in high organic solvent content was observed, which was directly dependent on the length of the PNA. Finally, it was shown that PNAs isothermally anneal rapidly in high DMF content solutions. Altogether, these findings highlight the potential of microcalorimetry to study duplex formation events in a wider range of solvent media and in the absence of a measurable spectrometric change. They also provide a good starting point for further applications of peptide nucleic acids as chemical templates in water-free media.

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- M. L. McKee, P. J. Milnes, J. Bath, E. Stulz, A. J. Turberfield and R. K. O'Reilly, *Angew. Chem., Int. Ed.*, 2010, **49**, 7948–7951.
- Y. He and D. R. Liu, *J. Am. Chem. Soc.*, 2011, **133**, 9972–9975.
- M. H. Hansen, P. Blakskjaer, L. K. Petersen, T. H. Hansen, J. W. Hoøjfeldt, K. V. Gothelf and N. J. V. Hansen, *J. Am. Chem. Soc.*, 2009, **131**, 1322–1327.
- W. Meng, R. A. Muscat, M. L. McKee, P. J. Milnes, A. H. El-Sagheer, J. Bath, B. G. Davis, T. Brown, R. K. O'Reilly and A. J. Turberfield, *Nat. Chem.*, 2016, **8**, 542–548.
- X. Li and D. R. Liu, *Angew. Chem., Int. Ed.*, 2004, **43**, 4848–4870.
- C. J. Krusemark, N. P. Tilmans, P. O. Brown and P. B. Harbury, *PLoS One*, 2016, **11**, e0154765.
- Z. J. Gartner, B. N. Tse, R. Grubina, J. B. Doyon, T. M. Snyder and D. R. Liu, *Science*, 2004, **305**, 1601–1605.
- B. N. Tse, T. M. Snyder, Y. Shen and D. R. Liu, *J. Am. Chem. Soc.*, 2008, **130**, 15611–15626.
- M. W. Kanan, M. M. Rozenman, K. Sakural, T. M. Snyder and D. R. Liu, *Nature*, 2004, **431**, 545–549.
- B. Seelig and J. W. Szostak, *Nature*, 2007, **448**, 828–831.
- D. D. Albergo and D. H. Turner, *Biochemistry*, 1981, **20**, 1413–1418.
- P. E. Nielsen, M. Egholm, R. H. Berg and O. Buchardt, *Science*, 1991, **254**, 1497–1500.
- C. Zambaldo, S. Barluenga and N. Winssinger, *Curr. Opin. Chem. Biol.*, 2015, **26**, 8–15.
- S. Barluenga and N. Winssinger, *Acc. Chem. Res.*, 2015, **48**, 1319–1331.
- A. Singhal, V. Bagnacani, R. Corradini and P. E. Nielsen, *ACS Chem. Biol.*, 2014, **9**, 2612–2620.
- S. Ficht, A. Mattes and O. Seitz, *J. Am. Chem. Soc.*, 2004, **126**, 9970–9981.
- A. Sen and P. E. Nielsen, *Nucleic Acids Res.*, 2007, **35**, 3367–3374.
- A. Sen and P. E. Nielsen, *Biophys. J.*, 2006, **90**, 1329–1337.
- L. E. Morrison, T. C. Halder and L. M. Stols, *Anal. Biochem.*, 1989, **183**, 231–244.
- B. G. Moreira, Y. You, M. A. Behlke and R. Owczarzy, *Biochem. Biophys. Res. Commun.*, 2005, **327**, 473–484.
- P. L. Privalov and A. I. Dragan, *Biophys. Chem.*, 2007, **126**, 16–24.
- T. Ratilainen, A. Holmén, E. Tuite, P. E. Nielsen and B. Nordén, *Biochemistry*, 2000, **39**, 7781–7791.
- P. E. Nielsen, G. Haaima, A. Lohse and O. Buchardt, *Angew. Chem., Int. Ed.*, 2004, **35**, 1939–1942.
- M. M. Rozenman and D. R. Liu, *ChemBioChem*, 2006, **7**, 253–256.
- M. Sundaralingam and P. K. Ponnuswamy, *Biochemistry*, 2004, **43**, 16467–16476.
- M. C. Chakrabarti and F. P. Schwarz, *Nucleic Acids Res.*, 1999, **27**, 4801–4806.

