



# Ambiphilicity of a mononuclear cobalt(III) superoxo complex†

 Cite this: *Chem. Commun.*, 2020, 56, 14821

 Received 5th August 2020,  
 Accepted 15th October 2020

DOI: 10.1039/d0cc05337f

rsc.li/chemcomm

 Ting-Yi Chen,<sup>ab</sup> Po-Hsun Ho,<sup>a</sup> Can-Jerome Spyra,<sup>b</sup> Franc Meyer,<sup>id</sup> \*<sup>b</sup>  
 Eckhard Bill,<sup>id</sup> <sup>c</sup> Shengfa Ye<sup>id</sup> \*<sup>de</sup> and Way-Zen Lee<sup>id</sup> \*<sup>af</sup>

**Addition of HOTf to a mixture of Co<sup>III</sup>(BDPP)(O<sub>2</sub>•) (1, H<sub>2</sub>BDPP = 2,6-bis((2-(S)-diphenylhydroxymethyl-1-pyrrolidinyl)methyl)pyridine) and Cp\*<sub>2</sub>Fe produced H<sub>2</sub>O<sub>2</sub> in high yield implying formation of Co<sup>III</sup>(BDPP)(OOH) (3), and reaction of Sc(OTf)<sub>3</sub> with the same mixture gave a peroxo-bridged Co<sup>III</sup>/Sc<sup>III</sup> 5. These findings demonstrate the ambiphilic property of Co<sup>III</sup>-superoxo 1.**

Metal-superoxo species are often believed to be the first intermediate following dioxygen (O<sub>2</sub>) association in the catalytic cycle of O<sub>2</sub> activating metalloenzymes.<sup>1</sup> Despite intensive work in the past, the chemistry of metal-superoxo complexes remains largely unexplored, and hence attracts significant attention from chemists and biochemists.<sup>2</sup> *Inter alia*, metal-superoxo intermediates can react with NO• or organic radicals to furnish metal-peroxynitrite<sup>3</sup> and -alkylperoxo<sup>4</sup> complexes *via* radical coupling. Furthermore, they exhibit considerable electrophilicity as indicated by their capability of performing hydrogen atom abstraction (HAA)<sup>5</sup> from weak C–H and O–H bonds and oxygen atom transfer<sup>6</sup> to triphenylphosphine or thiol anisoles. On the other hand, they can initiate deformylation processes when treated with 2-phenylpropionaldehyde, thereby revealing their nucleophilic character.<sup>7,8</sup> Besides the aforementioned well know activities, in a given elementary transformation metal-superoxo intermediates may function not only as an electrophile but also as a nucleophile. In fact, ambiphilicity

of metal-superoxo species has been postulated in a series of theoretical and experimental investigations including O<sub>2</sub> activation catalyzed by α-ketoglutarate dependent dioxygenases,<sup>9</sup> and by Cu, Fe and Co model complexes.<sup>10</sup> Only recently has such ambiphilic property been experimentally confirmed.<sup>8</sup> In our continuing efforts devoted to investigating reactivity of metal-superoxo intermediates, some of us succeeded in preparing a range of homologous Fe<sup>III</sup>-, Co<sup>III</sup>- and Mn<sup>III</sup>-superoxo species by reacting O<sub>2</sub> with the corresponding divalent precursors.<sup>11</sup> It has been shown that these trivalent metal-superoxo complexes can convert into the metal-hydroperoxo complexes *via* HAA. In particular, the reaction of Mn<sup>III</sup>(BDP<sup>B</sup>P)(O<sub>2</sub>•) (H<sub>2</sub>BDP<sup>B</sup>P = 2,6-bis((2-(S)-di(4-bromo)-phenylhydroxymethyl-1-pyrrolidinyl)methyl)pyridine) with trifluoroacetic acid (TFA) and Sc(OTf)<sub>3</sub> yields rare examples of Mn<sup>IV</sup>-hydroperoxo complexes, Mn<sup>IV</sup>(BDP<sup>B</sup>P)(OOH), and [Mn<sup>IV</sup>(μ-OO)Sc(OTf)<sub>n</sub>]<sup>(3-n)+</sup> as evidenced by the combined spectroscopic and computational studies (Scheme 1).<sup>8</sup> Obviously, these proton- and metal-coupled electron transfer processes provide the first experimental support for the proposed ambiphilicity of metal-superoxo species. In this regard, more examples are desired to fully understand how the ambiphilic property of metal-superoxo species affects their chemical reactivity. To this end, we examined the reaction of a Co<sup>III</sup>-superoxo complex, Co<sup>III</sup>(BDPP)(O<sub>2</sub>•) (1, H<sub>2</sub>BDPP = 2,6-bis((2-(S)-diphenylhydroxymethyl-1-pyrrolidinyl)methyl)pyridine) with TFA and Sc(OTf)<sub>3</sub> together with external electron donors.

Treating 1 with HOTf in THF at –90 °C gave a gray-green solution attributed to intermediate 2 having two weak absorption bands at 470 and 640 nm, which reached maxima when 1 equiv. of HOTf was added (the inset of Fig. 1). The existence of an isosbestic point at 590 suggested that no intermediate was formed in the course of conversion of 1 to 2 (Fig. 1). Conversely, complex 1 can be retrieved from deprotonation of 2 by 1 equiv. of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) with a yield of 80% with respect to 1 (Fig. S1, ESI†). Moreover, complex 2 can be obtained from one-electron oxidation of the hydroperoxo complex Co<sup>III</sup>(BDPP)(OOH) (3). Adding equimolar of tris(4-bromophenyl)ammoniumyl hexachloroantimonate, which is often referred to as magic blue, to a THF solution of 3 at –90 °C resulted in a gray-green solution, whose

<sup>a</sup> Department of Chemistry, National Taiwan Normal University, Taipei 11677, Taiwan. E-mail: wzlee@ntnu.edu.tw

<sup>b</sup> Universität Göttingen, Institut für Anorganische Chemie, D-37077 Göttingen, Germany. E-mail: franc.meyer@chemie.uni-goettingen.de

<sup>c</sup> Max-Planck-Institut für Chemische Energiekonversion, Mülheim an der Ruhr D-45470, Germany. E-mail: eckhard.bill@cec.mpg.de

<sup>d</sup> State Key Laboratory of Catalysis, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China. E-mail: shengfa.ye@dicp.ac.cn

<sup>e</sup> Max-Planck-Institut für Kohlenforschung, Mülheim an der Ruhr, D-45470, Germany. E-mail: shengfa.ye@kofo.mpg.de

<sup>f</sup> Department of Medicinal and Applied Chemistry, Kaohsiung Medical University, Kaohsiung 807, Taiwan

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0cc05337f





Scheme 1 Ambiphilicity of metal-superoxo complexes.

Fig. 1 UV-vis spectral changes of the reaction of **1** (1.0 mM, blue trace) with 1 equiv. of HOTf in THF at  $-90\text{ }^{\circ}\text{C}$ . Inset: Titration curve of **1** with HOTf. X-axis: the equivalents of HOTf, Y-axis: the conversion ratio of the absorption peak at 640 nm.

absorption spectrum displayed the same signature features as those found for **2** (Fig. S2, ESI<sup>†</sup>).

To identify the exact nature of the resulting species **2**, we have undertaken detailed spectroscopic characterization and DFT calculations. The EPR measurement of **2** exhibited a spectrum similar to that of **1** ( $A_{\text{Co}} = 18\text{ G}$ ) except for a slightly larger  $^{59}\text{Co}$  hyperfine coupling constant ( $A_{\text{Co}} = 24\text{ G}$ ) seen in Fig. 2, thus indicating that **2** still consists of a  $\text{Co}^{\text{III}}$  center coupled with a radical ligand yielding an overall doublet ground state. The radical ligand thus would be a hydroperoxyl radical or a superoxo having a strong hydrogen bonding interaction with the protonated BDPP<sup>2-</sup> ligand (Scheme 1) as

Fig. 2 X-band EPR spectra of (A) **1** ( $g_1 = 2.098$ ,  $A_{\text{Co}} = 18\text{ G}$ ); (B) **1** with 1 equiv. of HOTf ( $g_1 = 2.066$ ,  $A_{\text{Co}} = 24\text{ G}$ ); (C) **1** with 1 equiv. of DOTf ( $g_1 = 2.064$ ,  $A_{\text{Co}} = 24\text{ G}$ ) and (D) **1** with 1 equiv. of  $\text{Sc}(\text{OTf})_3$  ( $g_1 = 2.066$ ,  $A_{\text{Co}} = 24\text{ G}$ ). Measurement condition:  $T = 77\text{ K}$ ,  $f_{\text{mw}} = 9.6\text{ GHz}$ .

suggested by the crystal structure of **3**.<sup>11b</sup> However, the EPR spectrum of the product generated by reacting **1** with deuterated triflate acid (DOTf) is almost identical to that of **2** without discernable line broadening, which essentially rules out the possibility of the radical ligand being a hydroperoxyl radical (Fig. 2C). Repeated attempts to obtain the O–O vibrational frequencies of **2** from resonance Raman measurements did not accomplish, largely because **2** has only weak chromophores in the usual UV-vis region (Fig. 1). Consequently, the intensity of the O–O stretching signal is too low to be readily detected.

DFT calculations also suggested the O donor of the BDPP<sup>2-</sup> ligand to be the favored protonation site of **1**, consistent with experiment. Even when the starting geometry contained a OOH ligand in which the distal H atom forms a hydrogen bond with the BDPP<sup>2-</sup> ligand, the geometry optimizations invariably shifted the H atom back to the O atom of BDPP<sup>2-</sup> and eventually converged to **A** (Fig. 3). We also tested the initial geometry without the hydrogen bond by tilting the H atom upward. The computations indeed yielded a  $\text{Co}^{\text{III}}$  center bound to a hydroperoxyl radical ligand (**C**), but **C** lies  $23.2\text{ kcal mol}^{-1}$  higher in energy above **A** (Fig. 3). Moreover, formation of a hydrogen bond between the superoxo motif and the proton of the OH group of the protonated BDPP<sup>2-</sup> ligand stabilized **A** by  $10.4\text{ kcal mol}^{-1}$  relative to **B** (Fig. 3). Thus, **A** is best deemed as the most appropriate model for **2**.

Alternatively, to transform **1** into the corresponding peroxy product, we then added 1 equiv. of decamethylferrocene ( $\text{Cp}^*_2\text{Fe}$ ) or sodium naphthalenide ( $\text{NaC}_{10}\text{H}_8$ ) to THF solutions of **1** at  $-90\text{ }^{\circ}\text{C}$ , but UV-vis measurements suggested that no reactions occurred (Fig. S3 and S4, ESI<sup>†</sup>). Taken together, neither proton nor electron donors alone can realize the superoxo-to-peroxy conversion for **1**.

Interestingly, upon treating a mixture containing equimolar **1** and  $\text{Cp}^*_2\text{Fe}$  with 1 equiv. of HOTf, the color of the reaction solution gradually changed from gray-green to dark green then orange; meanwhile, characteristic features of decamethylferrocenium ( $\text{Cp}^*_2\text{Fe}^+$ ) emerged suggesting that  $\text{Co}^{\text{III}}$ -superoxo **1** was reduced in the presence of both HOTf and  $\text{Cp}^*_2\text{Fe}$  (Fig. 4A).





Fig. 3 Computed models for **2**.

During this process, we did not observe the formation of  $\text{Co}^{\text{III}}$ -hydroperoxo **3**. Instead, the reaction produced 19% of  $\text{H}_2\text{O}_2$  with respect to **1**, as determined by iodometric titration (Fig. S5, ESI<sup>†</sup>). When 2 equiv. of HOTf was added, 42% of  $\text{H}_2\text{O}_2$  was furnished (Fig. S6, ESI<sup>†</sup>). Thus, we reasoned that the aforementioned reaction indeed generates **3**; however, once formed, **3** further reacted with HOTf to produce  $\text{H}_2\text{O}_2$ . On the other hand,

treating **2** with 1 equiv. of  $\text{Cp}^*_2\text{Fe}$  (Fig. 4B) also generated 23% of  $\text{H}_2\text{O}_2$  (Fig. S7, ESI<sup>†</sup>). Therefore, all experimental findings revealed that transformation of **1** to **3** proceeds *via* concerted proton coupled electron transfer, which clearly demonstrated the ambiphilicity of **1**.

Reaction of **1** with  $\text{Sc}(\text{OTf})_3$  in THF at  $-90^\circ\text{C}$  generated a product, which showed nearly identical absorption and EPR spectra to those of **2** (Fig. S8 (ESI<sup>†</sup>) and Fig. 2D). On the basis of these observations, we tentatively suggest that the reaction furnished a  $\text{Co}^{\text{III}}$ -superoxo- $\cdots\text{Sc}^{\text{III}}$  species,  $[\text{Co}(\text{BDPP})(\text{O}_2^{\bullet-})\cdots\text{Sc}(\text{OTf})_n]^{(3-n)+}$  (**4**). Furthermore, addition of  $\text{Sc}(\text{OTf})_3$  to the mixture of **1** and  $\text{Cp}^*_2\text{Fe}$  in THF at  $-90^\circ\text{C}$  afforded a  $\text{Co}^{\text{III}}$ -peroxo- $\text{Sc}^{\text{III}}$  complex,  $[\text{Co}(\text{BDPP})(\mu\text{-OO})\text{Sc}(\text{OTf})_n]^{(2-n)+}$  (**5**), as depicted in Scheme 1 (Fig. S9, ESI<sup>†</sup>). Addition of 2 equiv. of HOTf to the THF solution of **5** produced 76%  $\text{H}_2\text{O}_2$  as quantified by iodometric titration experiments (Fig. S10, ESI<sup>†</sup>). Formation of  $\text{H}_2\text{O}_2$  thus strongly supports the identity of peroxo-bridged binuclear **5**. These findings further reinforce the ambiphilic property of **1**.

In comparison with the similar reaction found for  $\text{Mn}^{\text{III}}$  ( $\text{BDP}^{\text{BP}}(\text{O}_2^{\bullet-})$ ) (Scheme 1), the difference can be readily attributed to the much higher oxidation potential of  $\text{Co}^{\text{III}}$  to  $\text{Co}^{\text{IV}}$  than that of  $\text{Mn}^{\text{III}}$  to  $\text{Mn}^{\text{IV}}$ , which can ultimately be rooted back to the distinct effective nuclear charge of low spin  $\text{Co}^{\text{III}}$  compared to high spin  $\text{Mn}^{\text{III}}$  centers. Therefore, formation of an otherwise hydroperoxo O–H bond does not provide a sufficient driving force to trigger an electron transfer from the  $\text{Co}^{\text{III}}$  center to the superoxo ligand. Consequently, the superoxo motif is not electron rich enough to accommodate the incoming proton from HOTf and protonation of the supporting  $\text{BDPP}^{2-}$  ligand is preferred. Therefore, to effect surperoxo-to-peroxo conversion for **1**, an external electron source has to be provided in addition to Brønsted or Lewis acids.

In conclusion, treatment of  $\text{Co}^{\text{III}}$ -superoxo **1** with HOTf and  $\text{Sc}(\text{OTf})_3$  afforded the ligand-protonated  $\text{Co}^{\text{III}}$ -superoxo **2** with a hydrogen bond formed between the  $\text{O}_2^{\bullet-}$  motif and the protonated  $\text{BDPP}^{2-}$  ligand and a superoxo-bridged binuclear  $\text{Co}^{\text{III}}/\text{Sc}^{\text{III}}$  **4**, and  $\text{Co}^{\text{III}}$ -superoxo **1** can be regenerated from deprotonation of **2** by DBU. However, addition of 2 equiv. of HOTf into the reaction mixture of **1** and  $\text{Cp}^*_2\text{Fe}$  produced 42% of  $\text{H}_2\text{O}_2$  suggesting the formation of  $\text{Co}^{\text{III}}$ -hydroperoxo **3**, and the reaction of  $\text{Sc}(\text{OTf})_3$  with **1** in the presence of  $\text{Cp}^*_2\text{Fe}$  gave a peroxo-bridged binuclear  $\text{Co}^{\text{III}}/\text{Sc}^{\text{III}}$  **5**. These findings provided strong experimental support for the ambiphilic property of  $\text{Co}^{\text{III}}$ -superoxo **1**. Interestingly, the ligand-protonated  $\text{Co}^{\text{III}}$ -superoxo **2** can be prepared from one-electron oxidation of  $\text{Co}^{\text{III}}$ -hydroperoxo **3**. The unveiled results underline the critical property of ambiphilicity for metal-superoxo species and direct us to design further investigation strategies towards better understanding  $\text{O}_2$  activation processes carried out by metalloenzymes and related catalysts.

We are grateful for the financial supports from the Ministry of Science and Technology of Taiwan (MOST 108-2113-M-003-009-MY3 to W.-Z. L.) and the Max-Planck Society. W.-Z. L. and S. Y. also acknowledge the MOST-DAAD Project-Based Personnel Exchange Program (MOST 107-2911-I-003-502 and DAAD 57320810). Open Access funding provided by the Max Planck Society.



Fig. 4 UV-vis spectral changes of addition of (A) 1 equiv. of HOTf into a solution of **1** (1.0 mM) in the presence of 1 equiv. of  $\text{Cp}^*_2\text{Fe}$  and (B) 1 equiv. of  $\text{Cp}^*_2\text{Fe}$  into a solution of **2** (1.0 mM) in THF at  $-90^\circ\text{C}$ .



## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- (a) E. G. Kovaleva and J. D. Lipscomb, *Nat. Chem. Biol.*, 2008, **4**, 186–193; (b) M. Costas, M. P. Mehn, M. P. Jensen and L. Que, *Chem. Rev.*, 2004, **104**, 939–986; (c) A. J. Jasniowski and L. Que, Jr., *Chem. Rev.*, 2018, **118**, 2554–2592; (d) C. E. Elwell, N. L. Gagnon, B. D. Neisen, D. Dhar, A. D. Spaeth, G. M. Yee and W. B. Tolman, *Chem. Rev.*, 2017, **117**, 2059–2107; (e) K. Ray, F. F. Pfaff, B. Wang and W. Nam, *J. Am. Chem. Soc.*, 2014, **136**, 13942–13958; (f) S. Sahu and D. P. Goldberg, *J. Am. Chem. Soc.*, 2016, **138**, 11410–11428.
- (a) M. H. Dickman and M. T. Pope, *Chem. Rev.*, 1994, **94**, 569–584; (b) S. Fukuzumi, Y. M. Lee and W. Nam, *Dalton Trans.*, 2019, **48**, 9469–9489; (c) H. Noh and J. Cho, *Coord. Chem. Rev.*, 2019, **382**, 126–144; (d) X. Cai, S. Majumdar, G. C. Fortman, C. S. J. Cazin, A. M. Z. Slawin, C. Lhermitte, R. Prabhakar, M. E. Germain, T. Palluccio, S. P. Nolan, E. V. Rybak-Akimova, M. Temprado, B. Captain and C. D. Hoff, *J. Am. Chem. Soc.*, 2011, **133**, 1290–1293; (e) R. Huacuja, D. J. Graham, C. M. Fafard and C.-H. Chen, *J. Am. Chem. Soc.*, 2011, **133**, 3820–3823; (f) X. Zhang, G. R. Loppnow, R. McDonald and J. Takats, *J. Am. Chem. Soc.*, 1995, **117**, 7828–7829.
- (a) S. Herold and W. H. Koppenol, *Coord. Chem. Rev.*, 2005, **249**, 499–506; (b) R. D. Harcourt, *Coord. Chem. Rev.*, 2018, **358**, 178–180; (c) P. R. Gardner, A. M. Gardner, L. A. Martin and A. L. Salzman, *Proc. Natl. Acad. Sci. U. S. A.*, 1998, **95**, 10378–10383; (d) A. K. Das and M. Meuwly, *Angew. Chem., Int. Ed.*, 2018, **57**, 3509–3513; (e) S. K. Sharma, A. W. Schaefer, H. Lim, H. Matsumura, P. Moëne-Loccoz, B. Hedman, K. O. Hodgson, E. I. Solomon and K. D. Karlin, *J. Am. Chem. Soc.*, 2017, **139**, 17421–17430; (f) R. Cao, L. T. Elrod, R. L. Lehane, E. Kim and K. D. Karlin, *J. Am. Chem. Soc.*, 2016, **138**, 16148–16158; (g) J. J. Liu, M. A. Siegler, K. D. Karlin and P. Moëne-Loccoz, *Angew. Chem., Int. Ed.*, 2019, **58**, 10936–10940.
- (a) E. G. Kovaleva and J. D. Lipscomb, *Science*, 2007, **316**, 453; (b) P. Kumar, S. V. Lindeman and A. T. Fiedler, *J. Am. Chem. Soc.*, 2019, **141**, 10984–10987.
- (a) M. N. Blakely, M. A. Dedushko, P. C. Yan Poon, G. Villar-Acevedo and J. A. Kovacs, *J. Am. Chem. Soc.*, 2019, **141**, 1867–1870; (b) A. Kunishita, M. Kubo, H. Sugimoto, T. Ogura, K. Sato, T. Takui and S. Itoh, *J. Am. Chem. Soc.*, 2009, **131**, 2788–2789; (c) R. L. Peterson, R. A. Himes, H. Kotani, T. Suenobu, L. Tian, M. A. Siegler, E. I. Solomon, S. Fukuzumi and K. D. Karlin, *J. Am. Chem. Soc.*, 2011, **133**, 1702–1705; (d) J. Cho, J. Woo and W. Nam, *J. Am. Chem. Soc.*, 2010, **132**, 5958–5959; (e) H. Kelm and H.-J. Krüger, *Angew. Chem., Int. Ed.*, 2001, **40**, 2344–2348; (f) E. Tamanaha, B. Zhang, Y. Guo, W.-C. Chang, E. W. Barr, G. Xing, J. St Clair, S. Ye, F. Neese, J. M. Bollinger, Jr. and C. Krebs, *J. Am. Chem. Soc.*, 2016, **138**, 8862–8874; (g) N. Kindermann, C.-J. Günes, S. Dechert and F. Meyer, *J. Am. Chem. Soc.*, 2017, **139**, 9831–9834.
- (a) M. T. Kieber-Emmons, J. Annaraj, M. S. Seo, K. M. Van Heuvelen, T. Tosha, T. Kitagawa, T. C. Brunold, W. Nam and C. G. Riordan, *J. Am. Chem. Soc.*, 2006, **128**, 14230–14231; (b) S. Yao, E. Bill, C. Milsman, K. Wieghardt and M. Driess, *Angew. Chem., Int. Ed.*, 2008, **47**, 7110–7113; (c) J. Cho, J. Woo and W. Nam, *J. Am. Chem. Soc.*, 2012, **134**, 11112–11115; (d) K. Fujita, R. Schenker, W. Gu, T. C. Brunold, S. P. Cramer and C. G. Riordan, *Inorg. Chem.*, 2004, **43**, 3324–3326; (e) T. Tano, Y. Kubo, A. Kunishita, M. Kubo, H. Sugimoto, N. Fujieda, T. Ogura and S. Itoh, *Inorg. Chem.*, 2013, **52**, 10431–10437; (f) L.-L. Liu, H.-X. Li, L.-M. Wan, Z.-G. Ren, H.-F. Wang and J.-P. Lang, *Chem. Commun.*, 2011, **47**, 11146–11148.
- (a) P. Pirovano, A. M. Magherusan, C. McGlynn, A. Ure, A. Lynes and A. R. McDonald, *Angew. Chem., Int. Ed.*, 2014, **126**, 6056–6060; (b) W. D. Bailey, N. L. Gagnon, C. E. Elwell, A. C. Cramblitt, C. J. Bouchev and W. B. Tolman, *Inorg. Chem.*, 2019, **58**, 4706–4711.
- Y.-H. Lin, Y. Kutin, M. van Gastel, E. Bill, A. Schnegg, S. Ye and W.-Z. Lee, *J. Am. Chem. Soc.*, 2020, **142**, 10255–10260.
- (a) S. Ye, C. Riplinger, A. Hansen, C. Krebs, J. M. Bollinger, Jr. and F. Neese, *Chem. – Eur. J.*, 2012, **18**, 6555–6567; (b) T. Borowski, A. Bassan and P. E. M. Siegbahn, *Chem. – Eur. J.*, 2004, **10**, 1031–1041.
- (a) W. D. Bailey, D. Dhar, A. C. Cramblitt and W. B. Tolman, *J. Am. Chem. Soc.*, 2019, **141**, 5470–5480; (b) S. Hong, K. D. Sutherlin, J. Park, E. Kwon, M. A. Siegler, E. I. Solomon and W. Nam, *Nat. Commun.*, 2014, **5**, 5440–5547; (c) M. Sankaralingam, Y.-M. Lee, W. Nam and S. Fukuzumi, *Coord. Chem. Rev.*, 2018, **365**, 41–59; (d) A. R. Corcos, O. Villanueva, R. C. Walroth, S. K. Sharma, J. Bacsá, K. M. Lancaster, C. E. MacBeth and J. F. Berry, *J. Am. Chem. Soc.*, 2016, **138**, 1796–1799.
- (a) C. W. Chiang, S. T. Kleespies, H. D. Stout, K. K. Meier, P. Y. Li, E. L. Bominaar, L. Que, Jr., E. Munck and W. Z. Lee, *J. Am. Chem. Soc.*, 2014, **136**, 10846–10849; (b) C. C. Wang, H. C. Chang, Y. C. Lai, H. Fang, C. C. Li, H. K. Hsu, Z. Y. Li, T. S. Lin, T. S. Kuo, F. Neese, S. Ye, Y. W. Chiang, M. L. Tsai, W. F. Liaw and W. Z. Lee, *J. Am. Chem. Soc.*, 2016, **138**, 14186–14189; (c) Y. H. Lin, H. H. Cramer, M. van Gastel, Y. H. Tsai, C. Y. Chu, T. S. Kuo, I. R. Lee, S. Ye, E. Bill and W. Z. Lee, *Inorg. Chem.*, 2019, **58**, 9756–9765.

