


 Cite this: *Phys. Chem. Chem. Phys.*, 2024, 26, 11445

# IR spectroscopic characterization of $[M,C,2H]^+$ ( $M = Ru$ and $Rh$ ) products formed by reacting 4d transition metal cations with oxirane: Spectroscopic evidence for multireference character in $RhCH_2^{+\dagger}$

 Frank J. Wensink,<sup>a</sup> Corry E. Smink,<sup>a</sup> Brandon C. Stevenson,<sup>b</sup> Ryan P. Steele,<sup>b</sup> Joost M. Bakker<sup>\*a</sup> and P. B. Armentrout<sup>\*b</sup>

A combination of infrared multiple-photon dissociation (IRMPD) action spectroscopy and quantum chemical calculations was employed to investigate the  $[M,C,2H]^+$  ( $M = Ru$  and  $Rh$ ) species. These ions were formed by reacting laser ablated  $M^+$  ions with oxirane (ethylene oxide,  $c-C_2H_4O$ ) in a room-temperature ion trap. IRMPD spectra for the  $Ru$  species exhibit one major band and two side bands, whereas spectra for the  $Rh$  species contain more distinct bands. Comparison with density functional theory (DFT), coupled-cluster (CCSD), and equation-of-motion spin-flip CCSD (EOM-SF-CCSD) calculations allows assignment of the  $[M,C,2H]^+$  structures. For the spectrum of  $[Ru,C,2H]^+$ , a combination of  $HRuCH^+$  and  $RuCH_2^+$  structures reproduces the observed spectrum at all levels of theory. The well-resolved spectrum of  $[Rh,C,2H]^+$  could not be assigned unambiguously to any calculated structure using DFT approaches. The EOM-SF-CCSD calculations showed that the ground-state surface has multireference electronic character, and symmetric carbenes in both the  $^1A_1$  and  $^3A_2$  states are needed to reproduce the observed spectrum.

 Received 2nd January 2024,  
 Accepted 13th March 2024

DOI: 10.1039/d4cp00012a

rsc.li/pccp

## Introduction

The functionalization of C–H bonds is a longstanding topic in chemistry.<sup>1,2</sup> Functionalization reactions start with activation, which is a bottleneck for energy-efficient and selective functionalization processes involving C–H bonds because of their typically high bond energies. As the major component of natural gas and the simplest alkane molecule, methane is the most commonly studied alkane, often with the objective to

investigate how to selectively convert methane into liquid fuels like methanol that remain energy rich but are more easily transported. Methane activation catalysis is currently neither specific nor energy efficient enough for direct utilization of methane. Rather, current schemes rely on thermal cracking, followed by synthetic pathways to more valuable chemicals. For more efficient, direct utilization of methane, it is imperative to develop better C–H bond activation catalysts.

Rational design of more active and selective catalyst materials requires a detailed understanding of the interaction between potential catalyst materials and substrates. One way to obtain this information for methane is to study its interactions with model systems for the active site.<sup>3–5</sup> To obtain the most detailed information, the reaction can be isolated in the gas phase, which removes the perturbing effects of a support or solvent. This approach enables studies to focus on the intrinsic interactions. Once such interactions are understood in detail, support, solvent, and pressure effects can be considered.

Previously, studies have shown that methane is activated at room temperature by the third-row (5d) transition metal (TM) cations,  $Ta^+$ ,  $W^+$ ,  $Os^+$ ,  $Ir^+$ , and  $Pt^+$ , leading to dehydrogenation and the formation of  $[M,C,2H]^+$  products.<sup>6–17</sup> Infrared (IR) spectroscopic characterization of  $[M,C,2H]^+$  products showed

<sup>a</sup> Radboud University, Institute for Molecules and Materials, HFML-FELIX, Toernooiveld 7, 6525 ED Nijmegen, The Netherlands. E-mail: joost.bakker@ru.nl

<sup>b</sup> Department of Chemistry, University of Utah, 315 South 1400 East, Salt Lake City, Utah 84112, USA. E-mail: peter.armentrout@chem.utah.edu

† Electronic supplementary information (ESI) available: Figures showing the mass spectrum for reaction of  $Ru^+$  and  $Rh^+$  with oxirane, branching ratios for the photofragmentation of  $[Ru,C,2H]^+$  and  $[Rh,C,2H]^+$ , valence Kohn-Sham orbitals for  $RuCH_2^+$  ( $^4B_2$ ), EOM-SF-CCSD potential energy curves for  $RuCH_2^+$ , spectra of various superpositions of  $[Ru,C,2H]^+$  states, spin density of the  $RhCH_2^+$  ( $^1A_1$ ) state, comparison of the experimental spectrum of  $[Rh,C,2H]^+$  to the unscaled spectra calculated at the EOM-SF-CCSD level, a comparison of  $RhCH_2^+$  spectra calculated at different levels of theory, and  $\omega B97M-V$  spectra for  $[Ru,C,2H]^+$ . A table provides details regarding the calculated states of  $[M,C,2H]^+$  and a comparison to literature calculations. See DOI: <https://doi.org/10.1039/d4cp00012a>



that  $[\text{Pt,C,2H}]^+$  has a  $\text{PtCH}_2^+$  carbene structure with  $C_{2v}$  symmetry, whereas  $[\text{Ta,C,2H}]^+$  and  $[\text{W,C,2H}]^+$  are carbene structures that are distorted by agostic interactions.<sup>18–21</sup> In contrast,  $[\text{Os,C,2H}]^+$  and  $[\text{Ir,C,2H}]^+$  have a hydrido metal carbyne  $\text{HMCH}^+$  structure, and for  $\text{Ir}^+$ , experimental evidence was also found for the co-existence of a higher-energy symmetric  $\text{IrCH}_2^+$  carbene structure.<sup>18–20,22</sup> The observation of the  $\text{HIrCH}^+$  structure demonstrated the need for spectroscopic characterization, because the same structure had been discarded earlier on energetic grounds.<sup>23</sup>

In contrast to third-row TM cations, none of the second-row (4d) TM cations dehydrogenates methane exothermically,<sup>15,24–37</sup> although the reaction with  $\text{Zr}^+$  is only slightly endothermic.<sup>38</sup> To understand the underlying reasons why reactions of the lighter TM cations are endothermic, it is of interest to characterize the structures of  $[\text{M,C,2H}]^+$  products where M is a second-row TM. Because  $\text{Os}^+$  and  $\text{Ir}^+$  were shown to be the most reactive ions for methane dehydrogenation and also show the most diversity in their structures,<sup>13,16–19,22</sup> the present study focuses on the bonding nature and structures of  $[\text{M,C,2H}]^+$  with their second-row congeners: the elements of groups 8 and 9,  $\text{M} = \text{Ru}$  and  $\text{Rh}$ . Earlier studies investigated the  $\text{M}^+ - \text{CH}_2$  bond strengths for first-row TMs and found that the bond dissociation energy (BDE,  $D_0$ ) decreases with an increase in the promotion energy (required for promoting a metal atom in its ground state to an electron configuration where there is one electron in the valence  $s$  orbital that is spin-decoupled from the remaining metal  $d$  electrons) of the atomic metal cation.<sup>39</sup> It seems likely that a parallel trend is followed for the second-row TMs considered here, although this has not been quantified. The value of  $D_0(\text{Ru}^+ - \text{CH}_2)$  has been determined to be  $3.57 \pm 0.05$  eV,<sup>37</sup> such that formation of  $\text{RuCH}_2^+$  and  $\text{H}_2$  from  $\text{Ru}^+$  and  $\text{CH}_4$  requires  $1.17 \pm 0.05$  eV given  $D_0(\text{CH}_2 - \text{H}_2) = 4.743 \pm 0.001$  eV.<sup>40</sup> Analogously,  $D_0(\text{Rh}^+ - \text{CH}_2)$  has been determined as  $3.69 \pm 0.08$  eV in guided ion beam experiments<sup>41</sup> and as  $4.08 \pm 0.22$  eV by ion–molecule bracketing reactions.<sup>42</sup> Thus, formation of  $\text{RhCH}_2^+$  and  $\text{H}_2$  from  $\text{Rh}^+$  and  $\text{CH}_4$  requires an energy of  $1.03 \pm 0.08$  ( $0.67 \pm 0.22$ ) eV.<sup>32</sup>

Elsewhere, a B3LYP potential energy surface (PES) for the reaction between  $\text{Ru}^+$  and  $\text{CH}_4$  identified the first step as physisorption with an exothermicity of 0.78 eV.<sup>37</sup> This energy can be used to transfer one hydrogen atom from methane to  $\text{Ru}^+$ , forming  $\text{H-Ru}^+ - \text{CH}_3$ , which still lies 0.18 eV below the energy of the reactants. A further step toward  $(\text{H}_2)\text{RuCH}_2^+$  requires the crossing of endothermic energy barriers and a spin flip from quartet to doublet. Formation of the doublet-spin  $(\text{H}_2)\text{RuCH}_2^+$  intermediate is still exothermic by 0.09 eV, but loss of  $\text{H}_2$  requires over 1 eV, such that the overall dehydrogenation reaction to form  $\text{RuCH}_2^+ (^2\text{A}_2) + \text{H}_2$  is calculated to be endothermic by 1.08 eV. Energetically more favorable are the formation of the doublet-spin  $\text{HRuCH}^+$  (+0.97 eV) and quartet-spin  $\text{RuCH}_2^+$  (+0.94 eV) products. Formation of the former has no barriers exceeding the product energy, whereas progress along the quartet-spin surface requires passing over a tight-transition state barrier of 1.03 eV.<sup>37</sup>

For  $\text{RhCH}_2^+$ , multireference – single double configuration interaction – complete active space self-consistent field

(MR-SDCI-CASSCF)<sup>43</sup> and B3LYP<sup>44</sup> calculations have identified a  $^1\text{A}_1$ -state carbene as lowest in energy, followed by a  $^3\text{A}_2$  state lying 0.20 and 0.19 eV, respectively, higher in energy. (The relative energies of higher-lying states identified in the former study are included in Table S1, ESI†). Again, the only intermediate that can be formed exothermically on the PES of  $\text{Rh}^+(^3\text{F}) + \text{CH}_4$  is the physisorption step;<sup>43,44</sup> however, the CASSCF calculations found a barrier of about 0.4 eV for the reverse  $\text{RhCH}_2^+ + \text{H}_2$  reaction, whereas the later B3LYP calculations found no barrier. The latter result agrees with experiments for both the forward and reverse reactions.<sup>32,42</sup>

As described above, none of the metal cations we study here react exothermally with methane.<sup>26,31,32,37</sup> Therefore, in order to form the  $[\text{M,C,2H}]^+$  species of interest, we react the corresponding  $\text{M}^+$  ions with oxirane (ethylene oxide,  $\text{c-C}_2\text{H}_4\text{O}$ ). Oxirane readily reacts with many TM cations because of its ring strain and because  $[\text{M,C,2H}]^+$  formation is accompanied by the stable  $\text{CH}_2\text{O}$  neutral product. Thus, extraction of  $\text{CH}_2$  from oxirane requires much less energy than from methane,  $D_0(\text{CH}_2 - \text{CH}_2\text{O}) = 3.375 \pm 0.004$  eV, compared to the  $D_0(\text{CH}_2 - \text{H}_2) = 4.743 \pm 0.001$  eV already mentioned.<sup>40</sup> In contrast to the endothermic reactions with methane, the thermochemistry noted above indicates that formation of  $\text{MCH}_2^+ + \text{CH}_2\text{O}$  in the reaction of  $\text{M}^+$  with oxirane is exothermic by  $0.20 \pm 0.05$  and  $0.32 \pm 0.08$  eV for  $\text{M} = \text{Ru}$  and  $\text{Rh}$ , respectively. Although it has been shown that laser-ablated  $\text{Rh}^+$  reacts at thermal energies with oxirane to yield  $\text{RhCH}_2^+$ ,<sup>42</sup> product distributions for the reactions of  $\text{Ru}^+$  and  $\text{Rh}^+$  with  $\text{c-C}_2\text{H}_4\text{O}$  have not been published previously.

In the present work, we probe the structures of the  $[\text{M,C,2H}]^+$  product ions for  $\text{M} = \text{Ru}$  and  $\text{Rh}$  using a combination of IR multiple photon dissociation (IRMPD) action spectroscopy and DFT or coupled cluster with single and double excitations (CCSD) calculations. IRMPD spectra are recorded using the same Fourier transform ion cyclotron resonance (FTICR) mass spectrometer recently used to show that the reaction of  $\text{Pt}^+$  with two methane molecules leads to the formation of a  $\text{Pt}^+(\text{ethene})$  complex following two dehydrogenations and C–C coupling.<sup>21</sup> In that same work, we demonstrated that the spectrum of  $[\text{Pt,C,2H}]^+$  recorded with this instrument is consistent with that reported previously,<sup>18</sup> where  $[\text{Pt,C,2H}]^+$  was formed in a molecular beam environment without mass-isolation prior to irradiation. Earlier attempts to record IRMPD spectra of  $[\text{M,C,2H}]^+$  with  $\text{M} = \text{Ru}$  were unsuccessful in the molecular beam apparatus because the ion intensities were insufficient. Here, we take advantage of the possibility to react  $\text{TM}^+$  with  $\text{c-C}_2\text{H}_4\text{O}$  over longer times and to mass-isolate the formed  $[\text{M,C,2H}]^+$  species in order to obtain the desired spectra.

Critically, these small ions also form benchmark systems for the accuracy of quantum chemistry methods. Previously, we found that relatively standard DFT methods proved sufficient to accurately predict the IR fingerprints, both in terms of frequencies and IR intensities that are necessary for assigning the experimental IR spectra of  $[\text{M,C,2H}]^+$  species with 5d TM elements.<sup>18–22</sup> In contrast, in a parallel publication investigating the IR spectra of  $\text{FeCH}_2^+$  and  $\text{CoCH}_2^+$ ,<sup>45</sup> 3d congeners of the



species currently under investigation, we found that a large variety of DFT methods provide a significant underestimation of the IR absorption intensity for the M–C stretch vibration. We provided computational evidence suggesting that this relates to inaccurate charge separation of these DFT methods. One motivation of the present work is therefore to assess whether DFT treatments of this mode are as poor as they were for 3d elements, or whether they are accurate enough, like for the 5d elements studied earlier. We therefore also employ higher-level single- and multireference methods to interpret the observed spectra. One previous computational study,<sup>43</sup> based on CASSCF and MR-SDCI-CASSCF methods, noted only modest multireference contributions to the ground-state wavefunction of  $\text{RhCH}_2^+$ . The same analyses were unable to reproduce the experimentally observed  $\text{Rh}^+-\text{CH}_2$  binding energy and yielded relative electronic state energies that differ appreciably from the ones computed herein. We find that adequately treating the multireference character, while employing modern core potentials and basis sets, is needed to properly reproduce the observed spectrum of  $\text{RhCH}_2^+$ . In the case of  $[\text{Ru,C,2H}]^+$ , multi-reference character does not appear to be relevant, but here contributions from multiple structures and states, whose energies vary with the level of theory, are required to match the experimental spectrum.

## Methods

### Experimental

Atomic metal (Ru, Rh) cations were produced in a laser vaporization source.<sup>46,47</sup> In this source, a solid metal target disk was irradiated by a frequency doubled Nd:YAG laser at 30 Hz in the presence of helium gas injected by a pulsed valve. The helium cooled the metal ions through collisions and guided them toward the end of the source, where the gas mixture adiabatically expanded into vacuum, thereby further cooling the ions. They were then transferred, *via* a quadrupole mass filter in guidance mode, to a quadrupole ion trap with rectangular electrodes where they were trapped by dissipating the kinetic energy with  $5 \times 10^{-4}$  mbar of argon buffer gas. In the trap, the metal ions reacted with oxirane at a partial pressure of  $1 \times 10^{-6}$  mbar, which was mixed into the argon. After accumulating ions for approximately 400 ms, the voltage on the exit electrode of the ion trap was lowered. This released the ions from the ion trap after which they were guided into the cavity of the Free Electron Laser for IntraCavity Experiments (FELICE). Here, the ions were trapped in one of the cells of the FTICR mass spectrometer integrated within the laser cavity. This FTICR has four cells, with the first (cell 1) located in the laser focus and the fourth 30 cm from the focus, thereby lowering the photon fluence by a factor of 14. After ion capture, all unwanted masses were ejected from the FTICR cell using a combination of radio frequency (rf) excitation pulses (chirped and single-frequency),<sup>48</sup> so that only the species of interest were irradiated by IR light produced by FELICE. Resonant absorption of IR photons increased the internal energy of the ions and once the

fragmentation threshold was overcome, the ions dissociated, which indicates the presence of a vibrational band at the resonant IR frequency. The IRMPD spectrum was obtained by recording the intensities of precursor ( $I_p$ ) and fragment ( $I_{\text{frag}}$ ) ions present in the FTICR cell after irradiation and is expressed as the fragmentation yield  $Y$ :

$$Y = \ln\left(\frac{I_p + I_{\text{frag}}}{I_p}\right)$$

The IR light of FELICE is tunable in the 100–2000  $\text{cm}^{-1}$  spectral range, although for this work only the 300–2000  $\text{cm}^{-1}$  spectral range was used. The spectral bandwidth was near transform limited and, during these measurements, the full-width at half-maximum (FWHM) was approximately 0.7% of the central frequency. FELICE produced micropulses at 1 GHz, and the macropulses lasted between 6–10  $\mu\text{s}$  and were repeated at 5 or 10 Hz. All spectra were recorded using a single macropulse with the ions in FTICR cell 4 (low fluence) unless specifically indicated otherwise. FELICE macropulse energies used ranged from 0.3 J at lower frequencies to 0.5 J at higher frequencies, corresponding to macropulse fluences ranging from 0.9 to 6.6  $\text{J cm}^{-2}$  in FTICR cell 4.

### Computational

The DFT calculations were done using the Gaussian 16 software package,<sup>49</sup> where all molecular structures, closed- and open-shell, were optimized using the unrestricted B3LYP hybrid functional and the def2-TZVPPD basis set.<sup>50–52</sup> This combination was used because the  $\text{M}^+-\text{CH}_2$  BDEs calculated for the metals explored here matched the experimental values relatively well:  $D_0(\text{Ru}^+-\text{CH}_2) = 3.73$  eV (experimental  $3.57 \pm 0.05$  eV) and  $D_0(\text{Rh}^+-\text{CH}_2) = 3.60$  eV (experimental  $3.69 \pm 0.08$  eV). All electrons were treated explicitly for H and C, whereas an effective core potential (ECP) of 28 electrons (small core) was used for Ru and Rh with the 4s, 4p, 5s, and 4d orbitals treated explicitly by the basis set. To ascertain true minima and for comparison with the experimental spectra, harmonic vibrational frequencies were calculated. All relative energy values provided below include harmonic zero-point vibrational energies. For each vibrational mode, rovibrational envelopes were simulated assuming pure a-, b-, or c-type transitions on the basis of moments obtained from the computations.<sup>53</sup> For comparison with experiment, the simulated rotational substructures were shifted to the unscaled, calculated vibrational frequencies and the individual rotational transition intensities multiplied by the vibrational transition intensities. (For these calculations, no scaling factors were used to compare the theoretical spectra with experiment, which is unusual for these types of systems.<sup>18–22,54–56</sup> Nevertheless, such scaling factors were not found to improve the agreement between theory and experiment and therefore are not used throughout this work except as noted.) The resulting rovibrational transition lines were subsequently convoluted using a Gaussian line shape function with a FWHM of 0.9% of the central frequency, mimicking the FELICE spectral bandwidth. It should be noted



that because all calculated spectra correspond to absorption of a single IR photon, the predicted intensities can differ from the experimental multiple-photon spectra.

Additional geometry and frequency calculations were performed at the second-order Möller–Plesset, MP2,<sup>57</sup> and coupled-cluster with single and double excitations, CCSD,<sup>58</sup> levels using the same def2-TZVPPD basis set. These calculations were followed by single-point calculations with added perturbative triple excitations, *i.e.*, at the CCSD(T)/def2-TZVPPD level of theory.

Because of the large degree of spin contamination initially observed in DFT (and even CCSD) computations of the singlet  $[\text{Rh,C,2H}]^+$  and doublet  $[\text{Ru,C,2H}]^+$  species, additional calculations were performed with the equation-of-motion spin-flip<sup>59,60</sup> CCSD (EOM-SF-CCSD) level of theory using the cc-pVTZ-PP basis set<sup>61</sup> and the ECP28MDF pseudopotential<sup>61</sup> on Ru/Rh and the cc-pVTZ basis set<sup>62</sup> on C and H. To clarify relative energies, perturbative triples were added *via* single-point computations with the EOM-SF-CCSD using the perturbative Fock triples correction (fT)<sup>63</sup> on optimized structures. The Q-Chem software package was utilized for the MP2, CCSD, and EOM computations, and the 4s and 4p electrons on Ru/Rh and the 1s electrons on C were frozen for the electron-correlation calculations. The ground triplet state was used as the spin-flip reference for  $[\text{Rh,C,2H}]^+$ . For  $[\text{Ru,C,2H}]^+$ , a peculiar symmetry restriction required special treatment: the effectively doubly degenerate ground quartet states of  $B_1$  and  $B_2$  symmetry led to independent (and also nearly degenerate) manifolds of spin-flip states. Therefore, both quartets were used as spin-flip references, and properties of both manifolds will be presented herein. In the asymmetric HRuCH<sup>+</sup> configuration, the lowest  $^4A'$  state was used as the spin-flip reference. The structures of at least the two lowest-energy electronic states of each spin-flip transition symmetry were first optimized with the analogous cc-pVDZ-PP basis set, using a series of disparate starting structures. Stable structures were then re-optimized with the cc-pVTZ-PP basis, and harmonic zero-point corrections and spectra were computed. Total electronic densities, spin densities,  $\langle S^2 \rangle$  values, and Mulliken spin populations and charges were computed with the resulting EOM-SF-CCSD wavefunctions. Constrained potential energy curves along the metal–carbene coordinate were also computed by optimizing the structure of the lowest-energy electronic state at a series of Rh–C( $A_1$ )/Ru–C( $B_1/B_2$ ) distances between 1.5 and 2.5 Å. The energies of all other electronic states were computed as vertical single-points from these structures. Here, adequate comparison of the predicted IR spectrum for  $[\text{Rh,C,2H}]^+$ , but not  $[\text{Ru,C,2H}]^+$ , with experiment required the use of frequency scaling factors, which are detailed below.

## Results & discussion

Reactions between  $M^+$  ( $M = \text{Ru}, \text{Rh}$ ) and oxirane resulted in the ionic products  $MC^+$ ,  $[\text{M,C,2H}]^+$ ,  $[\text{M,O,2H}]^+$  for both  $M$ ,  $MO^+$  and  $[\text{M,2C,2H}]^+$  for  $M = \text{Ru}$ , and  $MH^+$ ,  $[\text{M,C,3H}]^+$ ,  $[\text{M,C,O}]^+$ , and

$[\text{M,C,O,H}]^+$  for  $M = \text{Rh}$ . These product mass distributions are shown in Fig. S1 of the ESI.† Relative product intensities are dependent on the metal ion; for example,  $MO^+$  is fairly intense for  $\text{Ru}^+$ , whereas it is barely visible for  $\text{Rh}^+$ . A low-intensity  $\text{RhH}^+$  ion was also observed, but the reaction of ground-state  $\text{Rh}^+$  with  $c\text{-C}_2\text{H}_4\text{O}$  to form  $\text{RhH}^+ + \text{CH}_3\text{CO}$  (the lowest energy isomer) is endothermic by  $0.95 \pm 0.04$  eV.<sup>40,64</sup> It seems possible that this species is formed by reactions with contaminants, by electronically excited  $\text{Rh}^+$ , or by trap loading at excess kinetic energy. Overall, we conclude that these reactions are likely to mainly generate species that can be formed exothermically from ground-state reactants, but some excess excitation energy may be available as well.

### IR spectroscopy of $\text{RuCH}_2^+$

Ruthenium has a complex isotopic distribution with natural abundances at 96 (5.5%), 98 (1.9%), 99 (13%), 100 (13%), 101 (17%), 102 (32%), and 104 (19%) amu. Fig. 1(a) shows the experimental IRMPD spectrum of  $[\text{Ru,C,2H}]^+$ . The spectrum was recorded using two FEL settings, indicated by the trace colors, with overlap between 900–1200  $\text{cm}^{-1}$ . It is constructed using only the  $^{101}\text{Ru}$  and  $^{102}\text{Ru}$  isotopes ( $m/z$  115, 116), using  $\text{RuC}^+$  as the only photofragment observed ( $m/z$  113, 114, Fig. S2a, ESI†), consistent with the dissociation thermochemistry in Table 1. Mass isolation of  $m/z$  115, 116 prior to IR irradiation leaves small amounts of  $^{104}\text{RuC}^+$  ( $m/z$  116),  $^{99}\text{RuO}^+$  ( $m/z$  115),  $^{100}\text{RuO}^+$  ( $m/z$  116), and  $^{98}\text{RuOH}_2^+$  ( $m/z$  116), all of which can be distinguished at the mass resolution of the FT-ICR. Because these species cannot fragment into mass channels  $m/z$  113, 114, their presence does not affect the spectrum of  $[\text{Ru,C,2H}]^+$ . The spectrum of  $[\text{Ru,C,2H}]^+$  in Fig. 1(a) shows one dominant band peaking at 637  $\text{cm}^{-1}$ . The band extends a bit further into the blue with positions of submaxima at 778 and 987  $\text{cm}^{-1}$ , identified from fitting the bands with Gaussian functions.

Fig. 1(b)–(h) shows the simulated IR spectra of different  $[\text{Ru,C,2H}]^+$  isomers and states calculated at the B3LYP level. Here, the  $^4B_2$  ground state (GS) is a symmetric ruthenium carbene geometry. With the convention that the  $z$ -axis is the symmetry axis and that the molecule lies in the  $yz$  plane, the  $^4B_2$  GS has a Ru–C  $\sigma$  bonding molecular orbital (MO) ( $a_1$  symmetry, using  $4d_{z^2}$  on Ru, double occupancy), a Ru–C  $\pi$  bond ( $b_1$ , Ru  $4d_{xz}$ , double), three non-bonding MOs localized on Ru: ( $a_1$ ,  $4d_{x^2-y^2}$ , single;  $a_2$ ,  $4d_{xy}$ , double;  $a_1$ , 5s, single), and an antibonding MO combining the Ru  $4d_{yz}$  and  $\text{CH}_2$  orbitals ( $b_2$ , single). Thus, the valence MO occupation (excluding the four  $\text{CH}_2$  bonding electrons) is  $(1a_1)^2(1b_1)^2(2a_1)^1(1a_2)^2(3a_1)^1(1b_2)^1$ . These valence molecular orbitals are shown in Fig. S3 of the ESI.†

A second electronic state,  $^4B_1$ , calculated to lie only 0.001 eV higher in energy, reverses the occupation of the  $4d_{x^2-y^2}$  ( $2a_1$ ) and  $4d_{xy}$  ( $1a_2$ ) non-bonding orbitals, explaining the small energy difference between these two states. In another low-lying state ( $^4A_2$ ), one electron in the non-bonding  $4d_{xy}$  ( $1a_2$ ) MO is transferred to the antibonding  $1b_2$  MO, leading to decreased stability, 0.15 eV above the GS. Additional carbene structures were located for different spin states. The most stable doublet-spin ruthenium carbene is  $^2A_2$ , 0.13 eV above the GS with the






Fig. 1 (a) Experimental IRMPD spectrum of  $[\text{Ru,C,2H}]^+$ , the black trace is the average of four scans, the red trace is a single scan; (b)–(h) B3LYP/def2-TZVPPD calculated IR spectra of different  $[\text{Ru,C,2H}]^+$  states and isomers with the harmonic vibrations in red and the rovibrational envelopes in black accompanied by geometric structure, relative energy, and electronic state.

Table 1 Fragmentation channels and dissociation energies ( $D_0$ ) of the lowest energy  $[\text{M,C,2H}]^+$  species

Species	Fragments	Bond dissociation energy $D_0$ (eV)	
		Theory <sup>a</sup>	Experiment
$\text{RuCH}_2^+$	$\text{RuC}^+ + \text{H}_2$	1.15	$0.72 \pm 0.09^{bcd}$
	$\text{Ru}^+ + \text{CH}_2$	3.73	$3.57 \pm 0.05^c$
$\text{RhCH}_2^+$	$\text{RhC}^+ + \text{H}_2$	1.29	$2.76 \pm 0.20^{be}$
	$\text{Rh}^+ + \text{CH}_2$	3.60	$3.69 \pm 0.08^e, 4.08 \pm 0.22^f$

<sup>a</sup> Theoretical values are calculated at the uB3LYP/def2-TZVPPD level of theory ignoring any barriers. <sup>b</sup> Uses  $D_0(\text{C-H}_2) = 3.320 \pm 0.001$  eV.<sup>40</sup> <sup>c</sup> Ref. 37. <sup>d</sup> Ref. 65. <sup>e</sup> Ref. 32,41. <sup>f</sup> Ref. 42.

MO occupation of  $(1a_1)^2(1b_1)^2(2a_1)^2(1a_2)^1(3a_1)^0(1b_2)^2$ . Moving an electron between the  $4d_{x^2-y^2}$  and  $4d_{xy}$  non-bonding orbitals results in a  $^2A_1$  state 0.16 eV above the GS. Energetics and

selected orbital configurations of all stable  $[\text{Ru,C,2H}]^+$  species located computationally are listed in Table S1 of the ESI.<sup>†</sup>

Interestingly, early generalized valence bond (GVB) calculations of  $\text{RuCH}_2^+$  by Carter and Goddard obtained a very different ordering of these states.<sup>66</sup> They found a ground state of  $^2A_2$  with a nearly degenerate  $^2A_1$  state, followed by the three quartet states ( $^4A_2$ ,  $^4B_2$ , and  $^4B_1$ ) lying about 0.41 eV above the ground state, with a  $^2B_2$  state at 0.87 eV (see also Table S1, ESI<sup>†</sup>). However, they calculate a bond energy ( $D_e$ ) at their most advanced level of theory of 2.95 eV and an estimated “exact” value of 3.19 eV, which is well below the experimentally observed values. In their analysis of the bonding of these states, they characterize the bonding in the doublet spin states as covalently bound “metal–methylidenes”, whereas the quartet states are viewed as “metal–carbenes” formed by  $\sigma$ -donor/ $\pi$ -acceptor interactions.

We checked for agostic carbene geometries in the DFT calculations, but these structures converged to symmetric carbenes. More interestingly, we did find a hydrido ruthenium carbyne geometry,  $\text{HRuCH}^+$ , with planar  $C_s$  symmetry. This species has a  $^2A'$  electronic state that lies only 0.029 eV above the GS, in agreement with previous theoretical results.<sup>37</sup> The  $^4A''$  hydrido carbyne was found at 1.32 eV above the GS. The lowest-energy dihydrido ruthenium carbide,  $\text{HHRuC}^+$ , is the  $^2A'$  state lying 1.72 eV above the GS. The lowest-energy dihydrogen ruthenium carbide,  $(\text{H}_2)\text{RuC}^+$ , is the  $^2A'$  state at 0.47 eV above the GS. The lowest-energy sextet species located is a  $^6A'$  state, 1.43 eV higher in energy than the GS. This species exhibits the ruthenium carbene geometry, but both hydrogen atoms are bent slightly out of plane (dihedral angle H–H–Ru–C = 8.8°).

The B3LYP-generated IR spectrum for the  $^4B_2$  state of  $\text{RuCH}_2^+$  is dominated by the  $\text{CH}_2$  rocking and wagging modes, which are calculated at 542 and 739  $\text{cm}^{-1}$ , respectively, Fig. 1(b). These vibrations are shifted to 441 and 865  $\text{cm}^{-1}$  for the  $^4B_1$  state, where the  $\text{CH}_2$  rocking-mode intensity is decreased by a factor of seven while that of the  $\text{CH}_2$  wagging mode is increased by a factor of three. Thus, the spectra of the  $^4B_1$  and  $^4B_2$  states of  $\text{RuCH}_2^+$  differ appreciably. Interestingly, MP2 calculations enhance the intensity of the Ru–C stretch, but CCSD calculations yield an intensity similar to that obtained by B3LYP. For example, for the  $^4B_2$  GS, the frequencies in  $\text{cm}^{-1}$  (intensities,  $\text{km mol}^{-1}$ ) for the stretch calculated at the B3LYP/CCSD levels are 655 (1)/660 (9) compared to MP2 at 648 (53). Other  $\text{RuCH}_2^+$  states have similar comparisons. It seems plausible that the experimental spectrum could be reproduced by a superposition of the five low-lying  $\text{RuCH}_2^+$  states ( $^4B_2$ ,  $^4B_1$ ,  $^2A_2$ ,  $^4A_2$ , and  $^2A_1$ ), all of which are accessible according to the B3LYP calculations (Fig. 1) because the reaction of  $\text{Ru}^+$  with oxirane is exothermic by  $0.20 \pm 0.05$  eV. At the CCSD(T)//CCSD level of theory, these states lie 0.045, 0.044, 0.150, 0.091, and 0.134 eV, respectively (Table S1, ESI<sup>†</sup>), above the ground-state  $\text{HRuCH}^+$  species (see next paragraph), such that all five states could be formed exothermically.

Complicating the assignment of the experimental spectrum is the low energy of the  $\text{HRuCH}^+$  ground state. For the  $\text{HRuCH}^+$  ( $^2A'$ ) species, B3LYP predicts four modes below 1500  $\text{cm}^{-1}$ : an



out-of-plane C–H bending mode ( $743\text{ cm}^{-1}$ , c-type), two in-plane H–M–C–H bending modes ( $611$  and  $965\text{ cm}^{-1}$ , both with mixed a- and b-type transition character), and the M–C stretch mode ( $1058\text{ cm}^{-1}$ , a-type). The M–H and C–H stretch modes are calculated to lie above  $2000\text{ cm}^{-1}$  and are thus out of the scope of our investigation. According to B3LYP theory, only the two low-frequency modes have appreciable intensities, and these transitions fall directly in line with the observed spectrum, Fig. 1(d). For this structure, MP2 calculations again enhance the intensities of these transitions and the other two modes in the experimentally observed window. In contrast, CCSD calculations yield results in good agreement with the relative intensities predicted by B3LYP, with frequencies of  $622$ ,  $751$ ,  $974$ , and  $1036\text{ cm}^{-1}$ , Table 2. Where the B3LYP method puts this HRuCH<sup>+</sup> state  $0.029\text{ eV}$  above the RuCH<sub>2</sub><sup>+</sup> (<sup>4</sup>B<sub>2</sub>) state, CCSD(T)//CCSD and EOM-SF-CCSD(ft)/EOM-SF-CCSD calculations make

it the ground state, lying  $0.045\text{ eV}$  and  $0.006\text{ eV}$ , respectively, below the <sup>4</sup>B<sub>2</sub> state. The role of perturbative triples corrections was found to be critical here, as both CCSD and EOM-SF-CCSD predicted the <sup>4</sup>B states to be lower in energy without these corrections. The potential energy landscape of the EOM-SF-CCSD computed states is further clarified in Fig. S4 of the ESI.†

Comparisons of the predicted CCSD and EOM-SF-CCSD spectra for the low-lying states with the experimental spectrum are shown in Fig. 2 and 3, respectively. We conclude that the HRuCH<sup>+</sup> species is present and captures the main feature at  $637\text{ cm}^{-1}$ , but does not capture the full shape of the  $637$  and  $778\text{ cm}^{-1}$  structure. CCSD, with the pure vibrational splitting of the bright transitions in the HRuCH<sup>+</sup> (<sup>2</sup>A') spectrum of  $622$  and  $751\text{ cm}^{-1}$ , can account for both features, but does not provide the peak intensity for the  $637\text{ cm}^{-1}$  band, whereas EOM ( $633$  and  $718\text{ cm}^{-1}$ ) is not sufficient to capture the peak at  $778\text{ cm}^{-1}$ .

**Table 2** Experimental band positions and strengths (s = strong, m = medium) accompanied by theoretical calculated frequencies, intensities, and descriptions for the assigned structures

	Experiment		Theory B3LYP/CCSD/EOM <sup>a</sup>		Mode description
	Frequency (cm <sup>-1</sup> )	Strength	Frequency (cm <sup>-1</sup> )	Intensity (km mol <sup>-1</sup> )	
RuCH <sub>2</sub> <sup>+</sup> ( <sup>4</sup> B <sub>2</sub> )	637	s	542/534/508	27/34/33	CH <sub>2</sub> rock
	778	s	655/660/649	1/9/10	M–C stretch
	987	m	739/731/765	42/49/62	CH <sub>2</sub> wag
			1313/1343/1336	0/0/2	CH <sub>2</sub> scissor
RuCH <sub>2</sub> <sup>+</sup> ( <sup>4</sup> B <sub>1</sub> )	637	s	441/516/489	4/22/23	CH <sub>2</sub> rock
	778	s	651/659/648	1/9/10	M–C stretch
	987	m	865/759/781	115/64/72	CH <sub>2</sub> wag
			1310/1341/1335	0/0/3	CH <sub>2</sub> scissor
RuCH <sub>2</sub> <sup>+</sup> ( <sup>2</sup> A <sub>2</sub> )	637	s	568/597/388	0/1/31	CH <sub>2</sub> rock
	778	s	734/739/745	8/0/3	M–C stretch
	987	m	933/924/947	59/65/67	CH <sub>2</sub> wag
			1397/1413/1360	3/2/0	CH <sub>2</sub> scissor
RuCH <sub>2</sub> <sup>+</sup> ( <sup>2</sup> A <sub>1</sub> )	637	s	566/593/470	0/1/1	CH <sub>2</sub> rock
	778	s	766/749/747	8/0/0	M–C stretch
	987	m	946/925/942	59/64/69	CH <sub>2</sub> wag
			1401/1415/1413	3/2/3	CH <sub>2</sub> scissor
RuCH <sub>2</sub> <sup>+</sup> ( <sup>4</sup> A <sub>2</sub> )	637	s	703/517/471	25/22/1	CH <sub>2</sub> rock
	778	s	667/647/490	1/8/30	M–C stretch
	987	m	980/766/689	104/58/0	CH <sub>2</sub> wag
			1407/1352/1334	4/1/13	CH <sub>2</sub> scissor
HRuCH <sup>+</sup> ( <sup>2</sup> A')	637	s	611/622/633	92/97/94	HMCH ip bend
	778	s	743/751/718	91/94/121	C–H oop bend
	987	m	965/974/1003	1/1/3	HMCH ip bend
			1058/1036/1102	5/3/3	M–C stretch
RhCH <sub>2</sub> <sup>+</sup> ( <sup>1</sup> A <sub>1</sub> )	774	s	606/624/600	0/1/1	CH <sub>2</sub> rock
	904	s	793/748/819	13/1/28	M–C stretch
	1029	s	964/971/964	57/65/83	CH <sub>2</sub> wag
	1269	m	1386/1408/1386	4/1/19	CH <sub>2</sub> scissor
	1783	s			Overtone
RhCH <sub>2</sub> <sup>+</sup> ( <sup>3</sup> A <sub>2</sub> )	774	s	646/627/657	8/21/6	CH <sub>2</sub> rock
	904	s	674/674/669	21/27/28	M–C stretch
	1029	s	942/933/948	89/110/125	CH <sub>2</sub> wag
	1269	m	1396/1431/1425	5/2/2	CH <sub>2</sub> scissor
	1783	s			Overtone

<sup>a</sup> B3LYP and CCSD calculations used the def2-TZVPPD basis set. EOM-SF-CCSD calculations used the cc-pVTZ-PP basis set and ECP28MDF core potential on the metal and cc-pVTZ on the C/H atoms.






Fig. 2 (a) Experimental IRMPD spectrum of  $[\text{Ru,C,2H}]^+$ , the black trace is the average of four scans, the red trace is a single scan; (b)–(g) CCSD/def2-TZVPPD calculated IR spectra of different  $[\text{Ru,C,2H}]^+$  states and isomers with the harmonic vibrations in red and the rovibrational envelopes in black accompanied by geometric structure, relative energy, and electronic state.

Moreover, neither method can explain the broad band observed at  $987\text{ cm}^{-1}$ .

We, therefore, consider additional contributors to the unique lineshape observed experimentally for  $\text{RuCH}_2^+$ . One option is that the  $^2\text{A}_1$  and  $^2\text{A}_2$  states account for the  $987\text{-cm}^{-1}$  transition. For the EOM computations, the combination of the spectra for these states with that for  $\text{HRuCH}^+$  ( $^2\text{A}'$ ) would lead to two bands, roughly centered at  $700$  and  $950\text{ cm}^{-1}$ , respectively, but a large drop in the predicted spectral brightness between them (see Fig. S5 in the ESI†). For CCSD results, a more reasonable, continuous band is formed, but the intensity for the  $978\text{ cm}^{-1}$  band is comparable to that near  $637\text{ cm}^{-1}$ . Energetically, inclusion of these A states is plausible given the  $0.20 \pm 0.05\text{ eV}$  experimental exothermicity of the  $\text{Ru}^+$  + oxirane reaction. A more reasonable solution is to include both  $^2\text{A}_1/^2\text{A}_2$  states and the  $^4\text{B}_1/^4\text{B}_2$  states. In both the EOM and CCSD computations, such a combination appears satisfactory to explain the observed spectral substructure, although a minor point of criticism is predicted intensity below  $500\text{ cm}^{-1}$




Fig. 3 (a) Experimental IRMPD spectrum of  $[\text{Ru,C,2H}]^+$ , the black trace is the average of four scans, the red trace is a single scan; (b)–(f) EOM-SF-CCSD/cc-pVTZ-PP calculated IR spectra of different  $[\text{Ru,C,2H}]^+$  states and isomers with the harmonic vibrations in red and the rovibrational envelopes in black accompanied by geometric structure, relative energy, and electronic state.

(originating from the  $^4\text{B}_1/^4\text{B}_2$  states), in contrast to the experimental spectrum (Fig. S5, ESI†). It seems likely that an enhanced contribution of the ground state  $\text{HRuCH}^+$  ( $^2\text{A}'$ ) would provide an even better reproduction of the experimental bands observed and also suppress the tail to the red of  $500\text{ cm}^{-1}$ . A final, unambiguous assignment of this spectrum remains challenging because of the rotational widths and possibility of multiple-photon effects.

### IR spectroscopy of $\text{RhCH}_2^+$

Upon photofragmentation of  $[\text{Rh,C,2H}]^+$  ( $m/z\ 117$ ),  $\text{RhC}^+$  was the only fragment observed, as shown in Fig. S2b (ES), consistent with the thermochemistry in Table 1. The experimental IRMPD spectrum of  $[\text{Rh,C,2H}]^+$  is shown in Fig. 4(a). The spectra were recorded in cell 4 (black trace, lowest IR fluence) and cell 1 (red trace, higher IR fluence). In contrast to the more poorly resolved, one-band-dominated spectrum for Ru, we observed five distinct IR bands: a triad of three strong bands at  $774$ ,  $904$ , and  $1029\text{ cm}^{-1}$ , with a relatively well-resolved satellite band of medium intensity at  $1269\text{ cm}^{-1}$  and a clear, relatively intense band at  $1783\text{ cm}^{-1}$ . The five experimental bands for the rhodium species are more than the three bands observed for the ruthenium species, which could potentially






Fig. 4 (a) Experimental IRMPD spectrum of  $[\text{Rh,C,2H}]^+$ , the black trace is recorded in FTICR cell 4 (low fluence), the red trace in FTICR cell 1 (high fluence); (b)–(f) B3LYP/def2-TZVPPD calculated IR spectra of different  $[\text{Rh,C,2H}]^+$  states and isomers with the harmonic vibrations in black and the rovibrational envelopes in grey accompanied by geometric structure, relative energy, and electronic state.

indicate the formation of multiple species. Also, the narrow width of the band observed at  $904\text{ cm}^{-1}$  is striking as it is much sharper than the bands observed for the Ru species. The structure observed under the lower fluence conditions (black trace) slightly broadens when using a significantly increased laser fluence (red trace) but remains clearly distinguishable, which is at odds with the Ru species where only drastic broadening was observed at higher fluence. In addition, the rhodium species has experimental bands at much higher frequencies than found for  $[\text{Ru,C,2H}]^+$ .

Fig. 4(b)–(f) shows the B3LYP-simulated IR spectra of different  $[\text{Rh,C,2H}]^+$  isomers and states. The  $^1\text{A}_1$  ground state has the symmetric rhodium carbene geometry with  $\text{C}_{2v}$  symmetry and a valence MO occupation of  $(1a_1)^2(1b_1)^2(2a_1)^2(1a_2)^2(3a_1)^0(1b_2)^2$ . This configuration is markedly different from the Ru case, as this species has a ground state with multiple unpaired electrons on the metal center, including an electron in the

$3a_1$  (5s-like) orbital. In contrast, the  $\text{RhCH}_2^+$  GS clearly has no unpaired electrons nor any electrons in the 5s orbital. Identification of the true ground state was complicated by finding another  $^1\text{A}_1$  state that differed from that shown because the  $2a_1$  ( $4d_{x^2-y^2}$ ) and  $1a_2$  ( $4d_{xy}$ ) orbitals were mixed, leading to a lower energy by  $0.020\text{ eV}$ . This outcome is an artifact of the B3LYP calculation, as demonstrated by the EOM calculations described below.

The lowest-energy triplet species for  $[\text{Rh,C,2H}]^+$  according to the B3LYP method is a symmetric carbene with a  $^3\text{A}_2$  state and lies only  $0.18\text{ eV}$  above the GS. This  $^3\text{A}_2$  state has one electron in the  $4d_{xy}$  ( $1a_2$ ) orbital, two electrons in the  $4d_{x^2-y^2}$  ( $2a_1$ ) orbital, and one electron in the 5s ( $3a_1$ ) orbital. A  $^3\text{A}_1$  state, slightly higher in energy at  $0.20\text{ eV}$  above the GS, has two electrons in the  $4d_{xy}$  orbital, one electron in the  $4d_{x^2-y^2}$  orbital, and one electron in the 5s orbital. We find two singlet carbene states with 5s occupancy at  $0.65\text{ eV}$  ( $^1\text{A}_2$ ) and  $0.66\text{ eV}$  ( $^1\text{A}_1$ ), Table S1 (ESI<sup>†</sup>), and the lowest-energy quintet at  $1.78\text{ eV}$  above the GS. Hydrido rhodium carbyne structures are found at  $0.83\text{ eV}$  ( $^1\text{A}'$ ) and  $1.53\text{ eV}$  ( $^3\text{A}'$ ) above the GS. The lowest-energy dihydrido rhodium carbide ( $^3\text{A}''$ ) lies  $3.42\text{ eV}$  above the GS, while the lowest energy dihydrogen rhodium carbide ( $^1\text{A}'$ ) is  $0.79\text{ eV}$  above the GS. The energetically preferred geometry is thus clearly a symmetric carbene.

The calculated IR spectra of the  $^1\text{A}_1$  and  $^3\text{A}_2$  states exhibit moderately intense bands for the Rh–C stretch (a narrow a-type transition,  $13$  and  $21\text{ km mol}^{-1}$ ) and strong bands for the out-of-plane  $\text{CH}_2$  wag (a broad b-type transition,  $57$  and  $89\text{ km mol}^{-1}$ ), whereas the in-plane  $\text{CH}_2$  rock and scissors modes are relatively weak with intensities  $<10\text{ km mol}^{-1}$ . The two states differ mainly in the frequency of the Rh–C stretch vibration, which is  $793\text{ cm}^{-1}$  for the  $^1\text{A}_1$  GS and  $674\text{ cm}^{-1}$  for the  $^3\text{A}_2$  state, Table 2. Both states have similar  $\text{CH}_2$  wag frequencies of  $964$  and  $942\text{ cm}^{-1}$ , respectively. In comparison to the Ru carbene cation, the Rh–C stretch vibration is calculated to have a significantly higher IR intensity. The three intense bands calculated at  $674$ ,  $793$ , and  $964/942\text{ cm}^{-1}$  could conceivably match the three main peaks observed experimentally, but they are systematically shifted to lower frequencies. The band observed at  $1269\text{ cm}^{-1}$  could be associated with the  $\text{RhCH}_2^+$  scissor vibration, but conceivably has components of overtones of the in-plane  $\text{CH}_2$  rock ( $606/646\text{ cm}^{-1}$ ). Likewise, the band observed at  $1783\text{ cm}^{-1}$  seems likely to be either a combination band or an overtone, as found earlier for a similar system.<sup>19</sup> An anharmonic frequency calculation, using the second-order vibrational perturbation theory (VPT2) method<sup>67,68</sup> and the same B3LYP/def2-TZVPPD potential energy surface, places the  $\text{CH}_2$  wagging overtone at  $1892\text{ cm}^{-1}$  with an intensity of  $1.3\text{ km mol}^{-1}$ , and the combination band of the  $\text{CH}_2$  rocking and scissoring modes at  $1918\text{ cm}^{-1}$  with a similar intensity. The strong intensity with which this band is experimentally found is remarkable, but it could be attributed to a higher IRMPD efficiency for higher photon energies.

Fig. 4(f) shows the spectrum calculated for the hydrido carbyne  $\text{HRhCH}^+$  species, which has a  $^1\text{A}'$  ground state. This species is found at fairly high energy,  $0.83\text{ eV}$  above  $\text{RhCH}_2^+$  ( $^1\text{A}_1$ ), which would make its formation unlikely. Further, it





exhibits two strong bands at 548 and 736  $\text{cm}^{-1}$  that do not match the experimental spectrum. From a spectroscopic point of view, the possibility that this species has a small population cannot be eliminated, in particular because the predicted low-frequency band could explain the small intensity near 500  $\text{cm}^{-1}$  observed in the high-intensity (red) IR spectrum. However, it is clear that the majority of the spectrum must be associated with low-energy carbene species. This conclusion is even more evident from the more advanced theoretical results found in the next section.

### EOM-SF-CCSD results for $\text{RhCH}_2^+$

Because of the unusual spectrum for the  $[\text{Rh,C,2H}]^+$  species and the difficulty in assigning the ground state computed with the B3LYP level of theory, we also explored more advanced EOM-SF-CCSD calculations. The potential energy curves along the Rh–C coordinate, shown in Fig. 5, indicate a manifold of electronic states for the carbene species with many curve crossings (with relative energies at the equilibrium Rh–C distances listed in Table 3). In particular, the  $^1\text{A}_1$  ground-state potential crosses the  $^3\text{A}_2$  potential near 1.9 Å, which also nearly coincides with the minimum in the latter state. Furthermore, the ground state also exhibits an avoided crossing with the next-lowest  $\text{A}_1$  state ( $^3\text{A}_1$ ) near 2.1 Å. Such interactions are likely to explain why the single-reference calculations encountered substantial difficulties in establishing the structures and electronic state of this species. Even the ground  $^1\text{A}_1$  state exhibited notable open-shell singlet character in these EOM-SF computations, with Mulliken spin populations (using the EOM-SF-CCSD wavefunctions) of  $-0.38$  and  $+0.38$  on the Rh and  $\text{CH}_2$  subunits, respectively. This


**Table 3** Energies of  $[\text{Ru,C,2H}]^+$  and  $[\text{Rh,C,2H}]^+$  species calculated at the EOM-SF-CCSD/cc-pVTZ-PP level of theory. EOM-SF-CCSD(T) energies are provided in square brackets

Structure	State	$E_{\text{rel}}$ (eV)	$E_{\text{rel}} + \text{ZPE}$ (eV)	$\langle S^2 \rangle$	
$\text{RuCH}_2^+$	$^4\text{B}_1$	0.000	0.000 [0.005]	4.098	
	$^4\text{B}_2$	0.001	0.001 [0.006]	4.101	
	$^2\text{A}_2$	0.010	0.012 [0.006]	0.967	
	$^2\text{A}_1$	0.021	0.036 [0.029]	0.938	
	$^4\text{A}_2$	1.036	1.018 [1.013]	2.835 <sup>a</sup>	
$\text{HRuCH}^+$	$^2\text{A}'$	0.119	0.072 [0.000]	0.967	
$[\text{Rh,C,2H}]^+$	$^1\text{A}_1$	0.000	0.000 [0.000]	0.244	
	$^3\text{A}_2$	0.296	0.295 [0.312]	2.251	
	$(\text{H}_2)\text{RhC}^+ (\text{C}_s)$	$^1\text{A}_1$	0.795	0.636 [0.570]	0.149
	$\text{RhCH}_2^+$	$^3\text{A}_1$	0.833	0.833 [0.764]	1.556 <sup>a</sup>
	$\text{HRhCH}^+$	$^1\text{A}'$	0.947	0.895 [0.785]	0.256
	$\text{RhCH}_2^+$	$^3\text{B}_2$	1.271	1.242 [1.185]	1.355 <sup>a</sup>
	$\text{RhCH}_2^+$	$^3\text{B}_2$	1.358	1.355 [1.272]	1.137 <sup>b</sup>
	$\text{RhCH}_2^+$	$^1\text{A}_2$	1.594	1.598 [1.569]	0.277
	$\text{RhCH}_2^+$	$^3\text{B}_1$	1.761	— <sup>c</sup>	1.546 <sup>a</sup>
	$\text{HRhCH}^+$	$^3\text{A}'$	2.080	2.012 [1.958]	2.011
	$\text{RhCH}_2^+$	$^3\text{B}_2$	2.430	2.392 [2.315]	1.493 <sup>a</sup>
$(\text{H}_2)\text{RhC}^+ (\text{C}_s)$	$^3\text{A}'$	3.189	3.020 [2.878]	1.144 <sup>a</sup>	

<sup>a</sup> Significant remnant spin contamination in EOM-SF-CCSD state. <sup>b</sup> Species corresponds to a transition state in  $\text{C}_{2v}$  symmetry. <sup>c</sup> Finite-difference frequency alternated electronic states; no ZPE correction available.

outcome is consistent with the spin density depicted in Fig. S6 of the ESI.† Notably, although previous computational studies of  $\text{RhCH}_2^+$  correctly identified the  $^1\text{A}_1$  ground state and  $^3\text{A}_2$  first excited state,<sup>43,44</sup> the role of multireference character in the ground state has not been fully appreciated, particularly as the  $\text{Rh}^+-\text{CH}_2$  bond is stretched.

The EOM results identified eleven distinct species (some of which are excited states of a given symmetry): several  $\text{RhCH}_2^+$  carbenes along with  $(\text{H}_2)\text{RhC}^+$  ( $^1\text{A}_1$ ) and  $\text{HRhCH}^+$  ( $^1\text{A}'$  and  $^1,^3\text{A}'$ ) structures (Table 3). On the basis of these calculations, the  $^1\text{A}_1$  and  $^3\text{A}_2$  carbenes seem to be the likely candidates for formation in the experiments conducted here. The vibrational spectra of these two states are similar to those found at the B3LYP level, although the Rh–C stretch intensity is notably enhanced, particularly in the ground-state singlet, as listed in Table 2. The combination of these two species results in a spectrum that reproduces the three strong bands observed experimentally, as shown in Fig. 6 where the frequencies have been scaled by a factor 1.09, resulting in a blue-shift of about 90  $\text{cm}^{-1}$ . (The scaling factor of 1.09 is chosen to best match the experimental spectrum. The comparison to the unshifted spectrum is shown in Fig. S7 of the ESI.†) The spectrum computed for the  $^3\text{A}_2$  carbene may even rationalize the larger spectral width of the band observed at 774  $\text{cm}^{-1}$  compared to the 904  $\text{cm}^{-1}$  band. This is because the Rh–C stretch for the  $^1\text{A}_1$  species is isolated whereas the same mode for the  $^3\text{A}_2$  state, at 668  $\text{cm}^{-1}$ , nearly overlaps with the more intense  $\text{CH}_2$  rocking mode. Cooperative excitation of these bands could lead to the broadening observed, especially in the high-fluence spectrum. The assignment of the additional bands observed experimentally at higher wavenumbers remains inconclusive but seems likely to be the overtones identified above, similar to the overtone band previously found in the spectrum of  $\text{PtCH}_2^+$ .<sup>18,19</sup>



**Fig. 5** Constrained potential energy curves along the metal–carbon distance in  $\text{RhCH}_2^+$  in  $\text{C}_{2v}$  symmetry, as calculated at the EOM-SF-CCSD/cc-pVTZ-PP level of theory. Solid (dashed) lines depict the lowest-energy (first excited) state of each symmetry. The structure was optimized in the lowest state of  $\text{A}_1$  symmetry, and all other state energies represent vertical excitations at the same structure. Spin and symmetry labels for each state, corresponding to the assignments at the  $^1\text{A}_1$  minimum geometry, are provided for each curve. [The symmetry labels for the  $\text{B}_1$  and  $\text{B}_2$  states are reversed from the Q-Chem outputs, in order to align with the symmetry-axis conventions of Gaussian computations in the remainder of this work.]






Fig. 6 (a) Experimental IRMPD spectrum of  $[\text{Rh,C,2H}]^+$ ; (b) harmonic stick spectrum (in red) of the  $\text{RhCH}_2^+$  ( $^1\text{A}_1$ ) structure at the EOM-SF-CCSD level and rovibrational simulations (in black) with frequencies scaled by 1.09; (c) same as (b) for  $\text{RhCH}_2^+$  ( $^3\text{A}_2$ ).

Although a systematic methodology comparison is not the focus of the present study, we note anecdotally that the more recently developed, range-separated hybrid functional  $\omega\text{B97M-V}^{69}$  was found to exhibit harmonic frequencies and intensities that are notably more closely aligned with the EOM-based spectra (Fig. S8 of the ESI<sup>†</sup>). This functional may serve as a more accurate, cost-efficient method for future studies of this complex, although the state selectivity of the EOM methods provides considerably more control over the targeted electronic states.

Given the relative success of this alternative functional for  $[\text{Rh,C,2H}]^+$ , the spectra of  $[\text{Ru,C,2H}]^+$  were also reexamined with  $\omega\text{B97M-V/cc-pVTZ-PP}$ . The density functional encouragingly exhibited harmonic spectra for  $[\text{Ru,C,2H}]^+$  (Fig. S9 of the ESI<sup>†</sup>) that were quite similar to those computed by CCSD (Fig. 2). However, the energy of the  $\text{HRuCH}^+$  ( $^2\text{A}'$ ) state was computed to be more than 0.6 eV above the ground  $^4\text{B}_2$  state, in stark contrast to the results of CCSD(T) and EOM-SF-CCSD(fT) where it is the ground state structure. Simultaneously capturing both the overall energy landscape and the nuanced shape of the potential energy surface near equilibrium evidently remains a daunting challenge for density functional theory for these ionic transition-metal complexes.

#### DFT performance for prediction of M–C stretch vibrational intensities

Judging from the comparison between the B3LYP-, CCSD-, and EOM-SF-CCSD-calculated spectra for  $\text{RuCH}_2^+$  (Table 2), the

drastic underestimation of the IR intensities identified for the  $\text{FeCH}_2^+$  and  $\text{CoCH}_2^+$  species<sup>45</sup> remains evident for this system. For the quartet spin carbenes, B3LYP predicts intensities for the Ru–CH<sub>2</sub> stretch vibration that are an order of magnitude smaller than the CCSD and EOM-SF-CCSD calculations. Only for the  $^2\text{A}_2$  state does B3LYP predict an intensity for this stretch that is comparable to the EOM-SF-CCSD result. However, the concluded presence of the  $\text{HRuCH}^+$  species makes it complex to test this observation on the basis of comparison with experiment. Because the latter species is identified from dominant bands associated with bending modes, which are less sensitive to potential underestimations of charge separation, the  $\text{RuCH}_2^+$  species cannot form a good theoretical benchmark.

On the other hand, comparison of the computed spectra for the  $^1\text{A}_1$  and  $^3\text{A}_2$   $\text{RhCH}_2^+$  species suggests that the single-reference B3LYP method does not underperform compared to the CCSD/EOM approaches for frequencies (Table 2). Comparison with experiment would require an anomalous scaling factor for the B3LYP spectra compared with previous work. For IR intensities, the enhanced intensities of the Rh–C stretch found by EOM-SF-CCSD method helped solidify the assignment of the spectra (in particular, the bands at 774 and 904  $\text{cm}^{-1}$ ). Of course, the vicinity of so many low-lying electronic states clearly complicates finding the experimental ground states.

#### Group 8 and 9 trends: carbene versus hydrido carbyne structures

In previous work, we have presented the experimental IRMPD spectra of  $[\text{Os,C,2H}]^+$  and  $[\text{Ir,C,2H}]^+$ .<sup>18–20,22</sup> We concluded that the osmium species had the hydrido carbyne  $\text{HOSCH}^+$  structure, as did the iridium species, which exhibited small amounts of a higher-energy, symmetric carbene, as well. If we put the current assignments for Ru and Rh in perspective with these previous findings, we note that both 5d elements prefer the formation of a hydrido carbyne structure, whereas the 4d Ru forms both this and a carbene species and Rh yields only the carbene. As shown in Table 4, the 5d hydrido carbynes are energetically preferred, but more strongly so for the group 8 Os case. This observation is consistent with the assignment of  $\text{HOSCH}^+$  ( $^2\text{A}'$ ), whereas part of the product for iridium (group 9) is assigned to a carbene structure (0.3 or 0.7 eV higher in energy). This trend appears to hold for 4d elements, where the  $\text{RuCH}_2^+$  ( $^4\text{B}_2$ ) carbene is competitive with the  $^2\text{A}'$   $\text{HRuCH}^+$  hydrido carbyne, whereas  $\text{HRhCH}^+$  ( $^1\text{A}'$ ) is significantly higher in energy than  $\text{RhCH}_2^+$  ( $^1\text{A}_1$ ). Intrinsically, the 5d TMs have stronger bonds to C and H than the 4d TMs, e.g., for group 9, see Fig. 4 in ref. 13. (For group 8, see ref. 16, 37, 70 and 65).

Table 4 Theoretical energies (eV) of  $\text{HMCH}^+$  species relative to the most stable  $\text{MCH}_2^+$  carbene

M	B3LYP	CCSD(T)//CCSD
Ru	0.03	−0.005
Rh	0.80	0.76
Os	−0.50	−0.82
Ir	−0.30	−0.69






Fig. 7 Potential energy surfaces of  $[M,C,2H]^+$  along the C–M–H angle calculated at the uB3LYP/def2-TZVPPD level of theory with steps of  $1^\circ$ . Open diamonds indicate calculated structures that successfully converged to true minima at the indicated C–M–H angle. Solid lines indicate  $A''$  states while  $A'$  states are indicated by dashed lines. Doublet spin surfaces are depicted in green, quartet in black, singlet in blue, and triplet in red. All energy values are relative to the GS symmetric carbene structures.

Because  $HMCH^+$  has four covalent bonds to the metal whereas  $MCH_2^+$  only has two covalent M–C bonds, the more highly coordinated hydrido carbyne structure becomes more favorable for 5d relative to 4d TMs.

The hydrido carbyne structures for 4d and 5d elements are found on the same or a lower spin surface as that of the lowest-energy carbene structure. To investigate whether we might have missed hydrido carbyne structures, the potential energy surfaces (PESs) connecting the  $MCH_2^+$  and  $HMCH^+$  structures were calculated for  $M = Ru$  and  $Rh$  (Fig. 7) with B3LYP. PESs were generated by starting from a symmetric carbene structure and systematically increasing the C–M–H angle while the rest of the molecule was allowed to relax. The PES scans were generated for both the  $A'$  and  $A''$  configurations in dashed and solid lines, respectively. The PESs for  $Ru$  and  $Rh$  show clear minima and appreciable barriers for H migration from the metal to carbon. Here again, the higher stability of carbyne structures for  $Ru$  in comparison to  $Rh$  is clearly visible.

## Conclusion

Reacting the 4d group 8 and 9 transition metal ions  $M^+$  ( $M = Ru$  and  $Rh$ ) with oxirane in a room temperature ion trap led to the

formation of  $[M,C,2H]^+$ . These species were mass-isolated in the FTICR mass spectrometer coupled to the infrared intracavity free-electron laser FELICE, where they were spectroscopically characterized. The IRMPD spectrum of  $[Ru,C,2H]^+$  contains one broad band around  $700\text{ cm}^{-1}$ , whereas the IRMPD spectrum of  $[Rh,C,2H]^+$  is characterized by much sharper, distinct bands. On the basis of B3LYP and CCSD(T)//CCSD energetics, the  $[Rh,C,2H]^+$  species is expected to be a carbene, and for  $[Ru,C,2H]^+$ , a competitive hydrido carbyne structure is also found. Comparison of the experimental IRMPD spectra with the predictions from B3LYP calculations offers moderate agreement. Calculations at the CCSD and EOM-SF-CCSD levels offer a modest improvement, but also show how sensitive an experimental constraint the spectroscopic fingerprint provides for these electronically complex systems. These comparisons suggest the presence of the  $HRuCH^+$  species, with contributions from  $RuCH_2^+$  carbenes. The case of  $[Rh,C,2H]^+$  offers a similar complexity. Despite the much more structured experimental IR spectrum, assignment based on either B3LYP or conventional CCSD simulations is difficult. Because population of the energetically close singlet and triplet spin surfaces could significantly influence the behavior of this species and because of moderately strong state mixing within the singlet manifold, EOM-SF-CCSD calculations were performed. These computational results suggest that a combination of the (partially open-shell)  $^1A_1$  and  $^3A_2$  carbenes can explain the complex structure of the  $[Rh,C,2H]^+$  spectrum. Notably, including the multi-reference character of this species was critical in properly assigning the spectrum, but a modest shift in the calculated harmonic frequencies was still required to match experiment.

## Conflicts of interest

There are no conflicts of interest to declare.

## Acknowledgements

This work was funded by NWO's Materials for Sustainability program (grant no. 739.017.008). We gratefully acknowledge the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO) for the support of the FELIX Laboratory and for CPU time on the Dutch National Supercomputers Cartesius and Snellius (project number 2021.055). Additional financial support was provided by the National Science Foundation (grant no. CHE-2313553) and by a generous grant of computer time from the Center for High-Performance Computing at the University of Utah. RPS acknowledges funding from the U.S. Department of Energy under Grant DE-SC0019405 and helpful conversations regarding EOM methods with Prof. Anna Krylov.

## References

- 1 R. H. Crabtree, Organometallic alkane CH activation, *J. Organomet. Chem.*, 2004, **689**, 4083–4091.



- 2 K. L. Stone and A. S. Borovik, Lessons from Nature: Unraveling Biological C–H Bond Activation, *Curr. Opin. Chem. Biol.*, 2009, **13**, 114–118.
- 3 D. K. Böhme and H. Schwarz, Gas-Phase Catalysis by Atomic and Cluster Metal Ions: The Ultimate Single-Site Catalysts, *Angew. Chem., Int. Ed.*, 2005, **44**, 2336–2354.
- 4 J. Roithová and D. Schröder, Selective Activation of Alkanes by Gas-Phase Metal Ions, *Chem. Rev.*, 2010, **110**, 1170–1211.
- 5 J. Sauer and H. J. Freund, Models in Catalysis, *Catal. Lett.*, 2015, **145**, 109–125.
- 6 S. W. Buckner, T. J. MacMahon, G. D. Byrd and B. S. Freiser, Gas-Phase Reactions of Nb<sup>+</sup> and Ta<sup>+</sup> with Alkanes and Alkenes. C–H Bond Activation and Ligand-Coupling Mechanisms, *Inorg. Chem.*, 1989, **28**, 3511–3518.
- 7 K. K. Irikura and J. L. Beauchamp, Osmium Tetroxide and Its Fragment Ions in the Gas Phase: Reactivity with Hydrocarbons and Small Molecules, *J. Am. Chem. Soc.*, 1989, **111**, 75–85.
- 8 K. K. Irikura and J. L. Beauchamp, Electronic Structure Considerations for Methane Activation by Third-row Transition-metal Ions, *J. Phys. Chem.*, 1991, **95**, 8344–8351.
- 9 K. K. Irikura and J. L. Beauchamp, Methane Oligomerization in the Gas Phase by Third-Row Transition-Metal Ions, *J. Am. Chem. Soc.*, 1991, **113**, 2769–2770.
- 10 U. Achatz, M. Beyer, S. Joos, B. S. Fox, G. Niedner-Schatteburg and V. E. Bondybey, The Platinum Hydrido-Methyl Complex: A Frozen Reaction Intermediate?, *J. Phys. Chem. A*, 1999, **103**, 8200–8206.
- 11 X.-G. Zhang, R. Liyanage and P. B. Armentrout, The Potential Energy Surface for Activation of Methane by Pt<sup>+</sup>: A Detailed Guided-Ion Beam Study, *J. Am. Chem. Soc.*, 2001, **123**, 5563–5575.
- 12 P. B. Armentrout, S. Shin and R. Liyanage, Guided-Ion Beam and Theoretical Study of the Potential Energy Surface for Activation of Methane by W<sup>+</sup>, *J. Phys. Chem. A*, 2006, **110**, 1242–1260.
- 13 F.-X. Li, X.-G. Zhang and P. B. Armentrout, The Most Reactive Third-row Transition Metal: Guided Ion Beam and Theoretical Studies of the Activation of Methane by Ir<sup>+</sup>, *Int. J. Mass Spectrom.*, 2006, **255–256**, 279–300.
- 14 L. G. Parke, C. S. Hinton and P. B. Armentrout, Experimental and Theoretical Studies of the Activation of Methane by Ta<sup>+</sup> and the Bond Energies of Ta<sup>+</sup>–CH<sub>x</sub> (x = 1–3), *J. Phys. Chem. C*, 2007, **111**, 17773–17787.
- 15 A. Shayesteh, V. V. Lavrov, G. K. Koyanagi and D. K. Böhme, Reactions of Atomic Cations with Methane: Gas Phase Room-Temperature Kinetics and Periodicities in Reactivity, *J. Phys. Chem. A*, 2009, **113**, 5602–5611.
- 16 P. B. Armentrout, L. Parke, C. Hinton and M. Citir, Activation of Methane by Os<sup>+</sup>: Guided Ion Beam and Theoretical Studies, *ChemPlusChem*, 2013, **78**, 1157–1173.
- 17 P. B. Armentrout, Methane Activation by 5d Transition Metals: Energetics, Mechanisms, and Periodic Trends, *Chem. – Eur. J.*, 2017, **23**, 10–18.
- 18 V. J. F. Lapoutre, B. Redlich, A. F. G. van der Meer, J. Oomens, J. M. Bakker, A. Sweeney, A. Mookherjee and P. B. Armentrout, Structures of the Dehydrogenation Products of Methane Activation by 5d Transition Metal Cations, *J. Phys. Chem. A*, 2013, **117**, 4115–4126.
- 19 C. J. Owen, G. C. Boles, V. Chernyy, J. M. Bakker and P. B. Armentrout, Structures of the Dehydrogenation Products of Methane Activation by 5d Transition Metal Cations Revisited: Deuterium Labeling and Rotational Contours, *J. Chem. Phys.*, 2018, **148**, 044307.
- 20 J. M. Bakker, C. J. Owen, S. W. Nooteboom, O. V. Lushchikova and P. B. Armentrout, Structural characterization of [M,C,2H]<sup>+</sup> products formed by reaction of 5d metal cations Pt<sup>+</sup> and Ir<sup>+</sup> with ethylene oxide and Ta<sup>+</sup> with methane using messenger spectroscopy, *J. Mol. Spectrosc.*, 2021, **378**, 111472.
- 21 F. J. Wensink, N. Roos, J. M. Bakker and P. B. Armentrout, C–H Bond Activation and C–C Coupling of Methane on a Single Cationic Platinum Center: A Spectroscopic and Theoretical Study, *Inorg. Chem.*, 2022, **61**, 11252–11260.
- 22 P. B. Armentrout, S. Kuijpers, O. V. Lushchikova, R. L. Hightower, G. C. Boles and J. M. Bakker, Spectroscopic Identification of the Carbyne Hydride Structure of the Dehydrogenation Product of Methane Activation by Osmium Cations, *J. Am. Soc. Mass Spectrom.*, 2018, **29**, 1781–1791.
- 23 J. K. Perry, G. Ohanessian and W. A. Goddard, Mechanism and Energetics for Dehydrogenation of Methane by Gaseous Iridium Ions, *Organometallics*, 1994, **13**, 1870–1877.
- 24 L. F. Halle, P. B. Armentrout and J. L. Beauchamp, Formation of Chromium Carbene Ions by Reaction of Electronically Excited Chromium Ions with Methane in the Gas Phase, *J. Am. Chem. Soc.*, 1981, **103**, 962–963.
- 25 N. Aristov and P. B. Armentrout, Methane Activation by V<sup>+</sup>: Electronic and Translational Energy Dependence, *J. Phys. Chem.*, 1987, **91**, 6178–6188.
- 26 R. H. Schultz, J. L. Elkind and P. B. Armentrout, Electronic Effects in C–H and C–C Bond Activation: State-specific Reactions of Fe<sup>+</sup>(<sup>6</sup>D, <sup>4</sup>F) with Methane, Ethane and Propane, *J. Am. Chem. Soc.*, 1988, **110**, 411–423.
- 27 L. S. Sunderlin and P. B. Armentrout, Methane Activation by Ti<sup>+</sup>: Electronic and Translational Energy Dependence, *J. Phys. Chem.*, 1988, **92**, 1209–1219.
- 28 R. Georgiadis and P. B. Armentrout, Translational and Electronic Energy Dependence of Chromium Ion Reactions with Methane, *J. Phys. Chem.*, 1988, **92**, 7067–7074.
- 29 L. S. Sunderlin and P. B. Armentrout, Periodic Trends in Chemical Reactivity: Reactions of Sc<sup>+</sup>, Y<sup>+</sup>, La<sup>+</sup>, and Lu<sup>+</sup> with Methane and Ethane, *J. Am. Chem. Soc.*, 1989, **111**, 3845–3855.
- 30 R. Georgiadis and P. B. Armentrout, Translational and Electronic Energy Dependence of the Reaction of Mn<sup>+</sup> with Ethane, *Int. J. Mass Spectrom. Ion Processes*, 1989, **91**, 123–133.
- 31 C. L. Haynes, Y.-M. Chen and P. B. Armentrout, The Potential Energy Surface for Activation of Methane by Co<sup>+</sup>: An Experimental Study, *J. Phys. Chem.*, 1995, **99**, 9110–9117.
- 32 Y.-M. Chen and P. B. Armentrout, Activation of Methane by Gas-Phase Rh<sup>+</sup>, *J. Phys. Chem.*, 1995, **99**, 10775–10779.



- 33 Y.-M. Chen, M. R. Sievers and P. B. Armentrout, Activation of CH<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, C<sub>3</sub>H<sub>8</sub>, and c-C<sub>3</sub>H<sub>6</sub> by Gas-phase Pd<sup>+</sup> and the Thermochemistry of Pd–ligand Complexes, *Int. J. Mass Spectrom. Ion Processes*, 1997, **167–168**, 195–212.
- 34 M. R. Sievers, Y.-M. Chen, C. L. Haynes and P. B. Armentrout, Activation of CH<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, and C<sub>3</sub>H<sub>8</sub> by Gas-Phase Nb<sup>+</sup> and the Thermochemistry of Nb–ligand Complexes, *Int. J. Mass Spectrom.*, 2000, **195–196**, 149–170.
- 35 F. Liu, X.-G. Zhang and P. B. Armentrout, Activation of CH<sub>4</sub> by Gas-phase Ni<sup>+</sup> and the Thermochemistry of Ni Ligand Complexes, *Phys. Chem. Chem. Phys.*, 2005, **7**, 1054–1064.
- 36 P. B. Armentrout, Activation of CH<sub>4</sub> by Gas-phase Mo<sup>+</sup> and the Thermochemistry of Mo–ligand Complexes, *J. Phys. Chem. A*, 2006, **110**, 8327–8338.
- 37 P. B. Armentrout and Y.-M. Chen, Activation of Methane by Ru<sup>+</sup>: Experimental and Theoretical Studies of the Thermochemistry and Mechanism, *Int. J. Mass Spectrom.*, 2016, **413**, 135–149.
- 38 P. B. Armentrout and M. R. Sievers, Activation of CH<sub>4</sub> by Gas-phase Zr<sup>+</sup> and the Thermochemistry of Zr ligand Complexes, *J. Phys. Chem. A*, 2003, **107**, 4396–4406.
- 39 P. B. Armentrout, L. S. Sunderlin and E. R. Fisher, Intrinsic Transition Metal–Carbon Double Bond Dissociation Energies: Periodic Trends in M<sup>+</sup>–CH<sub>2</sub> Bond Strengths, *Inorg. Chem.*, 1989, **28**, 4436–4437.
- 40 B. Ruscic and D. H. Bross, Active Thermochemical Tables (ATcT) values based on ver. 1.130 of the Thermochemical Network, available at ATcT.anl.gov (accessed 2/12/24).
- 41 Y.-M. Chen and P. B. Armentrout, Activation of C<sub>2</sub>H<sub>6</sub>, C<sub>3</sub>H<sub>8</sub>, and c-C<sub>3</sub>H<sub>6</sub> by Gas-phase Rh<sup>+</sup> and the Thermochemistry of Rh–ligand Complexes, *J. Am. Chem. Soc.*, 1995, **117**, 9291–9304.
- 42 D. B. Jacobson and B. S. Freiser, Transition-metal-carbene chemistry. Structure, thermodynamics, and reactivity of rhodium methylidene ion (RhCH<sub>2</sub><sup>+</sup>) in the gas phase, *J. Am. Chem. Soc.*, 1985, **107**, 5870–5876.
- 43 D. G. Musaev, N. Koga and K. Morokuma, Ab initio molecular orbital study of the electronic and geometric structure of RhCH<sub>2</sub><sup>+</sup> and the reaction mechanism: RhCH<sub>2</sub><sup>+</sup> + H<sub>2</sub> → Rh<sup>+</sup> + CH<sub>4</sub>, *J. Phys. Chem.*, 1993, **97**, 4064–4075.
- 44 J. Westerberg and M. R. A. Blomberg, Methane Activation by Naked Rh<sup>+</sup> Atoms. A Theoretical Study, *J. Phys. Chem. A*, 1998, **102**, 7303–7307.
- 45 F. J. Wensink, C. E. Smink, J. M. Bakker and P. B. Armentrout, IR spectroscopic characterization of 3d transition metal carbene cations, FeCH<sub>2</sub><sup>+</sup> and CoCH<sub>2</sub><sup>+</sup>: A failure of DFT approaches, *Phys. Chem. Chem. Phys.*, 2024, DOI: **10.1039/D4CP00026A**.
- 46 F. J. Wensink, M. G. Münst, J. Heller, M. Ončák, J. M. Bakker and C. Van Der Linde, IR Multiple Photon Dissociation Spectroscopy of MO<sub>2</sub><sup>+</sup> (M = V, Nb, Ta), *J. Chem. Phys.*, 2020, **153**(17), 171101.
- 47 C. Berg, T. Schindler, G. Niedner-Schatteburg and V. E. Bondybey, Reactions of Simple Hydrocarbons with Nb<sub>n</sub><sup>+</sup>: Chemisorption and Physisorption on Ionized Niobium Clusters, *J. Chem. Phys.*, 1995, **102**, 4870–4884.
- 48 A. G. Marshall, C. L. Hendrickson and G. S. Jackson, Fourier Transform Ion Cyclotron Resonance Mass Spectrometry: A Primer, *Mass Spectrom. Rev.*, 1998, **17**, 1–35.
- 49 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson and H. Nakatsuji, *et al.*, *Gaussian 16 Rev. C.01*, Wallingford, CT, 2016.
- 50 A. D. Becke, Density-functional Thermochemistry. III. The Role of Exact Exchange, *J. Chem. Phys.*, 1993, **98**, 5648–5652.
- 51 C. Lee, W. Yang and R. G. Parr, Development of the Colle-Salvetti Correlation-Energy Formula into a Functional of the Electron Density, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 1988, **37**, 785–789.
- 52 F. Weigend and R. Ahlrichs, Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305, DOI: **10.1039/B508541A**.
- 53 W. L. Meerts and M. Schmitt, Application of Genetic Algorithms in Automated Assignments of High-Resolution Spectra, *Int. Rev. Phys. Chem.*, 2006, **25**, 353–406.
- 54 P. B. Armentrout, B. C. Stevenson, F. Yang, F. J. Wensink, O. V. Lushchikova and J. M. Bakker, Infrared Spectroscopy of Gold Carbene Cation (AuCH<sub>2</sub><sup>+</sup>): Covalent or Dative Bonding?, *J. Phys. Chem. A*, 2019, **123**, 8932–8941.
- 55 O. W. Wheeler, M. Salem, A. Gao, J. M. Bakker and P. B. Armentrout, Sequential Activation of Methane by Ir<sup>+</sup>: An IRMPD and Theoretical Investigation, *Int. J. Mass Spectrom.*, 2019, **435**, 78–92.
- 56 O. W. Wheeler, M. Salem, A. Gao, J. M. Bakker and P. B. Armentrout, Activation of C–H bonds in Pt<sup>+</sup> + xCH<sub>4</sub> Reactions, Where x = 1–4: Identification of the Platinum Dimethyl Cation, *J. Phys. Chem. A*, 2016, **120**, 6216–6227.
- 57 C. Möller and M. S. Plesset, Note on an Approximation Treatment for Many-Electron Systems, *Phys. Rev.*, 1934, **46**, 618–622.
- 58 R. J. Bartlett, Many-body Perturbation Theory and Coupled Cluster Theory for Electron Correlation in Molecules, *Ann. Rev. Phys. Chem.*, 1981, **32**, 359–401.
- 59 A. I. Krylov, Size-consistent wave functions for bond-breaking: the equation-of-motion spin-flip model, *Chem. Phys. Lett.*, 2001, **338**, 375–384.
- 60 A. I. Krylov, Spin-Flip Equation-of-Motion Coupled-Cluster Electronic Structure Method for a Description of Excited States, Bond Breaking, Diradicals, and Triradicals, *Acc. Chem. Res.*, 2006, **39**, 83–91.
- 61 K. A. Peterson, D. Figgen, M. Dolg and H. Stoll, Energy-consistent Relativistic Pseudopotentials and Correlation Consistent Basis Sets for the 4d Elements Y–Pd, *J. Chem. Phys.*, 2007, **126**, 124101.
- 62 T. H. Dunning, Gaussian basis sets for use in correlated molecular calculations. I. The atoms boron through neon and hydrogen, *J. Chem. Phys.*, 1989, **90**, 1007–1023.
- 63 P. U. Manohar and A. I. Krylov, A noniterative perturbative triples correction for the spin-flipping and spin-conserving



- equation-of-motion coupled-cluster methods with single and double substitutions, *J. Chem. Phys.*, 2008, **129**.
- 64 Y.-M. Chen, J. L. Elkind and P. B. Armentrout, Reactions of  $\text{Ru}^+$ ,  $\text{Rh}^+$ ,  $\text{Pd}^+$ , and  $\text{Ag}^+$  with  $\text{H}_2$ , HD and  $\text{D}_2$ , *J. Phys. Chem.*, 1995, **99**, 10438–10445.
- 65 P. B. Armentrout and I. Kretzschmar, Guided Ion Beam and Theoretical Studies of the Reaction of  $\text{Ru}^+$  with  $\text{CS}_2$  in the Gas-phase: Thermochemistry of  $\text{RuC}^+$ ,  $\text{RuS}^+$ , and  $\text{RuCS}^+$ , *Phys. Chem. Chem. Phys.*, 2010, **12**, 4078–4091.
- 66 E. A. Carter and W. A. Goddard, Bonding in Transition-Metal-Methylene Complexes. 2.  $(\text{RuCH}_2)^+$ , a Complex Exhibiting Low-Lying Methylidene-like and Carbene-like States, *J. Am. Chem. Soc.*, 1986, **108**, 2180–2191.
- 67 V. Barone, Anharmonic vibrational properties by a fully automated second-order perturbative approach, *J. Chem. Phys.*, 2005, **122**, 014108.
- 68 V. Barone, Vibrational zero-point energies and thermodynamic functions beyond the harmonic approximation, *J. Chem. Phys.*, 2004, **120**, 3059–3065.
- 69 N. Mardirossian and M. Head-Gordon, Mapping the genome of meta-generalized gradient approximation density functionals: The search for B97M-V, *J. Chem. Phys.*, 2015, **142**, 074111.
- 70 P. B. Armentrout and Y.-M. Chen, Activation of  $\text{C}_2\text{H}_6$ ,  $\text{C}_3\text{H}_8$ ,  $\text{HC}(\text{CH}_3)_3$ , and  $\text{c-C}_3\text{H}_6$  by Gas-Phase  $\text{Ru}^+$  and the Thermochemistry of Ru-ligand Complexes, *J. Am. Soc. Mass Spectrom.*, 1999, **10**, 821–839.

