

Cite this: *RSC Mechanochem.*, 2025, 2, 370

Conquering the impossible: mechanochemistry as a tool for tackling coordination chemistry challenges

Huanxin Zhang, Nathan Davison* and Erli Lu *

Organic solvents are ubiquitously used in synthetic coordination chemistry, but these solution-based synthetic methods have severe drawbacks, such as the incapability of isolating highly reactive species which react with the solvent molecules, less-controllable heterogeneous reactions when using insoluble substrate(s), and unfavourable solvent molecule coordination. In recent years, coordination chemists have started employing mechanochemical methods (e.g., ball mill) to overcome these drawbacks. Herein, we offer our perspective about how mechanochemical methods have enabled coordination chemists to achieve what would otherwise be impossible. It should be noted that, this perspective should not be treated as a comprehensive review of mechanochemistry in coordination chemistry, instead, a “stepping stone” aiming at inspiring further endeavours. Also, due to the research background of the authors, the selection of examples herein may appear biased towards main-group chemistry, which we feel necessary to remind our readers.

Received 8th January 2025

Accepted 14th March 2025

DOI: 10.1039/d5mr00005j

rsc.li/RSCMechanochem

1. Introduction

Organic solvents are used in almost all chemical reactions – there is no exception for coordination chemistry. Yet, it is not unusual that the organic solvents used give rise to undesired outcomes, such as product decomposition and formation of unfavoured by- or side-products. For example, tetrahydrofuran (THF), one of the most-used organic solvents, is well-known to undergo deprotonation and C–O bond cleavage, induced by strong organometallic Brønsted bases and/or nucleophiles.¹ While the ethereal solvents (including THF) are not inert towards C–H and C–O activations,² arene solvents, especially benzene and toluene, are also known to be activated *via* multiple pathways, such as toluene C(sp³)–H deprotonation,³ arene reduction,⁴ and transition-metal mediated dearomatization activation.⁵ Even the most inert aliphatic hydrocarbon solvents, such as the linear hexanes⁶ and *cyclo*-hexane,⁷ were reported to undergo C–H activation with highly reactive organometallic species.^{8,9} On one hand, these reactivities are a blessing as they open up new territories in small molecule activation. On the other hand, they are a curse which may hamper the isolation and characterisation of targeted highly reactive coordination complexes.

Another inherent challenge of the solution-based synthetic methods is the heterogeneity caused by poorly soluble reactants. A classic example is reductions using potassium graphite (KC₈), Na/

K alloy, K or Na mirror, or sodium over sodium chloride (Na/NaCl). The heterogeneity presents kinetic challenges: on the solution-solid surface (where the reaction takes place), the reactant: reductant stoichiometric ratio could differ substantially from the ideal ratio. To overcome the challenge, very fine highly dispersed reductants and vigorous stirring are usually needed, such as demonstrated in divalent lanthanide chemistry.¹⁰ A similar challenge presents in reactions involving zero-valent metals (as reductant in most cases), which are insoluble in organic solvents. The metal vapour deposition (MVD) method was used to tackle the challenge, such as in the synthesis of the classic samarocenes [(C₅Me₅)₂Sm(THF)₂]¹¹/[(C₅Me₄Et)₂Sm(THF)₂]¹¹ their non-solvated analogue [(C₅Me₅)₂Sm],¹² and large-scale synthesis of dysprosium and neodymium diiodides.¹³

A solvent-free synthetic method would be ideal for tackling these solvent-related challenges. Mechanochemistry, defined by IUPAC as “chemical reaction[s] that [are] induced by the direct absorption of mechanical energy”¹⁴ and regarded as the “fourth way” of synthetic chemistry alongside solvothermal, electrochemical, and photochemical methods, offers a prime candidate.¹⁵ Moreover, the unique energy transfer mode of mechanochemistry (e.g. impact forces for milling), opens exciting opportunities for the exploration of entirely new complexes, which are unable to be synthesised *via* traditional solution-state methods.

In this perspective, from a viewpoint of synthetic coordination chemists, we will categorise and analyse the unique advantages offered by mechanochemical methods, which have unlocked new horizons in synthetic coordination chemistry.¹⁶ In other words, instead of a comprehensive review of the

School of Chemistry, University of Birmingham. Edgbaston, Birmingham, B15 2TT, UK.
E-mail: n.davison@bham.ac.uk; e.lu@birmingham.ac.uk; Web: <https://www.birmingham.ac.uk/staff/profiles/chemistry/lu-erli>; https://x.com/erli_lu



applications of mechanochemical methods in synthetic coordination chemistry, this perspective focuses on the cases which “achieve the impossible”, *i.e.*, cases where mechanochemistry succeeds, while other methods fail. We believe that the capability of “achieving the impossible” is a unique value of mechanochemistry, beside the sustainability merits of being solvent-free and energy-efficient.

2. Isolating highly reactive species

2.1 Low-valent main-group complexes

The synthesis, structure, and reactivity studies of low-valent complexes, especially those of main-group elements (*s*-¹⁷ and *p*-blocks¹⁸), lanthanide,¹⁹ and actinide²⁰ metals have emerged as a hot topic of modern chemistry, not only because of their roles in expanding our knowledge boundaries, but also because of their potential in inert small molecule and chemical bond activations.²¹ Due to their rich electron density in the valence shell orbitals, these low-valent complexes usually feature high reactivity, which often lead to reactions with solvents such as tetrahydrofuran (THF) and benzene.²²

A classic example is magnesium: Mg-containing species have attracted interest in extraterrestrial, organic synthesis and biological systems.²³ Mg(i) radical species have long been postulated to exist as fleeting species and reaction intermediates, which are important for understanding life processes in Earth and extraterrestrial²⁴ environments and also play crucial roles in the formation of Grignard reagents.²⁵ They have been spectroscopically detected in harsh conditions, for example, [•]MgF as a gaseous molecule is generated at 2300 °C and trapped in an inert-gas gas matrix²⁶ and [•]MgCH₃ was produced by a laser ablation/photodissociation technique.²⁷ Calculations have showed that Mg₂Cl₂ and other Mg–Mg compounds disproportionate exothermically into solid Mg(0) and the corresponding Mg(II) compound, which further indicated the difficulty of isolation of a Mg(i) monomeric radical species.²⁸

The first stable and isolable Mg(i) complexes were Mg(i)–Mg(i) dimers, isolated in 2007 by Jones and Stasch,²⁹ which are thermodynamically stabilised by forming a Mg(i)–Mg(i) bond, and kinetically protected by bulky Priso or Nacnac ligands (Priso = [(DippN)₂CN[†]Pr₂], (Dipp = 2,6-diisopropylphenyl); Nacnac = [ArNC(Me)₂CH]). Since then, this field has emerged as one of the most fruitful areas of main group chemistry. In the 2010s and early 2020s, Mg(i) dimers have developed significantly from a niche area into a library of soluble, selective, stoichiometric and safe reducing agents for organic and inorganic synthesis.³⁰

Despite the flourishing *status quo*, almost all the reported Mg(i) complexes are dimers: the Mg(i)–Mg(i) intermetallic bond plays an underpinning role in their stabilisation.

The defining structural feature of these Mg(i) dimers is the aforementioned 2-electron Mg(i)–Mg(i) σ-bond. In contrast, a hypothetical monomeric Mg(i) complex would feature an unpaired Mg-based single electron, effectively creating a Mg(i) radical. Such a monomeric radical would likely exhibit distinct, and potentially enhanced, reactivity compared to the dimeric Mg(i)–Mg(i) complexes.³¹ Consequently, the pursuit of such monomeric Mg(i) radicals has been a longstanding challenge in the main-group chemistry community for over a decade.

One possible strategy to isolate a Mg(i) monomeric complex is increasing the steric repulsion between the two Mg(i) centres in the dimer until the Mg(i)–Mg(i) bond homolyses and generates the desired Mg(i) monomer. Great efforts have been devoted to this route: several research groups adopted sterically bulky ligands, and external Lewis bases to increase the steric repulsion.²² But these efforts only led to limited successes. In most of the cases, Mg(i)–Mg(i) bond elongation was observed upon increasing the steric repulsion. The Mg(i)–Mg(i) bond is fairly “elastic” and can adopt a wide range of bond lengths (2.808 Å (ref. 32) to 3.196 Å (ref. 33)) without cleavage. Increasing the steric repulsion by increasing the steric bulkiness of the ligands can also lead to a change of the ligand coordination mode and hapticity (*e.g.*, for β-diketiminato ligands, from κ²-N, N to κ¹-N) to release the steric repulsion, instead of forming the desired Mg(i) monomer.³⁴

In 2021, a bulky bis-fluorenyl flanked ligand, namely hydrindacene, was introduced into Mg chemistry by the Tan group. Reducing the Mg(II)-hydrindacene precursor led to a ligand-reduced product, instead of the desired Mg(i) monomer.³⁵ It is worth noting that the hydrindacene ligand family has succeeded in isolating a number of low-valent mononuclear *p*-block complexes,³⁶ emphasising the challenge of isolating a Mg(i) monomer.

The first evidence of Mg(i)–Mg(i) homolysis emerged in 2021 by using the photochemical method. The Jones group used blue/UV light to activate Mg(i) dimers containing bulky ligands ([{(ArNacnac)Mg}₂] (ArNacnac = [HC-(MeCNAr)₂]; Ar = Dipp or 2,4,6-tricyclohexylphenyl (TCHP))) to generate transient Mg(i) radical intermediates, which underwent fast reactions with aromatic solvents, such as benzene, toluene and xylenes³⁷ (Fig. 1). The presence of Mg(i) radical intermediates was supported by both computational studies and their observed enhanced reactivity compared with the corresponding Mg(i)



Fig. 1 Photo-activated Mg(i)–Mg(i) homolysis.³⁷





Fig. 3 Synthesis of low-valent Mg(I) complexes by ball-milling.⁴²

substantial progress has been made, especially since the 2020s, and benefitted from the use of mechanochemistry. First, inspired by their successful isolation of the Mg(I) monomer **4**, in 2023 the Harder group adapted the mechanochemical synthesis into Ca chemistry. A Ca(II) precursor **10** and K/KI were ball-milled to produce a purple solid. Like the Mg(I) monomer **4**, the deep colour of the solid likely links to the formation of a radical species. While the purple solid was not characterised in solid state, it was subjected to reaction with benzene to yield a biphenyl dianion complex **11**, [(^DIPEP)BDI]Ca(THF)₂(biphenyl) (Fig. 4). The production of **11** is postulated to proceed *via* a Ca(I) radical species, [•]Ca(^DIPEP)BDI(THF), which could react with benzene to give the coupling products.⁴⁸ In 2024, in another attempt to use mechanochemistry to reduce Ca(II) to Ca(I), a new Ca-based room temperature stable electride (RoSE) was obtained, instead of the Ca(I) complexes.⁴⁹ This Ca-based RoSE will be discussed in the following electride Chapter.

From the abovementioned cases, it is obvious that the mechanochemical ball-milling method has exhibited unique

capability, especially in isolating solvent-incompatible species (whether due to reaction with solvents or solubility issues). Though still in its infancy, successes such as the first, and so far, the only Mg(I) monomeric complex should encourage more synthetic main-group chemists to adopt mechanochemistry into their work.

2.2 Electrides

An electride is a type of chemical compound in which electrons serve as the anion.^{50,51} In electrides, these electrons are localised in voids, cavities, or channels within the structure, such as in crystals, molecular clusters, or between atomic layers, instead of occupying atomic/molecular orbitals. The electrons are not associated with any specific atom but are instead confined within spaces created by their surrounding structures (*e.g.*, crystal lattice).⁵² Though it should be noted electride phase does not necessarily only exist in crystalline materials, but also in amorphous and liquid materials (*e.g.*, Na metal dissolved in liquid ammonia⁵³). Such properties make electrides ideal for



Fig. 4 Syntheses of Ca biphenyl complexes.⁴⁸





Fig. 5 Synthesis of a room-temperature-stable, Li/K heterobimetallic electride $\text{K}^+(\text{LiHMDS})\text{e}^-$ by ball-milling.⁶⁴

a wide range of applications. In synthesis, they serve as highly effective reductants or catalysts, facilitating processes such as organic synthesis, ammonia production, and CO_2 splitting.⁵⁴ Beyond synthesis, electrides also play crucial roles in materials science, functioning as electron emitters, superconductors, battery anodes, and components in optics, lamps, and even radioactive waste storage.⁵⁵

Since 1983, when the Dye group first reported the electride $\text{Cs}^+(\text{18-C-6})_2\text{e}^-$ (18-C-6: 18-crown-6), the synthesis of electrides has proven to be exceptionally challenging.⁵⁶ For instance, the initial synthesis of $\text{Cs}^+(\text{18-C-6})_2\text{e}^-$ required dissolving caesium metal in dimethyl ether–methylamine mixture, with a 2 : 1 stoichiometric amount of the crown ether to metal, all conducted at $-30\text{ }^\circ\text{C}$ until crystals were formed.⁵⁷ Since the 1983 inaugural report, the Dye group synthesised eight “organic electrides”, with a general structure $\text{AM}^+(\text{crown ether/cryptand})_n\text{e}^-$, where AM^+ is a Group-1 alkali metal cation.⁵⁸ All the organic electrides were synthesised *via* a similar cryogenic low-boiling-point solvent dissolved metal process, which is challenging, if not impossible, for a conventional synthetic chemistry laboratory. Most of Dye’s organic electrides are unstable at room temperature. It wasn’t until 2005 that the first room-temperature-stable organic electride was successfully synthesised.⁵⁹ This breakthrough was achieved by employing a specially designed aza-cryptand ligand, in order to avoid the C–O cleavage decomposition pathway, and employed a week-long, meticulously controlled process in liquid ammonia at $-78\text{ }^\circ\text{C}$, performed in a helium glovebox, underscoring the extreme difficulty of obtaining these materials.

In 2022, the Martin group reported a second room-temperature-stable organic electride, namely a magnesium electride $[(\text{THF})_4\text{Mg}_4(\mu^2\text{-bipy})_4][(\text{THF})_6\text{Mg}_2(\mu^2\text{-bipy})(\text{Cl})]$, (bipy = 2,2’-bipyridine) which was synthesised as a serendipitous product during their attempt to reduce a bipyridine–Ni(II) dichloride.⁶⁰ Indeed, the two organic RoSE from the Dye and Martin group are tremendous synthetic achievements, but their need for a specialised aza-cryptand ligand, impractical synthetic protocol, or their serendipitous nature, prevent their wider application as a general gateway to a library of organic RoSEs.

In parallel with the organic electrides, there are several “inorganic electrides”, which are derived from binary or ternary inorganic materials A_xB_y or $\text{A}_x\text{B}_y\text{C}_z$.⁶¹ In 2003, the Hosono group achieved a milestone by synthesising the first air- and room-temperature-stable electride, $[\text{Ca}_{24}\text{Al}_{28}\text{O}_{64}]^{4+}(4\text{e}^-)$.⁶² However, this synthesis was not straightforward. The electride was derived from $12\text{CaO}\cdot 7\text{Al}_2\text{O}_3$ crystals, which were obtained by melting the starting material (melting point $1415\text{ }^\circ\text{C}$), followed

by a careful reduction process, involving heating in a hydrogen atmosphere. Similarly, the synthesis of the first water-stable electride, Y_5Si_3 , involved melting yttrium and silicon ingots in a precise stoichiometric ratio.⁶³ The resulting silver-coloured ingot was then finely ground using an agate mortar under a strictly controlled argon atmosphere.

Both organic and inorganic electrides are typically synthesised under harsh and demanding conditions, underscoring the pressing need for a milder, more accessible synthetic method. In this context, mechanochemistry presents a tempting candidate. But it was not introduced into this area until very recently.

In 2023, the Lu group first utilised the mechanochemical method for electride synthesis, successfully producing a RoSE, namely $\text{K}^+(\text{LiHMDS})\text{e}^-$ (**13**) (HMDS = 1,1,1,3,3,3-hexamethylsilylazide), using commercial accessible starting materials *via* the mechanochemical ball-milling (Fig. 5).⁶⁴ A 1 : 1 molar ratio mixture of the starting materials, LiHMDS and potassium metal, were ball-milled at room temperature for 15 minutes, resulting in the formation of the electride **13** as a blue coloured powder. Remarkably, this method could be scaled up to 20 mmol without significant deterioration in yield, showcasing the efficiency and scalability of mechanochemistry in synthesising such reactive species. **13** is stable at room temperature under an inert atmosphere (N_2 or Ar) for several months, therefore, **13** is a RoSE by definition. As a highly reactive amorphous material, characterisation of **13** is inherently challenging. By combining electron paramagnetic resonance (EPR), magnetometry, DFT calculations and *ab initio* random structure searching **13** was proved to feature a unique three-dimensional helical delocalised electron density (Fig. 6).

13 exhibited versatile reactivity, including: (i) C–H activation and C–C coupling of benzene and pyridine; (ii) solvent-free Birch reductions, highlighting its potential as a strong reducing agent in synthetic chemistry. Moreover, upon treating with external ligands, the anionic electron in **13** can selectively re-combine with the Li^+ or K^+ , achieving the first selective chemical reduction of Group-1 metal cations.⁶⁵

In 2024, a new calcium-based RoSE **15**, $\text{K}\{[\text{Ca}[\text{N}(\text{Mes})(\text{SiMe}_3)_3(\text{e}^-)]_2\text{K}_3]$ (Mes = 2,4,6-trimethylphenyl), was synthesised by the mechanochemical ball-milling method (Fig. 7).⁴⁹ This process involved ball-milling a calcium trisamide **14**, $[\text{Ca}\{\text{N}(\text{Mes})(\text{SiMe}_3)_3\text{K}\}]$, with potassium metal at 30 Hz for 30 min (mixer mill) or 4000 rpm for 3 h (planetary mill). It is worth to mention that in a previous work, the reduction of **14** with KC_8 was attempted both with 18-C-6 in THF, and in benzene without any sequestering agent, however





Fig. 6 The AIRSS calculated structure of electride **13**, showing the helical anionic electron density topology.⁶⁴

no reaction was observed.⁶⁶ Also, in a control reaction, the precursor complex **14** was observed to undergo direct degradation in toluene, with no formation of the electride **15**. This observation clearly indicates that, in the presence of a solvent, electride **15** cannot be obtained. **15** reacted with pyridine and benzene, confirming its potential as a strong reductant.

2.3 Alkalides

The alkali metals are the most electropositive elements of the periodic table, with their Pauling-scale electronegativities spanning from 0.98 (Li) to 0.79 (Cs) (*cf.* C 2.55, Mg 1.31, H 2.20).⁶⁷ Hence, the zero-valent alkali metal atoms AM(0) easily lose one electron to form the corresponding AM(+1) cations. However, another possibility is accommodating one more electron into their valence-shell *s*-orbitals, resulting in AM(−1) anions with a fully occupied electron-paired *ns*² close-shell structure. Indeed, alkalides are a unique class of compounds where the alkali metals (such as sodium, potassium) exist as anions (*e.g.*, Na[−], K[−]).⁶⁸ In alkalides, the zero-valent alkali metal atoms gain an extra electron, resulting in a negatively charged species. This unusual electronic configuration makes alkalides highly reactive.⁶⁹

Despite being predicted in the 1960s,^{70–72} the first definite proof of alkali metal anions came in 1974 when Dye and co-workers isolated the first alkalide, a sodide (Na[−]), and by 1979 the group had reported Na[−], K[−], Rb[−] and Cs[−].⁷³ Their unique electronic properties, exceptional reducing power, and the potential to advance coordination chemistry and electron

transfer mechanisms have generated significant interest.⁷⁴ However, like the electrides, the synthesis of alkalides is exceptionally challenging, which limits their reactivity studies. Their high sensitivity to air, moisture, and in some cases, light, necessitates that the synthesis and storage must be conducted under rigorously controlled inert atmospheres, often requiring specialised techniques and equipment. However, the synthesis itself is not the only difficulty. While alkalides achieve some degree of thermodynamic stability with fully occupied *s*-orbitals, their negatively charged metal anions in solution can undergo comproportionation with the cations, *i.e.*, AM(−1) + AM(+1) → 2 AM(0). Additionally, equilibria between the alkali metal and ligands in solution mean that electrides and alkalides can coexist, further complicating their isolation.⁷⁵ Moreover, trace impurities that are difficult to eliminate can readily trigger decomposition, adding another layer of difficulty to an already complicated synthesis process.

The synthesis of alkalides was developed by the Dye group, requiring stringent purification of all reagents to remove trace impurities that could induce product decomposition. These synthetic protocols commonly demand meticulous attention to detail to ensure their success. Typically, the synthesis of alkalides is achieved by dissolving appropriate ratios of the alkali metal and the macrocyclic ligands, such as cryptands or crown ether, in low-boiling point solvents, such as dimethyl ether or methylamine, at low temperatures. The vast majority of alkalides synthesised contain macrocyclic ligands. Indeed, the macrocyclic effect is a major driving force for the isolation of alkalides, in which the alkali metal cation is sequestered and kinetically kept away from the alkalide anion, preventing comproportionation. Dissolving the alkali metal often takes a long time. For instance, it takes about 3 h of constant agitation at −30 °C to dissolve enough Na–K alloy in 10 mL of C222-isopropylamine solution to produce a 0.04 M solution of K⁺C222Na[−].⁷⁶ For Na⁺(C222)Na[−] (**16**), sodium metal is dissolved in ethylamine or methylamine with [2.2.2]-cryptand (C222) at −78 °C.⁷⁷ Although higher temperatures might accelerate the dissolving and reaction, they also significantly increase the risk of comproportionation and decomposition.

Once the reaction is completed, most of the solvent needs to be removed by evaporation, and a suitable co-solvent or mixture is introduced to promote crystallisation. The resultant crystalline powder or crystals are then repeatedly washed with diethyl ether and dried *via* residual ether evaporation, which stabilises the samples at low temperatures. However, a vacuum seal-off near the preparation bulb presents additional risk, as sealing under dynamic vacuum, even with cooling from liquid nitrogen, can cause rapid, irreversible decomposition.⁷⁸ Consequently, it is crucial to transfer the resultant powder products to a clean



Fig. 7 Synthesis of a room-temperature-stable, Ca-based electride by mechanochemistry methods.⁴⁹





Fig. 8 Experiment set-up of classic alkali synthesis.^{76,78,80} (a) Apparatus for preparing alkali using flame-sealed tubes and ampoules. This is the old design, used mostly before 1980. Adapted with permission from *J. Phys. Chem.* 1980, **84**, 1084. Copyright {1980} American Chemical Society. (b) An H-cell with J.-Young taps, used with high-vacuum line. In between the cells is a sinter filter pad. This is the new design, used after 1980 until the 2000s. A similar K-cell was used as well. Reprinted with permission from *J. Phys. Chem.* 1984, **88**, 3842. Copyright {1984} American Chemical Society. (c) Pyrex-quartz apparatus for the preparation of alkali by vapor deposition. This was used when the volatile solvents need to be avoided. Adapted with permission from *J. Phys. Chem.* 1982, **86**, 7. Copyright {1982} American Chemical Society.

vessel before final sealing, even for the relatively stable $\text{Na}^+(\text{C}222)\text{Na}^-$.⁷⁷ To meet these rigorous synthesis requirements, specially designed glassware, as illustrated in Fig. 8, is employed.^{76,79} Although the Dye group has also explored Direct Vapor Deposition (shown in Fig. 8(c)) as an alternative alkali synthetic method, this approach has achieved only modest improvements regarding its synthetic efficiency (*e.g.*, improving the yield) and easiness of handling.⁸⁰

It is quite obvious that the alkali synthesis is very challenging, which limits a full exploration of the alkali chemistry. An accessible and scalable synthetic route is needed. Mechanochemistry presents a promising candidate: the absence of solvents could effectively eliminate equilibria and comproportionation issues.

In 2024, the Lu group introduced the mechanochemical ball-milling method into alkali chemistry, reporting the first facile and scalable synthesis of an alkali.⁸¹ The prototypical alkali, the sodide $\text{Na}^+(\text{C}222)\text{Na}^-$ (**16**), was synthesised by a simple 10 minutes ball-milling reaction between 2 equivalents of Na metal and 1 equivalent of [2.2.2]-cryptand at room temperature and under argon atmosphere (Fig. 9). **16** can be obtained in 48% yield as a crystalline solid from its dry THF solution at $-35\text{ }^\circ\text{C}$. The sodide **16** features two distinct signals on its magic-angle spinning (MAS) solid-state ^{23}Na NMR for Na^+ (-11 ppm) and Na^- (-62 ppm), respectively.

Compared to the classical alkali synthesis methods mentioned earlier, this mechanochemical method offers multiple advantages. Firstly, the ball-milling method employs commercially available materials: both the Na metal and the [2.2.2]-cryptand were used without further purification from their commercial sources; especially, removal of the oxide layer on the Na metal surface is not necessary, which makes the synthesis substantially easier. Secondly, it operates at room temperature without stringent temperature control. Additionally, the reaction time is drastically shortened to 10 minutes, a substantial improvement over the previous method.⁷⁹ Last but not least, the Lu protocol can be scaled up, which is essential for the following reactivity studies. With such an accessible entry to the sodide **16**, the authors investigated the sodide reactivity. **16** delivers 2-electron reductive cleavages of C–N bonds in primary amines, and 1-electron reduction of azobenzene (Fig. 10).⁸¹

2.4 Miscellaneous

The Nobel Prize-winning metallocene is an essential class of organometallic complexes for all metals across the periodic table. The centre of the metallocene chemistry is the coordination hapticity between the cyclopentadienyl (Cp) ligand (and its derivatives) and the metal cation.



Fig. 9 Synthesis of sodide complex $\text{Na}^+(\text{2.2.2-Cryptand})\text{Na}^-$ by ball-milling.⁸¹





Fig. 10 Reactivity of the sodide (16) towards amines and azobenzene.⁸¹

As a part of the early metallocene “gold rush”, Fischer and co-workers reported the beryllocene, [BeCp₂], in 1959.⁸² Four decades later, Carmona and co-workers reported the Cp* version, [BeCp*₂], in 2000.⁸³ It is interesting that the sterically less bulky [BeCp₂] adopts an η⁵:η¹ hapticity for its two Cp ligands, but the bulkier [BeCp*₂] adopts an η⁵:η⁵ hapticity. While indenyl complexes had previously been reported for all other non-radioactive s-block metals, no such beryllium indenyl complex had been isolated. To investigate such a complex, in 2021 Hanusa and co-workers set out to synthesise the indenyl (Ind) version of beryllocene, namely [Be(Ind)₂].⁸⁴ However, the authors found that the conventional solution synthesis between K[Ind] and BeBr₂ in THF yielded an intractable mixture.⁸⁴ The authors then turned to the mechanochemical ball-milling method, which led to the successful isolation of the desired complex **23** (Fig. 11). Noting that in **23**, one indenyl ligand coordinates in η⁵-mode, while the other in η¹-mode. In this case, the mechanochemical method appeared superior to the conventional synthesis in solution, though the exact reason(s) for this phenomenon remains unclear.

3. Reactions involving reactants with limited solubility

Traditional reactions are mostly conducted in solution. However, solution-state reactions are often challenging when

one or more reactants are poorly soluble, particularly in heterogeneous systems. Under such conditions, limited surface interaction between the reactants can lead to sluggish reaction rates and incomplete conversion. Mechanochemical methods offer a promising solid-state alternative to the traditional solution-state synthesis, successfully facilitating reactions with no solvent required, thereby providing a pathway to bypass the limitations in heterogeneous reactions.


In 2023, the Gouverneur and Hayward groups reported a remarkable example of mechanochemistry's transformative potential in facilitating reactions involving insoluble reactants, in this case is fluorspar (calcium fluoride CaF₂).⁸⁵ All fluorochemicals originate from naturally occurring fluorspar.⁸⁶ However, CaF₂, a white solid with a high melting point at around 1420 °C, is poorly soluble in water (0.016 g L⁻¹ at 20 °C) and insoluble in organic solvents, making it unsuitable for direct use in fluorochemical synthesis. Traditionally, the production of fluorochemicals relies on the process first reported by C. W. Scheele in 1771.⁸⁷ involving the generation of hydrogen fluoride (HF) *via* the reaction of acid-grade (>97%) fluorspar with sulfuric acid (>98%) at 100–300 °C.⁸⁸ Given the toxicity and corrosive nature of the HF and the concentrated sulfuric acid, a method directly utilising CaF₂ and circumventing the HF stage is highly desirable.

The Gouverneur and Hayward groups reported such a HF-free direct CaF₂ utilisation by using mechanochemistry.⁸⁵



Fig. 11 Synthesis of di(indenyl)beryllium.⁸⁴



Fig. 12 The synthesis of fluoromix.⁸⁵

Acid-grade fluorspar (CaF₂, 1 eq.) was milled with anhydrous K₂HPO₄ (>98%, 2.5 eq.), giving a fluorinating reagent named fluoromix. The fluoromix, a white powder, was prepared *via* a stepwise process: 1 eq. of CaF₂ and 1 eq. of K₂HPO₄ were milled at 35 Hz for 3 hours, followed by the addition of another 1 eq. of K₂HPO₄ for a further 3 hours of milling, and finally, 0.5 eq. of K₂HPO₄ was added and milled for further 3 hours (Fig. 12). The resulting fluoromix 25, obtained after 9 hours of milling. Even after storing under inert atmosphere at room temperature for 9 months, the fluoromix was found still active.

As a nucleophilic fluorination reagent, the fluoromix showed impressive versatility, enabled a number of fluorination reactions listed in Fig. 13. Mechanistic studies revealed that the fluoromix contains crystalline constituents, including K₃(HPO₄)F and K_{2-x}Ca_y(PO₃F)_a(PO₄)_β, which serve as fluorinating agents. Studies to confirm its composition include ³¹P and ¹⁹F NMR for soluble species, PXRD analysis for crystalline species, and control experiments for amorphous species. These species facilitate the synthesis of sulfonyl fluorides, α-fluoroketones, fluoroesters, α-fluoroamides, benzylic fluorides, alkyl fluorides, and (hetero)aryl fluorides.

In this work, the mechanochemical method unlocked the direct utilisation of the insoluble CaF₂ as a fluoro-building block, circumventing the conventional HF stage, which is undesirable from the aspects of cost, health and safety and sustainability.

Very recently, the Gouverneur group further developed the mechanochemical CaF₂ activation methodology by reporting a one-step synthesis of alkali metal fluorides (AMF, AM = Li, Na, K) from CaF₂.⁸⁹ This was achieved by ball-milling CaF₂ with 2.0 equivalents of AMOH and 1.0 equivalent of TiO₂ at 35 Hz, room

temperature, for 3 hours. Byproducts are CaTiO₃ and H₂O. Further fluorination reactions using AMF were not reported in this work, though.

Zero-valent metals are good electron sources for reduction reactions. The ability to use zero valent metals directly and easily is undoubtedly desirable. However, due to their poor solubility in organic solvents, the direct use of zero valent metals is often difficult in traditional solution-state reactions. Zero-valent metals can be accessed as variety of forms (*e.g.* powder, lumps, *etc.*) and typically have an oxide coating on the outer layer, which needs to be removed to allow the reaction to happen. Typically, the zero valent metals must be activated prior to the reaction to increase the surface area, for example, using highly dispersed metal suspensions.⁹⁰ Reactions involving zero-valent metals are often difficult to predict and are sometimes difficult to reproduce, since the surface area, a determining factor, is difficult to control and could be different from batch to batch. In recent years, mechanochemistry offers an alternative pathway to use zero-valent metals, such as Li,⁹¹ Mg, Mn, Zn, Ag, and Bi, as reductants.⁹²

A demonstration of mechanochemistry's potential is a manganese metal-mediated dimerisation of electron-poor alkenes reported by the Browne group in 2021.⁹³ Ball-milling arylidene malonates 32 with manganese pieces, LiCl, and a minimal amount of THF (1 eq., liquid-assisted grinding, LAG)⁹⁴ yielded the dimerised products 33 (Fig. 14). The precise reaction mechanism remains unclear – whether the manganese metal participates as a concerted two-electron reduction or two sequential single-electron reductions. Nevertheless, control experiments unequivocally proved the necessity of the ball-milling method: stirring the reactants and manganese metal (powder or pellets) in bulk organic solvents did not work.

The Birch reduction has been widely used by organic chemists for over half a century, enabling the dearomatisation of arenes into 1,4-cyclohexadiene derivatives.⁹⁵ Traditionally, this transformation relies on alkali metals dissolved in liquid ammonia under cryogenic conditions. However, these classical procedures are fraught with operational challenges, including the need of liquid ammonia, which has to be slowly evaporated and carefully exhausted after the reaction. The restricted solubility of alkali metals limits the scope of solvents available, prompting significant interest in alternative, more practical methodologies. In recent years there has been an increasing interest in ammonia-free Birch reduction, including reports utilising electrochemical,⁹⁶ photochemical,⁹⁷ and enzymatic catalytic methods,⁹⁸ as well as employing lithium metal in an ammonia-free system.⁹⁹ However, all these methods still need bulk organic solvent, which is a sustainability culprit.

In 2023, the Lu group reported the first mechanochemical solvent-free Birch reduction by using the newly developed RoSE

Fig. 13 Summary of C–F and S–F bonds formation.⁸⁵



Fig. 14 Synthesis manganese mediated reductive dimerisation of aryldiene malonates by ball-milling.⁹³

reagent, $\text{K}^+(\text{LiHMDS})\text{e}^-$ (**13**),⁶⁴ previously discussed in the electrides chapter. As the first solvent-free Birch reduction, the Lu group's work was a conceptual breakthrough, but there are two technical drawbacks: (i) the reductant, $\text{K}^+(\text{LiHMDS})\text{e}^-$, is still a specialised reagent; (ii) the reactions need to operate under argon.

Shortly after, the Ito/Kubota group reported a Li metal mediated mechanochemical Birch–Benkeser reaction, which operates under air, overcoming the technical drawbacks.¹⁰⁰ As shown in Fig. 15, arenes **34** were ball-milled with Li metal and amine additives at room temperature under air for only 1 min, yielding the corresponding Birch-type reduction products **35** with good to excellent yield. The scope of substrates covers from carboxylic acids to electron-rich aromatic ethers, arene-bearing alcohols and several bioactive molecules.

This approach circumvents the need to dissolve lithium in liquid ammonia or other solvents. More importantly, the reaction is fast, air-tolerant, requires no specialised equipment or handling of cryogenic or hazardous materials, and demonstrated good functional group compatibility.

Following these works, several reports on mechanochemical Birch-type reductions emerged, underscoring the growing interest in this area. These include a Birch reduction by sodium lumps¹⁰¹ and calcium metal¹⁰² (the Ito/Kubota group), as well as calcium- and magnesium-metal mediated Birch reductions¹⁰³ (the Kananovich group). It appears that the mechanochemical Birch reduction is turning into a hot area.

Beside Birch reduction, Group-1 metals have also been used in other mechanochemical reactions. In 2023, the Lu group reported mechanochemical reduction and Li-ion doping of transition metal oxides, by ball-milling Li metal and nickel and cobalt oxides under argon (Fig. 16).¹⁰⁴

In 2025, the Ito/Kubota group reported mechanochemical synthesis of organolithium reagents, by ball-milling Li metal and organo-halides in air (Fig. 17).¹⁰⁵ It is noteworthy that Et_2O is needed to facilitate the organolithium formation, through a LAG mechanism. The organolithiums were not isolated, instead, they were used for further mechanochemical reactions with electrophiles (E^+), also in air. This method is significantly



Fig. 15 Lithium-based Birch reduction by ball-milling.¹⁰⁰



Fig. 16 Ball-milling reactions between Li metal and transition metal oxides, leading to the transition metal reduction and Li-ion doping.



Fig. 17 Mechanochemical synthesis of organolithium reagents.¹⁰⁵

easier and more practical in comparison to the traditional method to make organolithiums, *i.e.*, refluxing Li metal pieces and organo-halides in solution under inert gas atmosphere.

Mechanochemical methods not only offer new and versatile reaction conditions for the direct utilisation of metals but also provide immediate and significant improvements for substrates with poor solubility in conventional solvents.

In recent years, the Ito/Kubota group has reported several mechanochemical Suzuki–Miyaura cross-coupling reactions.^{106–111} A key benefit of the mechanochemical cross-couplings is their capability to activate insoluble electrophiles, especially the aryl halides¹¹² (Fig. 18). In this work, the use of 1,5-cod as a dispersant and a LAG reagent is also crucial.

In Fig. 19, aryl halides are classified by their solubility based on US Pharmacopoeia, 10^{-2} to 10^{-3} M as 'slightly soluble', 10^{-3} to 10^{-5} M as 'very slightly soluble' and 10^{-5} to 10^{-6} M as 'practically insoluble'.¹¹² For aryl halides in different catalogues, they compared the results of the mechanochemical method (5–90 minutes at 250 °C) with solution-based methods (24 hours,





Fig. 18 Cross-coupling reactions with insoluble aryl halides.¹¹²



Fig. 19 Classification of aryl halides based on their solubility.¹¹² (Reprinted with permission from *J. Am. Chem. Soc.*, 2021, 143, 6165. Copyright (2021) American Chemical Society.)

120 °C). Despite the much longer reaction time, the solution-state reactions afforded significantly lower yields. Of particular note is the successful reaction of “practically insoluble” (class 3) alkyl halides which gave no reaction in solution. The mechanochemistry strategy offers the first practical method for the cross-coupling of insoluble aryl halides. While the need for high temperature ball-milling may be a drawback, this method provides new opportunities to expand the structural diversity of organic molecules. In 2023, the Ito/Kubota group managed to lower the reaction temperature to 45 °C by using a bespoke phosphine ligand.¹¹³

Beyond metals, minerals,¹¹⁴ and organic compounds, the mechanochemical method has also been used to activate chalcogen elements. Element selenium is a cheap, commercially available, stable, odourless, and easy to handle starting material for organic selenation reactions.¹¹⁵ However, elemental

selenium is fairly inert, has low solubility in organic solvent, and has a tendency to form transition-metal selenium clusters, which make using it directly challenging.

In 2024, the Wei group reported a ball-milling-enabled C–Se bond formation from accessible organic halides, Mg metal and elemental selenium, with a wide substrate scope and good functional group tolerance.¹¹⁶ After ball-milling aryl halides **39**, Mg, LiCl, Se under LAG¹¹⁷ conditions (THF), the corresponding diselenides **40** were obtained in good yields (Fig. 20). If excess Mg was used, a magnesium-based selenium nucleophile intermediate forms, with the corresponding unsymmetrical mono-selenides **41** forming after *N*-phenylacrylamide was added (Fig. 20). Notably, this mechanochemical method also works for thiolation and telluration. The formation of **40** and **41** share the same magnesium selenide intermediate. Though not clearly pointed out in Wei’s paper, we hypothesise that the reaction can





Fig. 20 The synthesis of symmetrical diselenides and unsymmetrical monochalcogenides.¹¹⁶

proceed in two ways: (1) in air, the Mg selenide oxidises into selenium radical and dimerise into **40**; (2) with *N*-phenylacrylamide, the Mg selenide undergoes a magnesium-selenation towards the C=C bond, which is then protonated to form **41**. Another possibility is that, Se reacts with the Grignard reagents ArMgX , through a Mg-C bond selenation mechanism,¹¹⁸ to yield the Mg selenide intermediate.

Grignard reagents are some of the most widely used organometallic reagents in synthetic laboratories. Recent advancements in mechanochemistry have not only enhanced the synthesis of traditional magnesium-based Grignard reagents but also expanded the scope to heavier Group-2 elements such as calcium, strontium, and barium.¹¹⁹ From the textbook, higher atomisation energies and larger first ionisation energies (for alkaline earth metal, the second ionisation) is needed to prepare organometallics from the alkaline earth metals. As a result, the heavier alkaline earth metals need elevated temperatures or specialised activation methods to initiate the reactions to form their Grignard-analogues. But these heavier Grignard reagents are not only chemical curiosities, but could offer enhanced reactivity from their more ionic, more reactive metal-carbon bond.¹²⁰

Since the activation of the insoluble metals (Ca, Sr, Ba) is the barrier, mechanochemistry offers a prime solution. In 2024, the Ito/Kubota group demonstrated a direct arylation of alkyl fluorides through *in situ* generated Ca-based heavy Grignard reagents *via* mechanochemical ball-milling.¹²¹ In this work, aryl iodides **42** and alkyl fluorides **43** were ball-milled with commercially available calcium metal and a small amount of tetrahydropyran (THP) in air for 60 minutes at 80 °C, giving the C(sp³)-F bond arylation product **44** in good to high yields (Fig. 21).




Fig. 21 Arylation of alkyl fluorides using mechanochemically generated calcium-based heavy Grignard reagents.¹²¹

This mechanochemical method not only simplified the generation of reactive Grignard species but also bypassed the need for transition metal catalysts. The direct arylation in Fig. 21 of alkyl fluorides is particularly noteworthy. Additionally, the reaction conditions were air-tolerant and conducted at 80 °C (internal), making the process far more practical than traditional methods that require inert atmospheres and highly controlled environments.

In 2024, the Ito/Kubota group reported the mechanochemical generation of aryl barium nucleophiles from inactive barium metal.¹²² Barium metal is usually inert – its activation was reported *via* metal vapour synthesis.¹²³ However, in Ito/Kubota group's study, ball-milling commercial Ba metal and the substrates (**45–48**) yielded the silylation and nucleophilic addition products, **47** and **49**, respectively (Fig. 22). These reactions proceed *via* the barium organometallics ArBaI , which subsequently react with the electrophiles.

There are more examples of using the mechanochemical strategy for Mg-based Grignard reagents.^{124,125} Advantages of this strategy include faster reaction rates, reduction in safety hazards and increasing reactivity of metal.

An interesting example of a mechanochemical preparation of *bis*(*n*-propyltetramethylcyclopentadienyl)strontium (SrCp'_2 , **51**, $\text{Cp}' = n$ -propyltetramethylcyclopentadiene) was reported by Peters and Blair in 2014.¹²⁶ The common strontium starting material, SrI_2 , is poorly soluble in hydrocarbon and ethereal solvents. By using a bespoke reactor modified from commercial glass flasks, SrI_2 and $\text{Cp}'\text{K}$ were made into a suspension with stirring from six 12.7 mm stainless steel balls for 24 hours, to afford the Et_2O adduct **50** (Fig. 23). After a vacuum distillation of **50**, the solvent-free **51** was produced. Though this case is more akin to a classic suspension reaction, the authors did categorise it into a mechanochemical reaction.

4. Elimination of solvent coordination

Coordinated solvent molecules are ubiquitous in coordination chemistry. Solvent molecules can bond strongly to metal centres and influence the structure and reactivity. On one hand, they can stabilise reactive species by masking the exposed metal centre. But on the other hand, they could also block the reactive





Fig. 22 Reactions via direct generation of various aryl barium nucleophiles from commercially available inactivated barium metal and aryl halides.¹²²



Fig. 23 The synthesis of unsolvated $\text{Sr}(\text{Cp}')_2$.¹²⁶

centre. The coordination solvent effects are particularly pronounced in cationic polymerisations, which are rather sensitive to the coordination number and electronic environments of the metal centre.¹²⁷ The elimination of solvent coordination is also particularly important when the aim is to isolate highly reactive, base-free, or low-coordinate metal complexes.¹²⁸ The mechanochemical methods have inherent advantages in avoiding the coordination of solvent molecules.

Early in 2014, Hanusa and co-workers reported a mechanochemical synthesis of unsolvated tris(allyl)aluminum

complexes, $[\text{1,3-(SiMe}_3)_2\text{C}_3\text{H}_3]_3\text{Al}$, **54**, and their scandium analogue, $[\text{1,3-(SiMe}_3)_2\text{C}_3\text{H}_3]_3\text{Sc}$, **55** (Fig. 24).¹²⁹ In their study, attempts to synthesise the base-free tris(allyl)aluminum complex, **54**, in the presence of solvents such as THF or diethyl ether resulted in unidentified complex mixtures. However, by employing ball-milling, aluminum/scandium halides (AlX_3 , $\text{X} = \text{Cl, Br, I; ScCl}_3$) were successfully reacted with the potassium allyl salt **52** (1,3-bis(trimethylsilyl)allyl potassium) in solid-state, yielding the unsolvated **54** and **55**, respectively. The high reactivity of the unsolvated complex, **54**, is demonstrated by its



Fig. 24 Synthesis of unsolvated $[\text{1,3-(SiMe}_3)_2\text{C}_3\text{H}_3]_3(\text{Al, Sc})$.¹²⁹





Fig. 25 Synthesis of unsolvated $\{\text{Sn}[1,3-(\text{SiMe}_3)_2\text{C}_3\text{H}_3]_3\text{K}\}_\infty$ and $\text{Sn}[1,3-(\text{SiMe}_3)_2\text{C}_3\text{H}_3]_4$.¹³⁰



Fig. 26 Synthesis of non-solvated dialkylaluminum anion, $[\text{K}(\text{AlR}_2)]_2$ via mechanochemical method.¹³¹

immediate reaction with benzophenone in hexanes at -78°C . In contrast, the THF adduct $[(\text{C}_3\text{H}_5)_3\text{Al}(\text{THF})]$ required a longer reaction time of 10 minutes at room temperature, despite the larger steric bulk of $1,3-(\text{SiMe}_3)_2\text{C}_3\text{H}_3$ compared to C_3H_5 .¹²⁹

In 2018, the Hanusa group reported a transformation in unsolvated organotin complexes driven by mechanochemistry.¹³⁰ In this work, both unsolvated tris(allyl) stannate **56** and tetra(allyl)tin species **57** are synthesised by ball-milling. The Hanusa group first ball-milled potassium allyl salt **52** ($[1,3-(\text{SiMe}_3)_2\text{C}_3\text{H}_3]\text{K}$) with SnCl_2 in a 2:1 ratio for 5 minutes in a planetary mill, generating a partially hexane soluble brown powder, tris(allyl) stannate **56**, $\{\text{Sn}[1,3-(\text{SiMe}_3)_2\text{C}_3\text{H}_3]_3\text{K}\}_\infty$ (Fig. 25). To further prove the necessity of the mechanochemical method, the potassium allyl salt **52** and SnCl_2 was stirred in hexane with no reaction occurring. In addition, a ball-milling-triggered disproportionation reaction between **52** and SnCl_2 resulted in the formation of a novel unsolvated tetra(allyl)tin(IV) species, **57**. It is worth to mention that the tetra(allyl)tin(IV) species **57** will decompose in solution in 1 hour but is stable in solid-state for several months. Furthermore, during the synthesis of **57**, the stereochemical control of products can be realised without solvent, which shows possibility of stereochemical control during mechanochemical reaction. These

unsolvated complexes could serve as strong Lewis acid catalysts or be utilised in metal-exchange reactions.¹³⁰

In 2024, the Yamashita group reported the synthesis of a non-solvated dialkylaluminum anion, $[\text{K}(\text{AlR}_2)]_2$ (**60**) via the mechanochemical method (Fig. 26).¹³¹ **60** is an example of monovalent aluminium anionic complex, *i.e.*, an aluminyl anion, which is a recent topical area in p-block chemistry.¹³² The corresponding toluene adduct, $[\text{R}_2\text{Al}-\text{K}(\text{tol})_2]$ (**60-tol**), was reported by the same group in 2020,¹³³ which is highly reactive but quite unstable. Though **60-tol** was characterised by SCXRD and UV-vis spectroscopy,¹³³ its instability prevented its characterisation using X-ray photoelectron spectroscopy (XPS): a nuclear-specific characterisation to elucidate oxidation state. Unlike its toluene adduct, **60** is stable at room temperature, which allowed collecting its Al 2p XPS spectra, unequivocally proving its Al(I) oxidation state and the Al 2p-electron binding energies.

5. Conclusion and outlook

As a closing remark, we summarise a few situations where we believe that mechanochemical methods could offer unique synthetic advantages:



(1) The reactants/products are highly reactive and react with solvent molecules;

(2) One or more reactants are insoluble, resulting in slow or no reaction and which causes the stoichiometric ratio to deviate from the designed value, and lead to unwanted side reactions;

(3) A non-solvated product is desired, but the reaction has to be conducted in coordinative solvents.

Though not reported yet, we envisage that the mechanochemical methods could also be used in the following directions:

(1) Trap highly unstable intermediate. A unique benefit of mechanochemistry is, once the ball-milling stops, the reaction is “frozen” and allows a number of solid-state characterisations. Fuelled by recent advances in the instrument (precise temperature-control cryogenic ball mill is now commercially available), we are sure that in future, the low-temperature ball-milling method will allow coordination chemists to isolate and characterise more elusive species;

(2) *In situ* monitoring of mechanochemical reactions has been recently widely used in organic reactions.¹³⁴ But this invaluable tool has seen less use in coordination chemistry. The major spectroscopic handles currently used in the *in situ* studies are vibrational spectroscopies (IR and THz Raman¹³⁵), and powder X-ray diffraction.¹³⁶ It can be envisaged that more analytical probes will be developed. We are particularly interested in electron paramagnetic resonance (EPR) and metal-nuclear solid-state NMR – there are works underway in our group on these regards.

(3) Combining mechanochemistry with other energy inputs. While mechano-photochemistry¹³⁷ and mechano-piezoelectric chemistry¹³⁸ have been reported in organic reactions, these transformative tools have not been introduced into synthetic coordination chemistry yet. We are very interested in seeing how these tools will enable coordination chemists to achieve more and more impossibles—which we believe is only a matter of time.

Data availability

No primary research results, software or code have been included and no new data were generated or analysed as part of this review.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors thank the Leverhulme Trust (RPG-2022-231 and RPG-2023-159) and the University of Birmingham for their generous financial support.

References

- 1 For representative examples of THF fragmentation, see: (a) J. Clayden and S. A. Yasin, *New J. Chem.*, 2002, **26**, 191–192,

- DOI: [10.1039/B109604D](https://doi.org/10.1039/B109604D); (b) A. R. Kennedy, J. Klett, R. E. Mulvey and D. S. Wright, *Science*, 2009, **326**, 706–708, DOI: [10.1126/science.1178165](https://doi.org/10.1126/science.1178165); (c) R. E. Mulvey, V. L. Blair, W. Clegg, A. R. Kennedy, J. Klett and L. Russo, *Nat. Chem.*, 2010, **2**, 588–591, DOI: [10.1038/nchem.667](https://doi.org/10.1038/nchem.667); (d) E. Crosbie, P. García-Álvarez, A. R. Kennedy, J. Klett, R. E. Mulvey and S. D. Robertson, *Angew. Chem., Int. Ed.*, 2010, **49**, 9388–9391, DOI: [10.1002/anie.201005119](https://doi.org/10.1002/anie.201005119); (e) E. Tanuhadi, A. S. Bair, M. Johnson, P. Fontaine and J. Klosin, *Chem. Sci.*, 2025, **16**, 280–287, DOI: [10.1039/D4SC05983B](https://doi.org/10.1039/D4SC05983B).
- 2 (a) M. H. Holthausen, T. Mahdi, C. Schleppehorst, L. J. Hounjet, J. J. Weigand and D. W. Stephan, *Chem. Commun.*, 2014, **50**, 10038–10040, DOI: [10.1039/C4CC01922A](https://doi.org/10.1039/C4CC01922A); (b) J. Cornella, C. Zarate and R. Martin, *Chem. Soc. Rev.*, 2014, **43**, 8081–8097, DOI: [10.1039/C4CS00206G](https://doi.org/10.1039/C4CS00206G).
- 3 For recent representative examples, see: (a) D. E. Anderson, A. Tortajada and E. Hevia, *Angew. Chem., Int. Ed.*, 2023, **62**, e202218498, DOI: [10.1002/anie.202218498](https://doi.org/10.1002/anie.202218498); (b) N. Davison, P. G. Waddell, C. Dixon, C. Wills, T. J. Penfold and E. Lu, *Dalton Trans.*, 2022, **51**, 10707–10713, DOI: [10.1039/D1DT03532K](https://doi.org/10.1039/D1DT03532K).
- 4 For recent representative examples, see: (a) S. K. Thaakur, N. Roig, R. Monreal-Corona, J. Langer, M. Alonso and S. Harder, *Angew. Chem., Int. Ed.*, 2024, **63**, e202405229, DOI: [10.1002/anie.202405229](https://doi.org/10.1002/anie.202405229); (b) Y. Wang, Y. Zhang, J. Liang, B. Tan, C. Deng and W. Huang, *Chem. Sci.*, 2024, **15**, 8740–8749, DOI: [10.1039/D4SC02491E](https://doi.org/10.1039/D4SC02491E); (c) K. R. McClain, A. H. Vincent, A. Rajabi, D. X. Ngo, K. R. Meihaus, F. Furche, B. G. Harvey and J. R. Long, *J. Am. Chem. Soc.*, 2024, **146**, 32708–32716, DOI: [10.1021/jacs.4c12278](https://doi.org/10.1021/jacs.4c12278); (d) G. M. Richardson, T. Rajeshkumar, F. M. Burke, S. A. Cameron, B. D. Nicholls, J. E. Harvey, R. A. Keyzers, T. Butler, S. Granville, L. Liu, J. Langley, L. F. Lim, N. Cox, N. F. Chilton, J. Hicks, N. J. L. K. Davis, L. Maron and M. D. Anker, *Nat. Chem.*, 2025, **17**, 20–28, DOI: [10.1038/s41557-024-01688-6](https://doi.org/10.1038/s41557-024-01688-6).
- 5 For a recent example, see: M. Jakoobi, Y. Tian, R. Boulatov and A. G. Sergeev, *J. Am. Chem. Soc.*, 2019, **141**, 6048–6053, DOI: [10.1021/jacs.9b01562](https://doi.org/10.1021/jacs.9b01562).
- 6 (a) *n*-Hexane C–H activation: H. M. Hoyt, F. E. Michael and R. G. Bergman, *J. Am. Chem. Soc.*, 2004, **126**, 1018–1019, DOI: [10.1021/ja0385944](https://doi.org/10.1021/ja0385944); (b) iso-hexane C–H activation: J. K. Hoyano and W. A. G. Graham, *J. Am. Chem. Soc.*, 1982, **104**, 3723–3725, DOI: [10.1021/ja00377a032](https://doi.org/10.1021/ja00377a032).
- 7 A. H. Janowicz and R. G. Bergman, *J. Am. Chem. Soc.*, 1982, **104**, 352–354, DOI: [10.1021/ja00365a091](https://doi.org/10.1021/ja00365a091).
- 8 Even methane can undergo C–H activation with highly reactive coordination complexes, see a recent example: Y. Liu, W. Dong, Z. H. Li and H. Wang, *Chem*, 2021, **7**, 1843–1851, DOI: [10.1016/j.chempr.2021.03.010](https://doi.org/10.1016/j.chempr.2021.03.010).
- 9 J. A. Labinger and J. E. Bercaw, *Nature*, 2002, **417**, 507–514, DOI: [10.1038/417507a](https://doi.org/10.1038/417507a).
- 10 For a recent review of divalent lanthanide chemistry, see: S. Schäfer, S. Kaufmann, E. S. Rösch and P. W. Roesky,



- Chem. Soc. Rev.*, 2023, **52**, 4006–4045, DOI: [10.1039/D2CS00744D](https://doi.org/10.1039/D2CS00744D).
- 11 W. J. Evans, I. Bloom, W. E. Hunter and J. L. Atwood, *Organometallics*, 1985, **4**, 112–119, DOI: [10.1021/om00120a019](https://doi.org/10.1021/om00120a019).
- 12 W. J. Evans, L. A. Hughes and T. P. Hanusa, *J. Am. Chem. Soc.*, 1984, **106**, 4270–4272, DOI: [10.1021/ja00327a037](https://doi.org/10.1021/ja00327a037).
- 13 W. J. Evans, N. T. Allen, P. S. Workman and J. C. Meyer, *Inorg. Chem.*, 2003, **42**, 3097–3099, DOI: [10.1021/ic0300316](https://doi.org/10.1021/ic0300316).
- 14 Mechano-chemical reaction, in *IUPAC Compendium of Chemical Terminology, International Union of Pure and Applied Chemistry*, 2025, 5th edn, Online version 5.0.0, DOI: [10.1351/goldbook.MT07141](https://doi.org/10.1351/goldbook.MT07141).
- 15 L. E. Wenger and T. P. Hanusa, *Chem. Commun.*, 2023, **59**, 14210, DOI: [10.1039/d3cc04929a](https://doi.org/10.1039/d3cc04929a).
- 16 J. F. Reynes, F. Leon and F. García, *ACS Org. Inorg. Au*, 2024, **4**, 432–470, DOI: [10.1021/acsorginorgau.4c00001](https://doi.org/10.1021/acsorginorgau.4c00001).
- 17 M. Arrowsmith, H. Braunschweig, M. Celik, T. Dellermann, R. D. Dewhurst, W. C. Ewing, K. Hammond, T. Kramer, I. Krummenacher, J. Mies, K. Radacki and J. K. Schuster, *Nat. Chem.*, 2016, **8**, 890, DOI: [10.1038/nchem.2542](https://doi.org/10.1038/nchem.2542).
- 18 S. Yadav, S. Saha and S. S. Sen, *ChemCatChem*, 2016, **8**, 486, DOI: [10.1002/cctc.201501015](https://doi.org/10.1002/cctc.201501015).
- 19 A. Rajabi, R. Grotjahn, D. Rappoport and F. Furche, *Dalton Trans.*, 2024, **53**, 410, DOI: [10.1039/d3dt03221c](https://doi.org/10.1039/d3dt03221c).
- 20 N. Tsoureas and I. Vagiakos, *Inorganics*, 2024, **12**, 275, DOI: [10.3390/inorganics12110275](https://doi.org/10.3390/inorganics12110275).
- 21 R. Akhtar, K. Gaurav and S. Khan, *Chem. Soc. Rev.*, 2024, **53**, 6150, DOI: [10.1039/d4cs00101j](https://doi.org/10.1039/d4cs00101j).
- 22 B. Rösch and S. Harder, *Chem. Commun.*, 2021, **57**, 9354, DOI: [10.1039/D1CC04147A](https://doi.org/10.1039/D1CC04147A).
- 23 S. Petrie, *Aust. J. Chem.*, 2003, **56**, 259, DOI: [10.1071/CH03006](https://doi.org/10.1071/CH03006).
- 24 M. A. Anderson and L. M. Ziurys, *Astrophys. J., Lett.*, 1995, **439**, 25, DOI: [10.1086/187736](https://doi.org/10.1086/187736).
- 25 T. Kruczyński, N. Pushkarevsky, P. Henke, R. Kęppe, E. Baum, S. Konchenko, J. Pikies and H. Schnöckel, *Angew. Chem., Int. Ed.*, 2012, **51**, 9025, DOI: [10.1002/anie.201204997](https://doi.org/10.1002/anie.201204997).
- 26 L. B. Knight Jr, W. C. Easley, W. Weltner Jr and M. Wilson, *J. Chem. Phys.*, 1971, **54**, 322, DOI: [10.1063/1.1674610](https://doi.org/10.1063/1.1674610).
- 27 R. Rubino, J. M. Williamson and T. A. Miller, *J. Chem. Phys.*, 1995, **103**, 5964, DOI: [10.1063/1.470476](https://doi.org/10.1063/1.470476).
- 28 R. Köppe, P. Henke and H. Schnöckel, *Angew. Chem., Int. Ed.*, 2008, **47**, 8740, DOI: [10.1002/anie.200802960](https://doi.org/10.1002/anie.200802960).
- 29 S. P. Green, C. Jones and A. Stasch, *Science*, 2007, **318**, 1754, DOI: [10.1126/science.1150856](https://doi.org/10.1126/science.1150856).
- 30 A. Stasch and C. Jones, *Dalton Trans.*, 2011, **40**, 5659, DOI: [10.1039/c0dt01831g](https://doi.org/10.1039/c0dt01831g).
- 31 C. Jones, *Commun. Chem.*, 2020, **3**, 159, DOI: [10.1038/s42004-020-00408-8](https://doi.org/10.1038/s42004-020-00408-8).
- 32 S. J. Bonyhady, C. Jones, S. Nembenna, A. Stasch, A. J. Edwards and G. J. McIntyre, *Chem.–Eur. J.*, 2010, **16**, 938, DOI: [10.1002/chem.200902425](https://doi.org/10.1002/chem.200902425).
- 33 S. P. Green, C. Jones and A. Stasch, *Angew. Chem., Int. Ed.*, 2008, **47**, 9079, DOI: [10.1002/anie.200803960](https://doi.org/10.1002/anie.200803960).
- 34 B. Rösch, T. X. Gentner, J. Eyselien, A. Friedrich, J. Langer and S. Harder, *Chem. Commun.*, 2020, **56**, 11402, DOI: [10.1039/d0cc05200k](https://doi.org/10.1039/d0cc05200k).
- 35 J. Pan, L. Zhang, Y. He, Q. Yu and G. Tan, *Organometallics*, 2021, **40**, 3365, DOI: [10.1021/acs.organomet.1c00512](https://doi.org/10.1021/acs.organomet.1c00512).
- 36 (a) Q. Yu, L. Zhang, Y. He, J. Pan, H. Li, G. Bian, X. Chen and G. Tan, *Chem. Commun.*, 2021, **57**, 9268, DOI: [10.1039/D1CC03448K](https://doi.org/10.1039/D1CC03448K); (b) M. Janssen, T. Frederichs, M. Olaru, E. Lork, E. Hupf and J. Beckmann, *Science*, 2024, **385**, 318–321, DOI: [10.1126/science.adp496](https://doi.org/10.1126/science.adp496); (c) D. Wang, W. Chen, R. Sun, J.-L. Liu, B. Wang, S. Ye, G. Tan and S. Gao, *Chem*, 2025, **11**, 102392, DOI: [10.1016/j.chempr.2024.102392](https://doi.org/10.1016/j.chempr.2024.102392).
- 37 D. D. L. Jones, I. Douair, L. Maron and C. Jones, *Angew. Chem., Int. Ed.*, 2021, **60**, 7087, DOI: [10.1002/anie.202017126](https://doi.org/10.1002/anie.202017126).
- 38 L. Sun, G. Yuan, L. Gao, J. Yang, M. Chhowalla, M. H. Gharahcheshmeh, K. K. Gleason, Y. S. Choi, B. H. Hong and Z. Liu, *Nat. Rev. Methods Primers*, 2021, **1**, 5, DOI: [10.1038/s43586-020-00005-y](https://doi.org/10.1038/s43586-020-00005-y).
- 39 C. E. Knapp and C. J. Carmalt, *Chem. Soc. Rev.*, 2016, **45**, 1036, DOI: [10.1039/c5cs00651a](https://doi.org/10.1039/c5cs00651a).
- 40 D. Jędrzkiewicz, J. Mai, J. Langer, Z. Mathe, N. Patel, S. DeBeer and S. Harder, *Angew. Chem., Int. Ed.*, 2022, **61**, e202200511, DOI: [10.1002/anie.202200511](https://doi.org/10.1002/anie.202200511).
- 41 M. Gimferrer, S. Danés, E. Vos, C. B. Yildiz, I. Corral, A. Jana, P. Salvador and D. M. Andrada, *Chem. Sci.*, 2022, **13**, 6583, DOI: [10.1039/d2sc01401g](https://doi.org/10.1039/d2sc01401g).
- 42 D. Jędrzkiewicz, J. Langer and S. Harder, *Z. Anorg. Allg. Chem.*, 2022, **648**, e202200138, DOI: [10.1002/zaac.202200138](https://doi.org/10.1002/zaac.202200138).
- 43 J. Hicks, M. Juckel, A. Paparo, D. Dange and C. Jones, *Organometallics*, 2018, **37**, 4810, DOI: [10.1021/acs.organomet.8b00803](https://doi.org/10.1021/acs.organomet.8b00803).
- 44 S. Kriek, H. Görls, L. Yu, M. Reiher and M. Westerhausen, *J. Am. Chem. Soc.*, 2009, **131**, 2977, DOI: [10.1021/ja808524y](https://doi.org/10.1021/ja808524y).
- 45 W. Huang and P. L. Diaconescu, *Dalton Trans.*, 2015, **44**, 15360, DOI: [10.1039/c5dt02198g](https://doi.org/10.1039/c5dt02198g).
- 46 Y. Xie, H. F. Schaefer and E. D. Jemmis, *Chem. Phys. Lett.*, 2005, **402**, 414–421, DOI: [10.1016/j.cplett.2004.11.106](https://doi.org/10.1016/j.cplett.2004.11.106).
- 47 J. Mai, J. Maurer, J. Langer and S. Harder, *Nat. Synth.*, 2024, **3**, 368, DOI: [10.1038/s44160-023-00451-y](https://doi.org/10.1038/s44160-023-00451-y).
- 48 J. Mai, M. Morasch, D. Jędrzkiewicz, J. Langer, B. Rösch and S. Harder, *Angew. Chem., Int. Ed.*, 2023, **62**, e202212463, DOI: [10.1002/anie.202212463](https://doi.org/10.1002/anie.202212463).
- 49 A. W. J. Bowles, J. A. Quirk, Y. Liu, G. H. Morritt, M. Freitag, G. F. S. Whitehead, A. W. Woodward, A. Brookfield, C. A. P. Goodwin, D. Collison, F. Tuna, C. L. McMullin, J. A. Dawson, E. Lu and F. Ortu, *J. Am. Chem. Soc.*, 2024, **146**, 28914–28924, DOI: [10.1021/jacs.4c09408](https://doi.org/10.1021/jacs.4c09408).
- 50 J. L. Dye, *Science*, 1990, **247**, 663, DOI: [10.1126/science.247.4943.663](https://doi.org/10.1126/science.247.4943.663).
- 51 J. L. Dye, *Acc. Chem. Res.*, 2009, **42**, 1564, DOI: [10.1021/ar9000857](https://doi.org/10.1021/ar9000857).
- 52 C. Liu, S. A. Nikolaev, W. Ren and L. A. Burton, *J. Mater. Chem. C*, 2020, **8**, 10551, DOI: [10.1039/d0tc01165g](https://doi.org/10.1039/d0tc01165g).



- 53 T. Buttersack, P. E. Mason, R. S. McMullen, H. C. Schewe, T. Martinek, K. Brezina, M. Crhan, A. Gomez, D. Hein, G. Wartner, R. Seidel, H. Ali, S. Thürmer, O. Marsalek, B. Winter, S. E. Bradforth and P. Jungwirth, *Science*, 2020, 1086, DOI: [10.1126/science.aaz7607](https://doi.org/10.1126/science.aaz7607).
- 54 H. Hosono and M. Kitano, *Chem. Rev.*, 2021, **121**, 3121, DOI: [10.1021/acs.chemrev.0c01071](https://doi.org/10.1021/acs.chemrev.0c01071).
- 55 X. Zhang and G. Yang, *J. Phys. Chem. Lett.*, 2020, **11**, 3841, DOI: [10.1021/acs.jpcclett.0c00671](https://doi.org/10.1021/acs.jpcclett.0c00671).
- 56 A. Ellaboudy and J. L. Dye, *J. Am. Chem. Soc.*, 1983, **105**, 6490, DOI: [10.1021/ja00359a022](https://doi.org/10.1021/ja00359a022).
- 57 S. B. Dawes, D. L. Ward, R. H. Huang and J. L. Dye, *J. Am. Chem. Soc.*, 1986, **108**, 3534, DOI: [10.1021/ja00272a073](https://doi.org/10.1021/ja00272a073).
- 58 J. L. Dye, *Acc. Chem. Res.*, 2009, **42**, 1564, DOI: [10.1021/ar9000857](https://doi.org/10.1021/ar9000857).
- 59 M. Y. Redko, J. E. Jackson, R. H. Huang and J. L. Dye, *J. Am. Chem. Soc.*, 2005, **127**, 12416, DOI: [10.1021/ja053216f](https://doi.org/10.1021/ja053216f).
- 60 C. S. Day, C. D. Do, C. Odena, J. Benet-Buchholz, L. Xu, C. Foroutan-Nejad, K. H. Hopmann and R. Martin, *J. Am. Chem. Soc.*, 2022, **144**, 13109–13117, DOI: [10.1021/jacs.2c01807](https://doi.org/10.1021/jacs.2c01807).
- 61 W. Meng, J. Wang, X. Wang, W. Wang, X. Zhang, Y. Bandoe and Z. Cheng, *J. Mater. Chem. A*, 2024, **12**, 2583, DOI: [10.1039/d3ta06546d](https://doi.org/10.1039/d3ta06546d).
- 62 S. Matsuishi, Y. Toda, M. Miyakawa, K. Hayashi, T. Kamiya, M. Hirano, I. Tanaka and H. Hosono, *science*, 2003, **301**, 626, DOI: [10.1126/science.1083842](https://doi.org/10.1126/science.1083842).
- 63 Y. Lu, J. Li, T. Tada, Y. Toda, S. Ueda, T. Yokoyama, M. Kitano and H. Hosono, *J. Am. Chem. Soc.*, 2016, **138**, 3970, DOI: [10.1021/jacs.6b00124](https://doi.org/10.1021/jacs.6b00124).
- 64 N. Davison, J. A. Quirk, F. Tuna, D. Collison, C. L. McMullin, H. Michaels, G. H. Morritt, P. G. Waddell, J. A. Gould, M. Freitag, J. A. Dawson and E. Lu, *Chem*, 2023, **9**, 576, DOI: [10.1016/j.chempr.2022.11.006](https://doi.org/10.1016/j.chempr.2022.11.006).
- 65 N. Davison, P. G. Waddell and E. Lu, *J. Am. Chem. Soc.*, 2023, **145**, 17007, DOI: [10.1021/jacs.3c06066](https://doi.org/10.1021/jacs.3c06066).
- 66 A. W. J. Bowles, Y. Liu, M. P. Stevens, I. J. Vitorica-Yrezabal, C. L. McMullin and F. Ortu, *Chem.–Eur. J.*, 2023, **29**, e202301850, DOI: [10.1002/chem.202301850](https://doi.org/10.1002/chem.202301850).
- 67 M. Weller, T. Overton, J. Rourke, and F. Armstrong, *Inorganic Chemistry*, OUP, Oxford, 2018.
- 68 J. L. Dye, *J. Chem. Educ.*, 1977, **54**, 332, DOI: [10.1021/ed054p332](https://doi.org/10.1021/ed054p332).
- 69 J. L. Dye, *Philos. Trans. R. Soc., A*, 2015, **373**, 2037, DOI: [10.1098/rsta.2014.0174](https://doi.org/10.1098/rsta.2014.0174).
- 70 E. Clementi, A. D. McLean, D. L. Raimondi and M. Yoshimine, *Phys. Rev.*, 1964, **133**, A1274, DOI: [10.1103/PhysRev.133.A1274](https://doi.org/10.1103/PhysRev.133.A1274).
- 71 S. Matalon, S. Golden and M. Ottolenghi, *J. Phys. Chem.*, 1969, **73**, 3098, DOI: [10.1021/j100843a052](https://doi.org/10.1021/j100843a052).
- 72 S. Golden, C. Guttman and T. R. Tuttle Jr, *J. Am. Chem. Soc.*, 1965, **87**, 135, DOI: [10.1021/ja01079a036](https://doi.org/10.1021/ja01079a036).
- 73 J. L. Dye, C. W. Andrews and J. M. Ceraso, *J. Phys. Chem.*, 1975, **79**, 3076, DOI: [10.1021/j100593a058](https://doi.org/10.1021/j100593a058).
- 74 J. L. Dye, *Annu. Rev. Phys. Chem.*, 1987, **38**, 271, DOI: [10.1146/annurev.pc.38.100187.001415](https://doi.org/10.1146/annurev.pc.38.100187.001415).
- 75 S. G. Dale and E. R. Johnson, *Phys. Chem. Chem. Phys.*, 2017, **19**, 12816, DOI: [10.1039/C7CP00882A](https://doi.org/10.1039/C7CP00882A).
- 76 J. L. Dye, *J. Phys. Chem.*, 1980, **84**, 1084, DOI: [10.1021/j100447a003](https://doi.org/10.1021/j100447a003).
- 77 J. L. Dye, *Angew. Chem. Int. Ed. Engl.*, 1979, **18**, 587, DOI: [10.1002/anie.197905871](https://doi.org/10.1002/anie.197905871).
- 78 J. L. Dye, *J. Phys. Chem.*, 1984, **88**, 3842, DOI: [10.1021/j150661a031](https://doi.org/10.1021/j150661a031).
- 79 J. L. Dye, *Sci. Am.*, 1977, **237**, 92, DOI: [10.1038/scientificamerican0777-92](https://doi.org/10.1038/scientificamerican0777-92).
- 80 L. D. Le, D. I sea, B. V. Eck and J. L. Dye, *J. Phys. Chem.*, 1982, **86**, 7, DOI: [10.1021/j100390a004](https://doi.org/10.1021/j100390a004).
- 81 N. Davison, J. M. Hemingway, C. Wills, T. Stolar, P. G. Waddell, C. M. Dixon, L. Barron, J. A. Dawson and E. Lu, *Inorg. Chem.*, 2024, **63**, 15247, DOI: [10.1021/acs.inorgchem.4c02914](https://doi.org/10.1021/acs.inorgchem.4c02914).
- 82 E. O. Fischer and H. P. Hofmann, *Chem. Ber.*, 1959, **92**, 482–486, DOI: [10.1002/cber.19590920233](https://doi.org/10.1002/cber.19590920233).
- 83 M. del M. Conejo, R. Fernández, E. G. Puebla, Á. Monge, C. Ruiz and E. Carmona, *Angew. Chem., Int. Ed.*, 2000, **39**, 1949–1951, DOI: [10.1002/1521-3773\(20000602\)39:11<1949::AID-ANIE1949>3.0.CO;2-O](https://doi.org/10.1002/1521-3773(20000602)39:11<1949::AID-ANIE1949>3.0.CO;2-O).
- 84 R. F. Koby, N. D. Schley and T. P. Hanusa, *Angew. Chem., Int. Ed.*, 2021, **60**, 21174, DOI: [10.1002/anie.202107980](https://doi.org/10.1002/anie.202107980).
- 85 C. Patel, E. André-Joyaux, J. A. Leitch, X. M. Irujo-Labalde, F. Ibba, J. Struijs, M. A. Ellwanger, R. Paton, D. L. Browne, G. Pupo, S. Aldridge, M. A. Hayward and V. Gouverneur, *Science*, 2023, **381**, 302, DOI: [10.1126/science.adi1557](https://doi.org/10.1126/science.adi1557).
- 86 V. Gouverneur and K. Seppelt, *Chem. Rev.*, 2015, **115**, 563–565, DOI: [10.1021/cr500686k](https://doi.org/10.1021/cr500686k).
- 87 C. W. Scheele, *Kongl. Vetensk. Acad. Handl.*, 1771, **32**, 120.
- 88 J. Aigueperse, P. Mollard, D. Devilliers, M. Chemla, R. Faron, R. Romano, and J. P. Cuer, Fluorine compounds, inorganic, in *Ullmann's Encyclopedia of Industrial Chemistry*, Wiley-VCH, Weinheim, 2000.
- 89 T. Schlatzer, C. A. Goult, M. A. Hayward and V. Gouverneur, *J. Am. Chem. Soc.*, 2025, **147**(8), 6338–6342, DOI: [10.1021/jacs.4c16608](https://doi.org/10.1021/jacs.4c16608).
- 90 P. Lei, Y. Ding, X. Zhang, A. Adijiang, H. Li, Y. Ling and J. An, *Org. Lett.*, 2018, **20**, 3439, DOI: [10.1021/acs.orglett.8b00891](https://doi.org/10.1021/acs.orglett.8b00891).
- 91 K. Fujishiro, Y. Morinaka, Y. Ono, T. Tanaka, L. T. Scott, H. Ito and K. Itami, *J. Am. Chem. Soc.*, 2023, **145**, 8163, DOI: [10.1021/jacs.3c01185](https://doi.org/10.1021/jacs.3c01185).
- 92 A. C. Jones, J. A. Leitch, S. E. Raby-Buck and D. L. Browne, *Nat. Synth.*, 2022, **1**, 763, DOI: [10.1038/s44160-022-00106-4](https://doi.org/10.1038/s44160-022-00106-4).
- 93 W. I. Nicholson, J. L. Howard, G. Magri, A. C. Seastram, A. Khan, R. R. A. Bolt, L. C. Morrill, E. Richards and D. L. Browne, *Angew. Chem., Int. Ed.*, 2021, **60**, 23128, DOI: [10.1002/anie.202108752](https://doi.org/10.1002/anie.202108752).
- 94 (a) T. Friščić, A. V. Trask, W. Jones and W. D. S. Motherwel, *Angew. Chem., Int. Ed.*, 2006, **45**, 7546, DOI: [10.1002/anie.200603235](https://doi.org/10.1002/anie.200603235); (b) T. Friščić, A. V. Trask, W. D. S. Motherwel and W. Jones, *Cryst. Growth Des.*, 2008, **8**, 1605, DOI: [10.1021/cg700929e](https://doi.org/10.1021/cg700929e); (c) T. Friščić, S. L. Childs, S. A. A. Rizvi and W. Jones, *CrystEngComm*, 2009, **11**, 418, DOI: [10.1039/B815174A](https://doi.org/10.1039/B815174A).



- 95 A. J. Birch, *J. Chem. Soc.*, 1944, 430, DOI: [10.1039/JR9440000430](https://doi.org/10.1039/JR9440000430).
- 96 B. K. Peters, K. X. Rodriguez, S. H. Reisberg, S. B. Beil, Y. Kawamata, M. Collins, J. Starr, L. Chen, S. Udyavara, K. Klunder, T. J. Gorey, S. L. Anderson, M. Neurock, S. D. Minter and P. S. Baran, *Science*, 2019, **363**, 838, DOI: [10.1126/science.aav5606](https://doi.org/10.1126/science.aav5606).
- 97 J. P. Cole, D. Chen, M. Kudisch, R. M. Pearson, C. Lim and G. M. Miyake, *J. Am. Chem. Soc.*, 2020, **142**, 13573, DOI: [10.1021/jacs.0c05899](https://doi.org/10.1021/jacs.0c05899).
- 98 B. Thiele, O. Rieder, B. T. Golding, M. Müller and M. Boll, *J. Am. Chem. Soc.*, 2008, **130**, 14050, DOI: [10.1021/ja805091w](https://doi.org/10.1021/ja805091w).
- 99 J. Burrows, S. Kamo and K. Koide, *Science*, 2021, **374**, 741, DOI: [10.1126/science.abk3099](https://doi.org/10.1126/science.abk3099).
- 100 Y. Gao, K. Kubota and H. Ito, *Angew. Chem., Int. Ed.*, 2023, **62**, e202217723, DOI: [10.1002/anie.202217723](https://doi.org/10.1002/anie.202217723).
- 101 K. Kondo, K. Kubota and H. Ito, *Chem. Sci.*, 2024, **15**, 4452, DOI: [10.1039/d3sc06052g](https://doi.org/10.1039/d3sc06052g).
- 102 K. Kubota, Y. Fukuzawa, K. Kondo, Y. Gao and H. Ito, *Chem. Lett.*, 2024, **53**, 60, DOI: [10.1093/chemle/upae060](https://doi.org/10.1093/chemle/upae060).
- 103 J. V. Nallaparaju, R. Satsi, D. Merzhyievskiy, T. Jarg, R. Aav and D. G. Kananovich, *Angew. Chem., Int. Ed.*, 2024, **63**, e202319449, DOI: [10.1002/anie.202319449](https://doi.org/10.1002/anie.202319449).
- 104 N. Davison, T. Khatun, I. Arce-Garcia, J. A. Gould, J. A. Dawson and E. Lu, *Eur. J. Inorg. Chem.*, 2023, e202300344, DOI: [10.1002/ejic.202300344](https://doi.org/10.1002/ejic.202300344).
- 105 K. Kondo, K. Kubota and H. Ito, *Nat. Synth.*, 2025, DOI: [10.1038/s44160-025-00753-3](https://doi.org/10.1038/s44160-025-00753-3).
- 106 K. Kubota, K. Kondo, T. Seo and H. Ito, *Synlett*, 2022, **33**, 898, DOI: [10.1055/a-1748-3797](https://doi.org/10.1055/a-1748-3797).
- 107 R. Takahashi, T. Seo, K. Kubota and H. Ito, *ACS Catal.*, 2021, **11**, 14803, DOI: [10.1021/acscatal.1c03731](https://doi.org/10.1021/acscatal.1c03731).
- 108 T. Seo, K. Kubota and H. Ito, *J. Am. Chem. Soc.*, 2020, **142**, 9884, DOI: [10.1021/jacs.0c01739](https://doi.org/10.1021/jacs.0c01739).
- 109 K. Kubota, K. Kondo, T. Seo, M. Jin and H. Ito, *RSC Adv.*, 2023, **13**, 28652, DOI: [10.1039/d3ra05571j](https://doi.org/10.1039/d3ra05571j).
- 110 T. Seo, T. Ishiyama, K. Kubota and H. Ito, *Chem. Sci.*, 2019, **10**, 8202, DOI: [10.1039/c9sc02185j](https://doi.org/10.1039/c9sc02185j).
- 111 K. Kubota, T. Seo, K. Koide, Y. Hasegawa and H. Ito, *Nat. Commun.*, 2019, **10**, 111, DOI: [10.1038/s41467-018-08017-9](https://doi.org/10.1038/s41467-018-08017-9).
- 112 T. Seo, N. Toyoshima, K. Kubota and H. Ito, *J. Am. Chem. Soc.*, 2021, **143**, 6165, DOI: [10.1021/jacs.1c00906](https://doi.org/10.1021/jacs.1c00906).
- 113 T. Seo, K. Kubota and H. Ito, *J. Am. Chem. Soc.*, 2023, **145**, 6823, DOI: [10.1021/jacs.2c13543](https://doi.org/10.1021/jacs.2c13543).
- 114 F. Puccetti, C. Schumacher, H. Wotruba, J. G. Hernández and C. Bolm, *ACS Sustainable Chem. Eng.*, 2020, **8**, 7262–7266, DOI: [10.1021/acssuschemeng.0c02447](https://doi.org/10.1021/acssuschemeng.0c02447).
- 115 T. Guo, Z. Li, L. Bi, L. Fan and P. Zhang, *Tetrahedron*, 2022, **112**, 132752, DOI: [10.1016/j.tet.2022.132752](https://doi.org/10.1016/j.tet.2022.132752).
- 116 S. Chen, C. Fan, Z. Xu, M. Pei, J. Wang, J. Zhang, Y. Zhang, J. Li, J. Lu, C. Peng and X. Wei, *Nat. Commun.*, 2024, **15**, 769, DOI: [10.1038/s41467-024-44891-2](https://doi.org/10.1038/s41467-024-44891-2).
- 117 P. Ying, J. Yu and W. Su, *Adv. Synth. Catal.*, 2021, **363**, 1246–4271, DOI: [10.1002/adsc.202001245](https://doi.org/10.1002/adsc.202001245).
- 118 E. Lu, J. Chu, Y. Chen, M. V. Borzov and G. Li, *Chem. Commun.*, 2011, **47**, 743, DOI: [10.1039/C0CC03212C](https://doi.org/10.1039/C0CC03212C).
- 119 S. Harder and J. Langer, *Nat. Rev. Chem.*, 2023, **7**, 843, DOI: [10.1038/s41570-023-00548-0](https://doi.org/10.1038/s41570-023-00548-0).
- 120 M. Westerhausen, A. Koch, H. Gçrls and S. Krieck, *Chem.–Eur. J.*, 2017, **23**, 1456, DOI: [10.1002/chem.201603786](https://doi.org/10.1002/chem.201603786).
- 121 P. Gao, J. Jiang, Y. Fukuzawa, S. Maeda, K. Kubota and H. Ito, *RSC Mechanochem.*, 2024, **1**, 486, DOI: [10.1039/D4MR00067F](https://doi.org/10.1039/D4MR00067F).
- 122 K. Kubota, S. Kawamura, J. Jiang, S. Maeda and H. Ito, *Chem. Sci.*, 2024, **15**, 17453, DOI: [10.1039/d4sc05361c](https://doi.org/10.1039/d4sc05361c).
- 123 O. P. E. Townrow, C. Färber, U. Zenneck and S. Harder, *Angew. Chem., Int. Ed.*, 2023, **63**, e202318428, DOI: [10.1002/anie.202318428](https://doi.org/10.1002/anie.202318428).
- 124 J. M. Harrowfield, R. J. Hart and C. R. Whitaker, *Aust. J. Chem.*, 2001, **54**, 423, DOI: [10.1071/CH01166](https://doi.org/10.1071/CH01166).
- 125 J. V. Nallaparaju, T. Nikonovich, T. Jarg, D. Merzhyievskiy, R. Aav and D. G. Kananovich, *Angew. Chem., Int. Ed.*, 2023, **62**, e202305775, DOI: [10.1002/anie.202305775](https://doi.org/10.1002/anie.202305775).
- 126 D. W. Peters and R. G. Blair, *Faraday Discuss.*, 2014, **170**, 83, DOI: [10.1039/c3fd00157a](https://doi.org/10.1039/c3fd00157a).
- 127 L. Sian, A. Dall'Anese, A. Macchioni, L. Tensi, V. Busico, R. Cipullo, G. P. Goryunov, D. Uborsky, A. Z. Voskoboynikov, C. Ehm, L. Rocchigiani and C. Zuccaccia, *Organometallics*, 2022, **41**, 547–560, DOI: [10.1021/acs.organomet.1c00653](https://doi.org/10.1021/acs.organomet.1c00653).
- 128 N. R. Rightmire and T. P. Hanusa, *Dalton Trans.*, 2016, **45**, 2352, DOI: [10.1039/c5dt03866a](https://doi.org/10.1039/c5dt03866a).
- 129 N. R. Rightmire, T. P. Hanusa and A. L. Rheingold, *Organometallics*, 2014, **33**, 5952, DOI: [10.1021/om5009204](https://doi.org/10.1021/om5009204).
- 130 R. F. Koby, T. P. Hanusa and N. D. Schley, *J. Am. Chem. Soc.*, 2018, **140**, 15934, DOI: [10.1021/jacs.8b09862](https://doi.org/10.1021/jacs.8b09862).
- 131 S. Kurumada, R. Yamanashi, K. Sugita, K. Kubota, H. Ito, S. Ikemoto, C. Chen, T. Moriyama, S. Muratsugu, M. Tada, T. Koitaya, T. Ozaki and M. Yamashita, *Chem.–Eur. J.*, 2024, **30**, e202303073, DOI: [10.1002/chem.202303073](https://doi.org/10.1002/chem.202303073).
- 132 J. Hicks, P. Vasko, J. M. Goicoechea and S. Aldrich, *Angew. Chem., Int. Ed.*, 2020, **60**, 1702–1713, DOI: [10.1002/anie.202007530](https://doi.org/10.1002/anie.202007530).
- 133 S. Kurumada, S. Takamori and M. Yamashita, *Nat. Chem.*, 2020, **12**, 36–39, DOI: [10.1038/s41557-019-0365-z](https://doi.org/10.1038/s41557-019-0365-z).
- 134 A. L. Michalchuk and F. Emmerling, *Angew. Chem., Int. Ed.*, 2022, **61**, e202117270, DOI: [10.1002/anie.202117270](https://doi.org/10.1002/anie.202117270).
- 135 T. H. Borchers, F. Topić, M. Arhangelskis, M. Ferguson, C. B. Lennox, P. A. Julien and T. Friščić, *Chem*, 2025, **11**, 102319, DOI: [10.1016/j.chempr.2024.09.018](https://doi.org/10.1016/j.chempr.2024.09.018).
- 136 (a) T. Friščić, I. Halasz, P. J. Beldon, A. M. Belenguer, F. Adams, S. A. J. Kimber, V. Honkimäki and R. Dinnebier, *Nat. Chem.*, 2013, **5**, 66–73, DOI: [10.1038/nchem.1505](https://doi.org/10.1038/nchem.1505); (b) S. T. Emmerling, L. S. Germann, P. A. Julien, I. Moudrakovski, M. Etter, T. Friščić, R. E. Dinnebier and B. V. Lotsch, *Chem*, 2021, **7**, 1639–1652, DOI: [10.1016/j.chempr.2021.04.012](https://doi.org/10.1016/j.chempr.2021.04.012); (c) N. Y. Gugin, K. V. Yusenko, A. King, D. Al-Sabbagh, J. A. Villajos and F. Emerling, *Chem*, 2024, **10**, 3459–3473, DOI: [10.1016/j.chempr.2024.07.033](https://doi.org/10.1016/j.chempr.2024.07.033).
- 137 Two recent examples, see: (a) D. M. Baier, C. Spula, S. Fanenstich, S. Grätz and L. Borchardt, *Angew. Chem.*,



Int. Ed., 2023, **62**, e202218719, DOI: [10.1002/anie.202218719](https://doi.org/10.1002/anie.202218719); (b) X. Xin, J. Geng, D. Zhang, H. T. Ang, H. Wang, Y. Cheng, Y. Liu, R. W. Toh, J. Wu and H. Wang, *Nat. Synth.*, 2024, **4**, 177–187, DOI: [10.1038/s44160-024-00681-8](https://doi.org/10.1038/s44160-024-00681-8).

138 For representative examples, see: (a) K. Kubota, Y. Pang, A. Miura and H. Ito, *Science*, 2019, **366**, 1500–1504, DOI:

[10.1126/science.aay8224](https://doi.org/10.1126/science.aay8224); (b) X. Wang, X. Zhang, L. Xue, Q. Wang, F. You, L. Dai, J. Wu, S. Kramer and Z. Lian, *Angew. Chem., Int. Ed.*, 2023, **62**, e202307054, DOI: [10.1002/anie.202307054](https://doi.org/10.1002/anie.202307054), A review; (c) E. M. Marrero, C. J. Caprara, C. N. Gilbert, E. E. Blanco and R. G. Blair, *Faraday Discuss.*, 2023, **241**, 91–103, DOI: [10.1039/D2FD00084A](https://doi.org/10.1039/D2FD00084A).

