

RSC Advances




This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Hand held sensing kits for detecting nerve agents simulants

ARTICLE

BODIPY dyes functionalized with 2-(2-dimethylaminophenyl)ethanol moieties as selective OFF-ON fluorescent chemodosimeters for the nerve agents mimics DCNP and DFP

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x


www.rsc.org/

R. Gotor^a, P. Gaviña^a, Luis E. Ochando^b, K. Chulvi^a, A. Lorente^a, R. Martínez-Máñez^c and A. M. Costero^{a*}

Two OFF-ON fluorescent chemodosimeters based on a BODIPY core for the detection of nerve agents mimics have been synthesized. Their reactivity towards diethylcyanophosphonate (DCNP) and diisopropylfluorophosphate (DFP) has been tested in organic and aqueous phase. These chemodosimeters selectively detect the nerve agents with good LODs. The chemodosimeters hold their sensing properties on solid supports, allowing the preparation of a hand held sensing kit. The sensing in solid-liquid phase has been demonstrated. The X-Ray structure of compound **2** has been resolved.

Introduction

Sarin, Soman and Tabun are by far the most known chemical warfare among the nerve gases. Their extreme toxicity, together with their low cost and easy production makes these compounds a big concern for governments and civilians, especially in areas involved in war related conflicts. Chemically, nerve agents are organophosphorous derivatives with a good leaving group (see Scheme 1). These compounds irreversibly bind to a free serine residue of the acetylcholinesterase, inhibiting its enzymatic task¹.



Scheme 1. Chemical structures of nerve-agents Sarin, Soman, and Tabun, their simulants (DFP and DCNP) and some pesticides as potential interfering agents.

The resulting over-accumulation of acetylcholine in the synaptic junctions of the nerves has immediate effects such as muscle relaxation hindering, “hyperventilation” and death in the worst cases.

Monitoring these agents has been achieved by means of different technologies, such as the use of ion mobility spectroscopy², biosensors³, electrochemical methods⁴, microcantilevers⁵, photonic crystals⁶, optical-fiber arrays⁷, etc. but most of them show some limitations, such as low portability, high costs or the need of qualified personal to operate the devices.

In the last years, the use of colorimetric and/or fluorimetric chemosensors has gained importance. Colorimetric and/or fluorimetric chemosensors have proved to be a cost-effective method capable of performing quantitative analysis with widespread technologies, or even semiquantitative analysis with the “naked-eye”. Among the existing dosimeters for the nerve agent simulants, the transduction mechanism usually involves charge transfer (ICT) or photoinduced electron transfer (PET) processes, whereas the interaction with the analytes is performed by quaternization of nitrogen atoms⁸, phosphorylation of hydroximate groups⁹ or displacement assays¹⁰.


Our research group has recently explored the use of 2-(2-(dimethylamino)phenyl)ethanol moieties¹¹ in the selective detection of nervous gases. The detection mechanism starts with the phosphorylation of the hydroxyl group by the organophosphorous derivatives forming a phosphoester, which in a second step is displaced by the attack of the nitrogen atom to give a five membered ring quaternary ammonium salt.

ARTICLE

On the other hand, borondipyrromethene (BODIPY) fluorophores have gained a huge attention due to their remarkable optic properties, such strong extinction coefficients, high quantum yields of fluorescence and good photostability.¹² Herein we report the synthesis of two new fluorescent probes (**1** and **2**) based on 2-(2-dimethylaminophenyl)ethanol receptors connected to BODIPY fluorophores, and their use in the selective detection of the nerve agent simulants DCNP and DFP.

Results and discussion


The synthesized probes (**1** and **2**) are depicted in Scheme 2. It was expected that these chemodosimeters in polar solvents would have no fluorescence due to ICT processes involving the nonbonding electrons of the nitrogen atom.^{13,14} In the presence of nerve agents, the nitrogen atom should form the corresponding ammonium salt through a displacement reaction and thus an increase of the fluorescence was expected to occur.



Scheme 2. Chemical structures of the synthesized probes.


Synthesis of the probes

BODIPY-based probe **1** was synthesized in 63% yield by means of a Liebeskind-Srögl coupling between the boronic acid derivative **3** and thiomethylbodipy **4** in the presence of Cu(I)-2-thienylcarboxylate (CuTC), trifurylphosphine (TFP) and a catalytic amount of Pd₂(dba)₃¹⁴ (Scheme 3).




Scheme 3. Coupling reaction to form probe **1**

The synthesis of the boronic acid precursor **3** started with the reductive hydrogenation of the 2-(2-nitrophenyl)ethanol in the presence of formaldehyde to yield 2-(2-dimethylaminophenyl)ethanol **5**, followed by an aromatic bromination with NBS catalyzed with ammonium acetate to form **6**. Lithiation of this compound with two equivalents of BuLi followed by treatment with triethoxyborane and acidic water work-up gave rise to the building block **3** (Scheme 4).



Scheme 4. Synthetic procedures to form the boronic acid of sensing unit **3**.

On the other hand, the thiomethyl BODIPY derivative **4** was obtained by the condensation of pyrrole with thiophosgene, followed by methylation with MeI and reaction with BF₃·Et₂O following procedures described in the literature^{15a}. The tetramethylated thiomethyl BODIPY analogous was also obtained by using 2,4-dimethylpyrrole instead of pyrrole. Unfortunately, the cross-coupling between **3** and this derivative was not successful, probably due to steric hindrance. Probe **2** was synthesized by means of a Suzuki cross-coupling reaction between the boronic acid derivative **3** and the *meso*-(*p*-bromophenyl) BODIPY **7** in 86% yield (Scheme 5).



Scheme 5. Synthesis of probe **2**.

Compound **7** was prepared in a 19% yield by the condensation of 2,4-dimethylpyrrole with *p*-bromobenzaldehyde, followed by aromatization with DDQ and reaction with BF₃·Et₂O^{15b}.

X-Ray structure of compound **2**

Single dark red crystals of compound **2** were obtained by slow evaporation of acetonitrile solutions. X-ray diffraction analysis of several batches revealed a monoclinic P2₁/c crystal system, with two molecules in the asymmetric unit.

In general, the bond lengths and angles within the BODIPY core exhibit the same geometric parameters as those examples reported in the literature.¹⁶ Thus, the boron atom has the typical bond lengths (B-N 1.548 Å and B-F 1.394 Å) and the N1-B1-N2 and F1-B1-F2 angles indicate a tetrahedral BF₂N₂ configuration¹⁷. Moreover, the dihedral angle between the pyrrole rings is 2.16° indicating a high level of planarity of the BODIPY core. By contrast, the phenyl group at the *meso* position is almost perpendicular to the indacene plane with a dihedral angle of 81.53°. The two aromatic rings in the biphenyl moiety are also not coplanar (dihedral angle of 22.16°) following the trend observed in other related compounds¹⁴ (Figure 1). To our surprise, the nitrogen atom of the dimethylamino group shows a remarkable pyramidalization with average C19-N3-C19' bond angles of 112.8°. The C_{ar}-N3 bond length is 1.429 Å indicating that there is no conjugation

between the nitrogen lone pair and the π -system of the aromatic ring. Besides, the methyl groups of the dimethylaniline moiety are facing away from the *ortho* alkyl substituent, observation that has been previously spotted in other *o*-substituted anilines.¹⁸

The 2-ethanol moiety shows a high grade of disorder in one of the molecules of the asymmetric unit, with three different positions for the oxygen atom, one of them at the 50% of occupation and the other two at 25% each.




Figure 1. (Top) Asymmetric unit of compound **2**. Thermal ellipsoids indicate the atom positions at probability levels of 35%. (Bottom) Molecular packing in the unit cell of the crystal structure along the *b* axis. Hydrogen atoms have been omitted for clarity.

Spectroscopic properties

The spectroscopic properties of compounds **1** and **2** were evaluated in acetonitrile solution (10^{-6} M). Compound **1** shows an absorption band at 495 nm ($\epsilon = 61600 \text{ M}^{-1}\text{cm}^{-1}$) whereas compound **2**, in addition to the main absorption band at 498 nm ($\epsilon = 91400 \text{ M}^{-1}\text{cm}^{-1}$), shows two small bands at 240 and 290 nm, which can be attributed to the biphenyl moiety attached at *meso* position. The fluorescence emission spectra of these compounds in acetonitrile show weak emission bands at $\lambda_{\text{em}} = 515 \text{ nm}$ ($\lambda_{\text{exc}} = 470 \text{ nm}$) for **1** and $\lambda_{\text{em}} = 507 \text{ nm}$ ($\lambda_{\text{exc}} = 480 \text{ nm}$) for **2**. The fluorescence quantum yields (Φ_f) of the free species in acetonitrile are rather low (< 0.02 for **2** and < 0.001 for **1**).

The strong fluorescence quenching observed for these compounds presumably resides in an intramolecular charge transfer (ICT) from the donor dimethylaniline moiety to the acceptor BODIPY core. In fact, it is known for similar compounds that the initial locally excited state (LE), after solvent reorganization, leads to a ICT state which is highly stabilized in polar solvents such as acetonitrile¹³. BODIPYs showing an ICT processes use to have low quantum yields due to the non-radiative deactivation nature of the ICT processes. Moreover, in the case of **1**, the existence of a twisted intramolecular charge transfer (TICT) could be considered, as it has previously been observed in related compounds¹⁴.

Reducing the dimethylaniline electron-donating power (i.e: protonation or quaternization), would result in the cancellation of the ICT processes with the concomitant increase in the lifetime of the LE state thereby restoring the fluorescence emission¹³.

In fact, we could observe that after protonation with HClO_4 , compound **2** restored its fluorescence to emission quantum yields of $\Phi_f = 0.446$, while the quantum yield of $\mathbf{1} + \text{H}^+$ increased to $\Phi_f = 0.019$.

Table 1. Spectroscopic properties of the synthesized dyes, free and protonated. Concentration is 10^{-6} M in MeCN. Proton source is HClO_4 . $\lambda_{\text{exc}} = 470$ and 480 nm for **1** and **2** respectively. Φ_f was measured using *meso*-phenyl bodipy as standard.¹⁷ a_x = relative amplitude of species *x*.

Probe	λ_{abs} (nm)	λ_{em} (nm)	Φ_f	τ_1 (ps)	a_1 (%)	τ_2 (ps)	a_2 (%)
1	495	515	< 0.001	n.m. ^[a]			
$\mathbf{1} + \text{H}^+$	501	520	0.014	97			
2	498	507	0.019	76.3	99.3	2002	0.7
$\mathbf{2} + \text{H}^+$	499	508	0.446	643	16.3	2370	83.7

[a] not measured due to low emission

Detection experiments in solution

As a first approach, detection studies were carried in liquid phase using acetonitrile solutions (10^{-6} M) of compounds **1** and **2**. Both probes showed small changes in the UV spectrum after addition of different aliquots of DCNP or DFP. Only compound **2** showed a new UV band at 260 nm along with a decrease of the bands at 240 and 290 nm. This change on the UV region can be ascribed to a change in the dihedral angle of the biphenyl moiety^{19,20} (see Figure 2 for compound **2** and Figure S-1 for compound **1**). On the other hand, the addition of DCNP or DFP to acetonitrile solutions of **1-2** immediately gave rise to a large enhancement of the fluorescence emission.





Figure 2. UV-vis (left) and emission (right, $\lambda_{\text{exc}}=470$ nm) spectra of compound **2** in MeCN (1×10^{-6} M for emission, 1×10^{-5} M for UV-vis) titrated with DCNP (from 100 μM to 20 mM). Yellow: free ligand, red: ligand+DCNP 20 mM

The side products from the nerve agents decomposition by hydrolysis or other mechanisms (*i.e.*: formation of the pyrophosphate) include the formation of acids. Thus, in order to demonstrate that the previously described changes were due to the reaction with the simulant and not to a mere protonation process, buffered solutions of the probes were used in the detection experiments

Thus, 10^{-6} M solutions of sensors **1** and **2** in water:MeCN 3:1, buffered with 0.1M MES pH 5.5 were also studied¹¹. Addition of DCNP or DFP in the range of 100 μM to 20 mM also gave rise to a large enhancement of the corresponding emission bands.

The nature of the resulting product of the reaction of **1** or **2** with the simulants was studied by the synthesis of the expected product through a different pathway. Thus, cyclization reaction of **2** was carried out with one equivalent of tosyl chloride in acetonitrile in the presence of K_2CO_3 .



Scheme 6. Cyclization reaction of **2** in the presence of tosyl chloride to form compound **8**.

The isolated product was characterized by NMR and HRMS, and its UV-vis and emission spectrum ($\lambda_{\text{abs}}=499$ nm, $\lambda_{\text{em}}=509$ nm, $\Phi_f = 0.435$, $\tau_1 = 424$ ps, $\tau_2 = 2289$ ps with relative amplitudes of $a_1 = 14\%$ and $a_2 = 86\%$) perfectly matched with that resulting of **2** + DCNP (see supplementary information, figure S-17).

Kinetic studies

The kinetic of the reaction between compounds **1** and **2** and DCNP and DFP in buffered water:MeCN (3:1 v/v) solutions (pH 5.5, MES 0.1 M) was studied. For each solution, a battery of experiments was carried out using different amounts of simulant. The simulant was always in an excess to consider pseudo first-order conditions.

The intensity of the emission band was continuously monitored (520 nm and 510 nm for **1** and **2** respectively). The obtained data allowed us to determine the observed rate constants at different simulant concentrations (k_{obs}). Plotting k_{obs} vs the concentration of the simulant, let us evaluate the constant rates (k) and the corresponding half-life time ($t_{1/2} = \ln 2/k$) for the reaction of DCNP and DFP with **1** and **2**. Table 2 summarizes the results obtained. Moreover, as an example, figure 3 shows the increment in the emission intensity at 510 nm for probe **2** in the presence of 6, 12, 18, 24 mM of DCNP in buffered water/acetonitrile 3:1 v/v mixtures (for compound **1** see supplementary information, figure S-18).




Figure 3. Kinetic profile of the emission intensity at 510 nm of probe **2** after the addition of 6, 12, 18 and 24 mM DCNP. Probe **2** is 10^{-6} M in pH=5.5 buffered water/acetonitrile 3:1 v/v mixture. The inset shows the correlation between k_{obs} and concentration of DCNP.

Table 2. Observed reaction rates (k_{obs}) and constant rates (k) for the reaction of probes **1** and **2** with different concentration of nerve gas simulants DCNP and DFP. Sensors are 10^{-6} M in pH=5.5 buffered water/acetonitrile 3:1 v/v mixture

Probe	k_{obs}				k (10^{-3}s^{-1})	$t_{1/2}$ (s)
	6	12	18	24		
1	0.116	0.173	0.235	0.286	9.53	72
2	0.122	0.156	0.198	0.221	5.68	122
DCNP (mM)	6	12	18	24		

Limits of detection and interferents

Detection limits for compounds **1** and **2** are summarized in Table 3. For this determination, increasing amounts of the simulants were added to solutions containing probes **1** or **2**. The emission values for these determinations were measured two minutes after the addition of the simulant. The limit of detection was considered to be reached at concentrations of simulant that fulfilled the condition:

$$I_s = I_{\text{blank}} + 3\sigma_{\text{blank}}$$

where I_s is the sample emission intensity after 2 minutes of addition of the simulant, I_{blank} is the emission intensity mean of the blank and σ_{blank} is the standard deviation of the blank.

Table 3. Limits of detection (ppm) of DCNP and DFP for probes **1** and **2**, in MeCN and in buffered water:MeCN (3:1 v/v) solutions (pH 5.5, MES 0.1 mol dm⁻³)

(ppm)	DCNP	DFP	DCNP (water:MeCN)
1	5	3	54
2	7	8	4

Additionally, the reactivity of probes **1** and **2** towards other organophosphorous compounds (OP) and 4,4'-DDE, or 4,4'-DDD (see Scheme 1) was studied in buffered aqueous solutions (water/acetonitrile 3:1 v/v pH=5.5 MES 0.1M). Thus, 100 ppm of the interfering species were added to the probe solutions. In general, no increase of fluorescence emission was observed indicating that no reaction occurred in the presence of potential interferents that may be present in common environments (see Figure 4).




Figure 4. Relative (to DCNP) emission intensity of compounds **1** and **2** (measured at 520 and 510 nm respectively) with 1000 ppm_v of DCNP, OP compounds, gasoline and diesel.

Moreover, DCNP or DFP were added to the water/acetonitrile 3:1 v/v solutions of **1** and **2**, containing the tested interferents and, as expected, the detection event still was observed, indicating that these species do not ruin the sensing capabilities of the probes.

Sensing experiments over supporting material

In order to assess the possibility of using these chemosensors as on-field sensing devices, we decided to fabricate solid polymer strips doped with the probes **1** and **2**. The strips, which had no initial fluorescence, were dipped in distilled water contaminated with DCNP (from 2 to 2000 ppm). Clear enhancement of the fluorescence was observed by the naked eye from 20 to 2000 ppm for **2** (see figure 5).




Figure 5. Hydrogel coated polyethylene strips. Hydrogel was previously doped with compound **2**. The strips were dipped in distilled water contaminated with (from the left to the right): 0, 2, 20, 200, and 2000 ppm_v of DCNP. Values indicate the relative (to the rightmost strip) luminance of the measured squared areas. Measurement was carried out by meaning the CIE luminance of all the pixels inside the squared area with a custom-made image processing software. (Fluorescent changes are more easily appreciated on the real samples rather than on the paper)

Additionally, tap water was also used as solvent in the tests showing that this complicated matrix has no influence in the sensing properties of the system. After the test was performed, washing the strips with distilled water did not quench the fluorescence, indicating that the irreversible cyclization had occurred.

Gas phase sensing

As a test experiment, we hang the strips in the center of a round bottom flask, containing an air atmosphere. Then, this atmosphere was contaminated with 5 ppm of DCNP. In the period of 5 minutes, the strip was removed from the system, and a clear increase of the fluorescence emission could be observed under a hand held UV_{254nm} lamp. With this simple procedure, we demonstrated the possibility of detection of nerve agent simulants in the gas phase. At the moment, this part of the investigation is still in progress.

Experimental

General methods

All the synthetic manipulations were performed in a dry argon atmosphere using standard techniques. Compound **4** and **7** were prepared following the procedures described in the literature.^{15a, 15b}. Tetrahydrofuran was distilled over Na prior to use. The other materials were purchased and used as received. Silica gel 60 F254 (Merck) plates were used for TLC. ¹H and ¹³C NMR spectra were recorded using a Bruker DRX-500 spectrometer (500 MHz for ¹H and 126 MHz for ¹³C) and a Bruker Avance 400 MHz (400 MHz for ¹H and 100 MHz for ¹³C) with the deuterated solvent as the lock and residual solvent as the internal reference. HRMS were recorded using a Shimadzu QP5050A. Absorption spectra were recorded with a Shimadzu UV-2101PC spectrophotometer. Fluorescence spectra were carried out in a Varian Cary Eclipse fluorimeter.

Crystal structure data collection and refinement.

Intensity data were collected at 120(2) K with an Oxford Diffraction S Ultra diffractometer equipped with a Mo co-mounted ($\lambda = 0.71073 \text{ \AA}$) enhance X-ray sapphire CCD detector. Structures were solved by direct methods and refined by full-matrix least-squares using SHELX-97.^{20, 21} Molecular graphics of the structures (Fig. 1) were produced with Mercury and Ortep.²² Solution, refinement, geometrical calculations and graphics were performed with the WinGX package.²³ CCDC 982038 contain the supplementary crystallographic data for this paper.

Experimental procedures

Synthesis of compound 6. 2-(2-Nitrophenyl)ethanol (10 g, 61 mmol), 37 % aq formaldehyde (11.24 mL, 75.2 mmol), ethanol

(200 mL, 99 %), and Pd/C (500 mg, 10 %) were placed under an H₂ atmosphere at 60 PSI until the uptake of hydrogen ceased. After filtration through celite, the solvent was evaporated, and the residue was dissolved in EtOAc, and washed twice with water, and NaCl (sat.). The organic phase was dried using MgSO₄, and the solvent was evaporated to give 2-(2-(N,N-dimethylamino)phenyl)ethanol as crude oil. The oil (10.12 g, 61 mmol) and ammonium acetate (470 mg, 6.1 mmol) were dissolved in acetonitrile (300 mL) in a round bottom flask. Then, the mixture was cooled down to 0 °C, and a solution of N-bromosuccinimide (10.85 g, 61 mmol) in 10 ml of acetonitrile was added dropwise. After 1 h reaction, the solvent was evaporated and the mixture dissolved in EtOAc. The organic solvent was washed twice with 10% aq Na₂CO₃ and NaCl (sat.). The organic phase was dried with MgSO₄ and evaporated. The remaining oil was purified by silica column chromatography using EtOAc:hexane 4:6 as eluent to yield 10.61 g (71%) of **6** as brownish oil. R_f = 0.77 (EtOAc:Hexane 6:4). ¹H NMR (500 MHz, CDCl₃) δ 7.24 (dd, J = 8.5, 2.4 Hz, 1H), 7.21 (d, J = 2.3 Hz, 1H), 6.97 (d, J = 8.5 Hz, 1H), 3.75 (t, J = 5.7 Hz, 2H), 2.87 (t, J = 5.7 Hz, 2H), 2.59 (s, 6H). ¹³C NMR (125 MHz, CDCl₃) δ 151.36, 138.22, 133.64, 130.35, 121.77, 117.61, 77.00, 63.86, 44.81, 35.61. HR-MS: calcd for C₁₀H₁₄BrNO [M+H]⁺: 244.0337, found: 244.0341.

Synthesis of compound 3. Compound **6** (1.45 g, 5.94 mmol) was dissolved in dry THF under argon atmosphere and the solution was cooled to -78 °C. Then, 15.77 mL of 1.13 M BuLi (17.82 mmol) was slowly added. After 30 min, the reaction was warmed to 0 °C in an ice bath for 5 min, and then cooled again to -78 °C. At this point, triethylborane (2.55 ml, 14.84 mmol) was added. The reaction was allowed to warm up and was kept overnight at room temperature. Then, 75 ml of NH₄Cl 10% in water was added, and the mixture was stirred for other 5 min. The organic solvent was partially evaporated, the organic phase was extracted with dichloromethane (5x100ml) and the combined organic phases were dried over MgSO₄. The solvent was evaporated, and the solid was dissolved in the minimum amount of CHCl₃. A 1:1 mixture of EtOAc:hexane was slowly added in order to precipitate the boronic acid **3**. The compound was isolated by centrifugation (0.96 g, 77%). R_f = 0.59 (EtOAc:MeOH 9:1). ¹H NMR (500 MHz, DMSO-d₆) δ 7.81 (br. s, 2H), 7.62 (d, J = 1.2 Hz, 1H), 7.59 (dd, J = 8.0, 1.2 Hz, 1H), 7.06 (d, J = 7.5 Hz, 1H), 4.70 (s, 1H), 3.64 (t, J = 7.5 Hz, 2H), 2.82 (t, J = 7.5 Hz, 2H), 2.65 (s, 6H). ¹³C NMR (125 MHz, DMSO-d₆) δ 136.69, 133.44, 132.29, 118.49, 62.20, 45.16, 34.72. HR-MS: calcd for C₁₀H₁₇BNO₃ [M+H]⁺: 210.1296, found: 210.1301.

Synthesis of compound 1. Thiomethyl-bodipy **4** (36 mg, 0.153 mmol) and boronic acid **3** (96 mg, 0.459 mmol) were dissolved in 6 ml of dry THF. The solvent was purged with argon for 10 min. At this point, Pd₂dba₃ (3.5 mg, 3.8 μmol, 2.5 mol%), tri(2-furyl)phosphine (2.6 mg, 11.4 μmol, 7.5 mol%) and copper(I) thiophene-2- carboxylate (174 mg, 0.918 mmol, 6 equiv) were added followed by 5 min N₂-purge. Then, the mixture was heated at reflux for 24 h. After this time, TLC plate revealed quantitative consumption of starting materials. Then, solvent was evaporated and the mixture was dissolved in dichloromethane, and washed twice with NaHCO₃ 10% and NaCl(sat.). The organic layer was dried with MgSO₄ and the solvent was evaporated. The remaining mixture was purified by aluminium oxide column chromatography using dichloromethane as eluent to yield 48 mg (84%) of an orange

solid. R_f = 0.42 (dichloromethane, aluminum oxide). ¹H NMR (500 MHz, CD₂Cl₂) δ 7.93 (s, 2H), 7.53 – 7.50 (m, 2H), 7.32 (dd, J = 7.6, 1.0 Hz, 1H), 7.09 (d, J = 4.2 Hz, 2H), 6.62 (d, J = 4.2 Hz, 2H), 3.92 (t, J = 6.0 Hz, 2H), 3.09 (t, J = 6.1 Hz, 2H), 2.85 (s, 6H). ¹³C NMR (125 MHz, CD₂Cl₂) δ 155.81, 143.20, 134.95, 134.78, 134.53, 133.49, 131.48, 130.30, 129.02, 119.55, 118.25, 63.49, 44.43, 35.44. HR-MS: C₁₉H₂₁BN₃OF₂ [M+H]⁺: 356.1740, found: 356.1749.

Synthesis of compound 2. *meso*-bromophenyl-bodipy **7** (96.3 mg, 0.239 mmol), boronic acid **3** (150 mg, 0.718 mmol) and sodium carbonate (151 mg, 1.44 mmol) were dissolved in 10 mL of a DFM/water (4/1 v/v) mixture. The solvent was purged with argon for 10 min.

Tetrakis(triphenylphosphine)palladium(0) (13.7 mg, 12 μmol) was added, and the mixture was degassed for 5 more minutes. Then, the mixture was heated to 100 °C and allowed to react for 1 h. After this time, TLC plate revealed quantitative consumption of starting materials. Solvents were then evaporated and the mixture was dissolved in dichloromethane. The organic solvent was washed twice with water, and the aqueous layer was extracted with dichloromethane. The combined organic layers were washed with NaCl(sat.) and dried with MgSO₄. After solvent evaporation, the remaining mixture was purified by silica column chromatography using EtOAc:Hexane 7:3 as eluent to yield 99 mg (86%) of an orange solid. R_f = 0.4 (EtOAc:Hexane 4:6). ¹H NMR (500 MHz, CD₂Cl₂) δ 7.78 (d, J = 8.2 Hz, 2H), 7.60 (dd, J = 8.2, 2.4 Hz, 1H), 7.56 (d, J = 2.3 Hz, 1H), 7.39 (d, J = 8.2 Hz, 2H), 7.34 (d, J = 8.3 Hz, 1H), 6.08 (s, 2H), 3.91 (t, J = 5.8 Hz, 2H), 3.11 (t, J = 5.8 Hz, 2H), 2.78 (s, 6H), 2.57 (s, 6H), 1.52 (s, 6H). ¹³C NMR (125 MHz, CD₂Cl₂) δ 155.35, 152.50, 143.38, 141.88, 141.21, 136.42, 136.10, 133.46, 131.44, 129.52, 128.49, 127.31, 125.93, 121.12, 120.40, 64.05, 44.76, 36.06, 14.33. HR-MS: calcd for C₂₉H₃₃BN₃OF₂ [M+H]⁺: 488.2679, found: 488.2685.

Synthesis of compound 8. Compound **2** (29.2 mg, 59.9 μmol), tosyl chloride (13.7 mg, 71.8 μmol) and potassium carbonate (24.8 mg, 180 μmol) were placed in a round bottom flask and water traces were evaporated twice with 5 ml of toluene. Then, the mixture was dissolved in 5 ml of dry acetonitrile. The mixture was allowed to react overnight at room temperature. At this point, the orange precipitate was filtered off and washed with cold Et₂O, to yield 35.2 mg (92 %) of the desired product. ¹H NMR (500 MHz, CDCl₃) δ 7.99 (dd, J = 8.5, 2.1 Hz, 1H), 7.82 (d, J = 8.1 Hz, 2H), 7.74 – 7.67 (m, 4H), 7.43 (d, J = 8.1 Hz, 2H), 7.17 (d, J = 7.9 Hz, 2H), 6.02 (s, 2H), 4.57 (t, J = 7.2 Hz, 2H), 3.88 (t, J = 1.8 Hz, 6H), 3.49 (t, J = 7.3 Hz, 2H), 2.59 (s, 6H), 2.35 (s, 3H), 1.45 (s, 6H). ¹³C NMR (125 MHz, CDCl₃) δ 155.86, 146.70, 144.00, 143.56, 142.84, 140.67, 139.77, 139.19, 135.39, 133.54, 131.33, 129.03, 128.80, 128.63, 127.83, 125.94, 125.09, 121.40, 118.15, 68.59, 55.23, 27.33, 21.23. HR-MS: calcd for C₂₉H₃₀BN₃F₂ [M-TsO]⁺: 470.2574, found: 470.2578.

Synthesis of 1,3,5,7-tetramethyl-8-thiomethyl-BODIPY. The synthesis of the tetramethylated analog of compound **4** was carried out according to reference 24. The NMR data are in agreement with those previously reported. ¹H NMR (500 MHz, CDCl₃) δ 6.09 (s, 2H), 2.61 (s, 6H), 2.52 (s, 6H), 2.46 (s, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 155.32, 143.46, 141.68, 134.22, 122.15, 22.17, 16.90, 14.65. HR-MS: calcd for C₁₄H₁₈N₂F₂SB [M+H]⁺: 295.1246, found: 295.1243.

Preparation of the strips. 2.0 g of poliuretane based hydrogel solution (Hydromed D4®) was mixed by 1h rotation with 27.5 mL of EtOH and 2.4 mL of water. To 2.5 mL of the viscous solution, 500 uL of a 4.3 mM dioxane solution of probes **1** or **2** was added, and the resulting mixture was again rotated for an additional hour. Polyethylene strips (40 x 5 x 0.5 mm) were then dipped once in the solutions and then allowed to dry at room temperature for 2 hours

Conclusions

Two OFF-ON BODIPY-based probes have been prepared and their utility in the detection of nerve agent simulant (DCNP and CFP) has been demonstrated. The dimethylamino groups present in compounds **1** and **2** inhibit the typical fluorescence of the BODIPY core probably due to an ICT mechanism. In the presence of the target species, the fluorescence of the system is restored through the quaternization of the dimethylamino nitrogen atom. The intensity of the fluorescence is stronger in compound **2** than in **1**. Registered LOD were in the 50 ppm, and in the case of **1** with DFP, went down to 3 ppm. In addition, selectivity towards different potential interferents present in civilian or military settings has been demonstrated. The properties of the probes are still present in solid supported kits which are able to detect the studied simulant not only in solution but also in gas phase. The achieved LOD are even lower when these solid supports are used.

Acknowledgements

We thank the Spanish Government and the European FEDER funds (project MAT2012-38429-C04-02 and 01). SCSIE and ICMOL (Universidad de Valencia) are gratefully acknowledged for all the equipment employed. R.G. is grateful to the Spanish Ministry of Education for its FPU grant. Dr. Knut Rurack is gratefully acknowledged for instrumental support and fruitful discussion.

Notes and references

^aCentro de Reconocimiento Molecular y Desarrollo Tecnológico (IDM), Universidad de Valencia, Doctor Moliner 50 46100 Burjassot, Valencia (Spain) Fax: (+34) 963543831. E-mail: Ana.Costero@uv.es

^bCentro de Reconocimiento Molecular y Desarrollo Tecnológico, Unidad Mixta Universidad de Valencia - Universidad Politécnica de Valencia, Departamento de Geología, Facultad de Ciencias Biológicas, Universidad de Valencia, Doctor Moliner 50, 46100 Burjassot, Valencia, Spain

^cCentro de Reconocimiento Molecular y Desarrollo Tecnológico, Unidad Mixta Universidad Politécnica de Valencia - Universidad de Valencia Departamento de Química, Universidad Politécnica de Valencia Camino de Vera s/n, 46022, Valencia (Spain)

1. S. M. Somani, *Chemical Warfare Agents*, Academic Press, San Diego, 1992.
2. E. Steiner, S. J. Klopsch, W. A. English, B. H. Clowers and H. H. Hill, *Anal. Chem.*, 2005, **77**, 4792.
3. a) L. M. Eubanks, T. J. Dickerson and K. D. Janda, *Chem. Soc. Rev.*, 2007, **36**, 458; b) J. J. Gooding, *Anal. Chim. Acta*, 2006, **559**, 137; c) M. T. McBride, S. Gammon, M. Pitesky, T. W. O'Brien, T. Smith, J. Aldrich, R. G. Langlois, B. Colston and K. S. Venkateswaran, *Anal. Chem.*, 2003, **75**, 1924; d) A. J.

- Russell, J. A. Berberich, G. F. Drevon and R. R. Koepsel, *Annu. Rev. Biomed. Eng.*, 2003, **5**, 1; e) H. Wang, J. Wang, D. Choi, Z. Tang, H. Wu and Y. Lin, *Biosens. Bioelectron.*, 2009, **24**, 2377; f) A. Mulchandani, I. Kaneva and W. Chen, *Anal. Chem.*, 1998, **70**, 5042; g) S. Munchaldani, P. Mulchandani, I. Kaneva and W. Chen, *Anal. Chem.*, 1998, **70**, 4140.
4. a) M. A. K. Khan, Y.-T. Long, G. Schatte and H.-B. Kraatz, *Anal. Chem.*, 2007, **79**, 2877; b) O. V. Shulga and C. Palmer, *Anal. Bioanal. Chem.*, 2006, **385**, 1116; c) J.-C. Chen, J.-L. Shih, C.-H. Liu, M.-Y. Kuo and J.-M. Zen, *Anal. Chem.*, 2006, **78**, 3752; d) G. Liu and Y. Lin, *Anal. Chem.*, 2006, **78**, 835; e) K. A. Joshi, M. Prouza, M. Kum, J. Wang, J. Tang, R. Haddon, W. Chen and A. Mulchandani, *Anal. Chem.*, 2006, **78**, 331; f) G. Liu and Y. Lin, *Anal. Chem.*, 2005, **77**, 5894; g) J. Wang, L. Chen, S. Mulchandani, P. Mulchandani and W. Chen, *Electroanalysis*, 1999, **11**, 866.
5. a) G. Zuo, X. Li, P. Li, T. Yang, Y. Wang, Z. Chen and S. Feng, *Anal. Chim. Acta*, 2006, **580**, 123; b) C. Karnati, H. Du, H. -F. Ji, X. Xu and Y. Lvov, A. Mulchandani, P. Mulchandani, W. Chen, *Biosensors Bioelec.*, 2007, **22**, 2636; c) Q. Zhao, Q. Zhu, W. Y. Shih and W. -H. Shih, *Sens. Actuators B*, 2006, **117**, 74.
6. a) W. He, Z. Liu, X. Du, Y. Jiang and D. Xiao, *Talanta*, 2008, **76**, 698; J. P. Walker, K. W. Kimble and S. A. Asher, *Anal. Bioanal. Chem.*, 2007, **389**, 2115; b) J. P. Walker and S. A. Asher, *Anal. Chem.*, 2005, **77**, 1596.
7. M. J. Aernecke and D. R. Walt, *Sens. Actuators B*, 2009, **142**, 464.
8. a) K. A. Van Houten, D. C. Heath and R. S. Pilato, *J. Am. Chem. Soc.*, 1998, **120**, 12359; b) S.-W. Zhang and T. M. Swager, *J. Am. Chem. Soc.*, 2003, **125**, 340; c) T. J. Dale and J. Rebek, Jr., *J. Am. Chem. Soc.*, 2006, **128**, 4500; d) A. M. Costero, S. Gil, M. Parra, P. M. E. Mancini, R. Martínez-Máñez, F. Sancenón and S. Royo, *Chem. Commun.*, 2008, 6002.
9. K. J. Wallace, J. Morey, V. M. Lynch and E. V. Anslyn, *New J. Chem.*, 2005, **29**, 1469.
10. D. Knapton, M. Burnworth, S. J. Rowan and C. Weder, *Angew. Chem., Int. Ed.*, 2006, **45**, 5825.
11. A. M. Costero, M. Parra, S. Gil, R. Gotor, P. M. E. Mancini, R. Martínez-Máñez, F. Sancenón and S. Royo, *Chem. Asian J.*, 2010, **5**, 1573.
12. a) G. Ulrich, R. Ziessel and A. Harriman, *Angew. Chem. Int. Ed. Engl.*, 2008, **47**, 1184; b) A. Loudet and K. Burgess, *Chem. Rev.*, 2007, **107**, 4891; c) R. Ziessel, G. Ulrich and A. Harriman, *New. J. Chem.*, 2007, **31**, 496.
13. a) M. Kollmannsberger, K. Rurack, U. Resch-Genger and J. Daub, *J. Phys. Chem. A*, 1998, **102**, 10211; b) W. Qin, M. Baruah, M. Van der Auweraer, F. C. De Schryver and N. Boens, *J. Phys. Chem. A*, 2005, **109**, 7371.
14. E. Lager, J. Liu, A. Aguilar-Aguilar, B. Y. Tang and E. Pena-Cabrera, *J. Org. Chem.*, 2009, **74**, 2053.
15. a) E. Pena-Cabrera, A. Aguilar-Aguilar, M. Gonzalez-Dominguez, E. Lager, R. Zamudio-Vazquez, J. Godoy-Vargas and F. Villanueva-Garcia, *Org. Lett.*, 2007, **9**, 3985. b) A. Nierth, A. Y. Kobitski, G. U. Nienhaus and A. Jäschke, *J. Am. Chem. Soc.*, 2010, **132**, 2646-2654
16. a) J.-S. Lu, S.-B. Ko, N. R. Walters and S. Wang, *Org. Lett.*, 2012, **14**, 5660; b) H. Lu, Q. Wang, L. Gai, Z. Li, Y. Deng, X. Xiao, G. Lai and Z. Shen, *Chem. Eur. J.*, 2012, **18**, 7852.
17. Z. Shen, H. Röhr, K. Rurack, H. Uno, M. Spieles, B. Schulz, G. Reck and N. Ono, *Chem Eur. J.*, 2004, **10**, 4853.
18. a) J.-Y. Shin, B. O. Patrick and D. Dolphin, *Tet. Lett.*, 2008, **49**, 5515; b) I.G. Voigt-Martin, Z. X. Zhang, D. H. Yan, A.

- Yakimanski, R. Matschiner, P. Krämer, C. Grania, D. Schollmeyer, R. Wortmann and N. Detzer, *Colloid Polym Sci.*, 1997, **275**, 18.
19. A. M. Costero, J. Sanchis, S. Gil, V. Sanz and J. A. G. Williams, *J. Mat. Chem.*, 2005, **15**, 2848.
 20. G. M. Sheldrick, SHELX97, Programs for Crystal Structure Analysis (Release 97–2). University of Göttingen, Germany, 1997.
 21. G. M. Sheldrick, *Acta Crystallogr., Sect. A*, 2008, **64**, 112.
 22. a) C. F. Macrae, I. J. Bruno, J. A. Chisholm, P. R. Edgington, P. McCabe, E. Pidcock, L. Rodriguez-Monge, R. Taylor, J. van de Streek, P. A. Wood, *J. Appl. Crystallogr.*, 2008, **41**, 466. b) L. J. Farrugia, *J. Appl. Crystallogr.*, 1997, **30**, 565.
 23. L. J. Farrugia, *J. Appl. Crystallogr.* 1999, **32**, 837.
 24. C. A. Osorio-Martínez, A. Urías-Benavides, C. F. A. Gómez-Durán, J. Bañuelos, I. Esnal, I. L. Arbeloa, and E. Peña-Cabrera, *J. Org. Chem.*, 2012, **77**, 5434.