


Analyst

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.


1
2
3 **Real-world carbon nanoparticle exposures induce brain and gonadal alterations in**
4 **zebrafish (*Danio rerio*) as determined by biospectroscopy techniques**
5
6

7
8 *Junyi Li^a, Guangguo Ying^b, Kevin C. Jones^a and Francis L. Martin^{a*}*
9

10
11 ^aCentre for Biophotonics, Lancaster Environment Centre, Lancaster University, Lancaster
12 LA1 4YQ, UK; ^bState Key Laboratory of Organic Geochemistry, Guangzhou Institute of
13 Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China
14
15
16
17
18
19
20
21
22

23 **Corresponding author* email: f.martin@lancaster.ac.uk; Tel: +44 (0)1524 510206
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ToC graphic



Differential tissue-specific effects of nanoparticles: biospectroscopy as a toxicity-sensing tool

Abstract

Carbon-based nanoparticles (CNPs) have emerged as novel man-made materials with diverse applications, which may present significant risks to organisms. To bridge the gap in our knowledge of nanotoxicology, a number of *in vitro* or *in vivo* studies have been carried out. However, toxicity data remains limited. Herein, we employed a biospectroscopy approach to assess CNP-induced effects in zebrafish (*Danio rerio*). Zebrafish were exposed to Fullerene (C₆₀), long or short multi-walled carbon nanotubes (MWCNTs), or single-walled carbon nanotubes (SWCNTs) for 21 days at two concentrations: 0.1 mg/L or 0.001 mg/L. Following exposure, the brain, gills, gonads and liver from zebrafish were interrogated by attenuated total reflection Fourier-transform infrared (ATR-FTIR) or Raman spectroscopy. Computational analysis was then applied to the acquired infrared (IR) spectra, and distinct biochemical segregations between the exposed tissues vs. control were observed with spectral biomarkers of alterations identified. In addition, lipid-to-protein ratios in all four tissues were calculated by the IR spectra; unsaturated lipid levels in brain and gonad were assessed by Raman spectroscopy. Marked lipid alterations were observed. These findings show that biospectroscopy approaches have the potential to detect CNP-induced biochemical alterations in zebrafish.

1. Introduction

Nanotechnology has introduced a wide range of man-made materials into the environment. Fullerene (C_{60}) and carbon nanotubes (CNTs) are the most promising manufactured carbon-based nanomaterials. With their unique physico-chemical properties, these materials offer extraordinary opportunities for applications in industry, biomedicine or everyday life.¹⁻⁵ Increasing interest commercially or scientifically is leading to massive production and application of these materials. However, little is known of their potential toxicity, or even how these materials will behave during the manufacturing process and in the environment.^{6,7} Consequently, this emerging issue is drawing more and more attention worldwide from scientists and governments.

Increasingly, numerous investigators are examining the toxic interactions of carbon nanoparticles (CNPs) with cells. Jia *et al.* observed that carbon nanomaterials (SWCNTs, MWCNTs or C_{60}) with different geometric structures induce varying levels of cytotoxicity in alveolar macrophages.⁸ Additional studies showed that CNPs cause cytotoxicity in HaCat human keratinocytes⁹, human osteoblasts and fibroblasts¹⁰, and human T-lymphocytes¹¹. Other biological models such as bacteria^{12,13}, algae¹⁴, *Daphnia magna*¹⁵ and, even, fish¹⁶ or rats¹⁷ have been employed to study such adverse effects. However, *in vitro* observations do not always faithfully extrapolate to the *in vivo* situation. Also, sometimes conventional assays are limited in their ability to assess nanoparticles.¹⁸ To date, the understanding of CNPs' toxicity remains limited and a paradigm-shifting methodology is required to offset such challenges.

Spectroscopy is increasingly used as a powerful tool in biological research. Infrared (IR) spectroscopy, including attenuated total reflection Fourier-transform infrared (ATR-FTIR) spectroscopy, has been applied in disease diagnosis¹⁹ and toxic assessment of

1
2
3 environmental contamination within cells.^{20, 21} In ATR-FTIR instrumentation, a mid-IR beam
4 ($\lambda=2-20 \mu\text{m}$) is transmitted through an internal reflection element (*e.g.*, diamond, zinc
5 selenide, germanium, silicon) in contact with the samples and penetrates a small distance
6
7 beyond the sample surface, allowing the generation of biochemical fingerprint spectra (1800-
8 900 cm^{-1}). Raman spectroscopy is a complimentary spectral method, which can also detect a
9
10 wide range of chemical bonds in cells and tissues.²² Some excellent studies have already
11
12 shown that Raman spectroscopy can also be applied to assess nanotoxicity.^{23, 24} Such
13
14 biospectroscopy tools employed in biological investigations may generate a large amount of
15
16 spectral data, which requires computational analysis in order to extract information.²⁵
17
18
19
20
21
22
23

24 Previously, we employed ATR-FTIR spectroscopy coupled with computational
25
26 analysis to detect CNP-induced alterations in MCF-7 cells; dose-related effects were
27
28 observed.²⁶ This suggests biospectroscopy as a novel tool capable of identifying the effects of
29
30 CNPs.²⁷ Aquatic environments may act as a sink for environmental contaminants including
31
32 CNPs,²⁸ and some studies have already reported that single-walled CNTs and C_{60} exert toxic
33
34 effects in rainbow trout (*Oncorhynchus mykiss*),²⁹ and that MWCNTs induce toxicity in
35
36 zebrafish embryos.³⁰ Thus, it is essential to develop sensitive and reliable methods to assess
37
38 biological effects of CNPs in fish. Additionally, zebrafish are a well-established model
39
40 organism widely applied in biological medicine³¹ and toxicological assessment.³² In order to
41
42 address the limitations of *in vitro* tests and extrapolate to the *in vivo* situation, we conducted
43
44 tests on zebrafish (*Danio rerio*) following exposure to CNPs and interrogated the tissues
45
46 using spectroscopy. In this study, zebrafish were exposed to four CNPs (C_{60} , short and long
47
48 MWCNTs or SWCNTs) at concentrations of 0.1 mg/L or 0.001 mg/L. Following 21-day
49
50 exposure, alterations in tissues including brain, gills, gonads and liver were then assessed *via*
51
52 biospectroscopy with computational analysis. Our aim was to ascertain real-world CNP
53
54 effects in a model system.
55
56
57
58
59
60

2. Materials and Methods

Chemicals and carbon nanoparticles

Bovine serum albumin (BSA) obtained from Sigma was $\geq 98\%$. All CNPs were purchased from Sigma. Short MWCNTs were $>90\%$ pure being 10-15 nm in diameter and 0.1-10 μm in length. Long MWCNTs were $>90\%$ pure also, but were 110-170 nm in diameter and 5-9 μm in length. C_{60} had a purity $>99.5\%$ and particle size of 1 nm. SWCNTs were described as CarboLex AP-grade (the purity of AP-grade products ranges from 50% to 70% by volume); major impurities are carbon nanospheres and carbon-encapsulated catalyst nanoparticles - the diameter was 1.2-1.5 nm. All CNPs were analysed by Raman spectroscopy (Renishaw PLC, UK) with a 785 nm laser, and determined to be of high purity. Additionally, images of CNPs were taken using a scanning electron microscope (SEM) [JSM 5600 (JEOL)] [see Electronic Supporting Information (ESI) Figure S1]. CNPs were dispersed in 1% BSA solution with a 15-min ultrasonication and stock solutions were made at concentrations of 100 mg/L and 1 mg/L. CNT solutions were stable and well-dispersed, while C_{60} appeared to agglomerate.

Fish maintenance and experimental conditions

All experiments were carried out following approval from the local Institutional Review Board. Zebrafish were maintained in the Aquatic Toxicology Laboratory at the Guangzhou Institute of Geochemistry, Chinese Academy of Sciences. All fish were kept in 50-L flow-through tanks filled with dechlorinated tap water in a temperature-controlled room maintained at $27\pm 1^\circ\text{C}$. The room was on a 14:10 h light:dark cycle, and fish were fed once a day with a quantity of commercial food at 5% of the wet weight.

Fish exposures were conducted in 10-L glass tanks, and each experimental tank contains 5 L dechlorinated tap water and 4 fish (2 males, 2 females). Prior to exposure,

1
2
3 zebrafish in 50-L tanks were randomly transferred to the experimental tank for a 7-day
4
5 adaptive period. Following this, CNP exposure was initiated and run for 21 days. There were
6
7 nine randomly assigned tanks for each exposure (control and treatment with one of four
8
9 CNPs at 0.1 mg/L or 0.001 mg/L). Exposure concentrations were chosen based on previous
10
11 *in vitro* studies,²⁶ which showed that speculated real-world environmental levels induced
12
13 alterations in exposed cell populations detectable by biospectroscopy techniques. To
14
15 minimize contamination, fish were only fed in the morning every day and in the afternoon,
16
17 each tank was cleaned to eliminate fish faeces and food remains by siphoning the water out of
18
19 the tanks. Then all tanks were filled with fresh water and treatment. All fish were terminated
20
21 at the end of the exposure and were sacrificed within seconds by immersion in melting ice
22
23 prior to body size measurements (weight and length) (see ESI Figure S2). From each fish,
24
25 gills, brain, liver and gonads were independently harvested and fixed in 70% ethanol.
26
27
28
29
30 Exposure experiments were conducted in triplicate.
31

32 **Biospectroscopy analysis**

33
34
35 Each fixed fish tissue for spectral interrogation was thinly sliced by hand and mounted on IR-
36
37 reflective low-E slides (Kevley Technologies, USA), allowed to air-dry and stored in a
38
39 desiccator prior to spectral acquisition. All tissue samples were interrogated using a Bruker
40
41 TENSOR 27 FTIR spectrometer (Bruker Optics Ltd., UK) equipped with a Helios ATR
42
43 attachment containing a diamond internal reflection element (IRE). Instrument parameters
44
45 were set at 32 scans, 16 cm⁻¹ resolution. For each slide, 10 IR spectra were acquired at
46
47 different points across the sample. Prior to starting a new slide, the ATR crystal was cleaned
48
49 with deionized water and a background taken. Additionally, the brain and gonad tissues from
50
51 the control group and the high-dose treatment groups for the four CNPs were further
52
53 interrogated by Raman spectroscopy. Raman spectra were recorded using an *InVia* Renishaw
54
55 Raman spectrometer (Renishaw plc, UK) equipped with a 785 nm streamline focus laser and
56
57
58
59
60

1
2
3 a Renishaw-automated 100 nm encoded XYZ stage. The system parameters consisted of 50
4 μm entrance slit, 830 lines/mm ($\sim 1.69\text{ cm}^{-1}$ spectral resolution) diffraction grating dispersing
5 Raman signal onto a master Renishaw Pelletier cooled charge detector (CCD, 1024 pixel \times
6 pixel). The system calibration was carried out using an internal Renishaw silicon calibration
7 source for wavenumber shift. A white light camera mounted on the microscope was used to
8 visualize the locations of the spectral acquisition. Spectra were collected using a Leica $\times 50$
9 objective lens (numeral aperture 0.75) at 50% laser power ($\leq 100\text{ mW}$ prior to lens) of 20-
10 second exposure and spectral range covering 2000-500 cm^{-1} . Approximately 30 spectra were
11 obtained from different sites from each sample.
12
13
14
15
16
17
18
19
20
21
22

23 **Data analysis**

24
25 All spectral data acquired from both ATR-FTIR and Raman spectroscopy were processed
26 using our IRootLab toolbox (<http://irootlab.googlecode.com>)³³ running on MATLAB r2010a
27 (The MathWorks, Inc., US). IR spectra were pre-processed as follows: cut to 1800-900 cm^{-1}
28 (the biochemical fingerprint range), rubberband baseline correction and normalization to the
29 Amide I peak (1650 cm^{-1}). Computational analysis using multivariate techniques including
30 principal component analysis (PCA) and linear discriminant analysis (LDA) can efficiently
31 analyse such large spectral datasets. Following pre-processing, cross-calculated PCA-LDA
32 was applied to the dataset; information was extracted and visualised as scores plots and
33 cluster vectors. PCA is an unsupervised technique employed to reduce the dimensions of the
34 data, and then the PCA scores are used as input for the subsequent LDA, where the class'
35 information contributes to differentiation of the derived clusters. To eliminate the risk of
36 LDA overfitting, "cross-calculation" was applied to the PCA-LDA scores (see ESI Figure
37 S3). It utilizes leave-one-out cross-validation to train the PCA-LDA loadings using $n-1$
38 (n =number of spectra in the dataset) samples and subsequently calculates the scores for the
39 remaining sample, repeating this process until all scores are calculated. The loadings are
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 derived as averages between the n PCA-LDA loadings model. Additionally, the number of
4
5 principal components (PCs) for PCA-LDA was calculated by classification with an
6
7 optimization procedure. The output data derived from PCA-LDA can be then visualized as 1-
8
9 D, 2-D or 3-D scatterplots (“scores plots”). In scores plots, nearness between two groups
10
11 means similarity, while increasing distance indicates dissimilarity. To reveal the biochemical
12
13 alterations associated with each group in the dataset, PCA-LDA cluster vectors were
14
15 developed.²⁵ To simplify the identification of the main biochemical alteration of each group,
16
17 cluster vectors plots were used to indicate the first eight highest peaks; tentative assignments
18
19 of the peak wavenumbers are listed (see ESI, pages 25-46), ranked by peak height.
20
21
22
23

24 **Lipid accumulation and unsaturated levels in tissues exposed to CNPs**

25
26 Lipid accumulation in tissues was calculated by measuring the lipid-to-protein ratio using the
27
28 intensity absorbance at two vibration modes derived from the pre-processed IR spectra. The
29
30 intensity absorbance at 1400 cm^{-1} corresponding to C=O symmetric stretching of amino acid
31
32 was used as a protein marker, while the intensity absorbance at 2924 cm^{-1} was associated
33
34 with $\nu_{\text{as}}\text{CH}_2$ for lipid. In Raman spectra, the intensity at 1445 cm^{-1} was assigned to CH_2
35
36 bending for lipid and the intensity at 1670 cm^{-1} was assigned to $\nu(\text{C}=\text{C})$ in lipid. The ratio of
37
38 $(\text{C}=\text{C})/(\text{CH}_2)$ can be used to calculate the lipid unsaturated level.³⁴
39
40
41
42

43 Repeated-measures one-way analysis of variance (ANOVA) with Dunnett’s post hoc
44
45 tests were used to examine whether the alteration of the tissues observed in scores plots (LD1
46
47 and LD2), and the ratios of lipid/protein and $(\text{C}=\text{C})/(\text{CH}_2)$ differed significantly between the
48
49 exposure vs. control groups. It’s not necessary to transform data to meet the underlying
50
51 assumption of homogeneity between categories and normality of residuals. *P-values* below
52
53 0.05 were considered statistically significant. All ANOVA tests were conducted in GraphPad
54
55 Prism 4 (GraphPad Software, USA).
56
57
58
59
60

Results and discussion

When zebrafish were exposed to one of the four CNPs tested, the response for each tissue examined was different and varied with exposure. CNP-specific dose responses compared to the control group were observed in all tissues examined, including brain, gills, gonads and liver (see ESI Figures S4-S7). Additionally, ANOVA tests showed that these were significant for the majority of exposures (see ESI Tables S1-S2). In the scores plots along the LD1 dimension, most category separation was observed; these simply show the effects for each tissue resulting from two different CNP exposures (Figures 1-2). It is clearly noted that along LD1, the gills and brain exhibit a similar response pattern to the four different CNPs. In contrast, the gonads and liver display an inverse response to these exposures, with the lower dose inducing more pronounced effects than the higher. On a physiologically-based toxicokinetics (PBTk) model, the gills are considered the initial site for uptake and elimination of CNPs, while brain, gonads and liver connected with the gills *via* arterial blood are perturbed by CNPs.³⁵ It is noted that in zebrafish, blood circulating in veins from the gonads will reach the liver, which may explain why these tissues respond similarly to CNPs. To highlight the main biochemical alterations induced by CNPs, a cluster vectors method was employed following cross-calculated PCA-LDA. CNP exposures generate a range of biochemical alterations associated with lipid, protein, glycogen and DNA/RNA (see ESI).

To reveal the profile of effects induced by CNPs, a dataset was developed to profile alterations in zebrafish tissues following CNP exposures at a concentration of 0.1 mg/L compared to the control group. Segregation in 2-D cross-calculated PCA-LDA scores plots showed that long MWCNTs, possessing a relatively larger size, exert the most pronounced alterations in the four tissues studied compared to the other three CNPs (Figure 3). The cluster vectors plots indicate that long MWCNTs-induced effects in tissues are mainly to lipid and protein, with limited alterations in the DNA/RNA region (Figure 4). However, C₆₀

1
2
3 appears to induce a high level of alterations in the DNA/RNA region, especially in the brain.
4
5 SWCNTs- and short MWCNTs-induced alterations in zebrafish appear to be similar. To
6
7 investigate the brain and gonads further, Raman spectroscopy was employed to interrogate
8
9 these two tissues; significant post-exposure alterations were again observed.
10

11
12 IR spectra allow an estimation of the lipid-to-protein ratio in tissues. This showed that
13
14 female fish display a higher lipid-to-protein ratio, except in the gonads. Among all tissue
15
16 types, male gonads exhibited the highest lipid-to-protein ratio (Figure 5). Additionally, the
17
18 unsaturated levels of lipid in brain and gonads were assessed by calculating the ratio of
19
20 (C=C)/(CH₂) in Raman spectra. It showed that there was a higher unsaturated lipid level in
21
22 male fish than females (Figure 6).
23
24

25
26 As the initial site of uptake and elimination of contamination, the gills would be
27
28 expected to directly interact with CNPs. Therefore, CNPs would exert effects by direct
29
30 physical injury and indirectly by generating reactive oxidative species (ROS) and free
31
32 radicals,³⁶ resulting in gill inflammation.^{37, 38} Exposure to CNPs induced significant
33
34 alterations in Amide I, Amide II, lipid and protein in both female and male fish. Additionally,
35
36 slight alterations in DNA/RNA region ($\nu_{as}PO_2^-$, ~1225 cm⁻¹; $\nu_sPO_2^-$, ~1080 cm⁻¹) were
37
38 observed as well. The lipid-to-protein ratio was most highly elevated in gills exposed to long
39
40 MWCNTs. Short MWCNTs and SWCNTs seemed to elicit a lower lipid-to-protein ratio in
41
42 gills, while C₆₀ showed no significant effect ($P > 0.05$). This suggests that the size of the
43
44 CNPs plays a major role in inducing inflammation in the gill, with larger sizes causing higher
45
46 levels of damage.³⁹ Though it was reported that CNPs would be precipitated on gill mucus,⁴⁰
47
48 CNPs could still penetrate the gill and be transported into the fish circulation. Even if tight
49
50 junctions between gill cells dramatically reduce the permeability to CNPs, inflammation
51
52 factors at the site could still facilitate their transported into the circulation to exert further
53
54 effects on other organs.
55
56
57
58
59
60

1
2
3 Post-exposure to CNPs, the brain also showed significant alterations. Both IR and
4
5 Raman spectra showed that long MWCNTs caused the most pronounced alterations in brain
6
7 for both genders. In contrast, there appeared to be no significant effect of long MWCNTs on
8
9 the lipid-to-protein ratio compared with control, while C₆₀ induced a significant elevation of
10
11 lipid-to-protein ratio, as well as short MWCNTs and single-walled CNTs. Because of their
12
13 larger size, it may be difficult for long MWCNTs to penetrate through the tight junctions
14
15 between the gill cells, which could also limit their ability to cross the blood-brain barrier
16
17 (BBB).⁴¹ However, with high levels of oxidative stress and even inflammation factors
18
19 induced by long MWCNTs, they still dramatically altered brain spectral signatures. Raman
20
21 spectra showed that male fish display higher lipid unsaturated levels than females, which
22
23 could contribute to the fluidity of the cell membrane. Interestingly, C₆₀ caused contrasting
24
25 effects on unsaturated lipid levels of brain in both genders. In female fish, the unsaturated
26
27 lipid level was increased, while it was decreased in males, both significantly. As the brain in
28
29 male fish has a lower lipid-to-protein ratio but higher unsaturated levels compared with that
30
31 to female fish, it means that the male brain contains less lipid, but possibly greater fluidity
32
33 due to an elevated content of unsaturated lipid.⁴² Thus, it is more likely that the highly
34
35 lipophilic C₆₀ will penetrate further into the lipid region,⁴³ and exert oxidative damage not
36
37 only to lipid, but also to further biochemical constituents such as protein and DNA/RNA.
38
39 However, the three CNTs resulted in a significant increase of unsaturated lipid levels in the
40
41 female brain, while only SWCNTs caused a significant elevation in males. Due to the lower
42
43 lipid-to-protein ratio in male brain compared to female, possibly implicating that the male has
44
45 a thinner layer of lipid, CNPs may exert a wider range of effects in cells. This was exhibited
46
47 by the IR spectra indicating a trend for CNPs to induce more effects on the DNA/RNA
48
49 spectral region derived from male brain compared to female. Profiled by both the IR and
50
51 Raman spectra, all CNPs widely and highly induced alterations in the protein and lipid
52
53
54
55
56
57
58
59
60

1
2
3 spectral region. Complimentary to IR, Raman spectra also indicated that CNPs caused
4
5 significant effects in the spectral range associated with S-S ($\sim 524\text{ cm}^{-1}$), C-S ($\sim 662\text{ cm}^{-1}$) in
6
7 protein and C=C in lipid ($\sim 1655\text{ cm}^{-1}$). It is accepted that ROS generation is a key pathway in
8
9 nanoparticle toxicity. High ROS levels could increase the depletion of thiol groups in proteins,
10
11 especially glutathione (GSH), and increase the oxidized forms such as GSSG.^{44, 45}
12
13

14
15 The interference of CNPs in brain could perturb the hypothalamic-pituitary-gonadal
16
17 axis and affect the gonads further, in addition to direct effects on these tissues. It is evidenced
18
19 by IR spectra that CNP exposures resulted in significant alterations in lipids, proteins and
20
21 DNA/RNA. Raman spectra also confirmed this and showed that CNPs could induce
22
23 alterations associated with C-S ($\sim 662\text{ cm}^{-1}$) in female fish rather than in males. With both
24
25 higher lipid-to-protein ratios and higher unsaturated lipid levels, male fish are more likely to
26
27 protect protein from CNP-induced oxidative stress through the antioxidant function at the
28
29 lipid region. These gonadal alterations would possibly further affect reproductive activity.
30
31
32

33
34 As an important organ for active metabolism and detoxification, the liver seems quite
35
36 sensitive to CNP exposure. IR spectra show that the most pronounced alterations induced by
37
38 C_{60} were associated with DNA/RNA region ($\nu_s\text{PO}_2^-$, $\sim 1080\text{ cm}^{-1}$) in both genders, followed
39
40 by alterations located in the protein and lipid spectral regions. Limited effects in DNA/RNA
41
42 caused from short MWCNTs and SWCNTs were observed as well. In contrast, long
43
44 MWCNTs seemed to have a weak capability to affect the liver in the DNA/RNA region, but
45
46 caused significant alterations in lipids and proteins. The lipid-to-protein ratio in liver was
47
48 significantly elevated by long MWCNTs in both genders. C_{60} and short MWCNTs reduced
49
50 the lipid-to-protein ratio in female fish, in the absence of significant effects in males, while
51
52 SWCNTs only elevated that ratio in males. The change in lipid-to-protein ratio in liver could
53
54 be caused from the direct interference of CNPs, while the lipid accumulation was possibly
55
56
57
58
59
60

1
2
3 due to CNP-induced inflammatory stress; long MWCNTs especially appear to disrupt
4
5 cholesterol trafficking in liver tissue.⁴⁶
6
7

8 **Conclusion**

9
10
11 Herein, biospectroscopy is presented as a global approach to nanotoxicity assessment,
12
13 providing one with a profile of CNP-induced alterations. IR and Raman spectra show that
14
15 CNPs can induce significant alterations in fish, and also highlights the differing effects of
16
17 four CNPs. Although it could not be shown that the CNPs passed through the bio-barrier (*e.g.*,
18
19 BBB) and moved to further tissues in fish, such as brain and gonads, the fact that these
20
21 organs were affected is a key indicator from the perspective of biological effects. Possibly the
22
23 alterations observed in the brain and gonads demonstrate that CNPs do exert potential toxic
24
25 effects on both nervous and reproductive systems.⁴⁷⁻⁵⁰ In addition, the alterations in
26
27 DNA/RNA spectral regions caused by CNPs, especially by C₆₀, suggested that DNA
28
29 methylation could be perturbed following the CNP exposure. It is necessary to conduct
30
31 further investigations to assess the DNA methylation levels in tissues of exposed fish. These
32
33 effects may be extrapolated to other organisms.⁵¹ Furthermore, future studies need to examine
34
35 whether the effects of CNPs impact on reproductive activity and test if transgenerational
36
37 effects might arise.
38
39
40
41
42
43
44
45

46 **Acknowledgments**

47
48
49 We would like to thank Yanqiu Liang, Yuxia Jiang and Dr. Guoyong Huang for their kind
50
51 assistance in setting up experiments. Funding for the laboratory experiments in this study
52
53 from the Chinese Academy of Sciences is greatly acknowledged.
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Analyst Accepted Manuscript

References

1. A. W. Jensen, S. R. Wilson and D. I. Schuster, *Biorg. Med. Chem.*, 1996, 4, 767-779.
2. R. Bakry, R. M. Vallant, M. Najam-ul-Haq, M. Rainer, Z. Szabo, C. W. Huck and G. K. Bonn, *Int. J. Nanomed.*, 2007, 2, 639.
3. R. H. Baughman, A. A. Zakhidov and W. A. de Heer, *Science*, 2002, 297, 787-792.
4. J. M. Planeix, N. Coustel, B. Coq, V. Brotons, P. S. Kumbhar, R. Dutartre, P. Geneste, P. Bernier and P. M. Ajayan, *J. Am. Chem. Soc.*, 1994, 116, 7935-7936.
5. A. Bianco, K. Kostarelos and M. Prato, *Curr. Opin. Chem. Biol.*, 2005, 9, 674-679.
6. A. Nel, T. Xia, L. Mädler and N. Li, *Science*, 2006, 311, 622-627.
7. H. Meng, T. Xia, S. George and A. E. Nel, *ACS Nano*, 2009, 3, 1620-1627.
8. G. Jia, H. Wang, L. Yan, X. Wang, R. Pei, T. Yan, Y. Zhao and X. Guo, *Environ. Sci. Technol.*, 2005, 39, 1378-1383.
9. S. K. Manna, S. Sarkar, J. Barr, K. Wise, E. V. Barrera, O. Jejelowo, A. C. Rice-Ficht and G. T. Ramesh, *Nano Lett.*, 2005, 5, 1676-1684.
10. J. Chłopek, B. Czajkowska, B. Szaraniec, E. Frackowiak, K. Szostak and F. Béguin, *Carbon*, 2006, 44, 1106-1111.
11. M. Bottini, S. Bruckner, K. Nika, N. Bottini, S. Bellucci, A. Magrini, A. Bergamaschi and T. Mustelin, *Toxicol. Lett.*, 2006, 160, 121-126.
12. A. Simon-Deckers, S. Loo, M. Mayne-L'hermite, N. Herlin-Boime, N. Menguy, C. Reynaud, B. Gouget and M. Carrière, *Environ. Sci. Technol.*, 2009, 43, 8423-8429.
13. Z. Tong, M. Bischoff, L. Nies, B. Applegate and R. F. Turco, *Environ. Sci. Technol.*, 2007, 41, 2985-2991.
14. E. Navarro, A. Baun, R. Behra, N. Hartmann, J. Filser, A.-J. Miao, A. Quigg, P. Santschi and L. Sigg, *Ecotoxicology*, 2008, 17, 372-386.
15. S. B. Lovern and R. Klaper, *Environ. Toxicol. Chem.*, 2006, 25, 1132-1137.
16. S. Zhu, E. Oberdörster and M. L. Haasch, *Mar. Environ. Res.*, 2006, 62, Supplement 1, S5-S9.
17. J. K. Folkmann, L. Risom, N. R. Jacobsen, H. Wallin, S. Loft and P. Møller, *Environ. Health Perspect.*, 2009, 117, 703-708.
18. A. Kroll, M. H. Pillukat, D. Hahn and J. Schnekenburger, *Eur. J. Pharm. Biopharm.*, 2009, 72, 370-377.
19. D. I. Ellis and R. Goodacre, *Analyst*, 2006, 131, 875-885.
20. V. Llabjani, J. Trevisan, K. C. Jones, R. F. Shore and F. L. Martin, *Environ. Sci. Technol.*, 2010, 44, 3992-3998.
21. V. Llabjani, J. D. Crosse, A. A. Ahmadzai, I. I. Patel, W. Pang, J. Trevisan, K. C. Jones, R. F. Shore and F. L. Martin, *Environ. Sci. Technol.*, 2011, 45, 10706-10712.
22. J. G. Kelly, J. I. Trevisan, A. D. Scott, P. L. Carmichael, H. M. Pollock, P. L. Martin-Hirsch and F. L. Martin, *J. Proteome Res.*, 2011, 10, 1437-1448.
23. P. Knief, C. Clarke, E. Herzog, M. Davoren, F. M. Lyng, A. D. Meade and H. J. Byrne, *Analyst*, 2009, 134, 1182-1191.
24. J. Dorney, F. Bonnier, A. Garcia, A. Casey, G. Chambers and H. J. Byrne, *Analyst*, 2012, 137, 1111-1119.
25. J. Trevisan, P. P. Angelov, P. L. Carmichael, A. D. Scott and F. L. Martin, *Analyst*, 2012, 137, 3202-3215.
26. J. Li, R. Strong, J. Trevisan, S. W. Fogarty, N. J. Fullwood, K. C. Jones and F. L. Martin, *Environ. Sci. Technol.*, 2013, 47, 10005-10011.
27. M. J. Riding, F. L. Martin, J. Trevisan, V. Llabjani, I. I. Patel, K. C. Jones and K. T. Semple, *Environ. Pollut.*, 2012, 163, 226-234.
28. M. N. Moore, *Environ. Int.*, 2006, 32, 967-976.

- 1
2
3 29. T. W. K. Fraser, H. C. Reinardy, B. J. Shaw, T. B. Henry and R. D. Handy, *Nanotoxicology*, 2011,
4 5, 98-108.
5 30. J. Cheng, C. M. Chan, L. M. Veca, W. L. Poon, P. K. Chan, L. Qu, Y.-P. Sun and S. H. Cheng,
6 *Toxicol. Appl. Pharmacol.*, 2009, 235, 216-225.
7 31. S. A. Brittin, S. J. Duivesteyn, M. Belmamoune, L. F. M. Bertens, W. Bitter, J. D. de Bruijn, D. L.
8 Champagne, E. Cuppen, G. Flik, C. M. Vandenbroucke-Grauls, R. A. J. Janssen, I. M. L. de Jong,
9 E. R. de Kloet, A. Kros, A. H. Meijer, J. R. Metz, A. M. van der Sar, M. J. M. Schaaf, S. Schulte-
10 Merker, H. P. Spaink, P. P. Tak, F. J. Verbeek, M. J. Vervoordeldonk, F. J. Vonk, F. Witte, H.
11 Yuan and M. K. Richardson, *Int. J. Dev. Biol.*, 2009, 53, 835-850.
12 32. H. Segner, *Comp. Biochem. Physiol., Part C: Toxicol. Pharmacol.*, 2009, 149, 187-195.
13 33. J. Trevisan, P. P. Angelov, A. D. Scott, P. L. Carmichael and F. L. Martin, *Bioinformatics*, 2013,
14 29, 1095-1097.
15 34. H. Wu, J. V. Volponi, A. E. Oliver, A. N. Parikh, B. A. Simmons and S. Singh, *Proc. Natl. Acad.*
16 *Sci. U. S. A.*, 2011, 108, 3809-3814.
17 35. A. R. R. Péry, J. Devillers, C. Brochot, E. Mombelli, O. Palluel, B. Piccini, F. Brion and R.
18 Beaudouin, *Environ. Sci. Technol.*, 2013, 48, 781-790.
19 36. T. B. Henry, E. J. Petersen and R. N. Compton, *Curr. Opin. Biotechnol.*, 2011, 22, 533-537.
20 37. N. Lewinski, V. Colvin and R. Drezek, *Small*, 2008, 4, 26-49.
21 38. D. Crouzier, S. Follot, E. Gentilhomme, E. Flahaut, R. Arnaud, V. Dabouis, C. Castellarin and J.
22 C. Debouzy, *Toxicology*, 2010, 272, 39-45.
23 39. Y. Sato, A. Yokoyama, K.-i. Shibata, Y. Akimoto, S.-i. Ogino, Y. Nodasaka, T. Kohgo, K. Tamura,
24 T. Akasaka, M. Uo, K. Motomiya, B. Jeyadevan, M. Ishiguro, R. Hatakeyama, F. Watari and K.
25 Tohji, *Mol. Biosyst.*, 2005, 1, 176-182.
26 40. C. J. Smith, B. J. Shaw and R. D. Handy, *Aquat. Toxicol.*, 2007, 82, 94-109.
27 41. A. Pietroiusti, L. Campagnolo and B. Fadeel, *Small*, 2013, 9, 1557-1572.
28 42. C. Chatgililoglu, C. Ferreri, M. Melchiorre, A. Sansone and A. Torreggiani, *Chem. Rev.*, 2013,
29 114, 255-284.
30 43. E. Oberdörster, *Environ. Health Perspect.*, 2004, 112, 1058-1062.
31 44. M. L. Circu and T. Y. Aw, *Free Radic. Biol. Med.*, 2010, 48, 749-762.
32 45. O. Zitka, S. Skalickova, J. Gumulec, M. Masarik, V. Adam, J. Hubalek, L. Trnkova, J. Kruseova,
33 T. Eckschlager and R. Kizek, *Oncol. Lett.*, 2012, 4, 1247-1253.
34 46. K. L. Ma, X. Z. Ruan, S. H. Powis, Y. Chen, J. F. Moorhead and Z. Varghese, *Hepatology*, 2008,
35 48, 770-781.
36 47. G. Bardi, A. Nunes, L. Gherardini, K. Bates, K. T. Al-Jamal, C. Gaillard, M. Prato, A. Bianco, T.
37 Pizzorusso and K. Kostarelos, *PLoS ONE*, 2013, 8, e80964.
38 48. M. Ema, N. Kobayashi, M. Naya, S. Hanai and J. Nakanishi, *Reprod. Toxicol.*, 2010, 30, 343-
39 352.
40 49. Y. Bai, Y. Zhang, J. Zhang, Q. Mu, W. Zhang, E. R. Butch, S. E. Snyder and B. Yan, *Nat Nano*,
41 2010, 5, 683-689.
42 50. T.-T. Win-Shwe and H. Fujimaki, *Int. J. Mol. Sci.*, 2011, 12, 6267-6280.
43 51. S. Rhiem, M. J. Riding, W. Baumgartner, F. L. Martin, K. T. Semple, K. C. Jones, A. Schäffer
44 and H. M. Maes, *Environ. Pollut.*, 2014, 196C: 431-439.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Legends to Figures

Figure 1. One-dimensional (1-D) cross-calculated PCA-LDA scores plots in 1st space of infrared (IR) spectra derived from brain or gill of zebrafish [male (M) or female (F)] exposed to carbon nanoparticles (CNPs) at two concentrations compared to the vehicle control. The horizontal line in the centre of each plot represents the mean value. C60, C₆₀ (fullerene); Long, long MWCNTs; Short, short MWCNTs; Single, SWCNTs

Figure 2. One-dimensional (1-D) cross-calculated PCA-LDA scores plots in 1st space of infrared (IR) spectra derived from gonad or liver of zebrafish [male (M) or female (F)] exposed to carbon nanoparticles (CNPs) at two concentrations compared to the vehicle control. The horizontal line in the centre of each plot represents the mean value. C60, C₆₀ (fullerene); Long, long MWCNTs; Short, short MWCNTs; Single, SWCNTs

Figure 3. Cross-calculated PCA-LDA scores plots in 2 dimensions (90% confidence ellipsoids) derived from tissues of zebrafish exposed to carbon nanoparticles (CNPs) at concentration of 0.1 mg/L and the vehicle control. C60, C₆₀ (fullerene); Long, long MWCNTs; Short, short MWCNTs; Single, SWCNTs

1
2
3 **Figure 4.** Cluster vector peak plots indicating the wavenumber basis for segregation
4
5 corresponding to the tissues exposed to CNPs (0.1 mg/L). Each exposed tissue was compared
6
7 with the vehicle control. The size of the symbol in cluster vector peaks plot is proportional to
8
9 the height of the corresponding peaks in the cluster vector plots, which relative to the extent
10
11 of alteration compared with the vehicle control. The hint line represents a typical IR spectrum
12
13 of the biochemical-fingerprint region (1800 cm^{-1} - 900 cm^{-1}) of the tissue. C60, C₆₀
14
15 (fullerene); Long, long MWCNTs; Short, short MWCNTs; Single, SWCNTs
16
17
18
19
20
21

22 **Figure 5.** Ratio of lipid to protein derived from IR spectra, *Mean ± standard deviation*: (a)
23
24 Comparison of different tissues from female (F) and male (M) zebrafish without exposure;
25
26 and, (b) comparison of tissues exposed to CNPs with the control. C60, C₆₀ (fullerene); Long,
27
28 long MWCNTs; Short, short MWCNTs; Single, SWCNTs
29
30
31
32
33
34

35 **Figure 6.** Ratio of (C=C) to (CH₂) in lipid derived from Raman spectra, *Mean ± standard*
36
37 *deviation*: (a) Comparison of brain and gonads in female (F) and male (M) zebrafish without
38
39 exposure; and, (b) comparison of the tissues (brain and gonads) exposed to CNPs with the
40
41 control. C60, C₆₀ (fullerene); Long, long MWCNTs; Short, short MWCNTs; Single,
42
43 SWCNTs
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1





Figure 2



Analyst Accepted Manuscript

Figure 3




Figure 4




Figure 5



a Tissues from Zebrafish**b**

Figure 6

