

ChemComm

Chemical Communications

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Y. Hashimoto, H. Kubo, K. Kawauchi and D. Miyoshi, *Chem. Commun.*, 2024, DOI: 10.1039/D4CC03753G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

COMMUNICATION

NRAS DNA G-quadruplex-targeting molecules for
sequence-selective enzyme inhibition

Yoshiki Hashimoto, Hiroki Kubo, Keiko Kawauchi and Daisuke Miyoshi*

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Sequence-selective G-quadruplex ligands are valuable for controlling gene expression. Here, we established a new fluorescence displacement assay using a NRAS G-quadruplex selective fluorescent probe to identify sequence-selective DNA G-quadruplex ligands. These sequence-selective NRAS G-quadruplex ligands maintained their binding affinity even in the presence of excessive human telomeric DNA G-quadruplex and regulated enzymatic activities in a sequence-selective manner.

Guanine-rich nucleic acid sequences can fold into G-quadruplex (G4) structures^{1, 2} that play diverse roles in biological processes such as transcription, translation, and replication^{3, 4}. Since G4 forming sequences are localized in biologically important genetic regions such as oncogenes and telomeres, G4s have garnered attention as targets for development of novel cancer treatment strategies to regulate oncogene expression. Various small molecules targeting G4 structures, G4 ligands, have been developed⁵. These compounds are expected to stabilize G4s to control gene expression.

Since almost all genomic DNA folds into canonical duplexes in cells, the structure-selectivity of G4 ligands is crucial for their intracellular applications. There are several reports on the development of structure-selective G4 ligands such as pyridostatin and telomestatin^{6, 7}. We also previously developed a screening system to identify structure-selective G4 ligands⁸. However, to target a specific gene, not only structure-selectivity but also sequence-selectivity of G4 ligands is highly required, since approximately 700,000 potential G4-forming sequences exist in the human genome⁹. Here, we define a sequence-selective G4 ligand as a small compound which binds to the target G4 but does not bind to other G4s.

Benzothiazole hydrazone analogues reported by Pradeepkumar's group showed approximately 3-20 times higher affinity for *c-MYC* G4 compared to other G4s¹⁰. Madder's

group developed a method for sequence-selective alkylation to a target G4 by conjugating an alkylation compound with the complementary sequence of the target DNA¹¹. Some sequence-selective G4 targeting by conjugated oligonucleotides has also been achieved^{12, 13}. However, conjugation with nucleic acids could make G4 ligands difficult to deliver into living cells. Thus, G4 ligands with higher sequence-selectivity are still required. However, it is difficult to design such sequence-selective G4 ligands rationally. Here, we established a novel screening system for sequence-selective G4 ligands based on fluorescent intercalator displacement (FID)¹². By utilizing this system, we identified new NRAS DNA G4 sequence-selective ligands. Importantly, it was demonstrated that the G4 ligands stabilized the target G4 and inhibited enzymatic activity in a sequence-selective manner.

A fluorescent probe for the FID is essential to screen NRAS DNA sequence-selective G4 ligands. We previously reported that an anionic phthalocyanine coordinating with Zn²⁺ (ZnAPC) can sequence-selectively bind to the NRAS mRNA G4¹³, whereas it did not bind to telomeric DNA G4, which is the most abundant G4 in cells¹⁴. We thus hypothesized that ZnAPC would bind to NRAS DNA G4 but not to telomeric DNA G4. Since the majority of G4 structures in cells are formed by telomere sequences, we evaluated the affinity and selectivity of ZnAPC with NRAS G4 against the telomeric DNA G4 as a proof of concept. We used two oligonucleotides, NRAS and TELO, which were derived from human NRAS and human telomere, respectively (Table S1). Figure S1 shows normalized UV melting curves at 295 nm and circular dichroism (CD) spectra of 10 μM NRAS and TELO. The hypochromic melting transition and the CD spectrum showed that NRAS and TELO folded into a parallel G4 and a mixed G4, respectively. Figure S2A and B shows the absorption spectra of 20 μM ZnAPC with various concentrations of NRAS and TELO, respectively. An absorbance increase at 620 nm and a decrease at 680 nm were observed with increasing concentration of NRAS. Conversely, no significant change was observed with TELO, indicating the selective binding of ZnAPC with NRAS. In addition, we confirmed whether ZnAPC can maintain its affinity for NRAS even in the presence of 50 μM TELO (Fig. S2C). Figure S2D shows plots of Δabs₆₈₀, (abs. at 680 nm with DNA) - (abs. at

^a *Frontiers of Innovative Research in Science and Technology, Konan university, 7-1-20 Minatogima-minamimachi, Chuo-ku, Kobe, Hyogo 650-0047, Japan.*

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x



680 nm without DNA), *versus* concentration of NRAS in the absence or presence of TELO. ZnAPC showed K_d values of $5.8 \pm 0.7 \mu\text{M}$ and $6.0 \pm 0.6 \mu\text{M}$ in the absence and presence of $50 \mu\text{M}$ TELO, respectively, at 25°C (see SI for evaluation of the K_d values). These results demonstrate that the affinity of ZnAPC for NRAS was maintained even in the presence of excessive TELO. Higher concentrations of NRAS increased the fluorescence intensity of ZnAPC more than 20-fold with and without $50 \mu\text{M}$ TELO (Fig. S3A), further confirming the sequence-selectivity of ZnAPC for NRAS (Fig. S3B). It is not yet clear how ZnAPC can bind to the target NRAS G4 in a sequence-selective manner, although it is considered that not only the loop sequence but also the flanking sequence are important for the sequence selective binding of ZnAPC¹⁴. The increase in fluorescence intensity with sequence-selective binding shows that ZnAPC is a suitable probe for FID.

Figure 1A shows a schematic illustration of a sequence-selective G4 ligand screening by ZnAPC displacement. In this assay, the affinity index (I_{Affinity}), showing the degree of displacement by a tested compound, was evaluated from the F_{690} (fluorescence intensity of ZnAPC at 690 nm) decrease as the following: $(F_{690} \text{ with the compound}) / (F_{690} \text{ without the compound})$. Compounds showing lower I_{Affinity} are expected to have stronger affinity for NRAS. We screened about 3,000 compounds to identify sequence-selective ligands targeting NRAS. The obtained I_{Affinity} values in the absence or presence of TELO were organized as shown in Fig. 1B. Compounds observed in the blue area lack sequence-selectivity, while compounds in the red area may possess sequence-selectivity. As shown in Fig. 1C, the screening results identified three compounds highlighted in red: pacritinib¹⁵, osimertinib¹⁶, and JNK-IN-8¹⁷ (Fig. 2A), which exhibited the lowest I_{Affinity} in the presence of TELO. It was demonstrated that the quenching of ZnAPC is not due to a direct interaction with the ligand (Fig. S4). Of these three,




Fig. 1 (A) Schematic illustration of sequence-selective G4 ligand screening by ZnAPC displacement. (B) Plots of I_{Affinity} of ligands in the presence of $50 \mu\text{M}$ TELO *versus* I_{Affinity} of ligands in the presence of $0 \mu\text{M}$ TELO. Plots located within the red area have sequence-selectivity. (C) Plots of I_{Affinity} of compounds in the presence of $50 \mu\text{M}$ TELO *versus* I_{Affinity} of compounds in the absence of $50 \mu\text{M}$ TELO. (i) Pacritinib, (ii) Osimertinib, (iii) JNK-IN-8, and (iv) PDS.

pacritinib exhibited the lowest I_{Affinity} in the presence of TELO, suggesting that it has the highest sequence-selectivity.

To confirm the binding affinity and selectivity of these compounds, we observed changes in I_{Affinity} with varying concentrations of the compounds. Figure 2B and S4A show the fluorescence spectra of $20 \mu\text{M}$ ZnAPC and $1 \mu\text{M}$ NRAS with various concentrations of pacritinib in the absence and presence of $50 \mu\text{M}$ TELO. The observed K_d (K_i) values of the ligands were evaluated from fluorescence changes of ZnAPC in the absence and presence of $50 \mu\text{M}$ TELO (Fig. 2C). The K_i values of pacritinib were $0.74 \pm 0.11 \mu\text{M}$ and $0.88 \pm 0.09 \mu\text{M}$ in the absence and presence of $50 \mu\text{M}$ TELO, respectively. The lack of significant change in these K_i values confirms the high sequence-selectivity of pacritinib. Similar results were obtained for osimertinib and JNK-IN-8 (Fig. S5B and C). In contrast, the K_i values of pyridostatin (PDS), which was identified as a non-sequence-selective ligand (Fig. 1C (iv)), in the presence of $50 \mu\text{M}$ TELO ($4.1 \pm 0.1 \mu\text{M}$) was about 10 times larger than that in the absence of $50 \mu\text{M}$ TELO (0.47 ± 0.08), confirming the lower sequence selectivity of PDS (Fig. S5D). In addition, we carried out ThT displacement (TD) assays⁸ for TELO (see SI for evaluation of K_i , Fig. S6). The K_i values of pacritinib, osimertinib, and JNK-IN-8 were $77 \mu\text{M}$, $89 \mu\text{M}$, and $103 \mu\text{M}$, respectively, for TELO, at 25°C , which are about 100-fold larger than that for NRAS (Fig. S6C). The K_i values support the high sequence-selectivity of pacritinib, osimertinib, and JNK-IN-8. In comparison, the K_i value of PDS for TELO was 0.64 , which is almost the same as that for NRAS, confirming that PDS binds to G4s in a non-sequence-selective manner.

Next, we investigated the effect of these compounds on the structure and stability of NRAS. Fig. S6A shows CD spectra of $10 \mu\text{M}$ NRAS in the absence and presence of the three G4 ligands at $50 \mu\text{M}$, suggesting that these compounds did not alter the structure of NRAS G4. The thermal stability of NRAS was studied using CD melting curves at 265 nm (Fig. S7B). The T_m value of $10 \mu\text{M}$ NRAS was $41.0 \pm 0.9^\circ\text{C}$. Pacritinib, osimertinib, and JNK-IN-8 changed the T_m value to $72.7 \pm 0.6^\circ\text{C}$, $63.9 \pm 0.6^\circ\text{C}$, and $54.2 \pm 0.3^\circ\text{C}$, respectively (Fig. S7C). These results indicate that the ligands not only bind sequence-selectively but also strongly




Fig. 2 (A) Chemical structures of pacritinib, osimertinib, and JNK-IN-8. (B) Fluorescence spectra of $20 \mu\text{M}$ ZnAPC with 0, 0.1, 0.3, 1, 3, 10, 30, and $100 \mu\text{M}$ pacritinib in the buffer containing 50 mM MES-LiOH (pH 7.0), 100 mM KCl , and $1 \mu\text{M}$ NRAS at 25°C . (C) Plots of I_{Affinity} *versus* concentration of pacritinib.



stabilize NRAS.

With the identification of the sequence-selective G4 ligands, we performed T7 RNA polymerase (RNAP) stop assays to investigate the inhibitory effects of the G4 ligands on transcription. A template containing the NRAS G4-forming sequence (NRAS template, Table S1) was utilized for the transcription reaction. The transcription products with G4 ligands in concentrations ranging from 0 to 100 μM were visualized using denaturing polyacrylamide gel electrophoresis (PAGE). When the transcription reaction proceeds completely, completed transcripts of 66 nucleotides can be observed (Fig. 3A). When the transcription reaction is inhibited by G4 ligands, arrested transcripts of 36 nucleotides are produced. Figure 3B shows the results for pacritinib in the absence and presence of 50 μM TELO. A decrement of the completed transcript and an increment of the arrested transcript were observed with increasing concentration of pacritinib under both conditions. The IC_{50} values of pacritinib in the absence and presence of 50 μM TELO were $28 \pm 2 \mu\text{M}$ and $32 \pm 2 \mu\text{M}$, respectively, at 37 $^{\circ}\text{C}$ (Fig. 3C) (see SI for evaluation of IC_{50}). These results show that pacritinib can inhibit T7 RNAP activity in a sequence-selective manner. The significant difference between the values of the IC_{50} and the K_i was observed. Although the reason for the difference remains unclear, one of possible reasons is that the NRAS stabilized by the ligands competes the T7 RNAP binding to the substrate DNA, resulting in an unwinding of the DNA G4 structure. In addition, it was reported that not only their thermal stability but also the kinetics of the G4 formation and the ligand binding with G4 are also important for the inhibition of the polymerase activity¹⁸. The kinetic controlling of the T7 RNAP activity may lead to the difference between the IC_{50} and




Fig. 3 (A) Schematic illustration of T7 RNA polymerase stop assay with G4-forming template strand. (B) Transcripts from 1 μM NRAS template with 0, 0.1, 0.3, 1, 3, 10, 30, and 100 μM pacritinib in the absence (left) and presence (right) of 50 μM TELO. The transcription reactions were carried out in a buffer containing 1 mM KCl, 99 mM LiCl, 8 mM MgCl_2 , and 40 mM Tris-HCl (pH 7.2) at 37 $^{\circ}\text{C}$ for 2 hours. (C) Plots of relative amounts of arrested transcript with NRAS template versus concentration of pacritinib in the absence (circles with dotted line) or presence (triangles with continuous line) of 50 μM TELO.

the K_i values. The sequence-selective inhibition of T7 RNAP was

also observed for osimertinib and JNK-IN-8 (Fig. S8). These results demonstrate that the novel sequence-selective G4 ligands can sequence-selectively inhibit transcription activity mediated by stabilization of NRAS G4. Because of the direct inhibition of T7 RNAP activity by PDS at high concentrations in the absence of TELO, the experiments were conducted with PDS only up to 30 μM (Fig. S9). Although PDS exhibited a high transcription inhibitory effect in the absence of TELO, the effect was hardly observed in the presence of TELO. To confirm that the transcription inhibitory effects of these compounds were not unique for T7 RNAP, we performed the T7 RNAP assay with a mutNRAS template (see Table S1 for sequence), which does not form a G4 structure (Fig. S10). Pacritinib, osimertinib, and JNK-IN-8 did not affect the transcription activity with the mutNRAS template, confirming that their inhibitory effect on transcription was mediated by G4 formation. In contrast, PDS decreased the total amount of transcripts in the absence of TELO, suggesting that a higher concentration of PDS directly inhibits the T7 RNAP activity.




Fig. 4 (A) Results of tsTRAP assay with 0, 1, 10, and 100 μM pacritinib. Telomerase elongation reactions were carried out in a buffer containing 200 mM Tris-HCl (pH 8.3), 15 mM MgCl_2 , 630 mM KCl, 0.5% Tween 20, and 10 mM EGTA for 30 min at 30 $^{\circ}\text{C}$. (B) Plots of relative telomerase activity versus concentrations of pacritinib, osimertinib, JNK-IN-8, and PDS.

Given that the ligands showed high sequence-selectivity, it is expected that no inhibitory effects on enzymatic activity via formation of other G4s would be observed. We thus investigated the effects of pacritinib, osimertinib, JNK-IN-8, and PDS on telomere elongation by telomerase using the two-step telomeric repeat amplification protocol (tsTRAP) assay¹⁹. This method allows the study of inhibitory effects of G4 ligands on the telomerase extension reaction without affecting the subsequent PCR stop¹⁹. Figure 4A shows a non-denaturing PAGE of tsTRAP products with 0, 1, 10, and 100 μM pacritinib, demonstrating the inhibitory effect of pacritinib on telomere elongation. The amount of the internal control by addition of pacritinib was constant, showing that the addition of pacritinib did not interfere with the PCR reaction. More importantly, no significant decrement of telomerase activity was observed upon addition of pacritinib up to 100 μM (Fig. 4B). These results confirm that the sequence-selective G4 ligands targeting NRAS did not inhibit enzymatic activity depending on G4 formation other than NRAS (Fig. 4B). Similar to pacritinib, osimertinib and JNK-IN-8 did not inhibit telomerase activity with increasing concentration (Fig. 4B and S11A). In contrast, PDS strongly inhibited telomerase activity (Fig. 4B and S11B) because of its non-sequence-selective binding to G4s. The sequence-selective



NRAS G4 ligands, pacritinib, osimertinib, and JNK-IN-8, inhibited enzyme activity of the target G4 sequence, suggesting that developing target sequence-selective G4 ligands can control target enzymatic reactions. Such target selective enzyme inactivation is an important aspect for cellular applications of G4 ligands involving drug development.

In summary, we established a new screening method for obtaining NRAS sequence-selective G4 ligands. Pacritinib, osimertinib, and JNK-IN-8 were identified as novel sequence-selective G4 ligands that strongly stabilized NRAS. More importantly, the ligands showed sequence-selective regulation for enzymatic activity in the transcription reaction and telomere elongation. Although disease-related proteins have attracted attention as therapeutic targets, some of them, including NRAS, are often difficult to target with small molecule drugs²⁰⁻²². The use of sequence-selective G4 ligands enables the targeting of specific G4s, potentially allowing the control of specific biological reactions regardless of the protein structure. Although sequence-selective G4 ligands are useful for inhibit target enzyme inhibition, this screening method can be applied only for NRAS due to the properties of ZnAPC. For a screening of sequence-selective G4 ligands targeting other sequences, a sequence-selective fluorescent probe for the target G4 is required. As an alternative approach, performing of screenings for the target G4 and for the decoy G4 makes possible to obtain sequence-selective G4 ligands. The results obtained in this study may stimulate further studies for screening and designing sequence-selective G4 ligands.

This work was supported by JSPS KAKENHI, grant numbers 24K21801, 21H02062, and 20K21259, a Research Grant of the Asahi Glass Foundation, Japan, the Hirao Taro Foundation and Junzo Tateno Foundation of Konan Gakuen for Academic Research, Japan, and JST SPRING, grant number JPMJSP2117.

Author contributions

Conceptualization, D.M.; methodology, D.M.; formal analysis, Y.H. and H.K.; investigation, Y.H. and H.K.; data curation, Y.H.; writing—original draft preparation, Y.H.; writing—review and editing, K.K. and D.M.; visualization, D.M.; supervision, K.K. and D.M.; project administration, D.M.; funding acquisition, Y.H., K.K. and D.M.

Conflicts of interest

There are no conflicts to declare.

Data availability

The data supporting this article have been included as part of the ESI.

Notes and references

1. M. Gellert, M. N. Lipsett and D. R. Davies, *Proc. Natl. Acad. Sci. U S A*, 1962, **48**, 2013-2018.

2. D. Sen and W. Gilbert, *Nature*, 1988, **334**, 364-366.
3. S. Balasubramanian, L. H. Hurley and S. Neidle, *Nat. Rev. Drug. Discov.*, 2011, **10**, 261-275.
4. D. Rhodes and H. J. Lipps, *Nucleic Acids Res.*, 2015, **43**, 8627-8637.
5. Y. H. Wang, Q. F. Yang, X. Lin, D. Chen, Z. Y. Wang, B. Chen, H. Y. Han, H. D. Chen, K. C. Cai, Q. Li, S. Yang, Y. L. Tang and F. Li, *Nucleic Acids Res.*, 2022, **50**, D150-d160.
6. R. Rodriguez, S. Müller, J. A. Yeoman, C. Trentesaux, J. F. Riou and S. Balasubramanian, *J. Am. Chem. Soc.*, 2008, **130**, 15758-15759.
7. M. Y. Kim, H. Vankayalapati, K. Shin-Ya, K. Wierzba and L. H. Hurley, *J. Am. Chem. Soc.*, 2002, **124**, 2098-2099.
8. Y. Hashimoto, Y. Imagawa, K. Nagano, R. Maeda, N. Nagahama, T. Torii, N. Kinoshita, N. Takamiya, K. Kawauchi, H. Tatesishi-Karimata, N. Sugimoto and D. Miyoshi, *Chem. Commun. (Camb)*, 2023, **59**, 4891-4894.
9. V. S. Chambers, G. Marsico, J. M. Boutell, M. Di Antonio, G. P. Smith and S. Balasubramanian, *Nat. Biotechnol.*, 2015, **33**, 877-881.
10. S. P. Pany, P. Bommiseti, K. V. Diveshkumar and P. I. Pradeepkumar, *Org. Biomol. Chem.*, 2016, **14**, 5779-5793.
11. E. Cadoni, L. De Paepe, G. Colpaert, R. Tack, D. Waegeman, A. Manicardi and A. Madder, *Nucleic Acids Res.*, 2023, **51**, 4112-4125.
12. D. Monchard, C. Allain and M. P. Teulade-Fichou, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 4842-4845.
13. S. Kumari, A. Bugaut, J. L. Huppert and S. Balasubramanian, *Nat. Chem. Biol.*, 2007, **3**, 218-221.
14. K. Kawauchi, W. Sugimoto, T. Yasui, K. Murata, K. Itoh, K. Takagi, T. Tsuruoka, K. Akamatsu, H. Tateishi-Karimata, N. Sugimoto and D. Miyoshi, *Nat. Commun.*, 2018, **9**, 2271.
15. S. Hart, K. C. Goh, V. Novotny-Diermayr, C. Y. Hu, H. Hentze, Y. C. Tan, B. Madan, C. Amalini, Y. K. Loh, L. C. Ong, A. D. William, A. Lee, A. Poulsen, R. Jayaraman, K. H. Ong, K. Ethirajulu, B. W. Dymock and J. W. Wood, *Leukemia*, 2011, **25**, 1751-1759.
16. D. A. Cross, S. E. Ashton, S. Ghiorghiu, C. Eberlein, C. A. Nebhan, P. J. Spitzler, J. P. Orme, M. R. Finlay, R. A. Ward, M. J. Mellor, G. Hughes, A. Rahi, V. N. Jacobs, M. Red Brewer, E. Ichihara, J. Sun, H. Jin, P. Ballard, K. Al-Kadhimi, R. Rowlinson, T. Klinowska, G. H. Richmond, M. Cantarini, D. W. Kim, M. R. Ranson and W. Pao, *Cancer Discov.*, 2014, **4**, 1046-1061.
17. T. Zhang, F. Inesta-Vaquera, M. Niepel, J. Zhang, S. B. Ficarro, T. Machleidt, T. Xie, J. A. Marto, N. Kim, T. Sim, J. D. Laughlin, H. Park, P. V. LoGrasso, M. Patricelli, T. K. Nomanbhoy, P. K. Sorger, D. R. Alessi and N. S. Gray, *Chem. Biol.*, 2012, **19**, 140-154.
18. S. Takahashi, A. Kotar, H. Tateishi-Karimata, S. Bhowmik, Z. F. Wang, T. C. Chang, S. Sato, S. Takenaka, J. Plavec and N. Sugimoto, *J. Am. Chem. Soc.*, 2021, **143**, 16458-16469.
19. H. Yaku, T. Murashima, D. Miyoshi and N. Sugimoto, *J. Phys. Chem. B*, 2014, **118**, 2605-2614.
20. A. D. Cox, S. W. Fesik, A. C. Kimmelman, J. Luo and C. J. Der, *Nat. Rev. Drug Discov.*, 2014, **13**, 828-851.
21. C. V. Dang, *Cell*, 2012, **149**, 22-35.
22. L. Mabonga and A. P. Kappo, *Biophys. Rev.*, 2019, **11**, 559-581.



NRAS DNA G-quadruplex-targeting molecules for sequence-selective enzyme inhibition

Yoshiki Hashimoto, Hiroki Kubo, Keiko Kawauchi and Daisuke Miyoshi*

Data availability

All experiment procedures and the data supporting this article can be found in the ESI.

