
Volume 27 Number 11 21 March 2025 Pages 1513-1670

CrystEngComm

rsc.li/crystengcomm



ISSN 1466-8033



COMMUNICATION

In-Hyeok Park et al. An unsymmetrically sandwiched bis(O_3S_2 -macrocycle) lead($\mathfrak n$) complex via an endo/exo-coordination mode

CrystEngComm



COMMUNICATION

View Article Online



Cite this: CrystEngComm, 2025, 27, 1529

Received 12th October 2024, Accepted 24th November 2024 An unsymmetrically sandwiched bis(O₃S₂macrocycle) lead(II) complex via an endo/exocoordination mode†

Seulgi Kim, ** Kyunghye Ju, ** Taehun Kim ** and In-Hyeok Park ***

DOI: 10.1039/d4ce01045k

rsc.li/crystengcomm


An unsymmetrically sandwiched complex was isolated from the reaction of O₃S₂-macrocycle with lead(II) perchlorate. The lead(II) centre is bound to one macrocycle in an endocyclic (face) mode and the other in an exocyclic (edge) mode, resulting in a dicapped trigonal prismatic geometry. The edge-to-face sandwich complex might be associated with the hemispherical lead(II) coordination.

Metal-ligand interactions play a crucial role in the design and functional control of metal complexes, metal-organic frameworks and various macromolecular (MOFs), compounds. 1-11 The precise regulation of such interactions can impart unique chemical properties and reactivities, making crown ethers and their derivatives notable for their ability to selectively bind metal ions. 12,13 Crown ethers have attracted much attention due to the preferential endocyclic (metal-in-cavity) complexation with alkali and alkali earth metal ions in a good fit mode. 14 Sometimes, crown ether-type macrocycles form unusual stoichiometric complexes including sandwich types with a 1:2 (metal-to-ligand) ratio, club sandwiches (2:3) and higher-ordered ones. 15-23 In general, sandwich-type complexes are formed when the cavity is relatively small to fully accommodate the metal ion.


On the other hand, various modifications of crown ethers including donor atoms allow new reactivities functionalities. 24-26 Thiacrowns (or thiamacrocycles), for example, form not only endocyclic complexes but also exocyclic (a metal ion binds outside of the cavity) coordination products due to the preferential exo-orientation of sulfur donors.²⁷⁻³² Interestingly, the exo-coordination between sulfur donors (soft base) and thiaphilic metal ions (soft acid)³³ so often provides an opportunity to construct diverse types of metallosupramolecules from monomers and cyclic oligomers to polymers as reviewed by the Lee group. 34-36 In addition, some controllable endo- and exo-coordination modes^{37,38} and their applications in preparing Cu_nI_n (n = 2 or 4) clusters,³⁹ photochemical sensors^{40–43} and chiral inversion⁴⁴ have been reported.

Considering the contribution of the exo-coordination, the sandwich-type macrocyclic complexes could be extended to three types (A-C in Scheme 1, see Fig. S1†) depending on the combination of the endo- (face) and exo-coordination (edge). Undoubtedly, type A (face-to-face) is the most common because large alkali metal ions tend to form this symmetrical sandwich. 15-23 Types B and C involve the edge mode, which appears in the exo-coordinated soft metal complexes of thiamacrocycles. 34-38 In particular, type C (edge-to-face) complexes are unsymmetrical because one macrocycle has most of the donors coordinated in the same plane, while one part of the donors in the other macrocycle participates in the coordination environment. Thus, isolating type C complexes is challenging, especially in a one-pot reaction.

More recently, we have reported a 17-membered O₃S₂macrocycle (L in Scheme 2) as one of the ditopic model ligands together with its interdonor (sulfur-to-sulfur) distancedependent cis-palladium(II) complex and infinite onedimensional copper(II) complex. 45,46 Due to three consecutive




† Electronic supplementary information (ESI) available: Synthetic details, characterization, NMR titration, and crystal structure data. CCDC 2389314 (1). For ESI and crystallographic data in CIF or other electronic format see DOI: https://doi.org/10.1039/d4ce01045k



Scheme 1 Sandwich-type macrocyclic complexes with different coordination modes: (a) type A (face-to-face), (b) type B (edge-toedge) and (c) type C (edge-to-face).

Communication CrystEngComm



Scheme 2 Chemical structure of a ditopic O₃S₂-macrocycle (L).

oxygen donors, **L** has a hard base nature for *endo*-binding with hard or borderline metal ions (Scheme 2). Meanwhile, two sulfur donors could act as soft bases for the *exo*-binding sites with soft metal ions. In the present work, we employed the ligand **L** to gain a further understanding of non-soft metal complexations.

We have extended such sandwich-type complexes of macrocycles with hard and soft metal ions to borderline metal ions including lead(II) because their complexation behaviours are important in coordination chemistry, materials science and biological areas. A7,48 Lead(II) may bind to both oxygen and sulfur donors in L with variable coordination numbers and unusual coordination geometries. Practically, plenty of hemidirected lead(II) complexes involving both Pb–O and Pb–S bonds have been reported so far. In this work, we have isolated an unsymmetrically sandwiched lead(II) complex of L with an *edgeto-face* mode (type C in Scheme 1) as the first non-soft metal species in this category. The details of our investigations are presented below.

In 1, as shown in Fig. 1a, the complex part $[Pb(L)_2(ClO_4)]^+$ comprises an unsymmetrical sandwich structure in which both *endo*-dentate (L_A) and *exo*-dentate (L_B) macrocycles are bonded to the lead(II) centre in an *edge-to-face* mode (type C in Scheme 1c). The lead(II) centre is eight-coordinated with one macrocycle (L_A) bound to one side of the metal ion *via* the O_3S_2 donors $(Pb1-S1\ 2.902(13),\ Pb1-S2\ 2.990(12),\ Pb1-O1\ 2.854(3),\ Pb1-O2\ 2.806(3),\ Pb1-O3\ 2.951(1)$ Å). Interestingly, the coordination sphere is completed by two *exo*-dentate sulfur donors $(Pb1-S3\ 2.902(11),\ Pb1-S4\ 3.100(12)$ Å) from the other macrocycle L_B in an *edge* mode and one perchlorate oxygen atom $(Pb1-O8\ 2.781(9)\ \text{Å})$.

The eight-coordinated geometry of 1 can be best described as a distorted di-capped trigonal prismatic geometry (Fig. 1b). The trigonal faces of the prism are defined by S1–S2–S3 and O1–O2–O8. One sulfur atom (S4) and one oxygen atom (O3) cap the corresponding rectangular faces (S1–S3–O8–O1 and S2–S3–O8–O2). Thus, the capping bond lengths of




Fig. 1 An unsymmetrically sandwiched lead(\shortparallel) complex, [Pb(L)₂(ClO₄)] ClO₄·0.5CH₂Cl₂·0.5CH₃CN (1): (a) front view, (b) coordination environment of the lead(\shortparallel) centre, showing a distorted di-capped trigonal prismatic geometry, (c) side view and (d) space-filling structure. Non-coordinated solvent molecules are omitted. Each non-coordinated perchlorate ion in (b–d) is not shown.

Pb1–S4 (3.100(12) Å) and Pb1–O3 (2.951(1) Å) are slightly longer than those of other Pb1–S (2.902(11)–2.990(12) Å) and Pb1–O (2.781(9)–2.854(3) Å) bonds, respectively. Similar to the Pb–S bonds (2.902 (11)–3.100(12) Å), the Pb–O_{ether} bond lengths (2.806(3)–2.951(1) Å) fall in the longer part of the literature range for such bonds (2.5–2.9 Å). The elongated bond lengths seem to be caused by the steric hindrance between two macrocycles. Also, these are the typical signs of a hemidirected complex discussed in the latter part.

Notably, the lead(II) centre is located 1.00 Å above the average plane of the O₃S₂ donors in L_A with a perching manner (Fig. 1c). In general, the perching position in the sandwich-type macrocyclic complexes is mainly due to the larger size of metal ions compared to the macrocyclic cavity. In 1, however, the perching position of the lead(II) centre is mainly caused by the exo-coordination of the macrocycle L_B that lifts the lead(II) centre. Unlike the face-to-face mode, which shields the metal centre effectively from anions and solvent molecules, the edge-to-face mode in 1 allows some metal space between the two macrocycles (Fig. 1c and d). Due to this reason, one ClO₄ ion participates to the coordination environment (Pb1-O8 2.781(9) Å), with another ClO₄ ion remaining uncoordinated (Pb1···O14 3.243(6) Å) (Fig. 1a). The IR spectrum for 1 shows an intense ClO_4 peak at 1045 cm⁻¹ (Fig. S6†).

As mentioned, lead(II) often shows a hemidirected coordination sphere. Indeed, the eight-coordinated O_4S_4 coordination environment in 1 occupies only one hemisphere keeping the other hemisphere unoccupied except for the lone pair electron (Fig. 2). In our previous work, the

CrystEngComm Communication




Fig. 2 The hemispherical coordination mode of the eight-coordinated lead(11) centre in 1

assembly reactions of lead(II) and zinc(II) with a mixture of the rod-like 1,4-bis(4-pyridyl)piperazine and V-shaped 4,4'sulfonyldibenzoic acid generate a polyrotaxane and a polycatenane, respectively.⁵⁴ In this case, interestingly, the polyrotaxane formation is attributed to the hemisphere configuration of the lead(II) centre. The Penner-Hahn and Godwin groups reported the hemisphere coordination of lead(II) with thiol-rich peptides, which do not induce protein folding unlike the zinc(II)-analogue, as the origin of the lead poisoning.⁵⁸ Considering the coordination numbers, elongated bond lengths, anion coordination and steric configuration, the formation of unsymmetrical sandwich-type complex 1 might be responsible for the hemispherical coordination of the lead(II) centre. To the best of our knowledge, this is the first example of unsymmetrically sandwiched macrocyclic complexes with non-soft metal ions. When the analogue product with the soft metal ion shown in Fig. S1c† is involved, this is the second example in the Type C category (Scheme 1c).

For the understanding of the complexation behaviours of L with lead(II) perchlorate in solution, we attempted NMR and ESI-mass experiments. In the ¹H NMR titration, the addition of lead(II) (0-1.7 equiv.) to L induced all of the proton peaks to shift downfield, reflecting a stable complexation with an exchange rate being fast on the NMR time scale (Fig. 3a and




Fig. 3 (a) 1 H NMR titration of L (1.0 \times 10 $^{-3}$ M) with lead(II) perchlorate in CDCl₃/CD₃CN (v/v 1:1) and (b) titration curves for each proton in L.

S3†). The titration curves show that the magnitudes of the chemical shifts follow the order $H_4 > H_3 \gg H_2 > H_1 > H_{ar}$ (Fig. 3b). The larger shifts for H₄ and H₃ than those for H₂ and H₁ indicate the more favourable binding of lead(II) to sulfur donors than oxygens. Above 1.0 equiv., no significant chemical shifts were observed probably due to the predominant formation of a 1:1 product in this region.

The stability constant of the complexation was obtained by HyperNMR software (Fig. S4†).⁵⁹ This complexation could not be described using a mixed model of the 1:1 and 1:2 ratios (metal-to-ligand). Instead, a good fit of the data with the 1:1 model yields the $\log K$ value of 5.9 \pm (0.5), indicating the formation of a typical endocyclic mononuclear lead(II) complex, unlike the crystal structure. It is not surprising that the crystal structure differs from that in solution because of the solvation of the lead(II) with acetonitrile, a dipolar aprotic solvent.

Alternatively, ESI-mass spectroscopy is quite sensitive in monitoring labile or less stable supramolecules assembled in solution. The ESI-mass spectrum of L with 1.0 equiv. of lead(II) perchlorate was dominated by peaks for the 1:1 (metal-to-ligand) complexes such as $[Pb(L)(ClO_4)]^+$ (m/z 683) and $[Pb(L)(AN)_2(ClO_4)]^+$ (m/z 765, AN = CH₃CN) (Fig. 4). In the same spectrum, the formation of both 1:2 and 2:2 complexes is also confirmed by the peaks for $[Pb(L)_2(ClO_4)]^+$ $(m/z \ 1059)$ and $[Pb_2(L)_2(ClO_4)_3]^+$ $(m/z \ 1465)$, respectively. This result indicates that the 1:2 stoichiometric species is less stable than in the solid state probably due to the influence of the solvation.

In summary, an unsymmetrically sandwiched 1:2 (metalto-ligand) complex with an edge-to-face mode via exo/endocoordination was isolated from the reaction of a ditopic O₃S₂macrocycle L with lead(II) perchlorate. In its crystal structure, the lead(II) centre in a hemisphere configuration adopts an eight-coordinated di-capped trigonal prismatic geometry with the elongated Pb-O and Pb-S bond distances matched to satisfy the steric and electronic conditions required. In solution, the mononuclear bis(macrocycle) product was traced by ESI-mass spectroscopy but such evidence was not observed in the ¹H NMR titration probably due to the solvation effect. The knowledge of such a sandwich-type complex is expected to provide insight into the lead(II) coordination found in biosystems and materials.




Fig. 4 ESI-mass spectrum of a mixture of L and Pb(ClO₄)₂·3H₂O in CH2Cl2/CH3CN (1:1, v/v).

Data availability

Communication

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the NRF of Korea (2021R1C1C1006765, 2022R1A4A1022252 and RS-2023-00245420), COMPA (2024-24020010-11r, R&D Equipment Engineer Education Program), and the research fund of Chungnam National University.

References

- 1 G. Chakraborty, I.-H. Park, R. Medishetty and J. J. Vittal, Chem. Rev., 2021, 121, 3751–3891.
- 2 M. Kim, J. Yi, S.-H. Park and S. S. Park, Adv. Mater., 2023, 35, 2203791.
- 3 J. Park, A. Adhikary and H. R. Moon, Coord. Chem. Rev., 2023, 497, 215402.
- 4 C. Bae, M. Gu, Y. Jeon, D. Kim and J. Kim, *Bull. Korean Chem. Soc.*, 2023, 44, 112–124.
- 5 B. B. Rath, D. Kottilil, W. Ji and J. J. Vittal, ACS Appl. Mater. Interfaces, 2023, 15, 26939–26945.
- 6 Z. Zheng, A. H. Alawadhi, S. Chheda, S. E. Neumann, N. Rampal, S. Liu, H. L. Nguyen, Y.-H. Lin, Z. Rong, J. I. Siepmann, L. Gagliardi, A. Anandkumar, C. Borgs, J. T. Chayes and O. M. Yaghi, J. Am. Chem. Soc., 2023, 145, 28284–28295.
- 7 J. Miao, W. Graham, J. Liu, E. C. Hill, L.-L. Ma, S. Ullah, H.-L. Xia, F.-A. Guo, T. Thonhauser, D. M. Proserpio, J. Li and H. Wang, J. Am. Chem. Soc., 2024, 146, 84–88.
- 8 I.-H. Choi, J.-M. Gu, H.-C. Kim, Y. Kim and S. Huh, *Bull. Korean Chem. Soc.*, 2023, 44, 780–787.
- 9 A. H. Alawadhi, S. Chheda, G. D. Stroscio, Z. Rong, D. Kurandina, H. L. Nguyen, N. Rampal, Z. Zheng, L. Gagliardi and O. M. Yaghi, *J. Am. Chem. Soc.*, 2024, **146**, 2160–2166.
- 10 D. Kim, H. Yoo, K. Kim, D. Kim, K. T. Kim, C. Kim, J. Y. Kim, H. R. Moon and M. Kim, *Chem. Commun.*, 2022, 58, 5948–5951.
- 11 J. Kim, C. Na, Y. Son, M. Prabu and M. Yoon, *Bull. Korean Chem. Soc.*, 2023, 44, 507–515.
- 12 I. Hwang, S.-Y. Huang, S. Smith, V. Lynch, R. Custelcean, B. A. Moyer, N. Kumar, V. S. Bryantsev and J. L. Sessler, J. Am. Chem. Soc., 2023, 145, 14387–14394.
- 13 C.-H. Wang, Y.-C. Lin, S. Bhunia, Y. Feng, P. Kundu, C. L. Stern, P.-L. Chen, J. F. Stoddart and M. Horie, *J. Am. Chem. Soc.*, 2023, 145, 21378–21386.
- 14 C. J. Pedersen, J. Am. Chem. Soc., 1967, 89, 2495-2496.
- 15 R. Rencsok, K. A. Jackson, T. A. Kaplan, J. F. Harrison and M. R. Pederson, *Chem. Phys. Lett.*, 1996, **262**, 207–212.

- 16 R. W. Saalfrank, N. Löw, S. Kareth, V. Seitz, F. Hampel, D. Stalke and M. Teichert, Angew. Chem., Int. Ed., 1998, 37, 172–175.
- 17 E. N. Ushakov, S. P. Gromov, O. A. Fedorova, Y. V. Pershina, M. V. Alfimov, F. Barigelletti, L. Flamigni and V. Balzani, I. Phys. Chem. A, 1999, 103, 11188–11193.
- 18 J. Kim, M. Shamsipur, S. Z. Huang, R. H. Huang and J. L. Dye, J. Phys. Chem. A, 1999, 103, 5615–5620.
- 19 J. W. Steed, Coord. Chem. Rev., 2001, 215, 171-221.
- S. P. Gromov, A. I. Vedernikov, N. A. Lobova, L. G. Kuz'mina,
 S. S. Basok, Y. A. Strelenko, M. V. Alfimov and J. A. K. Howard, New J. Chem., 2011, 35, 724–737.
- 21 A. Bey, O. Dreyer and V. Abetz, *Phys. Chem. Chem. Phys.*, 2017, **19**, 15924–15932.
- 22 H.-R. Yu, J.-Q. Hu, X.-H. Lu, X.-J. Ju, Z. Liu, R. Xie, W. Wang and L.-Y. Chu, *J. Phys. Chem. B*, 2015, **119**, 1696–1705.
- 23 I. Oral and V. Abetz, *Soft Matter*, 2022, **18**, 934–937.
- 24 R. M. Izatt, J. S. Bradshaw, S. A. Nielsen, J. D. Lamb, J. J. Christensen and D. Sen, *Chem. Rev.*, 1985, 85, 271–339.
- 25 L. F. Lindoy, The Chemistry of Macrocyclic Ligand Complexes, Cambridge University Press, Cambridge, 1989.
- 26 G. W. Gokel, *Crown Ethers and Cryptands*, The Royal Society of Chemistry, 1991.
- 27 R. E. Wolf, Jr., J. R. Hartman, J. M. E. Storey, B. M. Foxman and S. R. Cooper, *J. Am. Chem. Soc.*, 1987, **109**, 4328–4335.
- 28 G. H. Robinson and S. A. Sangokoya, *J. Am. Chem. Soc.*, 1988, **110**, 1494–1497.
- 29 J. Buter, R. M. Kellogg and F. van Bolhuis, J. Chem. Soc., Chem. Commun., 1991, 910–912, DOI: 10.1039/C39910000910.
- 30 A. J. Blake, G. Reid and M. Schröder, J. Chem. Soc., Chem. Commun., 1992, 1074–1076, DOI: 10.1039/C39920001074.
- 31 Y. Jin, I. Yoon, J. Seo, J.-E. Lee, S.-T. Moon, J. Kim, S. W. Han, K.-M. Park, L. F. Lindoy and S. S. Lee, *Dalton Trans.*, 2005, 788–796, DOI: 10.1039/B415794J.
- 32 M. Heller, Z. Anorg. Allg. Chem., 2006, 632, 441-444.
- 33 R. G. Pearson, J. Am. Chem. Soc., 1963, 85, 3533-3539.
- 34 S. Park, S. Y. Lee, K.-M. Park and S. S. Lee, *Acc. Chem. Res.*, 2012, **45**, 391–403.
- 35 E. Lee, S. Y. Lee, L. F. Lindoy and S. S. Lee, *Coord. Chem. Rev.*, 2013, 257, 3125–3138.
- 36 S. Kim, L. F. Lindoy and S. S. Lee, *Coord. Chem. Rev.*, 2014, **280**, 176–202.
- 37 H. J. Kim and S. S. Lee, *Inorg. Chem.*, 2008, 47, 10807-10809.
- 38 E. Lee and S. S. Lee, *Inorg. Chem.*, 2011, 50, 5803-5807.
- 39 S. Kim, A. D. Siewe, E. Lee, H. Ju, I.-H. Park, K.-M. Park, M. Ikeda, Y. Habata and S. S. Lee, *Inorg. Chem.*, 2016, 55, 2018–2022.
- 40 S. J. Lee, J. E. Lee, J. Seo, I. Y. Jeong, S. S. Lee and J. H. Jung, *Adv. Funct. Mater.*, 2007, 17, 3441–3446.
- 41 C. S. Park, J. Y. Lee, E.-J. Kang, J.-E. Lee and S. S. Lee, *Tetrahedron Lett.*, 2009, **50**, 671–675.
- 42 H. Lee and S. S. Lee, Org. Lett., 2009, 11, 1393-1396.
- 43 E. Lee, H. Ju, I.-H. Park, S. Park, M. Ikeda, Y. Habata and S. S. Lee, *Analyst*, 2020, **145**, 1667–1676.
- 44 E. Lee, H. Ju, I.-H. Park, J. H. Jung, M. Ikeda, S. Kuwahara, Y. Habata and S. S. Lee, *J. Am. Chem. Soc.*, 2018, 140, 9669–9677.

CrystEngComm Communication

- 45 S. Kim, H. Ryu, J. K. Clegg, L. F. Lindoy and S. S. Lee, Inorg. Chem., 2020, 59, 15807-15812.
- 46 S. Kim, I.-H. Park, H.-B. Choi, H. Ju, E. Lee, T. S. Herng, J. Ding, J. H. Jung and S. S. Lee, Dalton Trans., 2020, 49, 1365-1369.
- 47 R. L. Davidovich, V. Stavila, D. V. Marinin, E. I. Voit and K. H. Whitmire, Coord. Chem. Rev., 2009, 253, 1316-1352.
- 48 R. L. Davidovich, V. Stavila and K. H. Whitmire, Coord. Chem. Rev., 2010, 254, 2193-2226.
- 49 R. Luckay, I. Cukrowski, J. Mashishi, J. H. Reibenspies, A. H. Bond, R. D. Rogers and R. D. Hancock, J. Chem. Soc., Dalton Trans., 1997, 901-908, DOI: 10.1039/A605068I.
- 50 L. Shimoni-Livny, J. P. Glusker and C. W. Bock, *Inorg. Chem.*, 1998, 37, 1853-1867.
- 51 M. L. Golden, J. H. Reibenspies and M. Y. Darensbourg, Inorg. Chem., 2004, 43, 5798-5800.
- 52 I. Persson, K. Lyczko, D. Lundberg, L. Eriksson and A. Płaczek, Inorg. Chem., 2011, 50, 1058-1072.

- 53 X. X. Wu, B. Ding, J. H. Li, P. Yang, Y. Wang and G. X. Du, Inorg. Chem. Commun., 2013, 33, 170-174.
- 54 H. Ju, E. Lee, S. Kim, I.-H. Park, J.-H. Lee and S. S. Lee, CrystEngComm, 2016, 18, 2621-2625.
- 55 S. Mirdya, S. Roy, S. Chatterjee, A. Bauzá, A. Frontera and S. Chattopadhyay, Cryst. Growth Des., 2019, 19, 5869-5881.
- 56 Y.-M. Yang, C. Feng, Y.-H. Jiang, D.-H. Du, H. Zhao, G.-N. Zhang and Y.-C. Wang, Transition Met. Chem., 2023, 48,
- 57 G. K. Kole, A. M. P. Peedikakkal, B. M. F. Toh and J. J. Vittal, Chem. - Eur. J., 2013, 19, 3962-3968.
- 58 J. S. Magyar, T.-C. Weng, C. M. Stern, D. F. Dye, B. W. Rous, J. C. Payne, B. M. Bridgewater, A. Mijovilovich, G. Parkin, J. M. Zaleski, J. E. Penner-Hahn and H. A. Godwin, J. Am. Chem. Soc., 2005, 127, 9495-9505.
- 59 https://www.hyperquad.co.uk/hypnmr.htm.