


Metallomics

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

CRITICAL REVIEW

Selenium, selenoproteins and neurodegenerative diseases

Cite this: DOI: 10.1039/x0xx00000x

Bárbara Rita Cardoso,^{*ab} Blaine R. Roberts,^a Ashley I. Bush^{†a} and Dominic J. Hare^{†*acd}

Received 00th January 2015,
Accepted 00th January 2015

DOI: 10.1039/x0xx00000x

www.rsc.org/

It is unsurprising that our understanding of the role of selenium in neurological function is somewhat immature, considering its relatively recent discovery as an essential element to human health. Selenocysteine, the 21st amino acid, is the defining feature of the 25 selenoprotein-encoding genes so far discovered within the human genome. The low abundance of these proteins in the brain belies the integral role they play in normal neurological function, from well-characterised antioxidant activity in the periphery to poorly understood mechanisms that modulate mitochondrial function and response to brain pathology. Selenium has been identified as playing a role in several neurodegenerative disorders, including Alzheimer's and Parkinson's disease, though its function as a 'cause or effect' of disease process remains unclear. This review discusses selenium metabolism in detail, specifically with regard to the role it plays within the central nervous system, and examines the most current literature investigating how selenium may be involved in chronic diseases of the central nervous system.

Introduction

The observation that selenium (Se) prevented necrotic liver damage in mice by Schwarz and Foltz in 1957 was the first direct experimental evidence that this element was essential to normal health.¹ Almost twenty years later, Flohe *et al.*² confirmed the importance of Se by identifying it as an essential cofactor of glutathione peroxidase (GPx). Twenty-five mammalian selenoprotein-encoding genes have since been identified,³ all containing selenocysteine (Sec) residues, which are analogous to cysteine (Cys) with a selenol moiety replacing the thiol group.⁴ From these genes, it is estimated that more than 30 unique selenoproteins are synthesised in mammals,⁵ and it is likely more will be characterised in the future. Selenoproteins have diverse roles throughout the body, acting as antioxidants, modulators of immune function, detoxification agents for heavy metals and xenobiotics, and participants in thyroid hormone metabolism.⁶

Selenium is essential to human health, and the antioxidant activity (or lack thereof) of specific selenoproteins is of particular interest in diseases of the central nervous system. The brain is the most metabolically active of all organs in the body, and although it represents only 2% of the total mass, cerebral metabolism accounts for about 20% of the oxygen and 25% of the glucose consumed by the human body.⁷ In concert with this high metabolic activity, the high concentration of polyunsaturated fatty acids in the brain makes it more

vulnerable to peroxidation. A selective vulnerability within certain neurons to oxidative stress may arise from deficiencies in antioxidant enzyme activity,⁸ which is a likely upstream event in the pathogenesis of several neurodegenerative diseases,⁹ including Alzheimer's disease (AD),¹⁰ Parkinson's disease (PD)¹¹ and other neurodegenerative disorders. Selenoproteins, with their inherent antioxidant activity, thus play essential roles as part of the free radical defence system, though the function of these proteins is not necessarily restricted to this singular classification. This review summarises the current data and important concepts concerning selenoproteins on brain function and the association between Se and diseases of the brain.

Selenoprotein synthesis

Dietary Se first must be converted to selenide (Se²⁻), which serves as the donor for the incorporation of Se into selenoproteins. Inorganic Se forms, such as selenite (SeO₃²⁻), are converted to Se²⁻ via the glutathione-glutaredoxin and thioredoxin pathways, while organic forms are metabolised to Se²⁻ by selenocysteine lyase (Sclly) or trans-selenation (analogous to trans-sulfuration).¹²

The incorporation of Se into proteins was described in detail by Turanov *et al.*,¹³ recently reviewed by Labunskyy *et al.*,¹⁴ and is summarised in Figure 1. After conversion of dietary Se to Se²⁻, selenophosphate (SePO₃³⁻) is produced in a reaction




Fig. 1 Selenoprotein synthesis pathway. Dietary Se intake, as either inorganic or organic species is metabolised to Se^{2-} , and the SeH_3PO_3 , which is incorporated into proteins via a unique tRNA pathway, where a seryl (Ser) residue is replaced with selenocysteine (Sec) prior to integration into mammalian proteins. Deficiencies in key metabolic factors involved in selenoprotein synthesis have been associated with specific neurological disorders, marked here in red. The inset in the dashed blue box is the simplified cloverleaf structure of mammalian tRNA[Sec] (adapted from Birringer *et al.*¹⁹⁶), showing variable (in blue) and modifiable (in red) bases. Abbreviations: Ψ = pseudouridine; EFSec = Sec elongation factor; eIF4a3 = eukaryotic initiation factor; $i^6\text{A}$ = N^6 -isopentenyladenosine; L30 = ribosomal protein L30; mcm^5U = 5-methoxycarbonylmethyluridine; m^1A = 1-methyladenosine; mcm^5Um = 5-methoxycarbonylmethyluridine-2'- O -methylribose; SBP2 = Sec insertion sequence (SECIS) binding protein-2.

catalysed by selenophosphate synthase 2.⁴ Selenocysteine (Sec) is incorporated into proteins via its own unique tRNA, (tRNA[Ser]Sec). tRNA[Ser]Sec decodes the UGA codon as Sec instead of a stop codon, as is typical of other amino acids, and thus prevents premature termination. This is possible due to the presence of a Sec insertion sequence (SECIS) element, a specific *cis*-acting stem-loop structure, in the 3'-untranslated region (UTR) of selenoprotein mRNA that works as a platform for RNA-binding proteins. SECIS forms complexes with two trans-acting elements: the specific elongation factor (EFSec) and the SECIS binding protein 2 (SBP2).¹⁵ EFSec interacts with SBP2 to recruit tRNA[Ser]Sec and mediates insertion of Sec into nascent protein chains in response to the UGA codon.¹⁵ Among the three distinct domains of SBP2, the Sec incorporation domain and a COOH-terminal RNA-binding domain are specifically involved in SECIS binding. Additional

SECIS binding proteins have been identified, including ribosomal L30, which is part of the basal Sec insertion machinery and has a similar RNA-binding domain to SBP2;¹⁶ a eukaryotic initiation factor (eIF4a3) that links Se status and differential selenoprotein expression.¹⁷ and nucleolin, which demonstrates a high affinity for SECIS, though its complete function is not currently known.¹⁸ Recently, Bröcker *et al.*¹⁹ used *Escherichia coli* formate dehydrogenase mutants to reassign 64 codons encoding a critical Sec residue, finding that 58 of these were able to site-specifically incorporate Sec. At 15 sense codons selenoprotein synthesis could be promoted compared to the 3 stop codons. This study provides critical new information about how selenoprotein production can be manipulated through genetic recoding, and provides exciting new opportunities for additional studies to further elucidate the

mechanisms involved in selenoprotein synthesis, which remains a relatively new frontier in molecular biology.

It is known that selenoprotein synthesis is critical for viability of life, as shown by tRNA[Ser]Sec (*Trsp*) knockout mice, which present an embryonically lethal phenotype. However, *Trsp* null mice allow tissue-specific manipulation of tRNA[Ser]Sec, and this model has been used to investigate the importance of selenoprotein synthesis to brain metabolism. Wirth *et al.*²⁰ showed that inactivation of selenoprotein synthesis in *Trsp* null neurons abolished the developmental appearance of parvalbumin-positive (PV) interneuron cells, characterising progressive neurodegeneration. This was later confirmed by additional studies that demonstrated this damage was restricted to specific brain areas, such as the somatosensory cortex and hippocampus, confirming the essential requirement of selenoprotein synthesis for maintenance of CNS metabolism.^{21,22} In humans, two mutations in the gene encoding *O*-phosphoseryl-tRNA[Sec]:selenocysteinyl-tRNA[Sec] synthase (SepSecS) were associated with progressive cerebellocerebral atrophy (PCCA), an autosomal-recessive phenotype of nondysmorphic profound mental retardation, severe spasticity and microencephaly.²³ These gene alterations abrogate SepSecS activity and thus block the synthesis of tRNA[Ser]Sec, and subsequent Se incorporation into selenoproteins.

In vivo studies have shown that Cys can replace Sec in the Sec biosynthetic organisation, generating Cys-tRNA[Ser]Sec and the subsequent insertion of Cys in some selenoproteins.²⁴ This is possible due to either substitution of Se²⁻ by sulfide (S²⁻) in the SPS2 catalysed reaction; or substitution of SePO₃³⁻ by thiophosphates in the reaction with the *O*-phosphoseryl-tRNA[Ser]Sec intermediate during Sec-tRNA synthesis. The same study showed that the incorporation of Cys in selenoproteins is dependent of Se availability to the cells. Tobe *et al.*²⁵ showed that some antibiotics could also interfere with Sec insertion in murine EMT6 cells. Cells treated with doxycycline, chloramphenicol and G418 caused lower Sec insertion in thioredoxin reductase 1 (Trxr1), and GPx1 and 4. However, Sec was also exchanged by arginine and tryptophan in addition to Cys. The specific mechanisms that are associated with mis-insertion of amino acids in place of Sec and how they are associated with changes in selenoproteins activity need to be clarified, but these reports show another way selenoprotein hierarchy is potentially regulated, which may also include as-yet undetermined environmental factors.

Selenoprotein P and brain selenium hierarchy

Selenoprotein expression is modulated by Se availability, and a hierarchy is observed in response to dietary Se intake. In cases of Se deficiency the synthesis of some proteins is maintained at a higher level than that of others, suggesting that insufficient available Se results in competition between transcripts of different selenoproteins.²⁶ Iodothyronine deiodinase 1 is atop this hierarchy and is comparatively the least affected by restricted dietary Se; glutathione peroxidases GPx2 and GPx4

are less vulnerable to Se deficiency than GPx1 and GPx3; and selenoprotein P (SelP) is in an intermediary position on this selenoprotein transcription ladder.²⁷

Some mechanisms underlying Se regulation of the selenoprotein transcriptome have been identified. During Se deficiency, non-essential selenoproteins are down regulated via nonsense-mediated mRNA decay, which decodes the UGA-Sec codon as a termination codon causing RNA degradation.^{28,29} Also, Se levels directly correlate with the degree of Um34 methylation (the single methyl group differentiating the two tRNA[Ser]Sec isoforms³⁰), one step of maturation of tRNA[Ser]Sec which can be altered as a result of some mutations. The Um34-containing isoform of tRNA[Ser]Sec; 5-methoxycarbonylmethyl-2'-O-methyluridine (mcm⁵Um), is recruited for efficient expression of the stress-related selenoproteins; and the non-Um34-containing isoform, 5-methoxycarbonylmethyluridine (mcm⁵U), is not required specifically. The distinct use of these tRNA[Ser]Sec isoforms shows that this mechanism contributes to Se regulation for the purpose of selenoprotein synthesis.²⁸

The hypothesis that explains the hierarchy between selenoprotein syntheses suggests that they are classified according to their importance to the body. However, it is important to note that Se concentration is highly variable within the periphery (*i.e.* liver, kidneys, etc), though in the brain Se levels remain relatively stable by comparison during periods of deficiency.³¹ Additionally, the ability of organs outside the CNS to retain Se is also variable. The CNS, together with reproductive and endocrine organs, has the highest priority for Se *both* uptake and retention during Se deficiency, although the Se concentration in the CNS is low in comparison with other organs.³² Though interrelated, supply and retention are not exclusively linked, and it is thus a testament to the importance of Se in brain function that rigorous mechanisms to ensure adequate Se levels are available in times of dietary shortage are in place. The brain's top position on a tissue-specific grading further indicates that Se is essential for its normal metabolism.³³ This is primarily due to the high expression of apolipoprotein E receptor-2 (apoER2) at the blood-brain barrier (see below) to facilitate Se uptake and strong SelP-apoER2 interaction within the CNS providing an adequate pool of available Se. Selenium deficient rats presented approximately 29% lower Se levels in the brain compared to animals with a sufficient Se supply, which is not nearly as dramatic as the 99% and 92% decrease in Se concentration in the liver and kidneys of deficient animals.³⁴ Se distribution is heterogeneous among different brain regions, and grey matter appears to have preference for greater uptake. Imaging using a ⁷⁵Se radiotracer combined with autoradiographic localisation and proteomic analysis to profile Se levels in different brain regions of rats fed a Se-replete diet revealed that Se was enriched in choroid plexus, putamen, parietal inferior lobule and occipital cortex.³⁵ This observation has been reflected in human brains, where the highest Se concentration occurs in grey matter, with the putamen, parietal




Fig. 2 a) SelP, the master protein driving Se bioavailability is synthesised in the liver from both inorganic and organoselenium compounds accessible through dietary sources. At the blood-brain barrier, SelP releases Se into the CNS via the ApoER2 receptor, which is in turn incorporated into newly-formed SelP in astrocytes (figure adapted from Burk *et al.*³⁵), or is transported directly to neurons (dashed arrow). SelP is made available to neurons via the same membrane-bound ApoER2, where additional selenoproteins essential to neurological function (blue box) are biosynthesised. b) In cases of deficient Se intake Um34 methylation is downregulated, causing premature termination of protein synthesis via failure of tRNA[Ser]Sec to decode the UGA codon as Sec instead of a stop codon, and misincorporation of selenocysteine (Sec) as cysteine (Cys; in red). These three pathological features may have downstream effects on a range of neurological disorders (red boxes).

inferior lobule and occipital cortex in particular, while the lowest levels are found in the cerebellum and medulla.³⁶

The mechanism that maintains Se levels in the brain at the expense of other organs, even during times of deprivation, is mainly orchestrated by SelP, which is recognised as the most important Se supply to different tissue types. The role of SelP to deliver Se to cells is facilitated by the presence of ten SECIS forms: one is situated in the larger N-terminal region, which acts as an enzyme; and nine are located in a Sec-rich C-terminal domain, providing a mechanism for transporting the highly reactive Se atom safely, incorporated into the protein as Sec.³⁷ Deletion of the conserved, non-SECIS containing 3'UTR of

SelP does not affect the efficiency of incorporation of multiple Sec residues into SelP *in vitro* whilst still remaining sensitive to ambient Se levels, indicating that the 3'UTR are not involved in Se incorporation *per se*.³⁸ The structure of plasma SelP remains unresolved, and there is some contention as to the number of circulating isoforms of the protein. Ballihaut *et al.*³⁹ recently reported three isoforms in a human plasma standard reference material, two of which incorporated Se, suggesting the third was in fact a truncated variant.

Burk *et al.*⁴⁰ recently proposed the mechanism by which Se is delivered to the brain (represented in Figure 2a). SelP from plasma is endocytosed following docking with apoER2 at the

1 blood-brain barrier (BBB) in brain capillary endothelial cells
2 (BCECs) and choroid plexus epithelial cells, which facilitates
3 Se transport, in an as-yet unknown chemical form, into the
4 brain. Selenium is then delivered to brain cells via SeLP
5 synthesised by astrocytes on the abluminal side of BCECs,
6 composing both the main transport mechanism and distinct pool
7 of Se. Direct transport of SeLP to neurons from BCECs has not
8 been excluded. Neurons regulate Se levels via apoER2-
9 mediated uptake of SeLP,⁴¹ which has obvious implications for
10 AD due its role in neurodevelopment and synaptic function. In
11 the periphery, apoER2 is widely expressed in the body and is
12 also involved in Se uptake in myoblasts⁴² and testis cells.⁴³ In
13 the CNS, apoER2 is also expressed in choroid plexus and on
14 PV interneurons in the hippocampus, inferior colliculus, medial
15 septum, red nucleus, reticular thalamus and cerebellum.^{40,44}
16 ApoER2 expression is regulated by physiological factors
17 related to developmental stage; fetal brain mRNA ApoER2 are
18 more than 9 times greater than that observed in adult brain, and
19 apparently not influenced by Se status.⁴⁰ Although deletion of
20 *apoER2* gene does not affect the whole-body Se levels,
21 *apoER2*^{-/-} and *SelP*^{-/-} mice have lower Se levels in brain,^{45,46}
22 and this may be associated with uptake of Se during specific
23 developmental windows.⁴⁰ In this study, fetal *apoER2*^{-/-} mice
24 brains contain 75% less Se levels compared to age-matched
25 *apoER2*^{+/+} mice, decreasing to 39% when reaching adulthood.
26 Moreover, *apoER2*^{-/-} mice had no SeLP in BCECs or choroid
27 plexus cells at the BBB, suggesting a key role of ApoER2 in
28 brain Se uptake, and also the presence of an unknown
29 redundancy mechanism for Se transport. It remains unclear if
30 other members of the lipoprotein receptor family may
31 participate of SeLP uptake, since Burk *et al.*⁴⁷ observed an
32 alternative mechanism that does not appear to depend directly
33 on selenoproteins participating on the transfer of Se from
34 pregnant dam to the fetus, and it is unclear if this mechanism is
35 conserved in the postnatal period.

36
37
38 Though primarily expressed in astrocytes, neurons have also
39 been identified as a source of endogenous SeLP^{48,49} in all the
40 areas of the brain, though particularly high expression was
41 identified in the putamen and substantia nigra.⁵⁰ In contrast to
42 *apoER2*^{-/-} animals, *SelP*^{-/-} mice present generalised Se
43 deficiency, and in brain this gene deletion causes severe
44 neurological dysfunction.^{47,51} *SelP*^{-/-} mice showed decreased
45 density of PV interneurons in different brain areas, most
46 marked in the inferior colliculus. This was associated with
47 higher oxidative damage and a behavioural phenotype,
48 characterised by impairments in contextual fear extinction,
49 latent inhibition and sensorimotor gating. These data suggest
50 that besides the effect on PV-expressing neurons causing
51 increased vulnerability to oxidative damage, the behavioural
52 deficits resulting from *SelP* deletion are likely associated with
53 impaired GABAergic function of the inferior colliculus.⁴⁴ Both
54 *SelP*^{-/-} and *apoER2*^{-/-} mice, when maintained on a Se-deficient
55 diet, are more vulnerable to neuronal damage and eventual
56 neurodegeneration. In this way, Burk *et al.*⁴⁰ reported that
57 extremely Se-deficient *apoER2*^{-/-} and *SelP*^{-/-} mice displayed
58 neurological dysfunction, while the littermate wild type mice

did not show equivalent disruption, despite a decrease in brain
Se levels. In another important study,⁴⁶ progressive
neurodegeneration was evaluated in brains of *SelP*^{-/-} postnatal
mice maintained on a Se-deficient diet after weaning. Higher
lipid peroxidation and decreased dendrite length and spine
density was observed, and the areas more susceptible to
neurodegeneration during this post-weaning developmental
window were the forebrain bundle, somatosensory cortex and
lateral striatum. In the hippocampus decreased dendrite length
and spine density was also present, which might be associated
with the disrupted hippocampal-dependent learning and
alteration to synaptic transmission observed in these mice.

As Sely is essential for breakdown of Sec into L-aniline and
Se²⁻, it has been proposed that this enzyme plays important role
in Se metabolism in different tissue, and interacts with SeLP to
maintain adequate selenoprotein function in the brain. Although
Sely^{-/-} mice do not demonstrate neurological dysfunction,⁵¹
deletion of both *Sely* and *SelP* exacerbates the neurological
phenotype of *SelP*^{-/-} mice, which display impaired motor
coordination and locomotion compared with *SelP* null mice
alone. Moreover, *Sely*^{-/-}*SelP*^{-/-} mice have lower expression of
Trxr1, GPx1 and selenoprotein M (SelM) than *SelP*^{-/-} mice,
showing that interplay of Sely and SeLP helps maintain
selenoprotein levels within the brain.⁵²

Selenium and other selenoproteins: role in brain function

Selenium appears to have a multifaceted role in the nervous
system, with a number of studies showing its importance to
normal brain physiological function. The antioxidant activity of
selenoproteins in the CNS is well established, and lower levels
of Se has been associated with brain injury.⁵³ Selenium
treatment as organic selenomethionine and inorganic SeO₃²⁻
resulted in a similar increased activity of antioxidant enzymes,
scavenging of reactive oxygen species (ROS) and reduced
protein carbonyl content of patulin-induced brain damaged
rats,⁵⁴ and reduced the pro-oxidant effects of glyceryl trinitrate
in a rat model of migraine headaches.⁵⁵ Administration of
SeO₃²⁻ to suckling pups whose dams had methimazole-induced
hypothyroidism decreased total oxidant status and reduced
degenerative changes in the granular cell layer of the dental
gyrus,⁵⁶ and increased levels of brain-derived neurotrophic
factor (BDNF) in the hippocampus and cerebellum of both
hypothyroid and control groups.⁵⁷ In a rat model of postnatal
protein malnutrition oral Se and zinc (Zn) supplementation
increased not only antioxidant enzyme activity, but also
corrected the neurobehavioral deficits caused by a protein-
deficient diet.⁵⁸

Besides the role of Se as essential component of the
antioxidant system in the brain, studies have also shown that Se
can act in response to oxidative stress through regulation of
Ca²⁺ channels and mitochondrial biogenesis. Mice dorsal
ganglion root cells treated with Se showed reduced Ca²⁺ release
induced by H₂O₂ when compared with non-treated cells, with

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

corresponding decreases in apoptosis and lipid peroxidation levels, and increased levels of GPx and glutathione (GSH) activity.⁵⁹ In a traumatic brain injury animal model exhibiting oxidative stress, increased apoptosis and activation of transient receptor potential vanilloid 1 (TRVP1) cation channels, Se treatment modulated Ca²⁺ entry through TRVP1 in neurons, reduced levels of ROS, prevented apoptosis, and reduced activity of caspases 3 and 9. Since brain injury induces Ca²⁺ influx and mitochondrial dysfunction in neurons, these data show that Se can increase cell viability not only as an antioxidant, but also by modulating apoptotic pathways and mitochondrial function.⁶⁰ Studies with murine hippocampal HT22 cells treated with SeO₃²⁻ clearly showed improved mitochondrial function, even after glutamate exposure and hypoxia. This was associated with activation of complexes I, II+III and IV, and increased mitochondrial respiratory rate. Selenium administration increased levels of peroxisome proliferator-activated receptor γ -coactivator-1 α (PGC-1 α) and nuclear respiratory factor 1 (NRF1), two important nuclear transcriptional factors associated with mitochondrial biogenesis. This role of Se appears to be mediated through activation of the cyclic adenosine monophosphate response element-binding protein (CREB) and activated protein kinase (AKT) cycles.⁶¹⁻⁶³

Synthetic organoselenium compounds have been studied as potential antioxidant therapies due their lower toxicity compared to inorganic forms of Se. Diphenyl diselenide [(PhSe)₂] mimics GPx activity, producing both beneficial or harmful effects, depending on dose and route of administration.⁶⁴ Acute treatment of rat hippocampal slices with (PhSe)₂ modified various proteins involved in glutamate signaling damage to astrocytes.⁶⁵ Conversely, (PhSe)₂ demonstrated a neuroprotective activity in animal models of autoimmune encephalomyelitis and hypothyroidism due its antioxidant and anti-inflammatory role.^{66,67} A derivative of (PhSe)₂, m-trifluoromethyl (m-CF₃-PhSe)₂ and the Se-containing salicylic acid derivative 5-5'-diselenediylbis-(2-hydroxybenzoic acid) both exhibit antinociceptive action by interaction with serotonergic system, as well as an anti-inflammatory role in mice.^{68,69} Selenium supplementation as a promoter of antioxidant activity does appear to be somewhat dependent on the chemical form; 3-methyl-1-phenyl-2-(phenylseleno)oct-2-en-1-one was less effective as an antioxidant compared to SeO₄²⁻, though it did reduce adenosine deaminase activity, suggesting a potential immunomodulatory property.⁵⁴ Musik *et al.*⁷⁰ examined two other Se-containing molecules; (4-(o-tolyl)-selenosemicarbazide of 2-chlorobenzoic acid; 3-(2-chlorobenzoylamino)-2-(o-tolylimino)-4-methyl-4-selenazoline) reporting differential impacts on brain antioxidant mechanisms, including GPx and superoxide dismutase-1 (SOD-1). These contradictory results regarding Se supplementation on brain function are likely due to the variety of chemical structures, dose, route of administration and the animal models used. More targeted studies regarding the role and the safety of these compounds are

required before they can be considered as potential antioxidant therapeutics in humans.

The apparent neuroprotective role of Se supplementation in the brain is not only directly associated with antioxidant effects of organic and inorganic Se species, but also suggests a role in *de novo* selenoprotein synthesis, as demonstrated by increased activity of certain selenoproteins following Se treatment (Figure 2; Table 1). Methamphetamine treatment decreased levels of GPx1 and GPx4 in SH-SY5Y cells; an effect that was arrested with addition of SeO₃²⁻ 24-hours after methamphetamine exposure.⁷¹ In their patulin-induced brain damaged rats, Song *et al.*⁷² showed that, although both SeO₃²⁻ and selenomethionine were able to offer some degree of increased antioxidant activity, the organic form of Se was more efficient in increasing the expression and activity of TrxR, GPx1 and GPx4.

The GPx family of enzymes are expressed in both neurons and glia,^{73,74} and are recognised as one of the most important mechanisms protecting against ROS-induced damage.⁷⁵ Among these isoenzymes, GPx4 protects cells against lipid hydroperoxide damage⁷⁶ and is the most expressed GPx in brain, found in neurons of cerebellum, hippocampus and hypothalamus.⁷⁷ Following brain injury, GPx4 is also observed in reactive astrocytes, suggesting that this GPx may have an important role in the protective cellular response to stress-induced oxidative damage and apoptosis.⁷⁸ Autooxidation of dopamine to the reactive dopamine quinone causes a decrease in GPx4 activity by covalently modifying this mitochondrial enzyme,⁷⁹ suggesting that antioxidant properties of GPx4 are restricted by dopamine breakdown in neurons. Wirth *et al.*²⁰ showed that mice with a neuron-specific deletion of *GPx4* had mild neurological dysfunction, and that GPx4 deficiency may contribute to dysfunction in other selenoproteins expressed in the brain, including TrxR, SelM and selenoprotein W (SelW).

TrxR is abundant throughout the CNS. In the mouse brain TrxR1 (the cytosolic isoform) is abundantly expressed in glial cells of the cerebellar white matter and TrxR2 (the mitochondrial isoform) is observed mainly in the cell bodies of neurons localised in the Purkinje and molecular cell layers in the cerebellum.⁸⁰ This selenoenzyme plays an important role in maintaining the redox balance inside cells. It forms, together with thioredoxin (Trx), a disulfide reductase system.⁸¹ Mitozo *et al.*⁷⁵ showed that TrxR inhibition resulted in increased DNA damage and poor cell viability following a H₂O₂ challenge in neuronal C6 cells, though it did not impede the ability of N2a cells to dispose of peroxides. However, specific inhibitors used to estimate the relative role of the Trx2, glutathione system and catalase for hydrogen peroxide detoxification in different preparations from rat brain showed that Trx2-system contributes to over 60% of the intra-mitochondrial H₂O₂ turnover, whereas the glutathione system contributes only about 20% to H₂O₂ detoxification reactions in the rat hippocampus.⁸² Recently, Cohen-Kutner *et al.*,⁸³ using two thioredoxin mimetic peptides (NAc-Cys-Pro Cys-amide; CB3 and Ac-Cys-Gly-Pro-Cys-amide; CB4) as a treatment for oxidative stress in SH-SY5Y human neuroblastoma cells, observed anti-inflammatory

1 and antioxidant effects of both peptides, as well as reduced
2 apoptosis markers. The potentially protective role of TrxR is
3 also important for dopaminergic cells, which exhibit high levels
4 of oxidative stress when undergoing parkinsonian degeneration.
5 In cells exposed to 6-hydroxydopamine (6-OHDA) and
6 paraquat, inhibition of TrxR potentiated mitochondrial
7 dysfunction, decreased maximal and reserve respiratory
8 capacity, and increased H₂O₂ levels and cell death through
9 oxidative stress. Also, TrxR2 deficient cells showed decreased
10 basal mitochondrial oxygen consumption rates.⁸⁴
11 Immunoreactivity of TrxR2 was increased in the spinal neurons
12 and hippocampus of aged dogs, suggesting that this
13 selenoenzyme is important for maintaining neuronal
14 homeostasis following the increase in oxidative stress
15 experienced during normal ageing.⁸⁵ Human astrocytomas,
16 which are the most common form of gliomas, were found to
17 express TrxR in a manner that correlated with both tumour
18 grading and Ki-67 labelling index (a marker of cell
19 proliferation), with both quantitative reverse transcription
20 polymerase chain reaction and immunostaining confirming
21 higher TrxR levels correspond to tumour severity.⁸⁶ Though no
22 clear causative link between TrxR activity and astrocytoma
23 proliferation has been identified, this association with cancer
24 progression and TrxR expression is a tantalising research
25 direction considering that, besides its antioxidant role, the TrxR
26 class of enzymes contributes to several physiological and
27 biochemical functions, including cell growth, maintenance and
28 programmed death.

29
30 Most selenoproteins have the conserved Cys-X-X-Sec (or
31 *CXXU*) redox motif, including selenoproteins H, W, T and O.
32 Other selenoproteins include Cys-X-X-Cys (*CXXC*) in TrxRs,
33 Cys-X-Sec (*CXU*) in selenoprotein-15,⁸⁷ Sec-X-X-Cys (*UXXC*)
34 in SelP,⁸⁸ as well as Sec-X-X-Ser (*UXXS*) and Sec-X-X-Thr
35 (*UXXT*), identified in a range of prokaryotic selenoproteins
36 with mammalian homologs.⁸⁹ Their role it is not well
37 elucidated, though it is thought they act as thiol-dissulfide
38 oxidoreductases that participate in the formation of disulfide
39 bonds.^{37,90-94} SelM has an endoplasmic reticulum (ER)/Golgi-
40 directing signaling peptide with a short *N*-terminal extension,
41 the *CXXU* motif, and a thioredoxin-like domain in the middle.⁹⁵
42 It is characterised by a two-layer $\alpha/\beta/\alpha$ sandwich with a
43 $\beta\alpha\beta\beta\alpha$ secondary structure pattern, and an ER retention signal
44 tetrapeptide in the *C*-terminal.^{96,97} In the mouse brain, SelM
45 expression is highest in the olfactory bulb and cerebellum, with
46 intermediate levels found in the hypothalamus and brain stem
47 and low expression in the hippocampus and cerebral cortex.⁹⁸
48 The precise role of SelM is not well understood; some studies
49 using mice overexpressing this protein have shown that it
50 promotes antioxidant activity in brain, in this case increasing
51 the activity of other antioxidant enzymes including SOD-1 and
52 GPx, which could be enhanced with further dietary
53 supplementation of Se.⁹⁹⁻¹⁰² SelM knockdown primary cortical
54 neurons show disturbed Ca²⁺ homeostasis compared to
55 controls.⁹⁹ Pitts *et al.*⁹⁸ observed that the deletion of *SelM* gene
56 in mice did not result in morphological changes to the brain,
57 nor did it result in motor or cognitive deficits and anxiety-like

behavior. However, these mice presented an obesity phenotype
accompanied by higher leptin levels and elevated hypothalamic
leptin resistance when compared with wild type controls. As
SelM is expressed in high quantities in the hypothalamus, the
authors hypothesised that deletion of this gene adversely
affected hypothalamic thyroxine balance and cause metabolic
dysfunction, indicating that SelM plays an important role in
maintaining redox balance in brain areas associated with high
metabolic output, though further investigation is obviously
required to confirm this supposition.

Yim *et al.*¹⁰⁰ showed that the association between SelM
overexpression and Se treatment *via* intraperitoneal injection
contributed to the activation of ERK MAPK signaling pathway,
an important mechanism involved in the modulation of protein
phosphorylation. Additionally, hippocampal mouse cells and
C8-D1A cerebellar cells overexpressing SelM exhibited
decreased cytosolic Ca²⁺ in response to oxidative stress, without
affecting basal Ca²⁺ levels, suggesting that SelM can modulate
the apoptotic pathway by mediating Ca²⁺ release from internal
cell stores.⁹⁹ Another proposed mechanism for the role of SelM
in providing neuroprotection was the observation that SelM
interacts with galectin-1,¹⁰³ a β -galactoside-binding lectin
associated with modulation of microglia activation in
neuroinflammation, neuronal myelination and neuronal stem
cell proliferation.^{104,105} Additionally, it was recently shown that
overexpression of SelM could change the expression of other
proteins; Kim *et al.*¹⁰² showed that, when compared with wild
type controls, mice overexpressing SelM upregulated
expression of several different genes, including creatine-kinase
B-type and *LDH-B*, essential for energy homeostasis; the E3
ubiquitin-protein ligase *RING1* that participates in cellular
defense against environmental insult; *eIF-4H*, which facilitates
the binding of initiator tRNA to ribosomes, regulating protein
synthesis; and *SylXV*, that regulates Ca²⁺-dependent membrane
trafficking, and thus is crucial for maintenance of adequate
neurotransmitter release. On the other hand, these mice
demonstrated downregulation of CENP-N, which mediates
mitotic progression, assembly of kinetochore proteins and
chromosome segregation; dihydropyrimidinase-related protein
2, that plays important role in neurite guidance; and proteasome
subunit K. This study strongly suggests that the protective role
of SelM may in fact be carried out indirectly through the
modulation of different genes that regulate cell viability.

SelW is abundant in brain of chickens,^{106,107} and is highly
stable in rat brain even during Se-deficiency. *In vivo*, where Se
supply is entirely dictated by the media in which cells are
grown, SelW expression is highly dependent on ambient Se
concentrations. Low levels (10⁻⁹ mol L⁻¹) of Se yielded the
highest levels of SelW mRNA expression, which progressively
increased over time, along with better general embryonic
chicken neuronal health. High levels of Se (up to 10⁻⁵ mol L⁻¹)
were obviously neurotoxic, with SelW mRNA expression
suppressed.¹⁰⁸ This clearly indicates that neurons depend on an
external Se pool to which it can draw from, and has a place
lower on the Se brain hierarchy. It is more likely that delivery
of Se from astrocytes as SelP to endogenous neuronal apoE2

receptors is the more controlled mechanism of neuronal Se uptake, as opposed to direct translocation of SelP immediately following BBB transit. This study, though clearly useful in investigating what, if any, buffering capacity neurons have for excessive Se, does however highlight a very pertinent point when considering *in vitro* studies of Se (or any other low-level essential element) metabolism: what conclusions can appropriately be drawn when the multicellular interplay that typically regulates Se hierarchy are removed and the system is viewed in isolation? As such, these studies should be also considered somewhat *in vacuo*; in this case purely as an example of how excessive Se levels impact upon SelW expression and neurotoxicity.

In murine brain tissue, SelW is expressed in axons and dendrites of neurons in several nuclei, mainly the cortex, hippocampus and cerebellum. It is highly abundant in apical dendrites of most pyramidal neurons in the hippocampus, cingulate cortex and piriform cortex, and is also expressed in synapses.^{51,109,110} However, as observed for other selenoproteins, SelW expression is dependent on the master regulator SelP, as SelP deleted mice present a reduced level of SelW mRNA.¹¹⁰ SelW is present in survival motor neuron protein (SMN) complexes, the major components of the spliceosome (the machinery that carries out pre-mRNA splicing).¹¹¹ This particularly interesting study showed that SelW is in part dependent of SMN-complexes due to its neuronal transport along neurites, since SelW was decreased by 17% in neurites of SMN-depleted NSC-34 cells. The role of SelW in the brain remains unclear: its redox motif suggests that this selenoprotein acts as antioxidant in different cells, though in muscle cell cultures depletion of SelW was compensated for by alternate antioxidant proteins, suggesting its major function is not antioxidant activity alone.¹¹² Also, because SelW is highly expressed in brain areas associated with synaptic plasticity, Raman *et al.*¹¹⁰ suggested that it is essential for maintaining redox homeostasis and mediates proper synaptic adaptation and development.

Several selenoproteins are not highly expressed in brain, such as selenoproteins S (SelS) and H (SelH), though recent studies are attempting to determine their role in CNS, suggesting that whilst they exist in low abundance, they may be essential participants in normal brain function. Speciation of selenoproteins is analytically challenging, and *in situ* hybridisation often lacks the necessary sensitivity to probe selenoprotein RNA where expression is very low. Selenoprotein S appears to have a neuroprotective effect, and is found mainly in neurons, sparingly in astrocytes; and animal studies have reported that it is typically confined to the hippocampus and cortex.¹¹³⁻¹¹⁵ SelS expression is induced in reactive astrocytes as part of an activation program triggered by astrogliosis-inducing stimuli.¹¹⁴ As a result, SelS overexpression decreased ER-stress and IL-6 release in astrocytes. These results were confirmed by Liu *et al.*,¹¹⁵ who observed increased SelS expression after 3-7 days of ischemic reperfusion, suggesting that SelS modulates astrocyte function by reducing the synthesis and release of inflammatory

cytokines and limiting degradation of unfolded proteins. Considering that previous studies have shown that SelS has an important anti-inflammatory role in different tissue types,^{116,117} other work regarding the role and therapeutic potential of this selenoprotein in brain are encouraged. In the same way, although SelH is only moderately expressed in the brain,¹¹⁸ overexpression in neuronal cells improved mitochondrial biogenesis and functional performance. Overexpression of SelH in HT-22 neurons increased the respiratory function of mitochondria and preserved UVB stress-induced mitochondrial membrane potential depolarisation. The study of neurons either overexpressing SelH or with the gene ablated revealed that this selenoprotein improved mitochondrial biogenesis signaling via PGC-1 α , NRF1 and mitochondrial transcription factor A (Tfam), likely mediated by modulation of CREB dependent PGC-1 α activation.^{61,62} The protective role of SelH is also possibly linked to its capacity to maintain genomic stability, protecting the cells against replicative senescence under chronic oxidative stress.¹¹⁹ According to this hypothesis, besides its direct antioxidant role arising from the presence of the CXXU motif, which prevents ribosomal DNA oxidation in the nucleoli, SelH may induce *de novo* synthesis of GSH and phase II detoxification enzymes. Although this theory was not evaluated in brain cells, it does pose one possibly important role of this selenoprotein in neurodegenerative diseases associated with ageing. The examples described here exemplify the ways by which selenoproteins can exert a neuroprotective role, even though they are expressed at relatively low levels.

Selenium in brain diseases

Alzheimer's disease

Alzheimer's disease (AD) is a neurodegenerative condition characterised by progressive loss of memory and cognition, and compromising of daily activities. Its key pathological features are neuron death, loss of synaptic connections, extracellular deposition of beta-amyloid (A β) protein aggregates (plaques), and intracellular precipitation of hyperphosphorylated tau protein, which manifest as neurofibrillary tangles.^{120,121} Oxidative stress is a central component of AD pathogenesis, with ROS adversely affecting mitochondrial function, synaptic transmission, axonal transport, and stimulating neuroinflammation.¹²³⁻¹²⁵

Studies in humans have reported negative correlations between cognitive decline and Se levels.¹²⁶ Selenium status in the elderly with mild cognitive impairment (MCI) and AD compared to healthy cognitively intact controls revealed a decrease in the Se content of erythrocytes that correlated with a decrease in cognitive function. However, plasma Se was not significantly different between healthy and MCI participants.¹²⁷ As plasma is a marker of recent exposure and erythrocytes tend to report longer-term Se status, these data suggest that chronic deficiency correlates with cognition decline. González-Domínguez *et al.*¹²⁸ reported lower Se levels in serum from AD patients in comparison to MCI elderly, although both groups

1 were not considered deficient. Even in “mild” AD patients
2 (defined as a Mini-Mental State Examination [MMSE] score
3 ≥ 20) Se levels in plasma were reported to be lower compared to
4 healthy age-matched elderly subjects.¹²⁹ These data are in
5 agreement with previous studies that suggest that a lack of Se
6 might increase the risk of dementia.^{127,130} Supplementation with
7 compounds containing Se has shown potential for stimulating
8 cognitive improvement, although this effect cannot be directly
9 attributed to Se alone.^{131,132} Cardoso *et al.*¹³³ showed that the
10 daily intake of one Brazil nut over 6 months, which provides
11 about 288 μg of Se, was associated with cognitive performance
12 improvement assessed by constructional praxis and animal
13 naming; both subtests of the CERAD battery. These data are
14 promising; though as Brazil nuts contain other antioxidant
15 components, more studies are required to elucidate the precise
16 effects of selenium intake on cognition via this dietary source.

17
18 Animal models and *in vitro* studies have also provided some
19 insight into the role of Se on AD pathogenesis. A Se deficient
20 diet was associated with increased A β plaque formation in the
21 brains of *Tg2576* transgenic mice.¹³⁴ Sodium selenate treatment
22 reduced tau phosphorylation in both cell culture and tau
23 transgenic animal models. In cells, the treatment antagonised
24 the protein phosphatase 2A inhibitor OA,¹³⁵ activated the
25 protein phosphatase 2A and dephosphorylated tau.¹³⁶ This
26 treatment prevented and reversed memory and motor deficits;
27 neurofibrillary tangles formation and neurodegeneration in
28 transgenic animals with tau dysfunction. Although SeO_3^{2-}
29 appears to be more toxic to cells, the treatment with this
30 inorganic Se compound to streptozotocin toxin model of mental
31 impairment improved tolerance to oxidative stress and
32 prevented cognitive decline.¹³⁷ In hippocampal and cortical
33 cells, it was associated with lower activity of α - and β -
34 secretases and corresponding reduced production of A β_{1-40} and
35 A β_{1-42} .¹³⁸ Treatment of triple-transgenic AD mice expressing
36 mutant forms of human APP_{SWE}, PS1_{M146V}, and tau_{P301L} with
37 selenomethionine resulted in reduced total and phosphorylated
38 tau, lower inflammatory biomarkers and improvement in
39 cognition.¹³⁹ Organoselenium compounds also display
40 interesting effects in improving AD pathology in murine
41 models: p,p'-methoxyl-diphenyl diselenide treatment to
42 streptozotocin-exposed mice slowed memory decline, reduced
43 oxidative stress and normalised acetyl-cholinesterase
44 activity.¹⁴⁰ Selenium-containing 8-hydroxyquinoline derivatives
45 bound to Cu^{2+} , Fe^{2+} and Zn^{2+} inhibited aggregation and
46 disassembled Cu^{2+} -induced aggregates of A β .¹⁴¹

47
48 Some studies have shown an association between certain
49 selenoproteins and pathological features of AD. In the post-
50 mortem brain SelP was found to colocalise with senile plaques
51 and neurofibrillary tangles,¹⁴² and was elevated in the choroid
52 plexus and cerebrospinal fluid of AD patients.¹⁴³ Considering
53 that oxidative stress is a hallmark of AD, these data could
54 suggest that SelP might act by protecting neurons against the
55 hallmark lesions damage of the disease *via* its prominent
56 antioxidant role, or by transporting Se to the synthesis of other
57 antioxidant selenoproteins. SelP encodes two His-rich regions
58 that act as high-affinity binding sites for transition metals,

potentially blocking metal-mediated A β_{1-42} aggregation and
subsequent ROS generation.^{144,145} This protein also appears to
modulate tau aggregation, with Du *et al.*¹⁴⁶ showing that the
His-rich domain of SelP inhibited aggregation and
disassembled formed aggregates of tau induced by the presence
of $\text{Cu}^+/\text{Cu}^{2+}$. These two His-rich regions associate with the
acidic tail of α -tubulin via a charge-charge interaction. This
suggests that SelP possibly participates in microtubule events,
which are associated with maintenance of cell polarity,
intracellular transport and cell division and migration, although
the exact role of SelP within microtubules has yet to be
elucidated.¹⁴⁷ Overexpression of SelM, as well as Se treatment
contributes to activation of ERK signalling, which induces a
decrease in tau phosphorylation, α -secretase and γ -secretase
activity and an increase in β -secretase.¹⁰⁰ Selenium treatment to
SelM-overexpressing mice resulted in significant ROS
inhibition, reduced mitochondrial damage, A β aggregation;¹⁰¹
and increased γ -secretase activity.¹⁰²

Parkinson's disease

Parkinson's disease (PD) is a neurodegenerative disorder
characterised by loss of pigmented dopaminergic neurons in the
substantia nigra (SN) pars compacta together with the presence
of intraneuronal proteinaceous inclusions of α -synuclein termed
Lewy bodies. Several mechanisms have been proposed that
induce toxicity inside the cytoplasm of neurons and also in the
extracellular space, including α -synuclein/neuromelanin
pathology,¹⁴⁸ ceruloplasmin dysfunction,¹⁴⁹ copper and
cuproprotein dyshomeostasis¹¹ and an iron/dopamine-mediated
oxidative stress cycle.¹⁵⁰ Dopaminergic neurons are particularly
vulnerable to oxidative damage, potentially due to their
propensity to accumulate iron with age,^{151,152} and thus oxidative
stress presents a central role on PD pathogenesis,¹⁵³⁻¹⁵⁵ and a
logical target of selenoprotein antioxidant activity.

An association between Se status and PD pathology is not
clear. Shahar *et al.*¹⁵⁶ measured plasma levels in 1,012 elderly
(≥ 65 years of age) Italian subjects and were able to associate
performance-based assessments of coordination with
circulating Se concentrations. Although there was no
measurable association between plasma Se and PD, older
subjects demonstrated a significant association between
circulating Se levels and decreased performance in neurological
coordination tests. In other smaller PD cohorts (< 100 PD
subjects) no difference in plasma Se levels compared to healthy
controls have been identified.^{157,158}

As noted by Shahar *et al.*,¹⁵⁶ causal influence between Se
and PD pathology is difficult to ascertain, particularly
considering motor function is often related to malnutrition, and
thereby circulating Se levels. Dietary Se may indeed be such a
confounding factor that any possible causal, or even indicative
relationship between circulating Se and PD (or any other
chronic illness, for that matter) impossible to elucidate.
Selenium in primary food sources is heavily dependent on the
Se content of soil, which is highly variable around the world;
soil Se levels in parts of Scandinavia, New Zealand and a wide
belt of land extending from Siberia through north-eastern China

are notably low, and are particularly replete in the US Midwest and Amazon regions of the Americas.¹⁵⁹ In contrast to Shahar's comments that PD pathology may cause malnutrition and Se deficiency, a large cohort of PD patients and controls from eastern China, (238 PD; 302 controls) reported that plasma Se (and iron) levels were significantly increased in patients with idiopathic PD.¹⁶⁰ Clearly, epidemiological studies with a higher statistical power are required to confirm or deny a possible association between circulating Se and PD. Additionally, it is advisable that several different biological matrices (e.g. erythrocytes, hair, nails etc.) should be used to evaluate Se levels. This would provide a more comprehensive overview of Se levels, and could potentially be used to identify times of exposure. Further, speciation of selenoproteins, as opposed to measuring total Se levels provides another avenue to identify specific Se functionality and its possible role in disease process.

Several studies have suggested a Se-deficient diet may contribute to dopaminergic cell vulnerability,^{161,162} particularly prior to exposure to parkinsonian neurotoxins.¹⁶³ Conversely, Se pretreatment somewhat mitigated 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced depletion of striatal dopamine in a dose dependent manner, though only a high dose of 3 mg kg⁻¹ of Se (as SeO₃²⁻) produced a statistically significant effect.¹⁶⁴ In a paraquat animal model of PD when treated with SeO₃²⁻ at a low dose (in drinking water at 11.8 μg L⁻¹) presented normalisation of motor activity when co-administered with the toxin.¹⁶⁵

SelP is localised within cell bodies, axons and presynaptic terminals of SN neurons, and in PD patients this selenoprotein co-localises within Lewy bodies.⁵⁰ Like SelP, GPx4 is also reduced in the SN of the PD brain, but is increased relative to cell density in surviving neurons.¹⁶⁶ Hauser *et al.*⁷⁹ showed that GPx4 is covalently modified by dopamine quinone, an oxidised and reactive metabolite of dopamine, and that this deleterious product of neurotransmitter breakdown may result in degradation and reduced activity of this antioxidant selenoprotein. More studies are necessary to elucidate the precise mechanisms by which GPx metabolism is altered during PD progression.

Other neurodegenerative diseases

A number of other progressive neurological disorders have been studied with regard to a possible association with Se metabolism and selenoprotein activity. Multiple sclerosis (MS) is one such chronic, progressive, inflammatory disease affecting the CNS with a possible link to Se. Although the aetiology of MS remains elusive, it is believed that disease onset and development depend on a genetic and environmental interaction. The pathogenesis of this disease is characterised by abnormal immune response against self-antigens, resulting in inflammation and massive neurodegeneration characterised by marked demyelination. Additionally, infiltration of pro-inflammatory and autoreactive leukocytes into the brain initiates a cascade of events including blood-brain-barrier disruption, microglial activation and excitotoxicity; all potentiating oligodendrocyte death and axon damage.¹⁶⁷⁻¹⁶⁹

There is growing evidence that the involvement of oxidative stress in MS pathogenesis as a crucial factor in initiating and perpetuating mechanisms involved in neurodegeneration. Glutathione peroxidase activity, as well other antioxidant enzymes, is decreased in MS patients when compared to the control subjects.^{170,171} In a proteomic study, among a set of 44 proteins that discriminated experimental autoimmune encephalomyelitis (EAE) rats from control groups, GPx3 was increased in EAE animals.¹⁷² Selenium levels tend to decrease in MS, as observed in a cohorts of Iranian¹⁷³ and Polish patients.¹⁷⁴ Complementing the suggestive data regarding a role of Se in MS pathogenesis, Chadanay *et al.*⁶⁶ observed that treatment with diphenyl diselenide to the EAE animal model reduced signs of the disease phenotype, partly by preventing the reduction in body weight that accompanies the disease progression and decreased the time and the severity of symptoms. In these animals, the treatment also improved immune system function by diminishing the extent of infiltrating macrophages and controlling mononuclear cell proliferation.

Amyotrophic lateral sclerosis (ALS) is another degenerative disease, in this case characterised by motor neuron loss in the brain and spinal cord, which leads to muscle atrophy, weakness and fasciculation.¹⁷⁵ Although the cause of ALS is unknown, several studies indicate a link between genetic and environmental factors, including excessive exposure to inorganic Se. This association was first suggested in 1977, when researchers observed a higher prevalence of ALS in a known seleniferous area.¹⁷⁶ In a northern Italian locale, where some inhabitants consumed drinking water with unusually high inorganic Se content, a higher risk of ALS was observed, with dose-response relation.^{177,178} ALS patients presented higher SeO₃²⁻ levels and lower organoselenium compounds, particularly SelP, in CSF compared with healthy control subjects.¹⁷⁹ This finding is something of a curiosity, as SeO₃²⁻ is not typically found at high levels in biological fluids; conversion to non-redox active forms occurs rapidly through a reaction with GSH, as SeO₃²⁻ is toxic,¹⁸⁰ though improved analytical technology has identified the presence of SeO₃²⁻ at low ng mL⁻¹ concentrations in urine¹⁸¹ and serum.¹⁸² It should be noted that the same method used in this study previously failed to identify SeO₃²⁻ in CSF samples.¹⁸³ Selenite is, however, toxic, though paradoxically it has been shown to offer some degree of neuroprotection against lead neurotoxicity in *Caenorhabditis elegans*,¹⁸⁴ though this may result from conversion to other selenoproteins rather than activity of the SeO₃²⁻ itself. Selenium in excess has neurotoxic effects to motor neuronal cells *in vitro*. Human neuroblastoma SKNB cells were more vulnerable to Se toxicity when compared with kidney and prostatic cells when treated with organic and inorganic Se, even at a low dose (0.1 μM).¹⁸⁵ Selenite induced degeneration of cholinergic and GABAergic neurons and impaired locomotor activity in *Caenorhabditis elegans*.¹⁸⁶⁻¹⁸⁸ The elevated SeO₃²⁻ in CSF of ALS patients may be due to excessive exposure to this compound from environmental

sources, alteration of the body's detoxification capacity or even genetic modifications.

Huntington's disease (HD) is an autosomal dominant neurodegenerative disease caused by increased repetition of CAG trinucleotide in exon 1 of the Huntington gene, resulting in polyQ-expanded sequence in N-terminal region of the Huntington protein. This genetic alteration makes the Huntington protein more prone to aggregation, which compromises proteasomal activity, impairs mitochondrial function, increases oxidative stress and contributes to neurodegeneration.¹⁸⁹⁻¹⁹¹ Clinically, HD is characterised by chorea, psychiatric disturbance and cognitive decline.^{192,193} Although Se levels in plasma were not different between HD patients and healthy controls, HD patients presented a decrease in regional brain Se content; specifically in the putamen, dorsolateral prefrontal cortex, primary visual cortex, cingulate gyrus and cerebellum.¹⁹⁴ Treatment of N171-82Q HD mice with SeO₃²⁻ significantly protected against decreased brain weight and reduced Huntington protein aggregation and oxidised glutathione.¹⁹⁴ In another HD mice model (3-nitropropionic acid exposure) treatment with (Z)-2,3-bis(4-chlorophenylselenanyl)prop-2-en-1-ol(bis selenide) improved locomotor activity and motor coordination, restored succinate dehydrogenase enzyme activity and protected against oxidative stress.¹⁹⁵ Although there are few studies investigating Se in HD, these findings suggest that Se may be a potential avenue of exploration with regard to prevention and treatment of HD, and more studies are therefore encouraged.

Conclusions

Selenium has an essential role in the brain; not only due to its direct antioxidant role, but also via its participation in maintaining redox balance, mitochondrial dynamics, regulation of Ca²⁺ channels and modulation of neurogenesis. The relevance of Se in the pathogenesis of different neurological diseases has been shown by several studies in humans, animal models and cell culture that all associate Se deficiency with a disturbance in brain metabolism. It is important to note that the strong hierarchical nature of Se regulation in the brain, and its ability to remain somewhat efficiently isolated from Se deficiency in the periphery, and thereby *in vitro* studies that control ambient Se levels in a manner not reflective of *in vivo* Se metabolism are suited only to investing a specific selenoprotein in isolation, as opposed to the interplay between the 30+ known selenoproteins *in toto*. Future research is required to fully elucidate the role of different selenoproteins in brain cells, and in turn this will provide potential uses of selenoproteins as biomarkers of normal or pathogenic biological processes in CNS. As recognition of the importance Se has in maintaining normal brain function, studies examining the benefits of Se supplementation are emerging, but literature is still unclear regarding the precise effects of different forms and doses of Se. Thus, additional experiments investigating the potential neurotoxicity of Se overload are also required to better assess the safe range of Se exposures.

Notes and references

^a *The Florey Institute of Neuroscience and Mental Health, The University of Melbourne, Parkville, Victoria, Australia*

^b *Faculty of Pharmaceutical Sciences, Department of Food and Experimental Nutrition, University of São Paulo, São Paulo, Brazil*

^c *Elemental Bio-imaging Facility, University of Technology Sydney, Broadway, New South Wales, Australia*

^d *Exposure Biology, Frank R. Lautenberg Environmental Health Sciences Laboratory, Department of Preventive Medicine, Icahn School of Medicine at Mount Sinai, New York, New York, United States of America.*

* Address correspondence to:

Bárbara Rita Cardoso, The Florey Institute of Neuroscience and Mental Health, The University of Melbourne, 30 Royal Parade, Parkville, Victoria, 3052, Australia. Ph +61 450 633 537; Email: barbara.rita@florey.edu.au

Dominic James Hare, Elemental Bio-imaging Facility, University of Technology Sydney, Thomas Street, Broadway, New South Wales, 2007, Australia. Ph +61 3 9035 9549; Email: dominic.hare@uts.edu.au

† Co-senior authors


1. K. Schwarz and C. M. Foltz, *J. Am. Chem. Soc.*, 1957, **79**, 3292-3292.
2. L. Flohe, W. Günzler and H. Schock, *FEBS Lett.*, 1973, **32**, 132-134.
3. G. V. Kryukov, S. Castellano, S. V. Novoselov, A. V. Lobanov, O. Zehtab, R. Guigó and V. N. Gladyshev, *Science*, 2003, **300**, 1439-1443.
4. M. Roman, P. Jitaru and C. Barbante, *Metallomics*, 2014, **6**, 25-54.
5. J. Gromadzińska, E. Reszka, K. Bruzelius, W. Wąsowicz and B. Åkesson, *Eur. J. Nutr.*, 2008, **47**, 29-50.
6. D. H. Holben and A. M. Smith, *J. Am. Diet. Assoc.*, 1999, **99**, 836-843.
7. M. Bélanger, I. Allaman and P. J. Magistretti, *Cell Metab.*, 2011, **14**, 724-738.
8. X. Wang and E. K. Michaelis, *Front. Aging Neurosci.*, 2010, **2**, 12.
9. A. Federico, E. Cardaioli, P. Da Pozzo, P. Formichi, G. N. Gallus and E. Radi, *J. Neurol. Sci.*, 2012, **322**, 254-262.
10. W. R. Markesbery, *Free Radic. Biol. Med.*, 1997, **23**, 134-147.
11. K. M. Davies, S. Bohic, A. Carmona, R. Ortega, V. Cottam, D. J. Hare, J. P. M. Finberg, S. Reyes, G. M. Halliday, J. F. B. Mercer and K. L. Double, *Neurobiol. Aging*, 2014, **35**, 858-866.
12. M. Navarro-Alarcon and C. Cabrera-Vique, *Sci. Total Environ.*, 2008, **400**, 115-141.
13. A. A. Turanov, A. V. Lobanov, D. L. Hatfield and V. N. Gladyshev, *Nucleic Acids Res.*, 2013, **41**, 6952-6959.
14. V. M. Labunskyy, D. L. Hatfield and V. N. Gladyshev, *Physiol. Rev.*, 2014, **94**, 739-777.
15. J. N. Gonzalez-Flores, N. Gupta, L. W. DeMong and P. R. Copeland, *J. Biol. Chem.*, 2012, **287**, 38936-38945.
16. A. L. Bifano, T. Atassi, T. Ferrara and D. M. Driscoll, *BMC Mol. Biol.*, 2013, **14**, 12.
17. M. E. Budiman, J. L. Bubenik, A. C. Miniard, L. M. Middleton, C. A. Gerber, A. Cash and D. M. Driscoll, *Mol. Cell*, 2009, **35**, 479-489.
18. A. C. Miniard, L. M. Middleton, M. E. Budiman, C. A. Gerber and D. M. Driscoll, *Nucleic Acids Res.*, 2010, **38**, 4807-4820.
19. M. J. Bröcker, J. M. L. Ho, G. M. Church, D. Söll and P. O'Donoghue, *Angew. Chem. Int. Ed.*, 2014, **53**, 319-323.
20. E. K. Wirth, M. Conrad, J. Winterer, C. Wozny, B. A. Carlson, S. Roth, D. Schmitz, G. W. Bornkamm, V. Coppola, L. Tessarollo, L. Schomburg, J. Köhrle, D. L. Hatfield and U. Schweizer, *FASEB J.*, 2010, **24**, 844-852.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
21. E. K. Wirth, B. S. Bharathi, D. Hatfield, M. Conrad, M. Briellemeier and U. Schweizer, *Biol. Trace Elem. Res.*, 2014, **158**, 203-210.
22. S. Seher, B. A. Carlson, A. C. Miniard, E. K. Wirth, Y. Mahdi, D. L. Hatfield, D. M. Driscoll and U. Schweizer, *Biochem. J.*, 2014, **462**, 67-75.
23. O. Agamy, B. Ben Zeev, D. Lev, B. Marcus, D. Fine, D. Su, G. Narkis, R. Ofir, C. Hoffmann, E. Leshinsky-Silver, H. Flusser, S. Sivan, D. Söll, T. Lerman-Sagie and O. S. Birk, *Am. J. Hum. Genet.*, 2010, **87**, 538-544.
24. X.-M. Xu, A. A. Turanov, B. A. Carlson, M.-H. Yoo, R. A. Everley, R. Nandakumar, I. Sorokina, S. P. Gygi, V. N. Gladyshev and D. L. Hatfield, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 21430-21434.
25. R. Tobe, S. Naranjo-Suarez, R. A. Everley, B. A. Carlson, A. A. Turanov, P. A. Tsuji, M.-H. Yoo, S. P. Gygi, V. N. Gladyshev and D. L. Hatfield, *J. Biol. Chem.*, 2013, **288**, 14709-14715.
26. L. Schomburg and U. Schweizer, *Biochim. Biophys. Acta, Gen. Subj.*, 2009, **1790**, 1453-1462.
27. R. Sunde, in *Selenium – Its molecular biology and role in human health*, eds D. L. Hatfield, M. Berry and V. N. Gladyshev, Springer, 2012, pp. 137-152.
28. M. T. Howard, B. A. Carlson, C. B. Anderson and D. L. Hatfield, *J. Biol. Chem.*, 2013, **288**, 19401-19413.
29. A. Seyedali and M. J. Berry, *RNA*, 2014, **20**, 1248-1256.
30. J. Y. Kim, B. A. Carlson, X.-M. Xu, Y. Zeng, S. Chen, V. N. Gladyshev, B. J. Lee and D. L. Hatfield, *Biochem. Biophys. Res. Commun.*, 2011, **409**, 814-819.
31. B. A. Zachara, H. Pawluk, E. Bloch-Boguslawska, K. M. Sliwka, J. Korenkiewicz, Z. Skok and K. Ryć, *Arch. Environ. Health*, 2001, **56**, 461-466.
32. D. Behne and W. Wolters, *J. Nutr.*, 1983, **113**, 456-461.
33. M. Haratake, K. Koga, M. Inoue, T. Fuchigami and M. Nakayama, *Metallomics*, 2011, **3**, 1019-1026.
34. M. Kühbacher, J. Bartel, B. Hoppe, D. Alber, G. Bukalis, A. U. Bräuer, D. Behne and A. Kyriakopoulos, *J. Neurochem.*, 2009, **110**, 133-142.
35. M. Kühbacher, J. Bartel, D. Alber, G. Bukalis and H. Kühn, 2014, http://www.laborjournal.de/editorials/ed424/Ms421_v425_web.pdf (Accessed 11 May 2015).
36. P. Ramos, A. Santos, N. R. Pinto, R. Mendes, T. Magalhães and A. Almeida, *Biol. Trace Elem. Res.*, 2014, **163**, 89-96.
37. L. R. Ferguson, N. Karunasinghe, S. Zhu and A. H. Wang, *Mutat. Res.*, 2012, **733**, 100-110.
38. S. P. Shetty, R. Shah and P. R. Copeland, *J. Biol. Chem.*, 2014, **289**, 25317-25326.
39. G. Ballihaut, L. E. Kilpatrick, E. L. Kilpatrick and W. C. Davis, *Metallomics*, 2012, **4**, 533-538.
40. R. F. Burk, K. E. Hill, A. K. Motley, V. P. Winfrey, S. Kurokawa, S. L. Mitchell and W. Zhang, *FASEB J.*, 2014, **28**, 3579-3588.
41. A. E. Clatworthy, W. Stockinger, R. H. Christie, W. J. Schneider, J. Nimpf, B. T. Hyman and G. W. Rebeck, *Neuroscience*, 1999, **90**, 903-911.
42. S. Kurokawa, K. E. Hill, W. H. McDonald and R. F. Burk, *J. Biol. Chem.*, 2012, **287**, 28717-28726.
43. G. E. Olson, V. P. Winfrey, S. K. NagDas, K. E. Hill and R. F. Burk, *J. Biol. Chem.*, 2007, **282**, 12290-12297.
44. M. W. Pitts, A. V. Raman, A. C. Hashimoto, C. Todorovic, R. A. Nichols and M. J. Berry, *Neuroscience*, 2012, **208**, 58-68.
45. I. Masiulis, T. A. Quill, R. F. Burk and J. Herz, *Biol. Chem.*, 2009, **390**, 67-73.
46. S. W. Caito, D. Milatovic, K. E. Hill, M. Aschner, R. F. Burk and W. M. Valentine, *Brain Res.*, 2011, **1398**, 1-12.
47. R. F. Burk, G. E. Olson, K. E. Hill, V. P. Winfrey, A. K. Motley and S. Kurokawa, *FASEB J.*, 2013, **27**, 3249-3256.
48. H. Steinbrenner, L. Alili, E. Bilgic, H. Sies and P. Brenneisen, *Free Radic. Biol. Med.*, 2006, **40**, 1513-1523.
49. M. Scharpf, U. Schweizer, T. Arzberger, W. Roggendorf, L. Schomburg and J. Köhrle, *J. Neural. Transm.*, 2007, **114**, 877-884.
50. F. P. Bellinger, A. V. Raman, R. H. Rueli, M. T. Bellinger, A. S. Dewing, L. A. Seale, M. A. Andres, J. H. Uyehara-Lock, L. R. White, G. W. Ross and M. J. Berry, *J. Parkinsons Dis.*, 2012, **2**, 115-126.
51. A. V. Raman, M. W. Pitts, A. Seyedali, A. C. Hashimoto, L. A. Seale, F. P. Bellinger and M. J. Berry, *Genes Brain Behav.*, 2012, **11**, 601-613.
52. C. N. Byrns, M. W. Pitts, C. A. Gilman, A. C. Hashimoto and M. J. Berry, *J. Biol. Chem.*, 2014, **289**, 9662-9674.
53. K.-M. Fang, F.-C. Cheng, Y.-L. Huang, S.-Y. Chung, Z.-Y. Jian and M.-C. Lin, *Biol. Trace Elem. Res.*, 2013, **152**, 66-74.
54. P. Bitencourt, L. Belle, G. Bonfanti, L. Cargnelutti, K. d. Bona, P. Silva, F. Abdalla, R. Zanette, R. Guerra, C. Funchal and M. Moretto, *Hum. Exp. Toxicol.*, 2013, **32**, 942-949.
55. M. Nazıroğlu, Ö. Çelik, A. C. Uğuz and A. Bütün, *Biol. Trace Elem. Res.*, 2015, **164**, 72-79.
56. I. Ben Amara, H. Fetoui, F. Guermazi and N. Zeghal, *Int. J. Dev. Neurosci.*, 2009, **27**, 719-726.
57. A. S. Abedelhafiez and A. Hassan, *Acta Physiol. Hung.*, 2013, **100**, 197-210.
58. O. L. Adebayo, G. A. Adenuga and R. Sandhir, *Nutr. Neurosci.*, 2014, **17**, 268-278.
59. A. C. Uğuz and M. Nazıroğlu, *Neurochem. Res.*, 2012, **37**, 1631-1638.
60. M. Nazıroğlu, N. Şenol, V. Ghazizadeh and V. Yürüker, *Cell. Mol. Neurobiol.*, 2014, **34**, 895-903.
61. N. Mendelev, S. L. Mehta, S. Witherspoon, Q. He, J. Z. Sexton and P. A. Li, *Mitochondrion*, 2011, **11**, 76-82.
62. S. L. Mehta, N. Mendelev, S. Kumari and P. Andy Li, *Int. J. Biochem. Cell Biol.*, 2013, **45**, 604-611.
63. S. Kumari, S. L. Mehta and P. A. Li, *PLoS One*, 2012, **7**, e39382.
64. A. S. de Freitas and J. B. T. Rocha, *Neurosci. Lett.*, 2011, **503**, 1-5.
65. A. P. Ardaís, G. G. Viola, M. S. Costa, F. Nunes, G. A. Behr, F. Klamt, J. C. F. Moreira, D. O. Souza, J. B. T. Rocha and L. O. Porciúncula, *Toxicol. Sci.*, 2010, **113**, 434-443.
66. N. L. Chanaday, A. F. de Bem and G. A. Roth, *Neurochem. Int.*, 2011, **59**, 1155-1162.
67. G. Roseni Mundstock Dias, R. Medeiros Golombieski, R. de Lima Portella, G. Pires do Amaral, F. Antunes Soares, J. B. Teixeira da Rocha, C. Wayne Nogueira and N. Vargas Barbosa, *Neuroendocrinology*, 2014, **100**, 45-59.
68. C. A. Brüning, B. M. Gai, S. M. Soares, F. Martini and C. W. Nogueira, *Pharmacol. Biochem. Behav.*, 2014, **125**, 15-20.
69. P. M. Chagas, S. G. Rosa, M. H. M. Sari, C. E. S. Oliveira, R. F. S. Canto, S. C. A. da Luz, A. L. Braga and C. W. Nogueira, *Pharmacol. Biochem. Behav.*, 2014, **118**, 87-95.
70. I. Musik, M. Kielczykowska and J. Kocot, *BioMetals*, 2013, **26**, 763-771.
71. S. M. Barayuga, X. Pang, M. A. Andres, J. Panee and F. P. Bellinger, *Neurotoxicology*, 2013, **37**, 240-246.
72. E. Song, C. Su, J. Fu, X. Xia, S. Yang, C. Xiao, B. Lu, H. Chen, Z. Sun, S. Wu and Y. Song, *Life Sci.*, 2014, **109**, 37-43.
73. T. Garcia, J. L. Esparza, M. R. Nogués, M. Romeu, J. L. Domingo and M. Gómez, *Hippocampus*, 2010, **20**, 218-225.
74. S. Zhang, C. Rocourt and W.-H. Cheng, *Mech. Ageing Dev.*, 2010, **131**, 253-260.
75. P. A. Mitozo, L. F. de Souza, G. Loch-Neckel, S. Flesch, A. F. Maris, C. P. Figueiredo, A. R. S. dos Santos, M. Farina and A. L. Daffe, *Free Radic. Biol. Med.*, 2011, **51**, 69-77.
76. M.-H. Yoo, X. Gu, X.-M. Xu, J. Y. Kim, B. A. Carlson, A. D. Patterson, H. Cai, V. N. Gladyshev and D. L. Hatfield, *Antiox. Redox. Signal.*, 2010, **12**, 819-827.
77. M. W. Pitts, C. N. Byrns, A. N. Ogawa-Wong, P. Kremer and M. J. Berry, *Biol. Trace Elem. Res.*, 2014, **161**, 231-245.
78. N. E. Savaskan, A. Borchert, A. U. Bräuer and H. Kuhn, *Free Radic. Biol. Med.*, 2007, **43**, 191-201.
79. D. N. Hauser, A. A. Dukes, A. D. Mortimer and T. G. Hastings, *Free Radic. Biol. Med.*, 2013, **65**, 419-427.
80. J. R. Godoy, M. Funke, W. Ackermann, P. Haunhorst, S. Oesteritz, F. Capani, H. P. Elsasser and C. H. Lillig, *Biochim. Biophys. Acta*, 2011, **1810**, 2-92.
81. D. Silva-Adaya, M. x. Gonsebatt, a. E and J. Guevara, *Oxid. Med. Cell. Longev.*, 2014, **2014**, 1-13.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
82. A. P. Kudin, B. Augustynek, A. K. Lehmann, R. Kovács and W. S. Kunz, *Biochim. Biophys. Acta, Bioenergetics*, 2012, **1817**, 1901-1906.
83. M. Cohen-Kutner, L. Khomsky, M. Trus, H. Ben-Yehuda, J. M. Lenhard, Y. Liang, T. Martin and D. Atlas, *Redox Biology*, 2014, **2**, 447-456.
84. P. Lopert, B. J. Day and M. Patel, *PLoS One*, 2012, **7**, e50683.
85. J. H. Ahn, J. H. Choi, J. M. Song, C. H. Lee, K.-Y. Yoo, I. K. Hwang, J. S. Kim, H.-C. Shin and M. H. Won, *Exp. Gerontol.*, 2011, **46**, 946-952.
86. H. Esen, F. Erdi, B. Kaya, B. Feyzioglu, F. Keskin and L. S. Demir, *J. Neurooncol.*, 2014, **121**, 451-458.
87. L. V. Papp, J. Lu, A. Holmgren and K. K. Khanna, *Antioxid. Redox Signaling*, 2007, **9**, 775-806.
88. R. F. Burk and K. E. Hill, *Annu. Rev. Nutr.*, 2005, **25**, 215-235.
89. Y. Zhang and V. N. Gladyshev, *PLoS Genet.*, 2008, **4**, e1000095.
90. A. Dikiy, S. V. Novoselov, D. E. Fomenko, A. Sengupta, B. A. Carlson, R. L. Cerny, K. Ginalski, N. V. Grishin, D. L. Hatfield and V. N. Gladyshev, *Biochemistry (Mosc.)*, 2007, **46**, 6871-6882.
91. F. Musiani, S. Ciurli and A. Dikiy, *J. Proteome Res.*, 2011, **10**, 968-976.
92. M. E. Moustafa and H. A. Antar, *Biochem. Genet.*, 2012, **50**, 736-747.
93. F. Li, P. B. Lutz, Y. Pepelyayeva, E. S. J. Arnér, C. A. Bayse and S. Rozovsky, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 6976-6981.
94. S.-J. Han, B. C. Lee, S. H. Yim, V. N. Gladyshev and S.-R. Lee, *PLoS One*, 2014, **9**, e95518.
95. S. Guariniello, G. Colonna, R. Raucchi, M. Costantini, G. Di Bernardo, F. Bergantino, G. Castello and S. Costantini, *Biochim. Biophys. Acta, Proteins Proteomics*, 2014, **1844**, 447-456.
96. A. D. Ferguson, V. M. Labunskyy, D. E. Fomenko, D. Araç, Y. Chelliah, C. A. Amezcua, J. Rizo, V. N. Gladyshev and J. Deisenhofer, *J. Biol. Chem.*, 2006, **281**, 3536-3543.
97. J. L. Martin, *Structure*, 1995, **3**, 245-250.
98. M. W. Pitts, M. A. Reeves, A. C. Hashimoto, A. Ogawa, P. Kremer, L. A. Seale and M. J. Berry, *J. Biol. Chem.*, 2013, **288**, 26121-26134.
99. M. A. Reeves, F. P. Bellinger and M. J. Berry, *Antiox. Redox. Signal.*, 2010, **12**, 809-818.
100. S. Y. Yim, K. R. Chae, S. B. Shim, J. T. Hong, J. Y. park, C. Y. Lee, H. J. Son, Y. Y. sheen and D. Y. Hwang, *Int. J. Mol. Med.*, 2009, **24**, 91-96.
101. P. Chen, R.-R. Wang, X.-J. Ma, Q. Liu and J.-Z. Ni, *Int. J. Mol. Sci.*, 2013, **14**, 4385-4399.
102. Y. Kim, J. S. Goo, I. Y. Kim, J. E. Kim, M. H. Kwak, J. Go, S. Shim, J. T. Hong, D. Y. Hwang and J. K. Seong, *Int. J. Mol. Med.*, 2014, **34**, 1688-1698.
103. X. Qiao, J. Tian, P. Chen, C. Wang, J. Ni and Q. Liu, *Int. J. Mol. Sci.*, 2013, **14**, 22233-22245.
104. S. C. Starosom, I. D. Mascanfroni, J. Imitola, L. Cao, K. Raddassi, S. F. Hernandez, R. Bassil, D. O. Croci, J. P. Cerliani, D. Delacour, Y. Wang, W. Elyaman, S. J. Khoury and G. A. Rabinovich, *Immunity*, 2012, **37**, 249-263.
105. K. Kajitani, Y. Kobayakawa, H. Nomaru, T. Kadoya, H. Horie and Y. Nakabeppu, *Neuroreport*, 2014, **25**, 171-176.
106. J.-L. Li, H.-F. Ruan, H.-X. Li, S. Li, S.-W. Xu and Z.-X. Tang, *Mol. Biol. Rep.*, 2010, **38**, 4015-4022.
107. B.-R. Ou, M.-J. Jiang, C.-H. Lin, Y.-C. Liang, K.-J. Lee and J.-Y. Yeh, *BioMetals*, 2011, **24**, 323-333.
108. J.-L. Li, H.-X. Li, S. Li, X.-J. Gao, S.-W. Xu and Z.-X. Tang, *BioMetals*, 2012, **25**, 459-468.
109. I. J. Cajigas, G. Tushev, T. J. Will, S. t. Dieck, N. Fuerst and E. M. Schuman, *Neuron*, 2012, **74**, 453-466.
110. A. V. Raman, M. W. Pitts, A. Seyedali, A. C. Hashimoto, F. P. Bellinger and M. J. Berry, *Brain Behav.*, 2013, **3**, 562-574.
111. F. Rage, N. Boulisfane, K. Rihaan, H. Neel, T. Gostan, E. Bertrand, R. Bordonne and J. Soret, *RNA*, 2013, **19**, 1755-1766.
112. W. Xiao-Long, Y. Chuan-Ping, X. Kai and Q. Ou-Jv, *Biochemistry (Mosc.)*, 2010, **75**, 201-207.
113. Y. Zhang, Y. Zhou, U. Schweizer, N. E. Savaskan, D. Hua, J. Kipnis, D. L. Hatfield and V. N. Gladyshev, *J. Biol. Chem.*, 2008, **283**, 2427-2438.
114. N. Fradejas, M. Del Carmen Serrano Pérez, P. Tranque and S. Calvo, *Glia*, 2011, **59**, 959-972.
115. L. X. Liu, X. Y. Zhou, C. S. Li, L. Q. Liu, S. Y. Huang and S. N. Zhou, *Neurol. Sci.*, 2013, **34**, 1671-1678.
116. A. A. Turanov, V. A. Shchedrina, R. A. Everley, A. V. Lobanov, S. H. Yim, S. M. Marino, S. P. Gygi, D. L. Hatfield and V. N. Gladyshev, *Biochem. J.*, 2014, **462**, 555-565.
117. X. A. Du, H. M. Wang, X. X. Dai, Y. Kou, R. P. Wu, Q. Chen, J. L. Cao, X. Y. Mo and Y. M. Xiong, *Osteoarthr. Cartilage*, 2015, **23**, 210-216.
118. S. V. Novoselov, G. V. Kryukov, X.-M. Xu, B. A. Carlson, D. L. Hatfield and V. N. Gladyshev, *J. Biol. Chem.*, 2007, **282**, 11960-11968.
119. R. T. Y. Wu, L. Cao, B. P. C. Chen and W.-H. Cheng, *J. Biol. Chem.*, 2014, **289**, jbc.M114.611970-634388.
120. P. H. Reddy, *Brain Res.*, 2011, **1415**, 136-148.
121. P. H. Reddy and P. H. Reddy, *Biochim. Biophys. Acta, Mol. Basis Dis.*, 2011, **1812**, 1359-1370.
122. Y. Yoshiyama, V. M. Y. Lee and J. Q. Trojanowski, *J. Neurol. Neurosurg. Psychiatry*, 2013, **84**, 784-795.
123. R. H. Swerdlow, J. M. Burns and S. M. Khan, *Biochim. Biophys. Acta, Mol. Basis Dis.*, 2014, **1842**, 1219-1231.
124. A. Picco, M. C. Polidori, M. Ferrara, R. Cecchetti, D. Arnaldi, M. Baglioni, S. Morbelli, P. Bastiani, I. Bossert, G. Fiorucci, A. Brugnolo, M. E. Dottorini, F. Nobili and P. Mecocci, *Eur. J. Nucl. Med. Mol. Imaging*, 2013, **41**, 764-775.
125. S. Haider, S. Saleem, T. Perveen, S. Tabassum, Z. Batool, S. Sadr, L. Liaquat and S. Madiha, *Age*, 2014, **36**, 1291-1302.
126. B. R. Cardoso, V. Silva Bandeira, W. Jacob-Filho and S. M. Franciscato Cozzolino, *J. Trace Elem. Med. Biol.*, 2014, **28**, 422-426.
127. B. R. Cardoso, T. P. Ong, W. Jacob-Filho, O. Jaluul, M. I. d. a. Á. Freitas and S. M. F. Cozzolino, *Br. J. Nutr.*, 2010, **103**, 803-806.
128. R. Gonzalez-Dominguez, T. Garcia-Barrera and J. L. Gomez-Ariza, *BioMetals*, 2014, **27**, 539-549.
129. M. G. M. Olde Rikkert, F. R. Verhey, J. W. C. Sijben, F. H. Bouwman, P. L. J. Dautzenberg, M. Lansink, W. M. W. Sipers, D. Z. B. van Asselt, A. M. J. van Hees, M. Stevens, B. Vellas and P. Scheltens, *J. Alzheimers Dis.*, 2014, **41**, 261-271.
130. H. Vural, H. Demirin, Y. Kara, I. Eren and N. Delibas, *J. Trace Elem. Med. Biol.*, 2010, **24**, 169-173.
131. P. Scheltens, P. J. G. H. Kamphuis, F. R. J. Verhey, M. G. M. Olde Rikkert, R. J. Wurtman, D. Wilkinson, J. W. R. Twisk and A. Kurz, *Alzheimers Dement.*, 2010, **6**, 1-10.e11.
132. E. Kesse-Guyot, L. Fezeu, C. Jeandel, M. Ferry, V. Andreeva, H. Amieva, S. Hercberg and P. Galan, *Am. J. Clin. Nutr.*, 2011, **94**, 892-899.
133. B. R. Cardoso, D. Apolinário, V. da Silva Bandeira, A. L. Busse, R. M. Magaldi, W. Jacob-Filho and S. M. F. Cozzolino, *Eur. J. Nutr.*, 2015, DOI: 10.1007/s00394-014-0829-2, 1-10.
134. M. Haratake, S. Yoshida, M. Mandai, T. Fuchigami and M. Nakayama, *Metallomics*, 2013, **5**, 479-483.
135. J. van Eersel, Y. D. Ke, X. Liu, F. Delerue, J. J. Kril, J. Götz and L. M. Ittner, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 13888-13893.
136. N. M. Corcoran, D. Martin, B. Hutter-Paier, M. Windisch, T. Nguyen, L. Nheu, L. E. Sundstrom, A. J. Costello and C. M. Hovens, *J. Clin. Neurosci.*, 2010, **17**, 1025-1033.
137. T. Ishrat, K. Parveen, M. M. Khan, G. Khuwaja, M. B. Khan, S. Yousof, A. Ahmad, P. Shrivastav and F. Islam, *Brain Res.*, 2009, **1281**, 117-127.
138. A.-R. Gwon, J.-S. Park, J.-H. Park, S.-H. Baik, H.-Y. Jeong, D.-H. Hyun, K. W. Park and D.-G. Jo, *Neurosci. Lett.*, 2010, **469**, 391-395.
139. G. Song, Z. Zhang, L. Wen, C. Chen, Q. Shi, Y. Zhang, J. Ni and Q. Liu, *J. Alzheimers Dis.*, 2014, **41**, 85-99.
140. S. Pinton, C. A. Brünning, C. E. Sartori Oliveira, M. Prigol and C. W. Nogueira, *J. Nutr. Biochem.*, 2013, **24**, 311-317.
141. Z. Wang, Y. Wang, W. Li, F. Mao, Y. Sun, L. Huang and X. Li, *ACS Chem. Neurosci.*, 2014, **5**, 952-962.
142. F. P. Bellinger, Q. P. He, M. T. Bellinger and Y. Lin, *J. Alzheimers Dis.*, 2008, **15**, 465-472.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
143. R. H. L. H. Rueli, A. C. Parubrub, A. S. T. Dewing, A. C. Hashimoto, M. T. Bellinger, E. J. Weeber, J. H. Uyehara-Lock, L. R. White, M. J. Berry and F. P. Bellinger, *J. Alzheimers Dis.*, 2015, **44**, 379-383.
144. X. Du, H. Li, Z. Wang, S. Qiu, Q. Liu and J. Ni, *Metallomics*, 2013, **5**, 861-870.
145. X. Du, Z. Wang, Y. Zheng, H. Li, J. Ni and Q. Liu, *Inorg. Chem.*, 2014, **53**, 1672-1678.
146. X. Du, Y. Zheng, Z. Wang, Y. Chen, R. Zhou, G. Song, J. Ni and Q. Liu, *Inorg. Chem.*, 2014, **53**, 11221-11230.
147. X. Du, S. Qiu, Z. Wang, R. Wang, C. Wang, J. Tian and Q. Liu, *Int. J. Mol. Sci.*, 2014, **15**, 10199-10214.
148. G. M. Halliday, A. Ophof, M. Broe, P. H. Jensen, E. Kettle, H. Fedorow, M. I. Cartwright, F. M. Griffiths, C. E. Shepherd and K. L. Double, *Brain*, 2005, **128**, 2654-2664.
149. S. Ayton, P. Lei, J. A. Duce, B. X. W. Wong, A. Sedjahtera, P. A. Adlard, A. I. Bush and D. I. Finkelstein, *Ann. Neurol.*, 2013, **73**, 554-559.
150. D. J. Hare, P. Lei, S. Ayton, B. R. Roberts, R. Grimm, J. L. George, D. P. Bishop, A. D. Beavis, S. J. Donovan, G. McColl, I. Volitakis, C. L. Masters, P. A. Adlard, R. A. Cherny, A. I. Bush, D. I. Finkelstein and P. A. Doble, *Chem. Sci.*, 2014, **5**, 2160.
151. L. Zecca, A. Stroppolo, A. Gatti, D. Tampellini, M. Toscani, M. Gallorini, G. Giaveri, P. Arosio, P. Santambrogio and R. G. Fariello, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 9843.
152. D. Hare, S. Ayton, A. Bush and P. Lei, *Front. Aging Neurosci.*, 2013, **5**, 34.
153. C. Venkateshappa, G. Harish, R. B. Mythri, A. Mahadevan, M. M. S. Bharath and S. K. Shankar, *Neurochem. Res.*, 2011, **37**, 358-369.
154. G. S. Gaki and A. G. Papavassiliou, *Neuromolecular Med.*, 2014, **16**, 217-230.
155. A. Recasens and B. Dehay, *Front. Neuroanat.*, 2014, **8**.
156. A. Shahar, K. V. Patel, R. D. Semba, S. Bandinelli, D. R. Shahar, L. Ferrucci and J. M. Guralnik, *Mov. Disord.*, 2010, **25**, 1909-1915.
157. S. Younes-Mhenni, M. Aissi, N. Mokni, A. Boughammoura-Bouatay, S. Chebel, M. Frih-Ayed, A. Kerkeni, M. Bost, G. Chazot, M. T. Sfar and M. H. Sfar, *Tunis. Med.*, 2013, **91**, 402-405.
158. L. K. Mischley, J. Allen and R. Bradley, *J. Neurol. Sci.*, 2012, **318**, 72-75.
159. G. F. Combs, *Br. J. Nutr.*, 2001, **85**, 517-547.
160. H.-W. Zhao, J. Lin, X.-B. Wang, X. Cheng, J.-Y. Wang, B.-L. Hu, Y. Zhang, X. Zhang and J.-H. Zhu, *PLoS One*, 2013, **8**, e83060.
161. M. L. Vizuete, V. Steffen, A. Machado and J. Cano, *Eur. J. Pharmacol.*, 1994, **270**, 183-187.
162. H.-C. Kim, W.-K. Jhoo, E.-J. Shin and G. Bing, *Brain Res.*, 2000, **862**, 247-252.
163. D. J. Hare, P. A. Adlard, P. A. Doble and D. I. Finkelstein, *Metallomics*, 2013, **5**, 91-109.
164. H. A. Khan, *Neurochem. Int.*, 2010, **57**, 489-491.
165. J. H. Ellwanger, P. Molz, D. R. Dallemole, A. Pereira Dos Santos, T. E. Muller, L. Cappelletti, M. Goncalves da Silva, S. I. Rech Franke, D. Pra and J. A. Pegas Henriques, *Nutrition*, 2015, **31**, 359-365.
166. F. P. Bellinger, M. T. Bellinger, L. A. Seale, A. S. Takemoto, A. V. Raman, T. Miki, A. B. Manning-Boğ, M. J. Berry, L. R. White and G. W. Ross, *Mol. Neurodegener.*, 2011, **6**, 8.
167. R. Gandhi, A. Laroni and H. L. Weiner, *J. Neuroimmunol.*, 2010, **221**, 7-14.
168. B. Broux, P. Stinissen and N. Hellings, *Crit. Rev. Immunol.*, 2013, **33**, 283-306.
169. G. G. Ortiz, F. P. Pacheco-Moisés, M. Á. Macías-Islas, L. J. Flores-Alvarado, M. A. Mireles-Ramírez, E. D. González-Renovato, V. E. Hernández-Navarro, A. L. Sánchez-López and M. A. Alatorre-Jiménez, *Arch. Med. Res.*, 2014, **45**, 687-697.
170. S. Ljubisavljevic, I. Stojanovic, T. Cvetkovic, S. Vojinovic, D. Stojanov, D. Stojanovic, V. Bojanic, D. Stokanovic and D. Pavlovic, *NeuroImmunoModulation*, 2014, **21**, 13-20.
171. I. Tasset, E. Agüera, F. Sánchez-López, M. Feijóo, A. I. Giraldo, A. H. Cruz, F. Gascón and I. Túnez, *Clin. Biochem.*, 2012, **45**, 440-444.
172. T. Rosenling, M. P. Stoop, A. Attali, H. van Aken, E. Suidgeest, C. Christin, C. Stingl, F. Suits, P. Horvatovich, R. Q. Hintzen, T. Tuinstra, R. Bischoff and T. M. Luider, *J. Proteome Res.*, 2012, **11**, 2048-2060.
173. M. Mehrpour, A. Kyani, M. Tafazzoli, F. Fathi and M. T. Joghataie, *Magn. Reson. Chem.*, 2013, **51**, 102-109.
174. K. Socha, J. Kochanowicz, E. Karpińska, J. Soroczyńska, M. Jakoniuk, Z. Mariak and M. H. Borawska, *Nutr. J.*, 2014, **13**, 62.
175. L. P. Rowland and N. A. Shneider, *N. Engl. J. Med.*, 2001.
176. A. W. Kilness and F. H. Hichberg, *JAMA*, 1977, **237**, 2843-2844.
177. M. Vinceti, F. Bonvicini, K. J. Rothman, L. Vescovi and F. Wang, *Environ. Health*, 2010, **9**, 77.
178. M. Vinceti, D. Guidetti, M. Pinotti, S. Rovesti, M. Merlin, L. Vescovi, M. Bergomi and G. Vivoli, *Epidemiology*, 1996, **7**, 529-532.
179. M. Vinceti, N. Solovyev, J. Mandrioli, C. M. Crespi, F. Bonvicini, E. Arcolin, E. Georgouloupoulou and B. Michalke, *Neurotoxicology*, 2013, **38**, 25-32.
180. C. M. Weekley and H. H. Harris, *Chem. Soc. Rev.*, 2013, **42**, 8870-8894.
181. S. Kokarnig, A. Tsirigotaki, T. Wiesenhofer, V. Lackner, K. A. Francesconi, S. A. Pergantis and D. Kuehnelt, *J. Trace Elem. Med. Biol.*, 2015, **29**, 83-90.
182. Ö. Palacios, J. R. Encinar, D. Schaumlöffel and R. Lobinski, *Anal. Bioanal. Chem.*, 2006, **384**, 1276-1283.
183. B. Michalke and A. Berthele, *J. Anal. At. Spectrom.*, 2011.
184. W.-H. Li, Y.-C. Shi, I.-L. Tseng and V. H.-C. Liao, *PLoS One*, 2013, **8**, e62387.
185. T. Maraldi, M. Riccio, L. Zamboni, M. Vinceti, A. De Pol and G. Hakim, *Neurotoxicology*, 2011, **32**, 180-187.
186. K. L. Morgan, A. O. Estevez, C. L. Mueller, K. L. Morgan, N. J. Szewczyk, A. Miranda-Vizuete, N. J. Szewczyk and M. Estevez, *Toxicol. Sci.*, 2010, **118**, 530-543.
187. A. O. Estevez, K. L. Morgan, N. J. Szewczyk, D. Gems and M. Estevez, *Neurotoxicology*, 2014, **41**, 28-43.
188. A. O. Estevez, C. L. Mueller, K. L. Morgan, N. J. Szewczyk, L. Teece, A. Miranda-Vizuete and M. Estevez, *Neurotoxicology*, 2012, **33**, 1021-1032.
189. J. P. Vonsattel, R. H. Myers, T. J. Stevens, R. J. Ferrante, E. D. Bird and E. P. Richardson, Jr., *J. Neuropathol. Exp. Neurol.*, 1985, **44**, 559-577.
190. A. Bush, W. Pettingell, G. Multhaup, M. d Paradis, J. Vonsattel, J. Gusella, K. Beyreuther, C. Masters and R. Tanzi, *Science*, 1994, **265**, 1464-1467.
191. T.-C. Ju, H.-M. Chen, Y.-C. Chen, C.-P. Chang, C. Chang and Y. Chern, *Biochim. Biophys. Acta, Mol. Basis Dis.*, 2014, **1842**, 1668-1680.
192. A. D. Ha and V. S. C. Fung, *Curr. Opin. Neurol.*, 2012, **25**, 491-498.
193. J. Paulsen, R. Ready, J. Hamilton, M. Mega and J. Cummings, *J. Neurol. Neurosurg. Psychiatry*, 2001, **71**, 310-314.
194. Z. Lu, E. Marks, J. Chen, J. Moline, L. Barrows, M. Raisbeck, I. Volitakis, R. A. Cherny, V. Chopra, A. I. Bush, S. Hersch and J. H. Fox, *Neurobiol. Dis.*, 2014, **71C**, 34-42.
195. C. F. Bortolatto, C. R. Jesse, E. A. Wilhelm, P. M. Chagas and C. W. Nogueira, *Neurotox. Res.*, 2012, **23**, 214-224.
196. M. Birringer, S. Pilawa and L. Flohé, *Nat. Prod. Rep.*, 2002, **19**, 693-718.

Table of Contents



A review of selenium’s essential role in normal brain function and its potential involvement in neurodegenerative diseases.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60