



Cite this: *Dalton Trans.*, 2015, **44**, 3540

Received 1st September 2014,  
Accepted 27th November 2014

DOI: 10.1039/c4dt02663b

www.rsc.org/dalton

## Unprecedented dinuclear silver(I)-mediated base pair involving the DNA lesion 1,*N*<sup>6</sup>-ethenoadenine†

Soham Mandal,<sup>a,b</sup> Alexander Hepp<sup>a</sup> and Jens Müller<sup>\*a,b</sup>


The DNA lesion 1,*N*<sup>6</sup>-ethenoadenine ( $\epsilon$ A) has been investigated with respect to its metal-binding properties. A synthetic DNA duplex comprising an  $\epsilon$ A :  $\epsilon$ A mispair readily forms doubly silver(I)-mediated base pairs  $\epsilon$ A–Ag(I)<sub>2</sub>– $\epsilon$ A, representing the first example for a dinuclear metal-mediated homo base pair of a purine derivative. It also constitutes the first example for a Hoogsteen-type metal-mediated homo base pair within a B-DNA duplex.

The incorporation of transition metal ions into nucleic acids by means of metal-mediated base pairs represents an elegant and efficient method for the site-specific introduction of metal-based functionality into the self-assembling biomacromolecules.<sup>1</sup> In metal-mediated base pairs, the hydrogen bonds between complementary nucleobases are formally replaced by coordinative bonds to a central metal ion. Numerous artificial nucleosides bearing ligands as aglycones have been introduced to broaden the scope of metal-mediated base pairing.<sup>2</sup> Experimental structure determinations both by X-ray crystallography and by NMR spectroscopy have clearly proven that the formation of metal-mediated base pairs is compatible with the well-known B-DNA conformation.<sup>3</sup> Some prominent possible applications for such DNA-metal conjugates include charge transfer,<sup>4</sup> expansion of the genetic code,<sup>5</sup> sensors of various types,<sup>1c</sup> and the use of DNA as a logic gate,<sup>6</sup> to name just a few.

While most metal-mediated base pairs contain one metal ion per base pair,<sup>7</sup> few examples are known of base pairs which comprise two metal ions. These include homo base pairs of various pyrimidine derivatives<sup>8</sup> as well as the hetero base pair 1,3-dideazaadenine–Ag(I)<sub>2</sub>–thymine.<sup>9</sup> Dinuclear metal-mediated base pairs represent a particular challenge as two positively charged entities need to be positioned at very

short distance. On the other hand, they enable a high functionalization of nucleic acids with metal ions and thereby significantly increase the scope of DNA-based nanotechnology.<sup>9</sup> We report here the first example of a doubly metal-mediated homo base pair derived from a purine nucleobase. The exocyclic etheno adduct of adenine, 1,*N*<sup>6</sup>-ethenoadenine ( $\epsilon$ A, Scheme 1), is a DNA lesion that is known to form in the presence of *e.g.* vinyl chloride,<sup>10</sup> although endogenous pathways for its formation *in vivo* have also been suggested.<sup>11</sup> Its metal binding properties are superior to those of the parent compound adenine.<sup>12</sup>

To investigate metal-mediated base pair formation, an oligonucleotide duplex with a central  $\epsilon$ A :  $\epsilon$ A mispair was synthesized (Scheme 2). This particular sequence was chosen because it had been used in various earlier studies on other metal-mediated base pairs and therefore allows a good comparison of the observations.<sup>7a,b,d,13</sup> The melting temperature  $T_m$  of the unmetalated duplex comprising the central  $\epsilon$ A :  $\epsilon$ A mispair amounts to 35.7 °C. It is significantly more stable than previously investigated other mispairs such as imidazole : imidazole, which showed  $T_m$  values in the range of 21.7–30.4 °C.<sup>7b</sup> Its stability is lower though than that of the corresponding duplexes with regular A : T (42.5 °C) or G : C



**Scheme 1** 1,*N*<sup>6</sup>-ethenodeoxyadenosine including atom numbering scheme.

5' – d (GAG GGA XAG AAA G)  
3' – d (CTC CCT XTC TTT C)


**Scheme 2** Oligonucleotide double helix under investigation (X =  $\epsilon$ A).

<sup>a</sup>Westfälische Wilhelms-Universität Münster, Institut für Anorganische und Analytische Chemie, Corrensstr. 28/30, 48149 Münster, Germany.

E-mail: mueller.j@uni-muenster.de; Fax: +49 251 8336007; Tel: +49 251 8336006

<sup>b</sup>Westfälische Wilhelms-Universität Münster, NRW Graduate School of Chemistry, Corrensstr. 28/30, 48149 Münster, Germany

†Electronic supplementary information (ESI) available: Experimental details, spectroscopic characterization, stack plot of <sup>1</sup>H NMR spectra. See DOI: 10.1039/c4dt02663b



**Fig. 1** Increase of the melting temperature  $T_m$  upon the addition of  $\text{Ag(I)}$ . Inset: UV melting profile of the duplex with one central  $\epsilon\text{A}:\epsilon\text{A}$  mispair in the absence (black) and presence (red) of two equivalents of  $\text{Ag(I)}$ .  $A_{\text{norm}} = (A - A_{\text{min}})/(A_{\text{max}} - A_{\text{min}})$ .


(45.4 °C) base pairs.<sup>13</sup> Hence, the metal-free duplex is destabilized with respect to the purely canonical duplex because of the  $\epsilon\text{A}:\epsilon\text{A}$  mispair formation, whereas the large  $\pi$  surface of  $\epsilon\text{A}$  is expected to be the main contributor towards its stabilization with respect to mispairs with a smaller  $\pi$  surface.

Upon the addition of  $\text{Ag(I)}$ , the melting temperature of the DNA duplex increases significantly until two equivalents of  $\text{Ag(I)}$  are present per base pair (Fig. 1). From a plot of  $T_m$  versus added equivalents of  $\text{Ag(I)}$  it can clearly be discerned that the first two  $\text{Ag(I)}$  ions induce the largest stabilization, whereas additional  $\text{Ag(I)}$  has only very minor effects on the thermal stability of the duplex. This weak additional stabilization by  $\text{Ag(I)}$  has previously been attributed to an electrostatic interaction with the negatively charged DNA backbone.<sup>13</sup> Taken together, this is a first good indication for the formation of a doubly metal-mediated base pair  $\epsilon\text{A-Ag(I)}_2\text{-}\epsilon\text{A}$ .

Additional spectroscopic experiments have been performed to validate in an independent manner the 2 : 1 stoichiometry of the base pair. Towards this end, CD and UV spectra of the DNA duplex have been recorded in the presence of various amounts of  $\text{Ag(I)}$ . Fig. 2a shows a plot of the CD spectra, indicating a constant positive Cotton effect around 275 nm. In


addition, two negative Cotton effects at 232 nm and 244 nm are observed in the absence of  $\text{Ag(I)}$ , merging into a single and more intense negative Cotton effect at 236 nm upon the addition of  $\text{Ag(I)}$ . The CD spectrum in the presence of two equivalents of  $\text{Ag(I)}$  is reminiscent of that of B-DNA.<sup>14</sup> The somewhat unusual CD spectrum in the absence of  $\text{Ag(I)}$  is likely to be due to the presence of the  $\epsilon\text{A}:\epsilon\text{A}$  mispair in which the nucleosides probably adopt a conformation not found in a regular B-DNA duplex (e.g. by stacking on top of one another or by extruding one base out of the base stack). A plot of the molar ellipticity at 236 nm versus added equivalents of  $\text{Ag(I)}$  (Fig. 2b) confirms that the distinct change of the ellipticity occurs only until two equivalents of  $\text{Ag(I)}$  are present. Excess  $\text{Ag(I)}$  only leads to minor additional changes in the CD spectrum. The same behaviour is found for the UV spectra of the duplex (Fig. 2c): upon adding the first two  $\text{Ag(I)}$  ions per  $\epsilon\text{A}:\epsilon\text{A}$  mispair a significant change of the UV spectrum is observed, whereas excess  $\text{Ag(I)}$  does not significantly change the spectrum. As the changes in absorbance are most prominent in the wavelength region where the aromatic nucleobases absorb, a binding of the  $\text{Ag(I)}$  ions to the nucleobases can be concluded. Previous experiments with same sequence comprising entirely canonical base pairs indicate no significant changes of the spectral properties upon the addition of  $\text{Ag(I)}$ .<sup>13</sup> Hence, as all experiments nicely show that the binding site with the highest  $\text{Ag(I)}$  binding affinity binds exactly two  $\text{Ag(I)}$  ions, the  $\epsilon\text{A}:\epsilon\text{A}$  mispair can be unambiguously identified as the metal binding site. Scheme 3 shows the proposed structure of this doubly metal-mediated base pair.

The model nucleobase 9-ethyl-1, $N^6$ -ethenoadenine was applied for another independent proof of the metal-binding stoichiometry by performing  $^1\text{H}$  NMR- and  $^{15}\text{N}$  NMR-spectroscopy-based titrations of the ligand with  $\text{Ag(I)}$  ions (Fig. 3 and ESI†). As can clearly be discerned from Fig. 3a, the chemical shifts of the aromatic protons change until one  $\text{Ag(I)}$  ion is present per ligand, corresponding to two  $\text{Ag(I)}$  ions per pair of ligands.  $^{15}\text{N}$  NMR spectroscopy unambiguously confirms N6 and N7 as the sites of metalation, as both nitrogen atoms experience an upfield shift of >40 ppm upon binding of  $\text{Ag(I)}$  whereas the chemical shifts of the other three nitrogen atoms




**Fig. 2** (a) CD spectra of the duplex with one central  $\epsilon\text{A}:\epsilon\text{A}$  mispair upon the addition of  $\text{Ag(I)}$ . (b) Change of molar ellipticity at 236 nm upon the addition of  $\text{Ag(I)}$ . (c) Change of absorbance at 257 nm upon the addition of  $\text{Ag(I)}$ .





**Scheme 3** Structure of the  $\epsilon\text{A-Ag(I)}_2\text{-}\epsilon\text{A}$  base pair ( $\text{R}, \text{R}' = \text{DNA backbone}$ ).



**Fig. 3** Chemical shifts of the model nucleobase 9-ethyl-1, $N^6$ -etheno-adenine in the presence of varying amounts of  $\text{Ag(I)}$ . (a)  $^1\text{H}$  NMR; (b)  $^{15}\text{N}$  NMR.

change by only 1–6 ppm (Fig. 3b). Taken together, these NMR data confirm the formation of a dinuclear complex also in solution. In contrast to the situation inside the DNA duplex, the dinuclear complex can adopt two distinct conformations in solution. In the case of transoid ethyl substituents, two identical  $\text{Ag(I)}$  binding sites are present ( $\text{N6-Ag(I)-N7}$ ), whereas a cisoid orientation of the ethyl substituents (Scheme 3) leads to two distinct  $\text{Ag(I)}$  environments ( $\text{N6-Ag(I)-N6}$  and  $\text{N7-Ag(I)-N7}$ ). Unfortunately, it is not possible to differentiate between these conformations based on the NMR data, as the  $\text{Ag(I)}$  ions bind in a labile fashion, leading to a fast exchange on the NMR time scale (as indicated by the smooth change of the  $^1\text{H}$  and  $^{15}\text{N}$  NMR chemical shifts upon metalation). As the B-DNA

geometry enforces a cisoid orientation of the glycosidic bonds, and the  $^{15}\text{N}$  NMR data show that both N6 and N7 are metalated, it can nonetheless be concluded that inside the DNA duplex the metal-mediated base pair adopts the structure shown in Scheme 3.

## Conclusions

The exocyclic etheno adduct of adenine, 1, $N^6$ -etheno-adenine ( $\epsilon\text{A}$ ), is capable of forming doubly  $\text{Ag(I)}$ -mediated homo base pairs. This is the first time that doubly metal-mediated homo base pairs are being reported for a purine derivative, and the first example for a Hoogsteen-type metal-mediated base pair within a regular B-DNA context. Particularly in view of the commercial availability of  $\epsilon\text{A}$ , this finding represents a significant step forward towards easily prepared, highly metal-functionalized nucleic acids.

Financial support by the NRW Graduate School of Chemistry (GSC-MS) and the DFG (SFB 858) is greatly acknowledged. This work was performed in the context of COST Action CM1105.

## Notes and references

- (a) J. Müller, *Eur. J. Inorg. Chem.*, 2008, 3749–3763; (b) D. A. Megger, N. Megger and J. Müller, *Met. Ions Life Sci.*, 2012, **10**, 295–317; (c) P. Scharf and J. Müller, *Chem-PlusChem*, 2013, **78**, 20–34.
- (a) G. H. Clever and M. Shionoya, *Met. Ions Life Sci.*, 2012, **10**, 269–294; (b) Y. Takezawa and M. Shionoya, *Acc. Chem. Res.*, 2012, **45**, 2066–2076; (c) G. H. Clever and M. Shionoya, *Coord. Chem. Rev.*, 2010, **254**, 2391–2402; (d) G. H. Clever, C. Kaul and T. Carell, *Angew. Chem., Int. Ed.*, 2007, **46**, 6226–6236; (e) A. Ono, H. Torigoe, Y. Tanaka and I. Okamoto, *Chem. Soc. Rev.*, 2011, **40**, 5855–5866.
- (a) S. Johannsen, N. Megger, D. Böhme, R. K. O. Sigel and J. Müller, *Nat. Chem.*, 2010, **2**, 229–234; (b) S. Kumbhar, S. Johannsen, R. K. O. Sigel, M. P. Waller and J. Müller, *J. Inorg. Biochem.*, 2013, **127**, 203–210; (c) H. Yamaguchi, J. Šebera, J. Kondo, S. Oda, T. Komuro, T. Kawamura, T. Dairaku, Y. Kondo, I. Okamoto, A. Ono, J. V. Burda, C. Kojima, V. Sychrovský and Y. Tanaka, *Nucleic Acids Res.*, 2014, **42**, 4094–4099; (d) Y. Kondo, T. Yamada, C. Hirose, I. Okamoto, Y. Tanaka and A. Ono, *Angew. Chem., Int. Ed.*, 2014, **53**, 2385–2388.
- (a) S. Liu, G. H. Clever, Y. Takezawa, M. Kaneko, K. Tanaka, X. Guo and M. Shionoya, *Angew. Chem., Int. Ed.*, 2011, **50**, 8886–8890; (b) T. Ehrenschröder, W. Schmucker, C. Wellner, T. Augenstein, P. Carl, J. Harmer, F. Breher and H.-A. Wagenknecht, *Chem. – Eur. J.*, 2013, **19**, 12547–12552.
- C. Kaul, M. Müller, M. Wagner, S. Schneider and T. Carell, *Nat. Chem.*, 2011, **3**, 794–800.



- 6 D.-L. Ma, H.-Z. He, D. S.-H. Chan and C.-H. Leung, *Chem. Sci.*, 2013, **4**, 3366–3380.
- 7 (a) T. Richters and J. Müller, *Eur. J. Inorg. Chem.*, 2014, 437–441; (b) T. Richters, O. Krug, J. Kösters, A. Hepp and J. Müller, *Chem. – Eur. J.*, 2014, **20**, 7811–7818; (c) K. Petrovec, B. J. Ravoo and J. Müller, *Chem. Commun.*, 2012, **48**, 11844–11846; (d) K. Seubert, C. Fonseca Guerra, F. M. Bickelhaupt and J. Müller, *Chem. Commun.*, 2011, **47**, 11041–11043; (e) F.-A. Polonius and J. Müller, *Angew. Chem., Int. Ed.*, 2007, **46**, 5602–5604; (f) D. Böhme, N. Düpre, D. A. Megger and J. Müller, *Inorg. Chem.*, 2007, **46**, 10114–10119.
- 8 (a) I. Okamoto, K. Iwamoto, Y. Watanabe, Y. Miyake and A. Ono, *Angew. Chem., Int. Ed.*, 2009, **48**, 1648–1651; (b) I. Okamoto, T. Ono, R. Sameshima and A. Ono, *Chem. Commun.*, 2012, **48**, 4347–4349; (c) H. Mei, I. Röhl and F. Seela, *J. Org. Chem.*, 2013, **78**, 9457–9463.
- 9 D. A. Megger, C. Fonseca Guerra, J. Hoffmann, B. Brutschy, F. M. Bickelhaupt and J. Müller, *Chem. – Eur. J.*, 2011, **17**, 6533–6544.
- 10 J. A. Hall, R. Saffhill, T. Green and D. E. Hathway, *Carcinogenesis*, 1981, **2**, 141–146.
- 11 F.-L. Chung, H.-J. C. Chen and R. G. Nath, *Carcinogenesis*, 1996, **17**, 2105–2111.
- 12 K. H. Scheller and H. Sigel, *J. Am. Chem. Soc.*, 1983, **105**, 3005–3014.
- 13 I. Sinha, J. Kösters, A. Hepp and J. Müller, *Dalton Trans.*, 2013, **42**, 16080–16089.
- 14 M. Vorlíčková, I. Kejnovská, K. Bednářová, D. Renčíuk and J. Kypr, *Chirality*, 2012, **24**, 691–698.

