Organic & Biomolecular Chemistry



PAPER

View Article Online



Cite this: *Org. Biomol. Chem.*, 2025, **23**. 8719

Modular synthesis of zinc(||)-bis(triazole) recognition sites for the conformational control of foldamers

Flavio della Sala, o a, Benedicte Doerner and Simon J. Webb **

Zinc(ii) bis(triazolyl)(pyridyl)amine (Zn(BTPA)) complexes on the end of α -amino-iso-butyric acid (Aib) foldamers are able to transfer chirality from bound anions to the helical foldamer body. Zn(BTPA) could be obtained by simple synthetic methodology that allowed a range of functional groups to be installed around the binding site, exemplified with a fluorophore, a macrocyclic bridge and Aib itself. Changing functional group did not prevent chiral ligands from controlling foldamer conformation, although differences in complexation kinetics and equilibria were observed. Addition of acetate gave a 2:1 foldamer: acetate intermediate at sub-stoichiometric acetate; a similar intermediate was implied during titration with Boc-Pro. A bulkier phosphate ligand or a more sterically hindered site did not form similar intermediates. The modular construction of Zn(BTPA)-capped foldamers will allow these conformational relays to be installed in a wide range of biomimetic constructs.

Received 29th July 2025, Accepted 28th August 2025 DOI: 10.1039/d5ob01226k

rsc.li/obc

Introduction

Metal ions play important roles in enzyme active sites and during ligand binding to proteins. Zinc(II) for example can control protein shape by coordinating to side chains on different secondary structures, as found in the zinc finger motif. The resulting structurally defined regions are critical for protein–DNA binding. Metal ions in proteins can also directly bind to cognate ligands, *e.g.* some calcium(II)-dependent animal lectins form direct coordination links between the sugar hydroxyls and bound calcium(II). Ligand binding to metal ions in proteins can then induce global conformational changes, with the binding of oxygen to haemoglobin one of the best known examples. Protein folding into geometrically defined pockets around the metal ions are important for these proteins to function.

Folded oligomers (foldamers) can coordinate to metal ions, ^{4,5} with some foldamers shown to mimic metalloprotein structure, including the zinc finger motif. ^{6,7} The metal ions can also become catalytic centres on the foldamers and/or provide locations for ligand binding. Ligand-induced conformational change, as observed in haemoglobin, can also be replicated. Upon binding, some ligands will perturb the conformational landscape of dynamic foldamers, ⁸⁻¹¹ which are a

type of foldamer that undergo rapid conformational interchange. α-Amino-iso-butyric acid (Aib) foldamers are rod-like dynamic foldamers that can undergo rapid long range (>1 nm) conformational change in response to external stimuli. Their chief conformational populations are 310 helices that have either a right-handed (P) or left-handed (M) screw-sense. Ligand binding at one terminus can cause these Aib foldamers to undergo end-to-end conformational change, which changes the proportion of P to M helices. These changes can be expressed as the helical excess, h.e., which is the fractional excess of P helix over M helix (h.e. = ([P] - [M])/([P] + [M])); this can be calculated from representative NMR spectra. 12 This simple P vs. M conformational landscape has led to Aib foldamers being used to mimic aspects of biological signal transduction, particularly how ligand recognition can initiate conformational change across multi-nanometre distances. To mediate ligand recognition, Zn(II) and Cu(II) complexes can be placed at one end of the Aib foldamers. 9,13-15 Like related complexes in the literature, 16-21 they bind chiral anions, including carboxylates.9a

The M(I)-bis(quinolyl)(pyridyl)amine (BQPA, Fig. 1) binding site is the best to date for turning ligand chirality into a change in the P:M screw-sense ratio of an Aib foldamer. Its effectiveness has been ascribed to the steric bulk of the quinolyl arms and the "propeller" conformation they adopt. ²² Both Zn(II) and Cu(II) complexes are effective. Foldamers capped with Zn(II)-BQPA could sense the enantiomeric excess (ee) of scalemic mixtures of chiral carboxylates, ²² due to rapid ligand exchange at the Zn(II) site. ¹⁵ A Cu(II)-(BQPA) recognition site

^aDepartment of Chemistry, University of Manchester, Oxford Road, Manchester M13

9PL. UK E-mail: S Webh@manchester ac uk

^bManchester Institute of Biotechnology, University of Manchester, 131 Princess Street, Manchester M1 7DN, UK

Old design: BQPA-capped foldamer Zn(1)·2ClO₄

H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O
H
N
O

Fig. 1 Reported foldamer $Zn(1)\cdot 2ClO_4$ bearing the bis(quinolyl)(pyridyl) amine (BQPA) binding pocket.¹⁵ Perchlorate anions omitted for clarity.

provided a synthetic receptor that responded to chiral carboxy-lates (the input signal) by undergoing a conformational change either in solution or deep into lipid bilayers. 9a

These previous Aib foldamers with BQPA have a metal ionchelating pocket that is symmetric and unfunctionalised, ^{13–15} unlike the naturally asymmetric binding pockets of proteins. We wished to retain the desirable recognition characteristics of BQPA but add functionality to the "arms" around the binding site. BQPA itself was difficult to modify and its relatively poor stability also required it to be added last during synthesis.

To better replicate metal ion containing binding pockets in proteins, simpler methods were needed to introduce functionality and decrease symmetry. To this end, we have explored the use of copper-catalysed alkyne–azide cycloaddition (CuAAC) reactions to create metal ion binding pockets^{23,24} (Fig. 2 and 3)




Fig. 2 Foldamers Zn(2–5)·2ClO₄ bearing the bis(triazolyl)(pyridyl)amine (BTPA) binding pocket, which is constructed in a modular fashion from different azides. Perchlorate anions are omitted for clarity.




Fig. 3 Foldamer $Zn((R)-6)\cdot 2ClO_4$ bearing the (pyridyl)(triazolyl)(pyridyl) (PTPA) binding pocket with different binding arms. Perchlorate anions are omitted for clarity.

that are flanked by selected substituents. The use of CuAAC allows simple modification of the synthetic route to provide either symmetrical or unsymmetrical metal ion binding pockets.

To study the effect of these binding sites at the N-terminus of Aib foldamers, we placed the recently reported (R)-1-(tri-fluoromethyl)-ethylamido ((R)-TFEA) reporter group at the C-terminus.²⁵ This group provides ¹⁹F NMR spectroscopic reports on changes in the conformational populations of Aib foldamers that are induced by chiral anionic ligands binding to the Zn(π) (such as carboxylates or phosphates, Fig. 4).²⁶

Results and discussion

Synthesis

Fluorinated motifs have great utility for both controlling the helical screw-sense in Aib foldamers and reporting on the helical excess of Aib foldamers. $^{8,27-29}$ The TFEA 19 F NMR reporter group allows the determination of the helical excess induced at the N-terminus of Aib foldamers (h.e.₀) both in organic solvents, micelles and when embedded in phospholipid vesicles. 25,26 The robustness of TFEA towards synthetic conditions made it attractive for the development of new binding pockets. Either the (R)- or the (S)-TFEA reporter were conjugated to the readily accessible foldamer N_3 (Aib)₄OH, then azide hydrogenation and elaboration at the N-terminus

Fig. 4 Chiral anionic ligands for zinc(n). Boc-Pro-O⁻, Boc-Pip-O⁻ and TiPSY⁻ are derived from N-(tert-butoxycarbonyl)-proline, N-(tert-butoxycarbonyl)-pipecolinic acid and 3,3'-bis(triphenylsilyl)-1,1'-binaphthyl-2,2'-diyl hydrogenphosphate respectively.

gave key bis-alkyne precursors (R)-7 and (S)-7 (Scheme 1). CuAAC reactions on these precursors had been shown to give BTPA-capped fluorescent foldamers (R)-2 and (S)-2. ²⁶ Applying the same CuAAC procedure but using N₃AibO^tBu, ¹³ 8-azido-3,6-dioxaoctanol or 8-azido-3,6-dioxaoctyl mesylate gave the other N-functionalised BTPA foldamers (S)-3, (R)-4 and (R)-8 in good yield (see SI, Section S2). Foldamer (R)-8 in turn gave access to (R)-5, which has a catechol/oligo(ethyleneglycol) bridge that was created through a Cs⁺ templated S_N2 reaction. This oligoether bridge is close to the tetrapodal Zn(π) chelating site and might enhance or otherwise alter ligand recognition at Zn(π). It could also permit the introduction of rotaxanated structures.

Installing a monoalkyne in the place of the dialkyne in 7 can give unsymmetrical metal ion binding sites (Fig. 3). The (pyridyl)(triazolyl)(pyridyl)amine (PTPA) moiety of (R)-6 was accessed by reductive amination of amine 9.³⁰ Mono-alkyne precursor 10 was hydrolysed then coupled to NH₂(Aib)₄((R)-TFEA). Finally a CuAAC reaction with N₃AibO^tBu¹³ provided PTPA-capped foldamer (R)-6.

a) O
$$R^1$$
 R^2 R^2 R^3 R^4 R^2 R^4 R^2 R^4 R^4

Scheme 1 (a) Synthesis of foldamers 2–5. Reagents: a. 6-Azido-2-ethyl-1H-benzo[de]isoquinoline-1,3(2H)-dione, CuSO₄·5H₂O, sodium ascorbate, DMF, rt. 26 b. N₃AibO^fBu, 13 CuSO₄·5H₂O, sodium ascorbate, DMF, rt. c. 8-Azido-3,6-dioxaoctyl mesylate \$18, CuSO₄·5H₂O, sodium ascorbate, DMF, rt. d. 8-Azido-3,6-dioxaoctanol, CuSO₄·5H₂O, sodium ascorbate, DMF, rt. e. Catechol, Cs₂CO₃, CH₃CN, reflux. (b) Synthesis of foldamers 6, 11. Reagents: f. Methyl 6-(bromomethyl)nicotinate \$15, DIPEA, CH₃CN, rt. g. KOH, CH₃OH, reflux. h. (Aib)₄((R)-TFEA), EDC·HCl, HOBt, Et₃N, CH₃CN, rt.

(R)-11

b

(R)-6, R^1 = H, R^2 = Me

Addition of zinc(II) perchlorate

Foldamer 2 was chosen to exemplify complexation of Zn(II) by the BTPA group. Foldamer (R)-2 was titrated with zinc perchlorate in CD₃CN (Fig. 5).³¹ H NMR spectroscopy showed a gradual, generally downfield, shift of aromatic peaks over the course of the titration, consistent with fast exchange between free and complexed foldamers at substoichiometric ratios of Zn(II). Concurrent resonance broadening was also initially observed, before sharpening and a decrease of chemical shift movement at ca. 0.7 eq. of zinc. After complexation, significant downfield shifts were observed for the ortho-pyridyl (Ho), parapyridyl (H_p), and triazole (H_t) proton resonances ($\Delta \delta$ = 187, 428 and 218 ppb respectively, Fig. 5 and Fig. S1 in the SI), shifts that are consistent with coordination to Zn(II). 32 Another diagnostic change was a downfield shift (280 ppb) and splitting of the triazole-methylene (Harm) protons from a singlet to four doublets (Fig. S1 in the SI), consistent with Zn(II) complexation stopping tertiary amine inversion and making the arms inequivalent with diastereotopic methylenes. 19F NMR spectroscopy showed that Zn(II) complexation at the N-terminus gave only a very small +8 ppb downfield shift for the C-terminal CF₃ reso-




Fig. 5 Addition of one equivalent of zinc($_{\rm II}$) perchlorate to foldamer (R)-2. (a) Labelling of selected protons around the Zn($_{\rm II}$) ion. (b) Partial stacked 1 H NMR spectra (CD $_{3}$ CN, 400 MHz, 298 K). (c) Partial stacked 19 F NMR spectra (CD $_{3}$ CN, 376 MHz, 298 K); spectra referenced with C $_{6}$ F $_{6}$ at -164.38 ppm. 33

nance of the reporter group (Fig. 5c), indicating little involvement with the newly installed $Zn(\pi)$.

Similar changes were observed for (S)-3, (R)-4 and (R)-5 (see SI, Section S3). However, the 1H NMR spectrum of $Zn((R)-5)\cdot 2ClO_4$ (see the SI, Fig. S4) showed an increase in the number and broadness of peaks from the crown-ether protons, suggesting additional conformational states for the macrocycle after Zn(II) addition. Interestingly, the catechol protons were downfield shifted ($\Delta\delta$ ca. 130 ppb, see SI Fig. S4) suggesting that the phenyl ring may be bent over the binding pocket. Analysis of model compound $Zn(S21)\cdot 2ClO_4$, which lacks the Aib foldamer, supported this suggestion as it showed a NOE between the catechol protons and a methylene at the other end of the crown ether macrocycle (the OCH_2CH_2N protons, see SI Fig. S6 and 7).

Although the binding pocket of $Zn((R)-6)\cdot 2ClO_4$ is less symmetric than the others, analogous behaviour was observed upon addition of $zinc(\pi)$ perchlorate (see SI Fig. S8 and 9). However, unlike the foldamers with symmetric binding pockets (*e.g.* (*R*)-5, Fig. 6a), the addition of $zinc(\pi)$ perchlorate




Fig. 6 (a and b) Complexation of Zn(II) to (R)-5 in CD₃CN gives a small change in δ_F (CF₃) in the ¹⁹F NMR spectrum. (c and d) Complexation of Zn(II) to (R)-6 in CD₃CN causes splitting of the CF₃ resonance in the ¹⁹F NMR spectrum due to the generation of a stereogenic centre at the N-terminus and the production of diastereomers.

to (R)-6 caused the ¹⁹F singlet to split into two overlapping singlets (Fig. 6d, $\Delta\delta$ = 29 ppb). This is consistent with complexation to Zn(II) generating a chiral centre at the N-terminus (Fig. 6c), which in conjunction with the chiral (R)-TFEA group leads to the formation of diastereomeric complexes with distinct CF₃ resonances. Addition of EDTA to sequester the zinc (II) supported this hypothesis, as the two peaks merged and returned to their original position (see SI Fig. S9).

Complexation studies with anions in CD₃CN

Binding to acetate. As a simple anion for initial binding studies, achiral tetra-n-butylammonium (TBA) acetate was used. TBA acetate (up to 2 eq.) was titrated into $Zn((R)-2)\cdot 2ClO_4$. Both 1H and ^{19}F NMR spectroscopy confirmed that acetate bound to the Zn(II) pocket. Its enantiomer, $Zn((S)-2)\cdot 2ClO_4$, showed the same changes.

The *ortho*-pyridyl proton (H_o) resonance splits into two. One resonance gradually moves downfield by ca. 0.213 ppm, suggesting the unbound state and this bound state are in fast exchange on the 1H NMR spectroscopy timescale. Another 1H NMR resonance for H_o also appears further downfield (by ca. 1 ppm) after 0.1 eq. ligand has been added; this resonance disappears after 1.4 eq. ligand has been added (Fig. 7b). This H_o




Fig. 7 (a) Acetate complexed to $Zn((R)-2)\cdot 2ClO_4$. (b) Partial 1H NMR spectra (400 MHz, 298 K) showing the *ortho*-CH resonance during the titration of $Zn((R)-2)\cdot 2ClO_4$ (4.25 mM, 550 μ L) in CD₃CN with TBA acetate (47 mM, up to 2 eq.). (c) Partial ^{19}F NMR spectra (376 MHz, 298 K) showing the TFEA reporter region during the titration of $Zn((R)-2)\cdot 2ClO_4$ (4.25 mM, 550 μ L) in CD₃CN with TBA acetate (47 mM, up to 2 eq.). Spectra referenced with C_6F_6 at -164.38 ppm. 33


signal seems to come from a new species that is in slow exchange with the other two species. A corresponding new ¹⁹F resonance also appears, upfield of the original reporter signal, mirroring the new signal in the ¹H NMR spectrum by appearing at 0.1 eq. and disappearing at 1.4 eq. This mirroring indicates that the new signals arise from a single species. Indeed after this new peak has disappeared, the ¹⁹F NMR spectrum showed no shift from the uncomplexed ¹⁹F peak (Fig. 7c), confirming that an achiral carboxylate has no effect over the P/Mratio. Diffusion ordered spectroscopy (DOSY) ¹H spectra of Zn ((R)-2)-2ClO₄ with and without 0.7 eq. of TBA acetate confirmed that these new peaks belong to a single separate species. The DOSY data also shows that this species is larger than uncomplexed $Zn((R)-2)\cdot 2ClO_4$, with a hydrodynamic radius of 13.6 Å compared to 10.2 Å for the acetate-free foldamer (see SI Section S4.3.3). These DOSY data suggest the new species may involve more than one foldamer.

Binding to chiral anions

In previous work, Boc-D-Pro, Boc-D-Pip and S-TiPSY, were all shown to produce an M screw-sense in $Zn((R)-2)\cdot 2ClO_4$ or Zn ((S)-2)-2ClO₄. ²⁶ The maximum h.e.₀ (the helical excess induced adjacent to the chiral group) of each was estimated by interpolation of $\Delta \delta_{\rm F}({\rm CF_3})$ into our previously reported calibration curve (see SI Section S5), 25,26 giving h.e. values of -21%, -7% and -23% respectively.34 Although these values are half (or less) of the h.e.0 values that these ligands induced in Zn (1)·2ClO₄,³⁵ this performance is better than other analogues of Zn(1)·2ClO₄ that had the quinolyl or pyridyl motifs replaced by pyridyl or triazolyl respectively; these gave no clear relays of chirality. Instead complex behaviour was observed, including equilibria with unfavourable exchange kinetics for NMR studies and low solubility. 14,35 To understand how replacing the quinolyl groups with triazolyl avoids these problems, these three chiral anions (Fig. 4) were titrated into Zn(BTPA)-capped foldamers and the equilibria monitored by ¹H and ¹⁹F NMR spectroscopy.

Both $Zn((R)-2)\cdot 2ClO_4$ and $Zn((S)-2)\cdot 2ClO_4$ were titrated with up to 2 eq. Boc-D-Pro or Boc-L-Pro in the presence of 2,6-lutidine (a non-coordinating base, 1.2 eq. with respect to Boc-Pro). The four combinations give pairs of enantiomeric mixtures, *e.g.* $Zn((R)-2)\cdot 2ClO_4/Boc-L-Pro$ is enantiomeric with $Zn((S)-2)\cdot 2ClO_4/Boc-D-Pro$ (Fig. 8) and gives identical NMR spectra during the titration.

No significant problems with exchange kinetics or solubility were observed. The 1H NMR spectra of $Zn((R)\text{-}2)\text{-}2ClO_4$ showed a downfield shift of the *ortho*-pyridyl (H_o) resonance with Boc-D-Pro and Boc-L-Pro (with 2,6-lutidine, see SI Fig. S14). As observed for acetate, this shift was gradual with increasing ligand concentration, which is indicative of fast exchange between the unbound and ligand-bound states on the 1H NMR spectroscopy time-scale at 298 K. However, unlike during the addition of TBA acetate, no additional H_o signal appeared further downfield in the 1H NMR spectrum. Instead the resonance became much weaker and quite broad (albeit still visible) between 0.3 and 1 eq. of Boc-Pro. The 1H NMR reso-






Fig. 8 (a) Partial ¹⁹F NMR spectra (376 MHz, 298 K) showing the CF₃ region during the titration of Zn((R)-2)·2ClO₄ (4.25 mM, 550 µL) in CD₃CN with 47 mM Boc-Pro (up to 2 eq. and 2.4 eq. 2,6-lutidine). C₆F₆ as an internal standard, referenced at –164.38 ppm. ³³ This internal standard gave δ_F (CF₃)₀ as –77.6648 ppm. (b) Representative data fitting of $\Delta\delta_F$ (CF₃) using Dynafit during the titration of Zn((S)-2)·2ClO₄ in CD₃CN with Boc-D-Pro. Binding model: 2:1 [Zn((S)-2)·2ClO₄]/[Boc-D-Pro] with $K_{11} = 1 \times 10^5$ M⁻¹ and $K_{21} = 2 \times 10^3$ M⁻¹. Conditions: [Zn((S)-2)·2ClO₄] = 8.36 mM, [Boc-Pro] = 0–18.9 mM, [2,6-lutidine] = 0–22.7 mM. CFCl₃ as an internal standard, referenced at –1.14 ppm. ^{31,33} This internal standard gave δ_F (CF₃)₀ as –77.6205 ppm.

nances from the (R)-TFEA reporter were little affected by the addition of carboxylate.

In the ¹⁹F NMR spectrum, the titration of up to 2 eq. Boc-D-Pro (with 2,6-lutidine) into $Zn((R)-2)\cdot 2ClO_4$ gave a gradual downfield shift in $\delta_F(CF_3)$, confirming an increase in the proportion of M helix (Fig. 8a). Conversely, titration of Boc-D-Pro into the enantiomer $Zn((S)-2)\cdot 2ClO_4$ gave a gradual upfield shift in $\delta_F(CF_3)$; this is also consistent with an increase in the proportion of M helix (Fig. 8b). Plotting $\Delta\delta_F(CF_3)$ against concentration revealed a clear sigmoidal profile. Above 2 eq. Boc-D-Pro, $\delta_F(CF_3)$ remaining constant. The sigmoidal titration

profiles for $\operatorname{Zn}((R)\text{-}2)\text{-}2\operatorname{ClO}_4$ and $\operatorname{Zn}((S)\text{-}2)\text{-}2\operatorname{ClO}_4$ are consistent with the formation of intermediates at sub-stoichiometric Boc-Pro that are not visible in the ¹⁹F NMR spectrum.

Titration with Boc-D-Pip produced similar effects to Boc-D-Pro. Addition of Boc-D-Pip/2,6-lutidine to $Zn((R)-2)\cdot 2ClO_4$ in CD_3CN resulted in strong broadening of the H_o resonance in the 1H NMR spectrum with the concurrent appearance of a new, downfield broadened peak at 9.46 ppm (see SI Fig. S18). Both H_o resonances were very weak between 0.3 and 1.0 equivalents of ligand (see SI Section S4.3.6) but the observation of two, albeit broadened, resonances during the titration suggests slower exchange on the 1H NMR spectroscopy timescale than Boc-Pro. In contrast, $δ_F(CF_3)$ in the ^{19}F NMR spectrum gradually shifts downfield, consistent with fast exchange between free and bound foldamer. $δ_F(CF_3)$ follows a sigmoidal profile during the titration but with extensive signal broadening between 0.4 and 1 eq. Boc-D-Pip (see SI Fig. S19).

Like carboxylates, phosphates are reported to complex to zinc-tetraamine complexes, ^{36–38} so the chiral TiPSY anion was also assessed (Fig. 4). TiPSY/2,6-lutidine however produced different behaviour upon titration into Zn((*R*)-2)-2ClO₄. The ¹H and ¹⁹F NMR spectra show the bound and unbound states are now in slow exchange (Fig. 9 and see SI Fig. S20–22), with integration of the signals providing no evidence for the formation of an intermediate species during the addition of TiPSY/2,6-lutidine. The TiPSY anion is a much larger ligand than acetate, Boc-Pro-O⁻ or Boc-Pip-O⁻, perhaps creating a steric block to the formation of intermediate complexes that involve more than one foldamer.




Fig. 9 Partial 19 F NMR spectra (376 MHz, 298 K) showing the CF₃ region for the titration of Zn((R)-2)·2ClO₄ (4.25 mM, 550 μ L) in CD₃CN with 47 mM TiPSY (up to 2 eq. and 2.4 eq. 2,6-lutidine). C₆F₆ as an internal standard, referenced at -164.38 ppm. 33

Changing from the flat aromatic moieties in $Zn((S)-2)\cdot 2ClO_4$ to the 2-isobutyrate of $Zn((S)-3)\cdot 2ClO_4$ was hoped to encumber the binding site and allow bound ligands to better control helical screw-sense. Adding Boc-D-Pro/2,6-lutidine to $Zn((S)-3)\cdot 2ClO_4$ induced an M-helical bias, the same as for $Zn((S)-2)\cdot 2ClO_4$ (Fig. S24). Indeed the induction of M helix by Boc-D-Pro-O⁻ (and *vice versa* for Boc-L-Pro-O⁻) was observed for all BTPA-capped Aib foldamers tested (see SI Section S5), which is the same screw-sense induced in BQPA-capped Aib foldamers by Boc-D-Pro-O⁻. 9a,14,15,39 The kinetics of exchange in this complex are slower and more comparable to Boc-Pip with Zn $((S)-2)\cdot 2ClO_4$. For $Zn((S)-3)\cdot 2ClO_4$, interpolating the induced chemical shifts for Boc-D-Pro (-81 ppb; +76 ppb for Boc-L-Pro) into the calibration curve showed the h.e. $_0$ (32%) is greater than for $Zn((S)-2)\cdot 2ClO_4$ (21%, see Table 1).

Very similar behaviour was observed for the titration of Zn ((R)-4)-2ClO₄ and Zn((R)-5)-2ClO₄ with Boc-Pro (either D or L) under analogous experimental conditions (see SI Sections S4.5 and S4.6 respectively). Both showed fast exchange between bound and unbound foldamer and both gave h.e.₀ values of +29% after addition of 2 eq. Boc-L-Pro/2,6-lutidine (Table 1). As observed for Zn(1)-2ClO₄, ¹⁵ fast carboxylate exchange at zinc(II) on the NMR spectroscopy timescale was confirmed and the non-coordinating base 2,6-lutidine only bound weakly (see SI Fig. S35). The former was confirmed by using different scale-mic mixtures of Boc-Pro; δ_F (CF₃) correlated with the ee of the mixtures (see SI Fig. S37). Notably, addition of a 1:1 mixture of Boc-(D/L)-Pro to Zn((R)-4)-2ClO₄ gave δ_F (CF₃) at the same position, within the experimental error, as that of uncomplexed Zn((R)-4)-2ClO₄ (*i.e.* h.e.₀ = 0 in both cases).

As expected, given that it exists as two diastereomers, titration of unsymmetrical $Zn((R)-6)\cdot 2ClO_4$ with Boc-Pro (either D- or L-) gave complex data with multiple CF_3 resonances observed (see SI Section S4.7). Nonetheless these changes were broadly similar to those observed with symmetric $Zn((R)-4)\cdot 2ClO_4$.

Binding model and estimation of carboxylate affinity for Zn (2–5)·2ClO₄ in CD₃CN

The equilibria between different Boc-Pro/foldamer complexes were modelled. The sigmoidal profiles in the ¹H NMR and ¹⁹F NMR data during titrations with Boc-Pro or Boc-Pip are similar

Table 1 Calculated helical excess (h.e.₀) values induced by excess Boc-Pro/2,6-lutidine (\geq 2 eq.) for Zn(2–6)·2ClO₄. Values calculated from shifts of resonances ($\Delta\delta_{\rm F}({\rm CF_3})$) in the ¹⁹F NMR spectra (downfield for Zn ((S)-3)·2ClO₄, upfield for the others)

Foldamer	h.e. ₀ ^a
Zn((R)-2)·2ClO ₄	+21% (with Boc-L-Pro)
Zn((S)-3)·2ClO ₄	+32% (with Boc-L-Pro) ^b
Zn((R)-4)·2ClO ₄	+29% (with Boc-L-Pro)
Zn((R)-5)·2ClO ₄	+29% (with Boc-L-Pro)
Zn((R)-6)·2ClO ₄	+21% (with Boc-L-Pro)

^a h.e.₀ values are estimated from $\Delta \delta_F$ (CF₃) values according to a nonlinear model²⁵ (see SI Section S5). ^b The spectroscopic changes are reversed but Boc-1-Pro still induces *P* helix. to that observed during the TBA acetate titration of Zn((*R*)-2)·2ClO₄ and suggest that an intermediate is formed. This is presumed to be a 2:1 foldamer:ligand complex. We believe that because of the exchange kinetics when Boc-Pro or Boc-Pip (each with 2,6-lutidine) are used as titrants, this intermediate species is not observable by NMR spectroscopy for these anions. The structure of this intermediate 2:1 complex is unknown but we speculate that the carboxylate is bridging between zinc centres, which has been reported for similar complexes.⁴⁰ This may be a coordination mode permitted by the relatively open face of the BTPA binding pocket compared to the encumbered BQPA binding pocket of Zn(1)·2ClO₄. This bridged species may be disfavoured for the large bulky TiPSY anion.

In order to estimate the affinity of Boc-Pro for the Zn(BTPA)-capped foldamers, we used SupraFit and Dynafit to calculate the binding constants (Fig. 8b and 10, also see the SI). ^{41,42} We used a 2:1 foldamer: anion binding model for data fitting, with the binding constant K_{11} representing the for-




Fig. 10 Representative data fitting with Suprafit. For fits, see the SI Sections S4.5 and S4.6. (a) Global fit of ^1H protons of $\text{Zn}((R)\text{-4})\text{-2ClO}_4$ titrated with Boc-D-Pro. (b) Fits of the ^{19}F peak of $\text{Zn}((R)\text{-4})\text{-2ClO}_4$ titrated with Boc-D-Pro (purple) or Boc-L-Pro (red). (c) Fit of the triazole proton of $\text{Zn}((R)\text{-5})\text{-2ClO}_4$ (H_t) titrated with Boc-D-Pro. (d) Fits of the ^{19}F peak of $\text{Zn}((R)\text{-5})\text{-2ClO}_4$ titrated with Boc-D-Pro (purple) or Boc-L-Pro (red). Binding model: 2:1 [foldamer]/[anion] (see eqn (1) and (2) in SI Section S4.2). Conditions: $[\text{Zn}((R)\text{-4})\text{-2ClO}_4] = 2$ mM, [Boc-Pro] = 0-6.7 mM, [2,6-lutidine] = 0-7.2 mM. $[\text{Zn}((R)\text{-5})\text{-2ClO}_4] = 2$ mM, [Boc-Pro] = 0-3.6 mM, [2,6-lutidine] = 0-4.3 mM. ^{19}F NMR spectra referenced with C_6F_6 at -164.38 ppm. $^{31.33}$ R. err.: residual error (in ppm).

mation of the 1:1 complex and the binding constant K_{21} representing the formation of the intermediate 2:1 complex from the 1:1 complex and another equivalent of foldamer. Since for acetate binding, uncomplexed $\operatorname{Zn}((R)\text{-}4)\text{-}2\operatorname{ClO}_4$ and the 1:1 complex are in fast exchange with each other but the 2:1 complex is not, we assume the 2:1 complex should not affect the chemical shift of the 1:1 complex and fitting of the latter's chemical shift data should give an estimate of both binding constants. Nonetheless fitting the data to two equilibria presents challenges, with small uncertainties in concentration leading to significant differences in the individual binding constants K_{11} and K_{21} .

The titration of $Zn((S)-2)\cdot 2ClO_4$ with Boc-D-Pro gave 19 F NMR data (Fig. 8b) and 1 H data (see SI Fig. S16) that fitted a 2:1 binding model adequately using $K_{11} = 1 \times 10^5 \text{ M}^{-1}$ and $K_{21} = 2 \times 10^3 \text{ M}^{-1}$. The titration of foldamers $\text{Zn}((R)\text{-}4)\text{-}2\text{ClO}_4$ and Zn((R)-5)-2ClO₄ with both Boc-L-Pro and Boc-D-Pro could also be adequately fitted to a 2:1 binding model (Fig. 10). Within the uncertainty associated with fitting the formation of multiple complexes, there is reasonable agreement (within an order of magnitude) between the binding constants calculated for foldamers $Zn((R)-4)\cdot 2ClO_4$ and $Zn((R)-5)\cdot 2ClO_4$ (approximate values: $K_{11} = 1$ to 4×10^6 M⁻¹ and $K_{21} = 1$ to 2×10^4 M⁻¹); these are an order of magnitude higher than the analogous values for $Zn((S)-2)\cdot 2ClO_4$. These values for K_{11} (ca. ~10⁶ M⁻¹) are similar to K_{11} for the complexation of Boc-Pro to Zn (1)·2ClO₄ (4 × 10⁶ M⁻¹) in CD₃CN, although in that case there was no indication of an intermediate complex.³⁵ The K_{21} binding constants (ca. 10⁴ M⁻¹) are approximately 100-fold smaller than K_{11} , which is consistent with a steric barrier inhibiting the formation of this 2:1 complex. The similarity of the Boc-Pro binding constants for $Zn((R)-4)\cdot 2ClO_4$ and $Zn((R)-4)\cdot 2ClO_4$ 5)·2ClO₄ indicate that changing the oligoethyleneglycol for a crown ether has little influence on the binding of carboxylates. Estimated K values for $Zn((R)-4-6)\cdot 2ClO_4$ are summarised in Tables S1-S5 (see the SI).

Conclusions

The modular nature of CuAAC allows the construction of bis(triazole)pyridyl (BTPA) metal ion chelation sites that are flanked by different substituents, e.g. by a fluorophore or a crown ether bridge. Carboxylate and phosphate ligands both bind tightly to the Zn(BTPA) group, with the Zn(BTPA) structure allowing bound ligands to induce local conformational changes in the adjacent 3_{10} -helical Aib foldamer body. Corresponding increases in the proportion of either the P or the M screw-sense were detected by 19 F NMR spectroscopy using the recently developed 1-(trifluoromethyl)-ethylamido (TFEA) reporter group. 25

The strength of helical induction by different ligands was lower than that reported for the same ligands when bound to a previously described bis(quinolyl)pyridyl (BQPA)-capped foldamer. The greater hindrance created by the quinolyl arms compared to the more open face presented by the triazoles in Zn

(BTPA) complexes is proposed to lead to the better performance of BQPA. Lower steric encumbrance around the Zn(BTPA) site is also proposed to permit additional coordination equilibria, with new intermediates observed at sub-stoichiometric carboxylate that are not found with Zn(BQPA) foldamers. DOSY and titration data suggest these intermediates are (foldamer)2(carboxylate) complexes.

Nonetheless, the new BTPA binding site does not suffer from the drawbacks observed in previous replacements for BQPA.¹⁴ Despite showing similar coordination geometries to Zn(1)·2ClO₄, either replacing the quinolyl arms with pyridyl arms or replacing the pyridyl link with a triazolyl link led to foldamers with undesirable characteristics, including poor solubility, unfavourable ligand exchange rates on the NMR timescale, and inefficient relays of conformational information from the bound carboxylate (i.e. Boc-Pro). In contrast, all tested Zn(BTPA)-capped foldamers were soluble, had clear resonances at room temperature and possessed an effective conformational relay. These properties show that the modular BTPA motif is a versatile alternative to BQPA.

The versatility of CuAAC chemistry also permitted the creation of an unsymmetrical binding site on an Aib foldamer. The unsymmetrical binding site generated a chirogenic centre at the N-terminus upon Zn(II) complexation. The conformational preference of each handedness of the chirogenic Zn(II) complex is relayed along the foldamer body to the chiral CF₃-containing reporter group, leading to different ¹⁹F NMR spectroscopic outputs from each diastereomer. The net effect is to allow an achiral messenger (Zn(II)) to produce a spectroscopic output from the remote chiral TFEA reporter group.

With the Zn(BTPA) ligand binding site shown to mediate ligand-induced conformational change, the modular assembly of functional binding sites by CuAAC should lend itself to the generation of more complex constructs that better mimic natural binding sites in proteins. In this way ligand-triggered conformational change in larger and more functional foldamers will become possible.

Author contributions

S. J. W. conceived the idea, acquired the funding, administered the project. F. D. S., B. D. and S. J. W. designed the experiments, analysed the data and wrote the manuscript. F. D. S. and B. D. and carried out the experimental work. S. J. W. provided resources and supervision.

Conflicts of interest

There are no conflicts to declare.

Data availability

The data supporting this article have been included as part of the SI. See DOI: https://doi.org/10.1039/d5ob01226k.

Acknowledgements

This work was supported by the EPSRC (SJW and FdS, grants EP/P027067/1 and EP/K039547) and the BBSRC University of Manchester DTP2 (BD, grant BB/M011208/1, project 2282955). thank the University of Manchester Mass Spectrometry Service Centre for high-resolution spectrometry.

References

- 1 S. Negi, M. Imanishi, M. Hamori, Y. Kawahara-Nakagawa, W. Nomura, K. Kishi, N. Shibata and Y. Sugiura, J. Biol. Inorg. Chem., 2023, 28, 249-261, DOI: 10.1007/s00775-023-01991-6.
- 2 (a) T. K. Dam and C. F. Brewer, in Comprehensive Natural Products II: Chemistry and Biology, ed. L. Mander and H.-W. Liu, Elsevier, Oxford, 2010, vol. 3, ch. 3.21. DOI: DOI: 10.1016/B978-044451967-2/00058-1; (b) K. Drickamer, Curr. Opin. Struct. Biol., 1995, 5, 612-616, DOI: 10.1016/0959-440x(95)80052-2.
- 3 A. Bellelli and M. Brunori, Biochim. Biophys. Acta, Bioenerg., 2011, 1807, 1262-1272, DOI: 10.1016/j.bbabio.2011.04.004.
- 4 (a) J. L. Algar, J. A. Findlay and D. Preston, ACS Org. Inorg. Au, 2022, 2, 464-476, DOI: 10.1021/acsorginorgau.2c00021; (b) A. Galan, K. Lulic, J. Wang, B. Wicher, I. Huc, J. Duhamel and V. Maurizot, Chem. Commun., 2023, 59, 5253-5256, DOI: 10.1039/d3cc00866e.
- 5 S. R. Rao, S. L. Schettler and W. S. Horne, ChemPlusChem, 2021, 86, 137-145, DOI: 10.1002/cplu.202000730.
- 6 K. L. George and W. S. Horne, J. Am. Chem. Soc., 2017, 139, 7931-7938, DOI: 10.1021/jacs.7b03114.
- 7 C. M. Lombardo, V. M. V. Kumar, C. Douat, F. Rosu, J.-L. Mergny, G. F. Salgado and G. Guichard, J. Am. Chem. Soc., 2019, 141, 2516-2525, DOI: 10.1021/jacs.8b12240.
- 8 M. De Poli, W. Zawodny, O. Quinonero, M. Lorch, S. J. Webb and J. Clayden, Science, 2016, 352, 575-580, DOI: 10.1126/science.aad8352.
- 9 (a) F. G. A. Lister, B. A. F. Le Bailly, S. J. Webb and J. Clayden, Nat. Chem., 2017, 9, 420-425, DOI: 10.1038/ nchem.2736; (b) F. G. A. Lister, N. Eccles, S. J. Pike, R. A. Brown, G. F. S. Whitehead, J. Raftery, S. J. Webb and J. Clayden, Chem. Sci., 2018, 9, 6860-6870, DOI: 10.1039/ C8SC02532K.
- 10 R. Wechsel, M. Žabka, J. W. Ward and J. Clayden, J. Am. Soc., 2018, 140, 3528-3531, DOI: 10.1021/ Chem. jacs.8b00567.
- 11 K. Gratzer, V. Diemer and J. Clayden, Org. Biomol. Chem., 2017, 15, 3585-3589, DOI: 10.1039/C7OB00660H.
- 12 (a) J. Solà, G. A. Morris and J. Clayden, J. Am. Chem. Soc., 2011, 133, 3712-3715, DOI: 10.1021/ja1097034; (b) M. De Poli, L. Byrne, R. A. Brown, J. Solà, A. Castellanos, T. Boddaert, R. Wechsel, J. D. Beadle and J. Clayden, J. Org. Chem., 2014, 79, 4659-4675, DOI: 10.1021/jo500714b.

- 13 A. D. Peters, S. Borsley, F. della Sala, D. F. Cairns-Gibson, M. Leonidou, J. Clayden, G. F. S. Whitehead, I. J. Vitórica-Yrezábal, E. Takano, J. Burthem, S. L. Cockroft and S. J. Webb, *Chem. Sci.*, 2020, 11, 7023–7030, DOI: 10.1039/ d0sc02393k.
- 14 N. Eccles, F. della Sala, B. A. F. Le Bailly, G. F. S. Whitehead, J. Clayden and S. J. Webb, *ChemistryOpen*, 2020, 9, 338–345, DOI: 10.1002/open.201900362.
- 15 N. Eccles, B. A. F. Le Bailly, F. della Sala, I. J. Vitórica-Yrezábal, J. Clayden and S. J. Webb, *Chem. Commun.*, 2019, 55, 9331–9334, DOI: 10.1039/C9CC03895G.
- 16 L. You, G. Pescitelli, E. V. Anslyn and L. Di Bari, J. Am. Chem. Soc., 2012, 134, 7117–7125, DOI: 10.1021/ja301252h.
- 17 F. A. Scaramuzzo, G. Licini and C. Zonta, *Chem. Eur. J.*, 2013, **19**, 16809–16813, DOI: **10.1002/chem.201302721**.
- 18 D. Zhang, B. Bousquet, J.-C. Mulatier, D. Pitrat, M. Jean, N. Vanthuyne, L. Guy, J.-P. Dutasta and A. Martinez, *J. Org. Chem.*, 2017, 82, 6082–6088, DOI: 10.1021/acs.joc.7b00559.
- 19 N. A. C. dos Santos, E. Badetti, F. Begato, K. Wurst, G. Licini and C. Zonta, Eur. J. Inorg. Chem., 2021, 2021, 2942–2946, DOI: 10.1002/ejic.202100392.
- 20 M. M. Wootten, S. Tshepelevitsh, I. Leito and J. Clayden, Angew. Chem., Int. Ed., 2025, 62, e202502437, DOI: 10.1002/ anie.202502437.
- 21 M. M. Wootten, B. A. F. Le Bailly, S. Tshepelevitsh, I. Leito and J. Clayden, *Chem. Sci.*, 2022, **13**, 2258–2269, DOI: **10.1039/D1SC06812A**.
- 22 L. A. Joyce, M. S. Maynor, J. M. Dragna, G. M. Da Cruz, V. M. Lynch, J. W. Canary and E. V. Anslyn, *J. Am. Chem. Soc.*, 2011, 133, 13746–13752, DOI: 10.1021/ja205775g.
- 23 Y. H. Lau, P. J. Rutledge, M. Watkinson and M. H. Todd, Chem. Soc. Rev., 2011, 40, 2848–2866, DOI: 10.1039/ C0CS00143K.
- 24 (a) L. Fang, G. Trigiante, R. Crespo-Otero, C. S. Hawes, M. P. Philpott, C. R. Jones and M. Watkinson, *Chem. Sci.*, 2019, 10, 10881–10887, DOI: 10.1039/C9SC04300D;
 (b) M. Chernobryva, M. Motevalli, C. S. Hawes and M. Watkinson, *J. Mol. Struct.*, 2022, 1259, 132736, DOI: 10.1016/j.molstruc.2022.132736.
- S. Wang, F. della Sala, M. J. Cliff, G. F. S. Whitehead,
 I. J. Vitórica-Yrezábal and S. J. Webb, *J. Am. Chem. Soc.*,
 2022, 144, 21648–21657, DOI: 10.1021/JACS.2C09103.
- 26 B. Doerner, F. della Sala, S. Wang and S. J. Webb, Angew. Chem., Int. Ed., 2024, 63, e202405924, DOI: 10.1002/ anie.202405924.
- 27 N. Picois, L. Bodero, P. Milbeo, T. Brigaud and G. Chaume, *Chem. Eur. J.*, 2024, **30**, e202400540, DOI: **10.1002/ chem.202400540**.

- 28 S. J. Pike, M. De Poli, W. Zawodny, J. Raftery, S. J. Webb and J. Clayden, *Org. Biomol. Chem.*, 2013, 11, 3168–3176, DOI: 10.1039/C3OB40463C.
- 29 L. Bodero, K. Guitot, N. Lensen, O. Lequin, T. Brigaud, S. Ongeri and G. Chaume, *Chem. – Eur. J.*, 2022, 28, e202103887, DOI: 10.1002/chem.202103887.
- 30 H. Struthers, B. Spingler, T. L. Mindt and R. Schibli, Chem. – Eur. J., 2008, 14, 6173–6183, DOI: 10.1002/ chem.200702024.
- 31 For clarity, downfield shifts of resonances $(\Delta \delta)$ in NMR spectroscopy are positive while upfield shifts are negative.
- 32 K. S. Iyer, J.-P. Laussac and B. Sarkar, *Int. J. Pept. Protein Res.*, 1981, **18**, 468–477, DOI: **10.1111/j.1399-3011.1981**. **tb03008.**x.
- 33 C. P. Rosenau, B. J. Jelier, A. D. Gossert and A. Togni, Angew. Chem., Int. Ed., 2018, 57, 9528–9533, DOI: 10.1002/ anie.201802620.
- 34 J. Brioche, S. J. Pike, S. Tshepelevitsh, I. Leito, G. A. Morris, S. J. Webb and J. Clayden, *J. Am. Chem. Soc.*, 2015, 137, 6680–6691, DOI: 10.1021/jacs.5b03284.
- 35 B. A. F. Le Bailly, Controlling macromolecular conformation for the nanoscale transmission of information, PhD thesis, The University of Manchester, 2015.
- 36 E. Y. Tirel, Z. Bellamy, H. Adams, V. Lebrun, F. Duarte and N. H. Williams, *Angew. Chem., Int. Ed.*, 2014, 53, 8246–8250, DOI: 10.1002/anie.201400335.
- 37 T. Koike and E. Kimura, *J. Am. Chem. Soc.*, 1991, **113**, 8935–8941, DOI: **10.1021/ja00023a048**.
- 38 N. Kumar, Roopa, V. Bhalla and M. Kumar, *Coord. Chem. Rev.*, 2021, 427, 213550, DOI: 10.1016/j.ccr.2020.213550.
- 39 The use of a ¹³C reporter at the C-terminus shows that Boc-L-Pro gives a *P* helical preference in Zn(1)·2ClO₄. See ref. 35.
- 40 Carboxylates and other oxoligands have been shown to bridge between metal centres in tri(pyridylmethyl)amine complexes. See: (a) R. L. Rardin, W. B. Tolman and S. J. Lippard, New J. Chem., 1991, 15, 417-430; (b) C. M. Smith and R. E. Norman, Acta Crystallogr., Sect. E: Struct. Rep. Online, 2007, 63, m2480-m2481, DOI: 10.1107/S1600536807042857; (c) B. Chakraborty, I. Ghosh, R. D. Jana and T. K. Paine, Dalton Trans., 2020, 49, 3463-3472, DOI: 10.1039/C9DT04438H; (d) H. Adams, N. A. Bailey, D. E. Fenton and Q.-Y. He, J. Chem. Soc., Dalton Trans., 1997, 1997, 1533-1539, DOI: 10.1039/A608406K.
- 41 C. Hübler, *Chem. Methods*, 2022, 2, e202200006, DOI: 10.1002/CMTD.202200006.
- 42 P. Kuzmic, *Anal. Biochem.*, 1996, 237, 260–273, DOI: 10.1006/abio.1996.0238.