

Scheme 1 Summary of SAR studies (see the ESI†) for the aplyronines and opportunities for molecular editing, leading to the design of simplified aplyronine analogues 7–9 containing macrocyclic and side chain modifications. Our planned synthesis invokes a modular strategy using side chain *E*-11, aldehyde 12 and phosphonate 13 as the key fragments.

improve the step economy while retaining the biological activity.¹¹ Herein, we report the concise total synthesis of a series of simplified aplyronine analogues in 24–26 steps LLS (37–39 total steps) and their preliminary biological evaluation.


Inspection of the aplyronine A–actin complex crystal structure,⁵ and taking previous SAR studies³ into account, indicated that the C16–C22 region of the macrolactone did not appear to form any significant interaction with the protein binding site (see the ESI† for discussion). As outlined in Scheme 1, we proposed that this region was primarily acting as a conformational constraint for the macrocycle and that its replacement by a simple (CH₂)₇ unit should preserve the 3D structure and critical C1–C13 binding region, and hence maintain its biological activity. Additionally, our proposed

aplyrologues involved replacing the C25 hydroxyl and C31 acetoxy groups with methyl ethers, which simplifies the protecting group strategy, as well as removing the propensity for the acetate to eliminate (Scheme 1).⁹ Moreover, these latter modifications mirror the functionalities present in the equivalent side chains (Fig. 1) present in reidispongolide A (4), scytopycin C (5) and rhizopodin (6). We envisaged that these designed aplyrologues¹² would preserve the core pharmacophore, while significantly improving the synthetic tractability for this promising class of anticancer agent.


Building on our recent total synthesis of aplyronines A and D,⁹ we planned a late-stage diversification to construct aplyrologues bearing the aplyronine D side chain, as in 7 (Scheme 1), as well as the scytopycin–aplyronine A/D (bearing the C7 TMSer: 8) and C (9) analogues. This analysis leads back to the common scaffold 10, which would be forged through an aldol coupling sequence with the C28–C34 ketone *E*-11 and a macrocyclic aldehyde. The macrocycle itself can then be constructed by a Horner–Wadsworth–Emmons (HWE) olefination with the simplified C15–C27 aldehyde 12 and the previously reported C1–C14 phosphonate 13.^{9a}

Construction of the C15–C27 aldehyde 12 (Scheme 2) commenced with a Ti(OiPr)₃-mediated aldol reaction¹³ between the Roche ester-derived ketone *S*-13¹⁴ and the aldehyde 14 to give the 1,4-*syn*, 3,4-*syn* adduct 15. An Evans–Tishchenko reduction¹⁵ then delivered the 1,3-*anti* configured alcohol 16, which was methylated, desilylated and oxidised to afford the aldehyde 12 in five steps from 13 and 14. Notably, this expedient sequence represented more than a three-fold reduction in the step count relative to the analogous fragment required for the aplyronines.⁹

Synthesis of the modified C28–C34 side chain (Scheme 3) commenced with an analogous titanium aldol reaction between the ketone *R*-13 and acetaldehyde to give 17. Adduct 17 was reduced to give alcohol 18 followed by methylation, PMB ether cleavage and ester reduction to afford the diol 19. At this point, a double oxidation under Swern conditions gave ketoaldehyde 20, which was directly subjected to a Wittig olefination¹⁶ with phosphonium salt 21 to generate the enamides *E*/*Z*-11. Final completion of the configurationally pure C28–C34 ketone *E*-11 was accomplished by I₂-mediated isomerisation.




Scheme 2 Synthesis of the C15–C27 aldehyde 12. Reagents and conditions: (a) *S*-13, Ti(OiPr)₃, DIPEA, CH₂Cl₂, –78 °C; 14, CH₂Cl₂, –78 °C, 85%, 18 : 1 dr, (b) SmI₂, EtCHO, THF, –20 °C, 95%, >20 : 1 dr, (c) Me₃O·BF₄, Proton Sponge[®], CH₂Cl₂, r.t., 89%, 99% brsm, (d) HCl (3 M)/THF/H₂O (5 : 2 : 1), r.t., 99%, (e) (COCl)₂, DMSO, CH₂Cl₂, –78 °C; Et₃N, –78 °C to r.t., 94%.



Scheme 3 Synthesis of the C28–C34 ketone *E*-11. Reagents and conditions: (a) *R*-13, Ti(OiPr)₄, DIPEA, CH₂Cl₂, –78 °C; acetaldehyde, CH₂Cl₂, –78 °C, 77%, 17:1 dr, (b) SmI₂, EtCHO, THF, –20 °C, 97%, >20:1 dr, (c) Me₃O·BF₄, Proton Sponge[®], CH₂Cl₂, r.t., 90%, (d) DDQ, CH₂Cl₂/pH 7 buffer (2:1), r.t., 93%, (e) DIBAL, CH₂Cl₂, –78 °C, 87%, (f) (COCl)₂, DMSO, CH₂Cl₂, –78 °C; Et₃N, –78 °C to r.t., (g) **21**, LiHMDS, THF, –78 °C; **20**, THF –78 to –20 °C, 60% over two steps, 3:1 Z/E, (h) I₂ (5 mol%), CH₂Cl₂, r.t., 86%, >20:1 E/Z.

With all three fragments in hand, attention now turned towards the assembly of the full carbon skeleton of the targeted analogues. Aldehyde **12** was first coupled with the phosphonate **13** (10 steps)^{9a} via an HWE olefination¹⁷ to generate the C1–C27 fragment **22** (Scheme 4). A CBS reduction¹⁸ of the C13 ketone was then followed by methylation to give **23**. Next, the C23 ester was cleaved and a DDQ-mediated deprotection/allylic oxidation¹⁹ afforded the *seco* acid **24**. The macrolactonisation reaction of **24** proceeded smoothly under Yamaguchi conditions²⁰ to give **25**. Moving forward, oxidative cleavage of the PMB ether in **25** followed by Dess–Martin oxidation of the resulting alcohol afforded the aldehyde **26**. This was then subjected to a three-step sequence involving a Cy₂BCL-mediated aldol coupling^{9b} with the C28–C34 ketone *E*-11, a Burgess elimination followed by a copper-catalysed conjugate reduction of the resulting enone gave **10**, corresponding to a protected aplyrologue bearing the scytophycin/reidispongolide-type side chain. Additionally, the C29 ketone in **10** could be reduced and acylated with DMGly under Keck conditions²¹ to give the 1,3-diol **27** after deprotection, which corresponded to the aplyrologue scaffold bearing the aplyronine-type side chain. The advanced intermediate **10** was also deprotected to give the scytophycin–aplyronine C hybrid **9**. In turn, this could be site-selectively acylated at C7 with the required enantiomer of TMSer to generate the scytophycin–aplyronine A hybrids *S*-**8** and *R*-**8**. Noting that aplyronine D bearing the C7 TMSer and C29 DMGly residues was reported to be the most active congener,^{1b} the corresponding aplyrologue D was targeted. Thus, the diol **27** was site-selectively acylated with *S*-TMSer to generate **7**.

With this first series of aplyrologues in hand, attention now turned towards determining their *in vitro* cytotoxicity. It was considered prudent to initially examine the activity of our previously synthesised samples of aplyronines A (**1**), C (**2**) and D (**3**), with discodermolide included as a control (Table 1).



Scheme 4 Synthesis of aplyrologues. Reagents and conditions: (a) Ba(OH)₂, THF/H₂O (40:1), r.t., 85%, >20:1 E/Z, (b) *R*-MeCBS, BH₃·SMe₂, –10 °C, 93%, >20:1 dr, (c) Me₃O·BF₄, Proton Sponge[®], CH₂Cl₂, r.t., 87%, (d) DIBAL, CH₂Cl₂, –78 °C, 99%, (e) DDQ, CH₂Cl₂/pH 9.2 buffer (4:1), 0 °C to r.t., 75%, 91% brsm, (f) NaClO₂, NaH₂PO₄·H₂O, 2-methyl-but-2-ene, H₂O/*t*BuOH, 0 °C to r.t., (g) TCBC, Et₃N; DMAP, PhMe, r.t., 70% over two steps, (h) DDQ, CH₂Cl₂/pH 7.0 buffer (2:1), r.t., 82%, (i) DMP, NaHCO₃, CH₂Cl₂, r.t., (j) *E*-11, Cy₂BCL, Et₃N, Et₂O, 0 °C; **26**, Et₂O, –78 to –10 °C, (k) Burgess reagent, THF, r.t., 65% over three steps, (l) Cu(OAc)₂, PPh₃, TMDS, PhMe, r.t., 89%, (m) HF·py/py (1:2), THF, 0 °C to r.t., 85%, (n) TCBC, *R*- or *S*-TMSer, Et₃N, DMAP, PhH, 62% for *S*-**8**, 73% for *R*-**8**, (o) Zn(BH₄), Et₂O, 0 °C, 58%, 2:1 dr, (p) DMGly, DCC, DMAP, DMAP·HCl, r.t., (q) HF·py/py (1:2), THF, 0 °C to r.t., 69% over two steps, (r) TCBC, *S*-TMSer, Et₃N, DMAP, PhH, 65%. CBS: Corey–Bakshi–Shibata catalyst.

While the expected sub-nanomolar potencies were found for aplyronines A and D, we did not find any significant difference between these congeners in the same HeLa cell line (Table 1, entries 1–3). However, our synthetic aplyrologues did not register the same exquisite potency compared with the native aplyronines. Nevertheless, the compounds *R*- and *S*-**8** (entries 5 and 6), bearing the scytophycin side chain, as well as **7** (entry 4) still gave inhibitory activities in the nM range. Interestingly, both the configuration of the TMSer residue and the nature of the C29 functionality as the ketone or DMGly ester did not appear to greatly influence the biological activity, suggesting that the deleterious effect may be attributable to an oversimplification of the southern hemisphere region of the macrolactone.

Table 1 Antiproliferative activity of synthetic aplyronines and aplyrologues against the HeLa cell line (discodermolide was included as a control)

Entry	Compound	IC ₅₀ ^a (nM)
1	Aplyronine A (1)	0.31
2	Aplyronine C (2)	3.67
3	Aplyronine D (3)	0.26
4	7 (aplyrologue D)	124
5	S-8 (aplyronine A, D/scytophycin C hybrid)	164
6	R-8 (aplyronine A, D/scytophycin C hybrid)	192
7	9 (aplyronine C/scytophycin C hybrid)	>1000
8	Discodermolide	3.40

^a *n* = 3. See the ESI for cytotoxicity curves and further experimental details.

In conclusion, the preliminary biological results obtained for this first series of aplyrologues suggests that the southern hemisphere of the macrocycle may play a subtle but vital role in reinforcing the proposed actin–tubulin interaction that the aplyronines uniquely mediate. Additionally, the biological data obtained for our synthetic aplyronines does not replicate the greater cytotoxicity previously reported for a natural sample of aplyronine D relative to aplyronine A,^{1b} signifying the scope for structural diversification of the side chain. In future work, further judicious molecular editing of the southern hemisphere region can be flexibly adapted towards the incorporation of more elaborate macrocyclic tethers. We envisage that our modular synthetic platform offers a means for enhancing the synthetic tractability of the aplyronines to facilitate the continued development of this promising class of anticancer agent.²²

We thank Trinity College (TRP), AstraZeneca (RJP and SJW), EPSRC (RJP and SJW), Downing College (MPH) and the Woolf Fisher Trust (NYSL) for financial support, Dr Nika Anžiček and Dr Simon Williams for helpful discussions, and the National Mass Spectrometry Centre (Swansea) for mass spectra.

Conflicts of interest

There are no conflicts to declare.

Notes and references

1 (a) K. Yamada, M. Ojika, T. Ishigaki, Y. Yoshida, H. Ekimoto and M. Arakawa, *J. Am. Chem. Soc.*, 1993, **115**, 11020; (b) M. Ojika,

- H. Kigoshi, K. Suenaga, Y. Imamura, K. Yoshikawa, T. Ishigaki, A. Sakakura, T. Mutou and K. Yamada, *Tetrahedron*, 2012, **68**, 982.
 2 (a) M. Ojika, H. Kigoshi, T. Ishigaki, I. Tsukada, T. Tsuboi, T. Ogawa and K. Yamada, *J. Am. Chem. Soc.*, 1994, **116**, 7441; (b) H. Kigoshi, M. Ojika, T. Ishigaki, K. Suenaga, T. Mutou, A. Sakakura, T. Ogawa and K. Yamada, *J. Am. Chem. Soc.*, 1994, **116**, 7443.
 3 (a) H. Kigoshi, K. Suenaga, M. Takagi, A. Akao, K. Kanematsu, N. Kamei, Y. Okugawa and K. Yamada, *Tetrahedron*, 2002, **58**, 1075; (b) M. Ojika, H. Kigoshi, Y. Yoshida, T. Ishigaki, M. Nisiwaki, I. Tsukada, M. Arakawa, H. Ekimoto and K. Yamada, *Tetrahedron*, 2007, **63**, 3138.
 4 K. Hirata, S. Muraoka, K. Suenaga, T. Kuroda, K. Kato, H. Tanaka, M. Yamamoto, M. Takata, K. Yamada and H. Kigoshi, *J. Mol. Biol.*, 2006, **356**, 945.
 5 (a) K.-S. Yeung and I. Paterson, *Angew. Chem., Int. Ed.*, 2002, **41**, 4632; (b) J. S. Allingham, V. A. Klenchin and I. Rayment, *Cell. Mol. Life Sci.*, 2006, **63**, 2119.
 6 M. Kita, Y. Hirayama, K. Yoneda, K. Yamagishi, T. Chinen, T. Usui, E. Sumiya, M. Uesugi and H. Kigoshi, *J. Am. Chem. Soc.*, 2013, **135**, 18089.
 7 M. Kita and H. Kigoshi, *Nat. Prod. Rep.*, 2015, **32**, 534.
 8 (a) R. V. J. Chari, M. L. Miller and W. C. Widdison, *Angew. Chem., Int. Ed.*, 2014, **53**, 3796; (b) K. C. Nicolaou and S. Rigol, *Angew. Chem., Int. Ed.*, 2019, **58**, 11206; (c) D. J. Newman, *Mar. Drugs*, 2019, **17**, 324.
 9 (a) N. Anžiček, S. Williams, M. P. Housden and I. Paterson, *Org. Biomol. Chem.*, 2018, **16**, 1343; (b) I. Paterson, S. J. Fink, L. Y. W. Lee, S. J. Atkinson and S. B. Blakey, *Org. Lett.*, 2013, **15**, 3118.
 10 I. Hayakawa, K. Saito, S. Matsumoto, S. Kobayashi, A. Taniguchi, K. Kobayashi, Y. Fujii, T. Kaneko and H. Kigoshi, *Org. Biomol. Chem.*, 2017, **15**, 124.
 11 P. A. Wender, R. V. Quiroz and M. C. Stevens, *Acc. Chem. Res.*, 2015, **48**, 752.
 12 (a) K. Futaki, M. Takahashi, K. Tanabe, A. Fujiedo, H. Kigoshi and M. Kita, *ACS Omega*, 2019, **4**, 8598; (b) T. Ohyoshi, A. Takano, M. Namiki, T. Ogura, Y. Miyazaki, Y. Ebihara, K. Takeno, I. Hawakawa and H. Kigoshi, *Chem. Commun.*, 2018, **54**, 9537.
 13 J. G. Solsona, J. Nebot, P. Romea and F. Urpi, *J. Org. Chem.*, 2005, **70**, 6533.
 14 I. Paterson, G. J. Florence, K. Gerlach, J. P. Scott and N. Sereinig, *J. Am. Chem. Soc.*, 2001, **123**, 9535.
 15 D. A. Evans and A. H. Hoveyda, *J. Am. Chem. Soc.*, 1990, **112**, 6447.
 16 I. Paterson, C. J. Cowden, V. S. Rahn and M. D. Woodrow, *Synlett*, 1998, 915.
 17 I. Paterson, K.-S. Yeung and J. B. Smaill, *Synlett*, 1993, 774.
 18 E. J. Corey, R. K. Bakshi, S. Shibata, C. P. Chen and V. K. Singh, *J. Am. Chem. Soc.*, 1987, **109**, 7925.
 19 I. Paterson, C. J. Cowden, V. S. Rahn and M. D. Woodrow, *Synlett*, 1998, 915.
 20 J. Inanaga, K. Hirata, H. Saeki, T. Katsuki and M. Yamaguchi, *Bull. Chem. Soc. Jpn.*, 1979, **52**, 1989.
 21 E. P. Boden and G. E. Keck, *J. Org. Chem.*, 1985, **50**, 2394.
 22 For other recent work from our group, see: (a) I. Paterson and N. Y. S. Lam, *J. Antibiot.*, 2017, **71**, 215; (b) N. Y. S. Lam and I. Paterson, *Eur. J. Org. Chem.*, DOI: 10.1002/ejoc.201901243; (c) N. Y. S. Lam, G. Muir, V. R. Challa, R. Britton and I. Paterson, *Chem. Commun.*, 2019, **55**, 9717; (d) M. Anketell, T. Sharrock and I. Paterson, *Angew. Chem., Int. Ed.*, DOI: 10.1002/acie.201914042.