

# Chemical Science

Volume 15  
Number 27  
21 July 2024  
Pages 10251-10672

rsc.li/chemical-science



ISSN 2041-6539

**EDGE ARTICLE**

Marcus W. Drover *et al.*  
Oxidatively-induced C(sp<sup>3</sup>)-C(sp<sup>3</sup>) bond formation  
at a tucked-in iron(III) complex

Cite this: *Chem. Sci.*, 2024, 15, 10359

All publication charges for this article have been paid for by the Royal Society of Chemistry

# Oxidatively-induced C(sp<sup>3</sup>)-C(sp<sup>3</sup>) bond formation at a tucked-in iron(III) complex†

Joseph A. Zurakowski,<sup>ab</sup> Connor S. Durfy,<sup>a</sup> Noah B. Stocck,<sup>c</sup> Giovanni Fanchini<sup>ac</sup> and Marcus W. Drover<sup>\*,a</sup>

Carbon-carbon (C-C) bond formation is a cornerstone of synthetic chemistry, relying on routes such as transition-metal mediated cross-coupling for the introduction of new carbon-based functionality. For  $\{[M]^{n+}-C\}$  ( $M = \text{metal}$ ) structural units, studies that offer well-defined relationships between metal oxidation state, hydrocarbon strain, and  $\{[M]^{n+}-C\}$  bond thermochemistry are thus informative, providing a means to reliably access new product classes. Here, we show that one-electron oxidation of the iron tucked-in complex  $[(\eta^6\text{-C}_5\text{Me}_4=\text{CH}_2)\text{Fe}(\text{dnppe})]$  ( $\text{dnppe} = 1,2\text{-bis}(\text{di-}n\text{-propylphosphino})\text{ethane}$ ) results in C(sp<sup>3</sup>)-C(sp<sup>3</sup>) bond formation giving unique  $\{\text{Fe}_2\}$  dimers. Freeze-quenched CW X-band EPR spectroscopy allowed for spectroscopic identification of the reactive  $[(\eta^6\text{-C}_5\text{Me}_4=\text{CH}_2)\text{Fe}(\text{dnppe})]^+$  intermediate. Density functional theory (DFT) calculations reveal a primarily Fe-centered radical and a weak  $\{[\text{Fe}]-C\}$  bond ( $\text{BDE}_{[\text{Fe}]-C} = 24.5 \text{ kcal mol}^{-1}$ , *c.f.*  $\text{BDE}_{C-C(\text{ethane})} = 90 \text{ kcal mol}^{-1}$ ). For comparison, a structurally analogous Fe(III) methyl complex was prepared,  $[\text{Cp}^*\text{Fe}(\text{dnppe})(\text{CH}_3)]^+$  ( $\text{Cp}^* = \text{C}_5\text{Me}_5^-$ ), where C(sp<sup>3</sup>)-C(sp<sup>3</sup>) coupling was not observed, consistent with a larger calculated  $\text{BDE}_{[\text{Fe}]-C}$  value of  $47.8 \text{ kcal mol}^{-1}$ . These data are analogized to the simple hydrocarbons ethane and cyclopropane, where a strain-induced  $\text{BDE}_{C-C}$  decrease of  $33 \text{ kcal mol}^{-1}$  is witnessed on cyclization.

Received 20th May 2024

Accepted 6th June 2024

DOI: 10.1039/d4sc03292f

rsc.li/chemical-science

## Introduction

Carbon-carbon (C-C) bond forming reactions are a cornerstone of synthetic chemistry. Most pharmaceutically active compounds contain a C-C bond, along with countless materials relevant to industry and everyday life. It thus comes as no surprise that transition metal mediated C-C bond forming reactions have been the focus of two Nobel prizes.<sup>1,2</sup> Using transition metals, one can form C-C bonds through cross-coupling – a sequence that proceeds *via* stepwise oxidative addition, transmetalation, and reductive elimination.<sup>3</sup> In recent years, there has been a surge of reports that have driven such transformations using electro-<sup>4-7</sup> or photochemistry,<sup>8-11</sup> many of these studies have focused on cheaper and more abundant base transition elements as feedstocks.<sup>12</sup>

Reductive elimination is the main step associated with C-C bond formation.<sup>13</sup> When reductive elimination from  $\{[M]^{n+}-C\}$

is unfavourable and prohibits C-C bond generation, oxidation of the metal-center can be used to induce so called oxidatively-induced reductive elimination (ORE; Fig. 1A).<sup>14-16</sup> In this way, oxidation increases the tendency to undergo elimination (*e.g.*, homolytic bond cleavage), thus returning the metal to a stable d-electron count.<sup>17,18</sup>

Sanford and co-workers have shown that one-electron oxidation of  $[(^t\text{Bu}_2\text{bpy})\text{Pd}^{\text{III}}(\text{CH}_3)_2]^+$  ( $^t\text{Bu}_2\text{bpy} = 4,4'\text{-di-}t\text{-tert-butyl-2,2'-bipyridine}$ ) results in C(sp<sup>3</sup>)-C(sp<sup>3</sup>) reductive elimination of ethane ( $\text{CH}_3\text{CH}_3$ ) from an intermediate  $\{[\text{Pd}^{\text{IV}}](\text{CH}_3)_3\}$  species (Fig. 1A).<sup>19</sup> Similarly, Baik, Chang, and co-workers showed that oxidation of an Ir(III) complex to Ir(IV) led to a  $19 \text{ kcal mol}^{-1}$  reduction in the barrier associated with reductive elimination in a C-H arylation reaction.<sup>20</sup> These examples illustrate the power of ORE in C-C bond formation, providing opportunities for otherwise unfavourable bond construction.

Of  $\{[M]^{n+}-C\}$  organometallics, sandwich complexes  $\{[\text{Cp}/\text{Cp}^*]_2\text{M}\}$ ;  $\text{Cp} = \text{C}_5\text{H}_5^-$ ;  $\text{Cp}^* = \text{C}_5\text{Me}_5^-$ ) represent a well-studied area owing to robust syntheses and product stability.<sup>21,22</sup> Lesser studied, however, are related “tucked-in” complexes, where a  $\text{H}^+/\text{H}'/\text{H}^-$  from the  $\text{Cp}^*-\text{CH}_3$  ligand has been removed, encouraging high reactivity.<sup>23</sup> This is especially true for first row late transition elements. As a salient example, it was reported that the ferrocenyl carbocation,  $[(\eta^6\text{-C}_5\text{Me}_4)=\text{CH}_2]\text{Fe}(\text{Cp}^*)^+$  is reactive at temperatures  $> -82 \text{ }^\circ\text{C}$ .<sup>24-26</sup>

<sup>a</sup>Department of Chemistry, Western University, 1151 Richmond Street, London, ON, N8K 3G6, Canada. E-mail: marcus.drover@uwo.ca


<sup>b</sup>Department of Chemistry and Biochemistry, University of Windsor, 401 Sunset Avenue, Windsor, ON, N9B 3P4, Canada

<sup>c</sup>Department of Physics and Astronomy, Western University, 1151 Richmond Street, London, ON, N6A 3K7, Canada

† Electronic supplementary information (ESI) available: Experimental procedures, spectroscopic data, computational details. CCDC 2341812. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4sc03292f>




## A. Concept: oxidatively-induced reductive elimination (r.e.)



- facile r.e. via a high-valent metal
- intermolecular or intramolecular r.e. possible

e.g.,



Sanford, 2009

## B. Reactivity and thermodynamics of a strained ring system




Fig. 1 (A) Overview of intra/intermolecular oxidatively induced reductive elimination with transition metals. (B) Decrease in  $BDE_{\text{C-C}}$  in cyclopropane upon oxidation, and this work.

Recently, we reported the first example of an Fe diphosphine tucked-in complex,  $[(\eta^6\text{-C}_5\text{Me}_4=\text{CH}_2)\text{Fe}(\text{dnppe})]$  ( $\text{dnppe} = 1,2$ -bis(di-*n*-propylphosphino)ethane) (**1**) (Fig. 1B) and its reactivity with electrophiles including  $\text{CO}_2$ ,  $\text{HBCy}_2$  ( $\text{Cy} = \text{cyclohexyl}$ ),  $\text{Br-Au}^1\text{-PPh}_3$ , benzaldehyde, and  $\text{B}(\text{C}_6\text{F}_5)_3$ .<sup>27</sup> Herein, we examine the redox behaviour of **1** using electrochemical, photochemical, and chemical means (Fig. 2A). Oxidation of the Fe-center in **1** led to rapid  $\text{C}(\text{sp}^3)\text{-C}(\text{sp}^3)$  formation between  $\text{CH}_2$  groups of the  $\eta^6\text{-C}_5\text{Me}_4=\text{CH}_2$  ligand. Model studies and theoretical calculations were used to rationalize iron-carbon  $\{[\text{Fe}]^{n+}\text{-C}\}$  bond strength (*i.e.*, bond dissociation energy (BDE)) as a function of both iron oxidation state and ring-strain. These observations are analogized to cyclic organic compounds (*e.g.*, cyclopropane) that also bear weak C-C bonds due to strain – a characteristic that has been leveraged for oxidatively-induced functionalization.<sup>28,29</sup>

## Results and discussion

An investigation into the oxidative chemistry of **1** was commenced by means of cyclic voltammetry (CV). Scanning from  $-1.90$  V to  $+0.50$  V in THF showed a quasi-reversible feature at  $E_{1/2} = -1.23$  V ( $i_p^{\text{red}}/i_p^{\text{ox}} = 0.89$  at  $200$   $\text{mV s}^{-1}$ ; Fig. S43†)<sup>30</sup> and an irreversible feature at  $E_{\text{p,a}} = +0.15$  V vs.  $\text{Fc}/\text{Fc}^+$  ( $\text{Fc} = \text{ferrocene}$ ) (Fig. 2B); these are assigned as  $\text{Fe}(\text{II}/\text{III})$  and  $\text{Fe}(\text{III}/\text{IV})$  events, respectively.<sup>31</sup> Varying the scan rate from  $50$  to  $500$   $\text{mV s}^{-1}$  did not change voltammogram shape (Fig. S44†). For other known  $\text{Fe}(\text{II})$  half-sandwich complexes,  $[\text{Cp}^*\text{Fe}^{\text{II}}(\text{dppe})(\text{X})]$  ( $\text{dppe} = 1,2$ -bis(diphenylphosphino)ethane;  $\text{X} = \text{H}, \text{CH}_3, \text{F}, \text{Cl}, \text{Br}, \text{I}$ ), the  $\text{Fe}(\text{II}/\text{III})$  couple was found to be  $-0.54 > E_{1/2} >$



A. Synthesis and diversification of  $\text{C}(\text{sp}^3)\text{-C}(\text{sp}^3)$  coupled productsB. Cyclic voltammetry studies on **1** and **2**

Fig. 2 (A) Synthesis of  $[\text{1}]^+$  and  $\text{C}(\text{sp}^3)\text{-C}(\text{sp}^3)$  coupled products. (B) Cyclic voltammograms for **1** and **2**.

$-0.88$  V, indicating the  $E_{1/2}$  value for **1** to be comparably cathodically shifted.<sup>32,33</sup> To probe the effect of hydrocarbon strain on electrochemical oxidation, an analogous  $\{[\text{Fe}^{\text{II}}]\text{-CH}_3\}$  complex  $[\text{Cp}^*\text{Fe}^{\text{II}}(\text{dnppe})(\text{CH}_3)]$  (**2**) (Fig. 2B) was examined.<sup>34</sup> Compound **2** showed a reversible feature at  $E_{1/2} = -1.16$  V with  $i_p^{\text{red}}/i_p^{\text{ox}}$  near  $1.00$  at  $200$   $\text{mV s}^{-1}$  (Fig. S45†), and an irreversible feature at  $E_{\text{p,a}} = +0.35$  V vs.  $\text{Fc}/\text{Fc}^+$  in THF, indicating that the tucked-in  $\eta^6\text{-C}_5\text{Me}_4=\text{CH}_2$  ligand contributes to a  $70$  mV cathodic shift in the  $\text{Fe}(\text{II}/\text{III})$  couple. For both **1** and **2**, the  $\text{Fe}(\text{III}/\text{IV})$  couple was irreversible – a reported outcome for  $[\text{Cp}^*\text{Fe}(\text{diphosphine})(\text{CH}_3)]^+$ -type complexes.<sup>32</sup>

As a result of our observations from the CV data, complex **1** was next exposed to **1** equiv. of  $[\text{Cp}_2\text{Fe}]\text{B}(\text{C}_6\text{F}_5)_4$  at  $25$  °C (Fig. 2A)



in CH<sub>3</sub>CN. Monitoring this reaction by <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy showed the formation of diamagnetic products in the range of *ca.* δ<sub>P</sub> = 78 ppm, consistent with complexes of the type [Cp\*Fe(diphosphine)(NCCH<sub>3</sub>)]<sup>+</sup>.<sup>33</sup> This hypothesis was further supported by FT-IR spectroscopy, providing a band at ν(C≡N) = 2252 cm<sup>-1</sup> (*c.f.*, ν(C≡N) = 2234 cm<sup>-1</sup>) for [Cp\*Fe(dnppe)(NCCH<sub>3</sub>)]<sup>+</sup>, prepared independently (*vide infra*; [7]<sup>+</sup>) (Fig. S28†).

Supposing the main product from oxidation of **1** to be the C(sp<sup>3</sup>)-C(sp<sup>3</sup>) coupled {μ-H<sub>2</sub>CCH<sub>2</sub><sup>-</sup>} dimer, [{η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>-(μ-CH<sub>2</sub>)}Fe<sup>II</sup>(dnppe)(NCCH<sub>3</sub>)<sub>2</sub>]<sup>2+</sup> ([5]<sup>2+</sup>), the redox behaviour of **1** was probed in the presence of 4 mL of THF with 100 μL CH<sub>3</sub>CN added.<sup>35</sup> Consistent with an electrochemical-chemical (EC) reaction (onwards reactivity), scanning oxidatively from -2.00 to +1.00 V showed the previously discussed quasi-reversible couple at E<sub>1/2</sub> = -1.23 V, with a decreased value of *i*<sub>p</sub><sup>red</sup>/*i*<sub>p</sub><sup>ox</sup> = 0.62 (*c.f.*, 0.89 in THF at 200 mV s<sup>-1</sup>). In addition to the observed signature for **1**, an additional wave at E<sub>1/2</sub> = -0.16 V was assigned to the Fe(II/III) couple of [5]<sup>2+</sup>, confirmed through independent preparation (*vide infra*; Fig. S51†).

Outlining an alternative preparation of [5]<sup>2+</sup>, we envisaged treatment of **1** with a halide-based oxidant to initiate C(sp<sup>3</sup>)-C(sp<sup>3</sup>) coupling, followed by halide abstraction in CH<sub>3</sub>CN (Fig. 2A). Diiodoethane (ICH<sub>2</sub>CH<sub>2</sub>I), previously shown to oxidize {Fe(diphosphine)<sub>x</sub>} complexes,<sup>36</sup> was selected as the candidate oxidant, supplying iodide groups that could later be removed. In accord, exposure of **1** to 1 equiv. of ICH<sub>2</sub>CH<sub>2</sub>I showed an immediate colour change from orange to dark red (Fig. 2A). Analysis by <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy, however, showed an absence of signals, suggestive of paramagnetism. Growth of crystals from a THF solution layered with pentane at -35 °C overnight provided material appropriate for an Evans method measurement, giving μ<sub>eff</sub> = 2.66 μ<sub>B</sub> (298 K, CDCl<sub>3</sub>), falling within the expected range of a molecule containing two unpaired electrons. Further analysis by single-crystal X-ray diffraction (scXRD) revealed a connectivity map that supported a dimeric structure of [{η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>-(μ-CH<sub>2</sub>)}Fe<sup>III</sup>(dnppe)(I)]<sub>2</sub>(I)<sub>2</sub> ([3](I)<sub>2</sub>) (Fig. S54†). This molecule features a bridging {μ-H<sub>2</sub>CCH<sub>2</sub><sup>-</sup>} unit that links two {(C<sub>5</sub>Me<sub>4</sub>-(CH<sub>2</sub>))Fe<sup>III</sup>(dnppe)(I)} moieties. Consistent with the μ<sub>eff</sub> value obtained from Evans method and the two exogenous iodide counter-anions observed by scXRD, both Fe-centers are formally Fe(III). High-resolution electrospray ionization mass-spectrometry (HRMS-ESI) additionally provides a signal for [3]<sup>+</sup> at *m/z* = 1158.299 (calcd. *m/z* = 1158.293).

Formation of [3](I)<sub>2</sub> is a four-electron process involving two successive oxidations of **1**. Considering the Fe(II) resonance contributor for **1** (Fig. 1B), oxidation leads to homolytic bond cleavage, forming an *S* = 1 Fe(II) center<sup>37</sup> and a {C<sub>5</sub>Me<sub>4</sub>-(·CH<sub>2</sub>)} radical (*vide infra*). Recombination of two {C<sub>5</sub>Me<sub>4</sub>-(·CH<sub>2</sub>)} radicals results in C(sp<sup>3</sup>)-C(sp<sup>3</sup>) bond formation. Iodide coordination and subsequent oxidation to Fe(III) (by excess ICH<sub>2</sub>CH<sub>2</sub>I) thus provides the observed product [3](I)<sub>2</sub> (Fig. 2A). To provide the Fe(II) complex [{η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>-(μ-CH<sub>2</sub>)}Fe<sup>II</sup>(dnppe)(I)]<sub>2</sub> (**4**), 0.5 equiv. of ICH<sub>2</sub>CH<sub>2</sub>I was added to **1** (to preclude oxidation from Fe(II) to Fe(III)). However, this reaction resulted in mixtures of products, with at least two <sup>31</sup>P{<sup>1</sup>H} resonances appearing in

the range of **4** (δ<sub>P</sub> = *ca.* +80 ppm), unreacted **1** (δ<sub>P</sub> = +89.6 ppm), and free dnppe (δ<sub>P</sub> = -28.3 ppm) (Fig. S30†).

As an alternative route towards **4**, reduction of [3](I)<sub>2</sub> was explored using cobaltocene ([Cp<sub>2</sub>Co]). Addition of 2 equiv. of [Cp<sub>2</sub>Co] to a THF solution of [3](I)<sub>2</sub> produced an insoluble yellow precipitate and a purple supernatant over 3 h. Extraction of the purple product into toluene and analysis by NMR spectroscopy revealed a diamagnetic product. By <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy, one signal at δ<sub>P</sub> = +78.5 ppm was observed, standard of [Cp\*Fe<sup>II</sup>(dnppe)(halide)]-type compounds (*e.g.*, for halide = Cl, δ<sub>P</sub> = +79.4 ppm).<sup>34</sup> Analysis of the <sup>1</sup>H NMR spectrum also revealed a characteristic signal for the bridging {μ-H<sub>2</sub>CCH<sub>2</sub><sup>-</sup>} moiety (δ<sub>H</sub> = 2.57 ppm) as well as two pairs of signals for the unsymmetric methyl {η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>} groups (δ<sub>H</sub> = 1.89 and 1.82 ppm). Crystals of **4** were grown from slow evaporation of a benzene solution overnight; analysis by scXRD confirmed its structural identity as [{η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>-(μ-CH<sub>2</sub>)}Fe<sup>II</sup>(dnppe)(I)]<sub>2</sub> (**4**) (Fig. 3). Offering context, a Cambridge Structural Database (CSD) search provides fewer than 5 examples of structurally-authenticated Cp\*-based compounds featuring a bridging {μ-H<sub>2</sub>CCH<sub>2</sub><sup>-</sup>} unit.<sup>24,38-43</sup> Using pre-formed ligand constructs, Wang and co-workers reported a set of polymeric Ir complexes bridged by {C<sub>5</sub>Me<sub>4</sub>-(CH<sub>2</sub>)<sub>*n*</sub>-C<sub>5</sub>Me<sub>4</sub>} (*n* = 2, 3, 4),<sup>40</sup> while Brintzinger and co-workers synthesized monometallic complexes of the type [(C<sub>5</sub>Me<sub>4</sub>-(CH<sub>2</sub>)<sub>2</sub>-C<sub>5</sub>Me<sub>4</sub>)MCl<sub>2</sub>] (M = Ti, Zr).<sup>41</sup> Another example has been accessed by Cp\*{H<sup>+</sup>} removal – Severin and co-workers utilized [OAl(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>]<sup>-</sup> in the presence of Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> to perform H-atom abstraction of [Cp\*Fe], generating a [(η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>)(·CH<sub>2</sub>)Fe(Cp\*)] radical,<sup>38</sup> which dimerized to give [{η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>-(μ-CH<sub>2</sub>)}Fe<sup>II</sup>(Cp\*)]<sub>2</sub>. This example, however, does not require Fe oxidation nor does it inform on the relationship between iron-ring-strain and C-C coupling.

Compound **4** is a viable candidate for halide abstraction in CH<sub>3</sub>CN. Reaction of a CH<sub>3</sub>CN solution of **4** with 2 equiv. K [B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] (Fig. 2A) produced an immediate colour change from purple to orange. After workup, FT-IR (ATR) spectroscopic analysis showed a stretch at ν(N≡C) = 2252 cm<sup>-1</sup>, corroborating formation of [{η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>-(μ-CH<sub>2</sub>)}Fe<sup>II</sup>(dnppe)(NCCH<sub>3</sub>)<sub>2</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sub>2</sub> ([5][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sub>2</sub>). The <sup>1</sup>H NMR spectrum of [5]<sup>2+</sup> also showed the expected inequivalent {η<sup>5</sup>-C<sub>5</sub>Me<sub>4</sub>} signals




Fig. 3 Molecular structure of **4** with ellipsoids drawn at 50% probability. Hydrogen atoms omitted except for those on C(1) and C(2). Solvent molecule (benzene) has been omitted for clarity.



( $\delta_{\text{H}} = 1.63$  and  $1.48$  ppm), a  $\{\mu\text{-CH}_2\}$  signal ( $\delta_{\text{H}} = 2.22$  ppm), and a peak for  $[\text{Fe}]\text{-NCCH}_3$  ( $\delta_{\text{H}} = 2.41$  ppm; *c.f.*,  $\delta_{\text{H}} = 1.91$  ppm for  $[\text{Cp}^*\text{Fe}(\text{dnppe})(\text{NCCH}_3)]^+$  ( $[7]^+$ )). Comparison of the FT-IR spectra for the reaction between **1** and  $[\text{Cp}_2\text{Fe}]\text{B}(\text{C}_6\text{F}_5)_4$ ,  $[5]^{2+}$ ,  $[7]^+$ , and neat  $\text{CH}_3\text{CN}$ , confirmed that  $[5]^{2+}$  was formed from oxidation of **1** (Fig. 4) with an observed stretch at  $\nu(\text{N}\equiv\text{C}) = 2252\text{ cm}^{-1}$ . Notably, the combination band for neat  $\text{CH}_3\text{CN}$  ( $\nu = 2300\text{ cm}^{-1}$ ) also disappears upon  $\text{CH}_3\text{CN}$  coordination.<sup>44,45</sup> In addition to our observations from IR spectroscopy, a THF solution of **4** exposed to  $100\text{ }\mu\text{L}$  of  $\text{CH}_3\text{CN}$  and analyzed by CV displayed a predominant wave at  $E_{1/2} = -0.16\text{ V}$ , akin to the voltammogram acquired from the addition of  $100\text{ }\mu\text{L}$   $\text{CH}_3\text{CN}$  to **1**, noted above (Fig. S51<sup>†</sup>).

Compound **4** is also a valuable synthon for clean access to dimeric iron hydrides. For example, exposure of a THF solution of **4** to 2 equiv. of  $\text{K}[\text{HBET}_3]$  led to clean formation of the bis-hydride complex  $[\{\eta^5\text{-C}_5\text{Me}_4\text{-}(\mu\text{-CH}_2)\text{Fe}^{\text{II}}(\text{dnppe})(\text{H})_2\}]$  (**6**) (Fig. 2A).<sup>46</sup> By  $^1\text{H}$  NMR spectroscopy, the bridging  $\{\mu\text{-H}_2\text{CCH}_2\}$  group was located at  $\delta_{\text{H}} = 2.62$  and the inequivalent  $\{\eta^5\text{-C}_5\text{Me}_4\}$  signals were observed at  $\delta_{\text{H}} = 2.05$  and  $1.96$  ppm. Successful  $[\text{Fe}]\text{-H}$  formation was further established by locating the  $[\text{Fe}]\text{-H}$  signal as a triplet at  $\delta_{\text{H}} = -17.81$  ppm ( $^2J_{\text{H-P}} = 70.2$  Hz) (*c.f.*,  $\delta_{\text{H}} = -17.90$  ppm ( $^2J_{\text{H-P}} = 70.2$  Hz) for the monomeric  $[\text{Cp}^*\text{Fe}(\text{dnppe})(\text{H})]$ ).<sup>34</sup> FT-IR spectroscopy also revealed a characteristic  $\nu(\text{Fe-H})$  stretch at  $1847\text{ cm}^{-1}$ .

Having characterized the terminal  $\text{C}(\text{sp}^3)\text{-C}(\text{sp}^3)$  coupled product, we next sought to characterize the reactive oxidized precursor,  $[1]^+$ . Oxidation of **1** was thus carried out in 2-MeTHF using  $[\text{Cp}_2\text{Fe}]^+$  at  $-78\text{ }^\circ\text{C}$ . Following 5 min of reaction time, freeze-quenched continuous wave (CW) X-band EPR spectroscopy provided a spectrum with rhombic symmetry ( $g = [2.0004, 2.0526, 2.2134]$ ) (Fig. 5A) – this represents the first example of a spectroscopically characterized  $\text{Fe}(\text{III})$  tucked-in complex by EPR. Compound  $[1]^+$  can likewise be photochemically generated in a 2-MeTHF glass using **1** and a 405 nm laser ( $350\text{ mW mm}^{-2}$ ) following 10 min of irradiation (Fig. 5B). To provide




Fig. 4 Enhanced view of the region between  $2400\text{--}2150\text{ cm}^{-1}$  for the FT-IR (ATR) spectra for the reaction between **1** and 1 equiv. of  $[\text{Cp}_2\text{Fe}]\text{B}(\text{C}_6\text{F}_5)_4$  (red),  $[5]^{2+}$  (black),  $[7]^+$  (dark grey), and neat  $\text{CH}_3\text{CN}$  (light grey).




Fig. 5 A) Freeze-quenched CW X-band EPR spectra (8.885 GHz) recorded in 2-MeTHF glass at 77 K for  $[1]^+$  (black) and  $[2]^+$  (red). (B) Freeze-quenched CW X-band EPR spectra for  $[1]^+$  (top) and a photo-oxidized sample of **1** under 405 nm ( $350\text{ mW mm}^{-2}$ ) irradiation (bottom). (C) DFT-calculated spin density plots for  $[1]^+$  and  $[2]^+$  (all C and H atoms from the *dnppe* ligand, except those C atoms directly attached to P, have been omitted for clarity).  $\text{Fc} = \text{Cp}_2\text{Fe}$ .

a comparative basis, the  $\{\text{Fe}^{\text{III}}\text{-CH}_3\}$  complex  $[2]^+$  was also prepared by reaction of **2** with 1 equiv. of  $[\text{Cp}_2\text{Fe}]^+$  in 2-MeTHF at  $-78\text{ }^\circ\text{C}$ . By CW X-band EPR spectroscopy, a rhombic signal was also observed ( $g = [1.9827, 2.0450, 2.3968]$ ) (Fig. 5A), again consistent with an  $S = 1/2$   $\text{Fe}(\text{III})$  radical species ( $[2]^+$ ).<sup>47</sup> The  $g_{\text{max}}$  shift between compounds  $[1]^+$  and  $[2]^+$  is attributed to a difference in geometry about Fe. For  $[1]^+$ ,  $g$ -values are all greater than 2.0 with a spread of 0.05 to 0.16 – characteristic of the unpaired electron predominately residing in a  $d_{z^2}$  orbital.<sup>48</sup> For  $[2]^+$ ,



a larger value of  $g_{\max}$  is consistent with an unpaired electron residing in a  $d_{x^2-y^2}$  orbital.<sup>49</sup> Calculated spin density plots for  $[1]^+$  and  $[2]^+$  additionally corroborate that the unpaired electron predominately resides on Fe, with populations of  $0.917e^-$  for  $[1]^+$  and  $0.987e^-$  for  $[2]^+$  (Fig. 5C). The marked geometry change noted above (between  $[1]^+$  and  $[2]^+$ ) is best visualized from an overlay plot of their DFT-optimized structures (with coincident Fe and P atoms) (Fig. 6A). This data also showcases an elongated Fe–C bond in  $[1]^+$  ( $d(\text{Fe–C}) = 2.24 \text{ \AA}$ ) as compared to  $[2]^+$  ( $d(\text{Fe–C}) = 2.01 \text{ \AA}$ ), indicating weaker Fe–C overlap in  $[1]^+$ . The ring-strain inherent to  $[1]^+$  is additionally described by an Fe–C–C triangle with angles of  $38.4^\circ$ ,  $60.5^\circ$ , and  $81.1^\circ$  (Fig. 6A).

Mayer bond order (MBO) and natural bond orbital (NBO) analysis provide further insight that helps to rationalize  $\text{C}(\text{sp}^3)\text{--}\text{C}(\text{sp}^3)$  coupling in  $[1]^+$  (Fig. 6B).<sup>50</sup> To determine the change in bond order upon oxidation, MBOs were calculated. For  $1/[1]^+$ , the MBO decreased by 0.06 on oxidation ( $1: 0.66/[1]^+: 0.60$ ), while for  $2/[2]^+$ , the MBO was found to slightly increase ( $2: 0.73/[2]^+: 0.80$ ). NBO calculations provided second order perturbation energies ( $E^{(2)}$ ) which showed a similar trend. For the  $\text{C}(\text{lp}) \rightarrow \text{Fe}(\text{lv})$  ( $\text{lp} = \text{lone pair}$ ,  $\text{lv} = \text{lone valence}$ ) interaction, energies were found to be  $43.4$  and  $53.8 \text{ kcal mol}^{-1}$  for  $[1]^+$  and  $[2]^+$ , respectively (Fig. 6B). This data indicates that (1) the Fe–C bond becomes weaker on going from  $1 \rightarrow [1]^+$  and (2) that the Fe–C

bond of  $1/[1]^+$  is weaker than  $2/[2]^+$  – a consequence of ring-strain.

The radical character intrinsic to  $[1]^+$  is also evidenced by its chemical reactivity with 10 equiv. of the radical trap, (2,2,6,6-tetramethylpiperidin-1-yl)oxyl (TEMPO), which gives  $[\{\eta^5\text{-C}_5\text{Me}_4\text{-(CH}_2\text{-ONC}_9\text{H}_{18}\text{)}\}\text{Fe}^{\text{II}}(\text{dmppe})(\text{NCCH}_3)]\text{B}(\text{C}_6\text{F}_5)_4$  ( $[8]$ )  $\text{B}(\text{C}_6\text{F}_5)_4$  (Fig. 6C). Compound  $[8]^+$  was characterized by signals in the  $^1\text{H}$  NMR spectrum corresponding to  $\{\text{CH}_2\text{-ONC}_9\text{H}_{18}\}$  at  $\delta_{\text{H}} = 4.18 \text{ ppm}$  and Fe-bound  $\text{NCCH}_3$  at  $\delta_{\text{H}} = 2.40 \text{ ppm}$ . Confirmation of an Fe-bound  $\text{NCCH}_3$  group was determined *via* FT-IR spectroscopy, with a characteristic  $\nu(\text{N}\equiv\text{C})$  stretch at  $2247 \text{ cm}^{-1}$ . The  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum also showed a signal at  $\delta_{\text{C}} = 71.0 \text{ ppm}$ , corresponding to the TEMPO-bound  $\text{CH}_2$  from the  $\eta^5\text{-C}_5\text{Me}_4\text{-(CH}_2\text{-ONC}_9\text{H}_{18}\text{)}$  ligand. HRMS-ESI also provided a signal corresponding to  $[8]^+$  at  $649.407$  (calcd;  $649.407$ ). These





Fig. 6 (A) Structural overlay (capped sticks) of  $[1]^+$  (black) and  $[2]^+$  (red). Select angles ( $^\circ$ ) and select bond lengths ( $\text{\AA}$ ) are provided. (B) Mayer bond order (MBO) and natural bond orbital (NBO) analysis providing  $E^{(2)}$  values between  $\text{Fe}\text{--}\text{C}$ . (C) Reaction of  $[1]^+$  with (2,2,6,6-tetramethylpiperidin-1-yl)oxyl (TEMPO) forming  $[8]^+$ .

**A.** Thermodynamics for  $\text{C}(\text{sp}^3)\text{--}\text{C}(\text{sp}^3)$  bond formation in  $[1]^+$



**B.** Thermodynamics for ethane elimination in  $[2]^+$  (not observed)




Fig. 7 (A) Thermodynamic calculations showing energies associated with the dimerization of  $[1]^+$ . (B) Analogous calculations for the elimination of ethane ( $\text{CH}_3\text{CH}_3$ ) from  $[2]^+$ . All energies are in  $\text{kcal mol}^{-1}$ .



data corroborate a radical mechanism, whereby the generated  $\{C_5Me_4-(\cdot CH_2)\}$  group has been trapped by TEMPO.<sup>51</sup>

To assess the effect of ring-strain on  $\{[Fe]-C\}$  bond homolysis, we next sought information regarding the  $BDE_{[Fe]-C}$  of  $[1]^+$ . For reference, the  $BDE_{C-C}$  associated with the simplest cyclical saturated hydrocarbon (cyclopropane;  $(CH_2)_3$ ) is substantially lower than ethane due to ring strain ( $BDE_{C-C(cyclopropane)} = 57 \text{ kcal mol}^{-1}$  vs.  $BDE_{C-C(ethane)} = 90 \text{ kcal mol}^{-1}$ ).<sup>52–54</sup> Given this, and assuming  $CH_3CH_3$  and  $RCH_2CH_2R$  ( $R = C_5Me_4^-$ ) to have similar bond strengths, C–C coupling should be favoured for  $\{[Fe]-CH_2R\}$  when  $BDE_{[Fe]-C} \leq 45 \text{ kcal mol}^{-1}$ .

Computations show the difference in  $BDE_{[Fe]-C}$  for  $[1]^+$  and  $[2]^+$  to be  $23.3 \text{ kcal mol}^{-1}$  (Fig. 7). The absolute values of  $BDE_{[Fe]-C}$ :  $24.5$  and  $47.8 \text{ kcal mol}^{-1}$  for  $[1]^+$  and  $[2]^+$ , respectively are also consistent with observation of  $C(sp^3)-C(sp^3)$  coupling for the former. These calculated values are benchmarked against ethane, which provides  $BDE_{C-C(calcd)} = 88.4 \text{ kcal mol}^{-1}$  *c.f.*, the experimental  $BDE_{C-C} = 90 \text{ kcal mol}^{-1}$ .<sup>52</sup> Notwithstanding differences in  $BDE_{C-C}$ , both processes were calculated to be mildly exergonic (Fig. 7):  $\Delta G^\circ = -2.1 \text{ kcal mol}^{-1}$  and  $-4.3 \text{ kcal mol}^{-1}$  for C–C coupling from  $[1]^+$  or  $[2]^+$ , respectively, suggesting a difference in kinetic barrier. Based on the above discussion, the  $C(sp^3)-C(sp^3)$  coupling associated with  $[1]^+$  can thus be ascribed to a minimum of two features: (1) a strained three-membered  $\{-Fe-CH_2-C-\}$  ring and (2) a weakened  $\{[Fe^{III}]-CH_2R\}$  bond *c.f.*, **1**.<sup>32</sup> This is contrasted with  $[2]^+$ , where experimentally,  $CH_3CH_3$  is not evolved during synthesis (Fig. 7B).

## Conclusions

Metal-mediated cross-coupling has emerged as a powerful means to construct C–C bonds. In recent years, such studies have focused predominantly on base transition elements, such as iron, cobalt, and nickel, as cheaper and more sustainable sources. Here, we have studied an oxidized iron tucked-in complex  $[(\eta^6-C_5Me_4=CH_2)Fe^{III}(dnppe)]^+$  ( $[1]^+$ ), which due to both ring-strain and iron oxidation state, promotes  $C(sp^3)-C(sp^3)$  bond coupling. The product  $\{Fe_2\}$  diphosphine dimers invite reactions to give functionalized  $\{Fe_2^{III}\}$  or  $\{Fe_2^I\}$  products having coordinated halide, hydride, or solvent equivalents. Characterization of the reactive oxidized “tucked-in”  $\{Fe^{III}\}$  precursor was performed using CW X-band EPR spectroscopy, with computations revealing a weak  $\{[Fe^{III}]-C\}$  bond with  $BDE_{[Fe]-C} = 24.5 \text{ kcal mol}^{-1}$ . The observed  $C(sp^3)-C(sp^3)$  coupling, however, is not simply an artifact of iron oxidation state. The model,  $[Cp^*Fe^{III}(dnppe)(CH_3)]^+$   $[2]^+$ , for instance, does not undergo  $C(sp^3)-C(sp^3)$  coupling, consistent with a higher calculated  $BDE_{[Fe]-C}$  of  $47.8 \text{ kcal mol}^{-1}$ . This pair of compounds and their contrasting thermochemistry/reactivity pave the way towards better understanding the intricacies of  $\{M-C\}$  strain and metal oxidation state on reductive elimination, offering insight into an important elementary transformation of direct relevance to C–C bond coupling.

## Data availability

The data supporting this article have been included as part of the ESI.† Crystallographic data for **4** has been deposited at the

CCDC under 2341812. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

## Author contributions

Joseph A. Zurawski: investigation, methodology, formal Analysis, visualization, writing – original draft, writing – review & editing. Connor S. Durfy: investigation, validation. Noah B. Stoczek: investigation, methodology. Giovanni Fanchini: resources, methodology, funding acquisition. Marcus W. Drover: project administration, funding acquisition, visualization, supervision, formal analysis, conceptualization, methodology, writing – review & editing.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

The authors are grateful to Western University, the Council of Ontario Universities for a John C. Polanyi award to M. W. D., the Canadian Foundation for Innovation (LOF-212442), and the Natural Sciences and Engineering Research Council of Canada (Discovery Grant, RGPIN-2020-04480 (M. W. D.), RGPIN-2020-05008 (G. F.), Discovery Launch Supplement, DGEGR-2020-00183), and graduate award (CGS-D/NSERC Vanier to J. A. Z.) for funding.

## References

- R. H. Grubbs, *Angew. Chem., Int. Ed.*, 2006, **45**, 3760–3765.
- C. C. C. Johansson Seechurn, M. O. Kitching, T. J. Colacot and V. Snieckus, *Angew. Chem., Int. Ed.*, 2012, **51**, 5062–5085.
- A. Biffis, P. Centomo, A. Del Zotto and M. Zecca, *Chem. Rev.*, 2018, **118**, 2249–2295.
- B. Zhang, Y. Gao, Y. Hioki, M. S. Oderinde, J. X. Qiao, K. X. Rodriguez, H.-J. Zhang, Y. Kawamata and P. S. Baran, *Nature*, 2022, **606**, 313–318.
- Y. Liu, P. Li, Y. Wang and Y. Qiu, *Angew. Chem., Int. Ed.*, 2023, **62**, e202306679.
- P. Li, G. Kou, T. Feng, M. Wang and Y. Qiu, *Angew. Chem., Int. Ed.*, 2023, **62**, e202311941.
- Y. Wang, S. Dana, H. Long, Y. Xu, Y. Li, N. Kaplaneris and L. Ackermann, *Chem. Rev.*, 2023, **123**, 11269–11335.
- C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.*, 2013, **113**, 5322–5363.
- C.-S. Wang, P. H. Dixneuf and J.-F. Soulé, *Chem. Rev.*, 2018, **118**, 7532–7585.
- M. De Abreu, P. Belmont and E. Brachet, *Eur. J. Org. Chem.*, 2020, **2020**, 1327–1378.
- L. K. G. Ackerman, J. I. Martinez Alvarado and A. G. Doyle, *J. Am. Chem. Soc.*, 2018, **140**, 14059–14063.
- M. L. Clapson, C. S. Durfy, D. Facchinato and M. W. Drover, *Cell Rep. Phys. Sci.*, 2023, **4**, 101548.



- 13 X. Chen, K. M. Engle, D. Wang and J. Yu, *Angew. Chem., Int. Ed.*, 2009, **48**, 5094–5115.
- 14 M. B. Watson, N. P. Rath and L. M. Mirica, *J. Am. Chem. Soc.*, 2017, **139**, 35–38.
- 15 J. Kim, K. Shin, S. Jin, D. Kim and S. Chang, *J. Am. Chem. Soc.*, 2019, **141**, 4137–4146.
- 16 A. Cizikovs, E. E. Basens, P. A. Zagorska, A. Kinens and L. Grigorjeva, *ACS Catal.*, 2024, **14**, 1690–1698.
- 17 A. Pedersen and M. Tilset, *Organometallics*, 1993, **12**, 56–64.
- 18 W. Lau, J. C. Huffman and J. K. Kochi, *Organometallics*, 1982, **1**, 155–169.
- 19 M. P. Lanci, M. S. Remy, W. Kaminsky, J. M. Mayer and M. S. Sanford, *J. Am. Chem. Soc.*, 2009, **131**, 15618–15620.
- 20 K. Shin, Y. Park, M.-H. Baik and S. Chang, *Nat. Chem.*, 2018, **10**, 218–224.
- 21 T. J. Kealy and P. L. Pauson, *Nature*, 1951, **168**, 1039–1040.
- 22 P. J. Chirik, *Organometallics*, 2010, **29**, 1500–1517.
- 23 P. Preethalayam, K. S. Krishnan, S. Thulasi, S. S. Chand, J. Joseph, V. Nair, F. Jaroschik and K. V. Radhakrishnan, *Chem. Rev.*, 2017, **117**, 3930–3989.
- 24 A. Z. Kreindlin, F. M. Dolgushin, A. I. Yanovsky, Z. A. Kerzina, P. V. Petrovskii and M. I. Rybinskaya, *J. Organomet. Chem.*, 2000, **616**, 106–111.
- 25 E. I. Fedin, A. L. Blumenfeld, P. V. Petrovskii, A. Z. Kreindlin, S. S. Fadeeva and M. I. Rybinskaya, *J. Organomet. Chem.*, 1985, **292**, 257–268.
- 26 A. Z. Kreindlin, S. S. Fadeeva and M. I. Rybinskaya, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1984, **2**, 403–406.
- 27 J. A. Zurakowski and M. W. Drover, *Chem. Commun.*, 2023, **59**, 11349–11352.
- 28 J. P. Dinnocenzo, W. P. Todd, T. R. Simpson and I. R. Gould, *J. Am. Chem. Soc.*, 1990, **112**, 2462–2464.
- 29 P. Peng, X. Yan, K. Zhang, Z. Liu, L. Zeng, Y. Chen, H. Zhang and A. Lei, *Nat. Commun.*, 2021, **12**, 3075.
- 30 T. Krämer, M. R. Gyton, I. Bustos, M. J. G. Sinclair, S. Tan, C. J. Wedge, S. A. Macgregor and A. B. Chaplin, *J. Am. Chem. Soc.*, 2023, **145**, 14087–14100.
- 31 Given the similarity of  $E_{1/2}$  for both **1** and **2**, along with the lower reduction potentials ( $E_{1/2} \ll -1.23$  V vs.  $\text{Fc}/\text{Fc}^+$ ) observed for related Fe(0) species, a formal Fe(II) oxidation state seemingly better describes the observed electrochemical behaviour of **1**.
- 32 M. Tilset, I. Fjeldahl, J.-R. Hamon, P. Hamon, L. Toupet, J.-Y. Saillard, K. Costuas and A. Haynes, *J. Am. Chem. Soc.*, 2001, **123**, 9984–10000.
- 33 D. Patel, A. Wooles, A. D. Cornish, L. Steven, E. S. Davies, D. J. Evans, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Dalton Trans.*, 2015, **44**, 14159–14177.
- 34 J. A. Zurakowski, K. R. Brown and M. W. Drover, *Inorg. Chem.*, 2023, **62**, 7053–7060.
- 35 Complex **1** is insoluble in pure acetonitrile and was found to react with  $\text{CH}_2\text{Cl}_2$ .
- 36 F. S. Pick, D. B. Leznoff and M. D. Fryzuk, *Dalton Trans.*, 2018, **47**, 10925–10931.
- 37 P. Hamon, L. Toupet, J.-R. Hamon and C. Lapinte, *Organometallics*, 1996, **15**, 10–12.
- 38 Y. Liu, E. Solari, R. Scopelliti, F. Fadaei Tirani and K. Severin, *Chem.–Eur. J.*, 2018, **24**, 18809–18815.
- 39 M. Kessler, S. Hansen, C. Godemann, A. Spannenberg and T. Beweries, *Chem.–Eur. J.*, 2013, **19**, 6350–6357.
- 40 X. Tan, B. Li, S. Xu, H. Song and B. Wang, *Organometallics*, 2013, **32**, 3253–3261.
- 41 F. Wochner, L. Zsolnai, G. Huttner and H. H. Brintzinger, *J. Organomet. Chem.*, 1985, **288**, 69–77.
- 42 D. V. Muratov, A. S. Romanov, M. Corsini, A. R. Kudinov, F. F. de Biani and W. Siebert, *Chem.–Eur. J.*, 2017, **23**, 11935–11944.
- 43 X. Wang, M. Sabat and R. N. Grimes, *Organometallics*, 1995, **14**, 4668–4675.
- 44 Y. Kwon, C. Lee and S. Park, *Chem. Phys.*, 2014, **445**, 38–45.
- 45 B. Dereka, N. H. C. Lewis, J. H. Keim, S. A. Snyder and A. Tokmakoff, *J. Phys. Chem. B*, 2022, **126**, 278–291.
- 46 E. S. Wiedner, M. B. Chambers, C. L. Pitman, R. M. Bullock, A. J. M. Miller and A. M. Appel, *Chem. Rev.*, 2016, **116**, 8655–8692.
- 47 J. Morrow, D. Catheline, M. H. Desbois, J. M. Manriquez, J. Ruiz and D. Astruc, *Organometallics*, 1987, **6**, 2605–2607.
- 48 D. Chong, W. E. Geiger, N. A. Davis, A. Weisbrich, Y. Shi, A. M. Arif and R. D. Ernst, *Organometallics*, 2008, **27**, 430–436.
- 49 F. Paul and C. Lapinte, *Coord. Chem. Rev.*, 1998, **178–180**, 431–509.
- 50 A. J. Bridgeman, G. Cavigliasso, L. R. Ireland and J. Rothery, *J. Chem. Soc., Dalton Trans.*, 2001, 2095–2108.
- 51 Of note, complexes **1** and **2** exhibited no reactivity with TEMPO.
- 52 B. Ruscic, *J. Phys. Chem. A*, 2015, **119**, 7810–7837.
- 53 F. H. Seubold, *J. Phys. Chem.*, 1954, **22**, 945–946.
- 54 X. Lei, X. Cao, J. Wang and X. Li, *Combust. Flame*, 2022, **237**, 111881.

