



Cite this: *Analyst*, 2018, **143**, 6113

## Tandem fluorescence and Raman (fluoRaman) characterisation of a novel photosensitiser in colorectal cancer cell line SW480†

Julia Gala de Pablo,<sup>†a</sup> David R. Chisholm,<sup>†b,c</sup> Andreas Steffen,<sup>d</sup> Amanda K. Nelson,<sup>d</sup> Christoph Mahler,<sup>d</sup> Todd B. Marder,<sup>d</sup> Sally A. Peyman,<sup>†a,e</sup> John M. Girkin,<sup>†f</sup> Carrie A. Ambler,<sup>†b,g</sup> Andrew Whiting<sup>†\*b,c</sup> and Stephen D. Evans<sup>†\*a,e</sup>

The development of new imaging tools, molecules and modalities is crucial to understanding biological processes and the localised cellular impact of bioactive compounds. A small molecule photosensitiser, DC473, has been designed to be both highly fluorescent and to exhibit a strong Raman signal in the cell-silent region of the Raman spectrum due to a diphenylacetylene structure. DC473 has been utilised to perform a range of novel tandem fluorescence and Raman (fluoRaman) imaging experiments, enabling a thorough examination of the compound's cellular localisation, exemplified in colorectal cancer cells (SW480). This multifunctional fluoRaman imaging modality revealed the presence of the compound in lipid droplets and only a weak signal in the cytosol, by both Raman and fluorescence imaging. In addition, Raman microscopy detected the compound in a cell compartment we labelled as the nucleolus, whereas fluorescence microscopy did not detect the fluoRaman probe due to solvatochromatic effects in a local polar environment. This last finding was only possible with the use of tandem confocal Raman and fluorescence methods. By following the approach detailed herein, incorporation of strong Raman functional groups into fluorophores can enable a plethora of fluoRaman experiments, shedding further light on potential drug compound's cellular behaviour and biological activity.

Received 31st July 2018,  
Accepted 1st November 2018

DOI: 10.1039/c8an01461b

rsc.li/analyst

## Introduction

Imaging and tracing bioactive compounds *in vivo* is essential to understanding their mechanism of action. However, tagging them while retaining their inherent biological activity and function is very difficult, in particular when large fluorescent molecules such as the 238 residue Green Fluorescent Protein

(GFP) are employed as labels.<sup>1</sup> An inherently fluorescent bioactive molecule that does not require labelling is, therefore, highly desirable. However, there are other issues to consider when using fluorescence for quantitative molecular tracking, including photobleaching during the experiment,<sup>1</sup> solvatochromatic or fluorogenic effects in different locations within the cell and non-linear responses from the fluorophore due to localised concentrations or self-absorption.<sup>2</sup> For this reason, novel and complementary techniques for the detection and characterisation of a bioactive molecule within a cellular environment are needed.

Raman spectroscopy uses the inelastic scattering of monochromatic light to provide information on the vibrational modes of a sample, allowing the detection of chemical functional groups.<sup>3</sup> It is a label-free and non-destructive technique, with the ability to diagnose disease states or track the biochemical changes occurring in a cell over time.<sup>4</sup> Raman spectroscopy can be a valuable tool, not only for imaging a compound within a cell in a semi-quantitative manner, but also for obtaining information on the molecules with which they co-locate.<sup>5</sup> However, Raman signal intensity is several orders of magnitude weaker than fluorescence or elastic

<sup>a</sup>Molecular and Nanoscale Physics Group, School of Physics and Astronomy, University of Leeds, Leeds, UK. E-mail: S.D.Evans@leeds.ac.uk

<sup>b</sup>LightOx Limited, Wynyard Park House, Wynyard Avenue, Wynyard, Billingham, TS22 5TB, UK

<sup>c</sup>Department of Chemistry, Durham University, South Road, Durham, DH1 3LE, UK. E-mail: andy.whiting@durham.ac.uk

<sup>d</sup>Institut für Anorganische Chemie, Julius-Maximilians-Universität Würzburg, Am Hubland, 97074 Würzburg, Germany

<sup>e</sup>Welcome Trust Brenner Building, St James's University Hospital, Faculty of Medicine and Health, University of Leeds, Leeds, UK

<sup>f</sup>Biophysical Sciences Institute, Department of Physics, Durham University, South Road, Durham, DH1 3LE, UK

<sup>g</sup>Department of Biosciences, Durham University, South Road, Durham, DH1 3LE, UK

†Electronic supplementary information (ESI) available. See DOI: 10.1039/c8an01461b

‡Joint first authors.



scattering signals making the technique inherently slow and hence, lacking in sufficient sensitivity.<sup>4</sup>

To overcome these drawbacks, and to obtain orthogonal information provided by Raman measurements, multimodal imaging systems can be used.<sup>4</sup> Fluorescence is usually a problem in Raman spectroscopy, as it can cause significant background signal. Much effort has, therefore, gone into the design of Raman systems to reduce interference from fluorescence.<sup>4</sup> Since 2003, several groups have adopted the multimodal approach for increasing the tissue-diagnosis sensitivity of Raman probes by combining Raman with tissue auto-fluorescence.<sup>4,6,7</sup> Fluorescence tends to be used either as a confirmatory technique for the Raman mapping, or as a quick mapping technique to choose the areas of interest prior to Raman imaging in a large sample.<sup>4</sup> Some researchers have used both techniques in a complementary manner, such as the work of Carney *et al.* on extracellular vesicles using laser tweezers Raman spectroscopy and fluorescence labelling with CD9+ antibodies.<sup>8</sup> Raman and fluorescence have also been shown to provide complementary information for studying pH effects of doxorubicin uptake,<sup>9</sup> for tissue imaging using surface-enhanced Raman scattering (SERS) and fluorescence based endoscopy<sup>10</sup> and for dual drug release studies, using both surface-enhanced resonance Raman scattering (SERRS) and energy transfer to probe length scales.<sup>11</sup> Other groups that have combined fluorescence and Raman include Zeng *et al.* who tracked autofluorescence lifetime and stimulated Raman scattering of naturally occurring lignin molecules in poplar cells,<sup>12</sup> Manen *et al.* tracked both the resonant Raman signal of flavocytochrome b<sub>558</sub> and the fluorescence of labelled quantum dots in single live neutrophils using the same excitation laser,<sup>13</sup> and Uzunbajakava *et al.* investigated DNA using spontaneous Raman and two-photon excited fluorescence of the DNA dye Hoechst 33342.<sup>14</sup> Designing a molecular probe that can be imaged by both tandem fluorescence and Raman scattering could give additional information in the location and environment of the probe. Following this approach, Li *et al.* designed a mitochondrial probe for both stimulated Raman scattering (SRS) and aggregation induced emission (AIE) fluorescence in HeLa cells.<sup>15</sup> However, to our knowledge, designing a potential chemotherapeutic compound that can be imaged by both fluorescence and specific spontaneous Raman scattering (RS) in tandem (fluoRaman) has not been shown previously.

The vibrational signals associated with chemotherapeutic drugs often occur within the so-called “fingerprint” region of the spectrum, that also contains the vibrational information of both cells and tissue.<sup>16</sup> Consequently, the reliable and confident analysis of drug distribution in cells can be extremely challenging. Designing bioactive compounds with Raman signatures in the cell-silent region between 1800–2800 cm<sup>-1</sup> can enable detection of the compound more accurately in cellular samples.<sup>17</sup> Alkynes are one such example, and typically exhibit a vibrational frequency of *ca.* 2125 cm<sup>-1</sup> with intrinsically sharp peaks (fwhm ~14 cm<sup>-1</sup>).<sup>18</sup> Alkyne labels for Raman spectroscopy were introduced in the last decade by Yamakoshi *et al.*<sup>19,20</sup> and alkyne palettes have been developed by Chen

*et al.* using C<sup>13</sup> and C<sup>12</sup> isotope combinations<sup>18</sup> and by Hu *et al.* with their, so called, Carbow.<sup>21</sup> These probes show significant potential for diagnostic purposes in cancer therapy, using Raman spectroscopy.

Colorectal cancer (CRC) is the third most common cancer worldwide, with 1.4 million new cases diagnosed in 2012, with numbers worldwide predicted to rise to 1.36 million for men and 1.08 million for women by 2035.<sup>22</sup> Around 50% of the patients develop metastasis to the liver<sup>23</sup> and resistance to standard chemotherapy.<sup>24</sup> Novel therapies for the treatment of CRC are, therefore, needed.

Photodynamic therapy utilises photosensitisers to induce the production of large quantities of reactive oxygen species (ROS) for the targeted destruction of diseased tissues, and has been used to treat a variety of cancers.<sup>25</sup> However, current photosensitisers approved for therapeutic use suffer from a wide variety of issues, including poor pharmacokinetic properties due to their high molecular weight, often porphyrin structures, and ineffective targeting leading to the death of healthy tissue.<sup>25–27</sup> We have developed a new low molecular weight, small photosensitiser, DC473 (Fig. 1A), that elicits cell death through the stimulated production of ROS when activated by UV-A, violet or corresponding near-infrared two-photon wavelengths.<sup>28</sup> To improve this photosensitising activity, a precise understanding of the compound's cellular localisation is necessary. DC473 exhibits strong solvatochromatic fluorescence and possesses two alkynes suitable for Raman spectroscopy. It was anticipated that this unique combination of structural and photophysical properties would enable a range of novel tandem fluorescence and Raman (fluoRaman) imaging methodologies and experiments. Indeed, herein we report a detailed investigation into the cellular localisation of DC473 in the SW480 CRC cell line using confocal fluorescence and Raman (fluoRaman) microscopy, proving co-location of the compound signal with both techniques, and obtain the Raman signature of the organelles with which it spatially correlates.

## Materials and methods

### General photophysical information

UV-visible absorption spectra were obtained on an Agilent 1100 Series Diode Array spectrophotometer using standard 1 cm path length quartz cells. Excitation and emission spectra were recorded on an Edinburgh Instruments FLSP920 spectrophotometer, equipped with a 450 W xenon arc lamp, double monochromators for the excitation and emission pathways, and a red sensitive photomultiplier (PMT-R928) and a near-IR PMT as detectors, or on a Horiba Jobin-Yvon Fluoromax 3 spectrophotometer with single monochromators for the excitation and emission pathways. The excitation and emission spectra were corrected using the standard corrections supplied by the manufacturers for the spectral power of the excitation source and the sensitivity of the detector. The quantum yields were measured by use of integrating spheres with either the





**Fig. 1** A. Chemical structure and summary of the properties of photosensitiser, DC473. B. Raman spectrum of DC473, in the solid state, taken with the 40 $\times$  and 785 nm laser excitation; the inset shows the C $\equiv$ C band. The Raman signals corresponding to the alkyne and phenyl groups are indicated in red and blue on the peak labels, respectively. The main Raman bands tracked in this study are the 1597  $\text{cm}^{-1}$  phenyl C=C stretch and the 2197  $\text{cm}^{-1}$  C $\equiv$ C stretch vibrations. C. Absorption and D. emission spectra of DC473 in a variety of solvents MeCy = methylcyclohexane, Tol = toluene and DCM =  $\text{CH}_2\text{Cl}_2$ .

Edinburgh Instruments FLSP920 spectrophotometer or the Horiba Jobin-Yvon Fluoromax 3 spectrophotometer. The luminescence lifetimes were measured using a TCSPC module on an FLSP980 spectrometer equipped with a high-speed photomultiplier tube positioned after a single emission monochromator and operating with pulsed laser diodes (376 or 274 nm, repetition rate 1–5 MHz, pulse width *ca.* 200 ps, instrument response function *ca.* 500 ps). Decays were recorded to  $10^4$  counts in the peak channel with a record length of at least 1000 channels. The band-pass of the monochromator was adjusted to give a signal count rate of <20 kHz. Iterative deconvolution of the IRF with one decay function and nonlinear least-squares analysis were used to analyse the data. The quality of all decay fits was judged to be satisfactory, based on the calculated values of the reduced  $\chi^2$  and Durbin-Watson parameters and visual inspection of the weighted and autocorrelated residuals.

### Cell culture

SW480 cells were provided by Dr N. West from Saint James's University Hospital and were cultured in Dulbecco's modified Eagle medium (DMEM, Gibco) supplemented with 10% fetal bovine serum (Sigma), 2 mM glutamax (Thermo Fisher

Scientific) and penicillin 100 units per mL streptomycin 100  $\mu\text{g mL}^{-1}$  (Sigma). All experiments were done with passage numbers below 50. Quartz slides (UQG Optics,  $75 \times 25 \times 1$  mm) and coverslips ( $25.4 \times 25.4 \times 0.15$ – $0.25$  mm Alfa Aesar) were sonicated with acetone (VWR Chemicals), 2–5% decon 90 (VWR Chemicals) and rinsed with milliQ. Hydrogen peroxide 30% (Thermo Fisher) and sulphuric acid >95% (Thermo Fisher) were mixed in a 3 : 7 proportion (Piranha solution) and used to clean the slides for 20 minutes. Slides and coverslips were rinsed and stored in milliQ and dried under a stream of nitrogen immediately before using. Quartz coverslips were coated with poly-L-Lys (Sigma) and basement membrane extract (BME, Cultrex) to increase cell adherence. Treatment with poly-L-lysine was done covering the coverslip with 10  $\mu\text{g mL}^{-1}$  poly-L-lysine solution and incubating for 1 h, storing at 4  $^\circ\text{C}$  for 1 week. BME 12–18  $\text{mg mL}^{-1}$  was thawed and diluted to  $\sim 150 \mu\text{g mL}^{-1}$ . Coverslips were rinsed with DPBS and BME was added and incubated for 1 h. Cells (viability 96% Trypan blue) were seeded at a concentration of  $3.5 \times 10^5$  cells per well 2 days in advance. After two days, cells were treated with either DC473 4  $\mu\text{M}$  or the equivalent volume of dimethyl sulfoxide (DMSO, sterile, New England Biolabs) in complete DMEM for 4 hours. After incubation, cells were washed with DPBS and








laser after baseline subtraction, with the characteristic presence of the diphenylacetylene triple bond stretching vibration at  $2197\text{ cm}^{-1}$ . Alkyne peaks usually appear at *ca.*  $2125\text{ cm}^{-1}$ ;<sup>18</sup> however, the electron-rich tetrahydroquinoline donor group causes a shift to  $2197\text{ cm}^{-1}$ .<sup>21</sup> *N*-Propargyl alkynes Raman bands are typically much less intense and of lower wavenumber than their diphenylacetylene counterparts and it is, therefore, likely that this alkyne stretch is present as one of the weaker signals in the  $2080\text{--}2160\text{ cm}^{-1}$  region. A strong signal assigned as phenyl ring vibrations was also evident at  $1597\text{ cm}^{-1}$ . Both the phenyl vibrations at  $1597\text{ cm}^{-1}$  and the  $2197\text{ cm}^{-1}$  diphenylacetylene alkyne peak were used for the detection of the compound *via* Raman spectroscopy in single cells.

### Cellular imaging

With the fluorescence and Raman signatures of DC473 understood, we next sought to characterise the compound's cellular localisation and behaviour using the primary adenocarcinoma colorectal cancer cell line, SW480. First, the Raman spectra were acquired with a  $785\text{ nm}$  laser at individual points in fixed SW480 cells that had been incubated with DC473, and compared with the confocal fluorescence images of the same cells. Fig. 2C shows high-quality raw Raman spectra taken from different parts of a fixed cell. DC473 accumulation was detected when measuring in vesicles and in membranes, but not in the nucleus. The fluorescence image in Fig. 2B shows accumulation of the compound mainly in vesicles and weakly in the cytosol, but not in the nucleus. Raman measurements in the indicated points (V1–3) confirmed the presence of DC473 in particles which correlate with the strong  $1455\text{ cm}^{-1}$

band associated with lipids, indicating that they were likely lipid droplets, previously described in this cell line.<sup>43,44</sup> Cytosol measurements showed weak signals of DC473, in agreement with the fluorescence characterisation. Single point measurements in the nucleus showed only a very weak signal from the compound.

When exciting with the  $532\text{ nm}$  laser, cytochrome *c*, a small, soluble heme-protein localised in mitochondria, shows a resonant Raman signal.<sup>45,46</sup> Fig. 2D and E show the bright field and confocal fluorescence images of an SW480 cell following incubation with DC473. Fig. 2F shows individual points probed with RS with a  $532\text{ nm}$  laser in the same cell in vesicles, the nucleus, cytosol and filopodia. Similar to the results obtained with the near-infrared excitation, DC473 was detected strongly in vesicles and weakly in the cytosol. The cytosol regions examined exhibited the characteristic  $749$ ,  $1128$ ,  $1310$  and  $1582\text{ cm}^{-1}$  cytochrome peaks, indicating that the DC473 in this compartment could be localized near, or inside, the mitochondria. Mitochondria are an important target for photodynamic therapy agents and it is, therefore, important to understand the behaviour of DC473 with respect to co-localisation with the mitochondria signal. Ideally, higher resolution studies should be undertaken to confirm this co-localisation. However, given the large membranous area associated with the mitochondria and the ability of DC473 to accumulate in lipid rich environments, this seems reasonable. Indeed, given that the DC473 signal is not resonantly enhanced whilst that of the cytochrome *c* is, would imply significant accumulations of DC473 in the mitochondria, perhaps as high as 3–4 orders of magnitude greater than that of cytochrome *c*.




**Fig. 2** A and D Show the bright field and B and E show the confocal fluorescence images of fixed SW480 cells following incubation with DC473 whose point spectra were measured in C and F, respectively, where V stands for vesicle, N for nucleus, F for filopodia and C for cytosol. C. Raw Raman  $785\text{ nm}$  spectra at selected points of the SW480 cell showed in A and B. The green bands show the main peaks associated with the presence of the compound in different cellular locations, with strong presence in vesicles. F.  $532\text{ nm}$  Raman baseline subtracted spectrum of selected points of the SW480 cell showed in D and E. The green bands show the main DC473 peaks, showing a similar trend to Fig. 1B. Measurements in the cytosol show weak signal of DC473 and resonant Raman bands of cytochrome *c*, indicated with red bands.




### Mapping of DC473 accumulation using confocal fluorescence and Raman spectroscopy

Raman mapping was undertaken on fixed SW480 cells incubated with the compound, or with an equivalent concentration of DMSO, as a control (Fig. 3) using 785 nm excitation. This excitation was chosen due to a small but non-zero overlap of

the fluorescence excitation spectrum of the molecule with the 532 nm excitation line. Bright field and fluorescence images for the DC473-treated cell are shown together with the Raman maps associated with the main spectral peaks. The  $1338\text{ cm}^{-1}$  band, usually assigned as a mixed contribution of  $\text{CH}_2$  wagging, C–O stretching, phenyl stretching and N–H deformation,<sup>47–49</sup> appears most strongly here in the nucleus.



**Fig. 3** Peak integrated intensity maps for SW480 fixed cells either incubated with DC473 or with an equivalent concentration of DMSO as a control. The  $1597$  and  $2197\text{ cm}^{-1}$  peaks show DC473 accumulation mainly in lipid droplets, as indicated by the correlation with the  $1445\text{ cm}^{-1}$  intensity map mainly due to lipid vibrations. The  $1338\text{ cm}^{-1}$  intensity map shows mixed contributions of proteins and nucleic acids, and the  $1656\text{ cm}^{-1}$  map shows Amide I intensity as a map of proteins. The  $1455$  and  $1434\text{ cm}^{-1}$  maps mainly show contributions of lipids.



**Fig. 4** A. Bright field and confocal fluorescence images of the fixed SW480 cell incubated with DC473. The maps of each of the scores of the 4 main PCs are shown. B. Coefficients for the first 4 PCs whose scores are mapped in A. C. Bright field, fluorescence and score maps of PCA coefficients with DC473 associated Raman bands. The cell main features have been overlaid with the images as a visual aid. The presence of PC4 suggests that Raman could be detecting presence of DC473 in a region in the nucleus that does not show in the fluorescence image.





The 1434  $\text{cm}^{-1}$  band associated with methylene stretching<sup>50,51</sup> shows a strong contribution in lipid droplets rich regions and exhibits a strong correlation with the 1455  $\text{cm}^{-1}$  band associated with lipids and fatty acids<sup>50</sup> as well as the 1597  $\text{cm}^{-1}$  phenyl peak associated with DC473. The 1656  $\text{cm}^{-1}$  Amide I band<sup>50</sup> due to proteins is evident over the whole cell area. The two main DC473 peaks, 1597  $\text{cm}^{-1}$  (phenyl) and 2197  $\text{cm}^{-1}$  (alkyne), show a strong correlation with the lipid peaks and a weaker contribution in the cytosol, indicating localisation of the compound in lipid-rich environments. Whilst the 1597  $\text{cm}^{-1}$  DC473 peak is very strong and provides good contrast in the Raman images, its presence on the side of the Amide I peak makes it difficult to separate from that of protein. In contrast, the 2197  $\text{cm}^{-1}$  band, whilst weaker, occurs in the 'silent' region of the spectrum giving unequivocal detection of the compound.

PCA analysis of one of the maps of a cell incubated with DC473 is shown in Fig. 4A, along with the coefficients of these components in Fig. 4B. PC1 shows mainly average cell signal with DC473 contribution, where the whole cell shows higher scores than the background and thus gives similar information to the fluorescence image. PC3 peaks in the nucleus, with negative contributions for both DC473 and the broad lipid band around 1448  $\text{cm}^{-1}$ . Similarly to the intensity maps, PC2 shows positive contributions for the region around 1338  $\text{cm}^{-1}$ . PC2 and PC4 show the main DC473-specific components. PC2 shows very strong DC473 presence and a positive 1434  $\text{cm}^{-1}$  methylene band indicating accumulation in the elongated structures in the region below the nucleus in Fig. 4A/C.

PC4 shares some similarities with PC2, but with lower DC473 content and with stronger contribution of a band at 1627  $\text{cm}^{-1}$ . This component shows the presence of the compound in different regions of the cell that do not correlate with the fluorescence signal, as shown in Fig. 4. The PC4 component indicates the presence of DC473 in a compact area inside the nucleus, suggesting that a low concentration of the compound could be present inside the nucleolus. The absence of fluorescence in this region might be explained by a solvatochromatic reduction in quantum yield when the compound is localised in a more polar environment. Hence, the use of fluoRaman microscopy can not only identify the presence of the compound in different cell compartments but also provide information about the polarity of the environment of the compound.

## Conclusions

A continuing and major requirement in the life sciences is for small molecules that can be specifically targeted to sites within cells, which report back on local conditions within the cell, and can then be externally activated to alter cell activity using a minimally invasive method, such as light. We have designed and synthesised a novel photosensitiser, DC473, which is both fluorescent and Raman active in the silent region of the spectra and can, therefore, be tracked within the cell, with the additional advantage of its solvatochromatic fluorescence properties changing due to local conditions. By

exploiting the unique structural and spectroscopic properties of DC473, we have utilised fluoRaman microscopy to characterise and understand the cellular localisation of the compound in CRC SW480 cells. These novel imaging experiments have allowed us to detect the compound mainly in lipid droplets and in the cytosol, where it correlates with the resonant Raman signal of cytochrome C from mitochondria, suggesting that it could be located in or near this cellular compartment. Furthermore, we have acquired maps from single SW480 CRC fixed cells where Raman imaging strongly detects the compound in lipid droplet rich areas and weakly in the cytosol. However, further analysis using PCA also shows weak signal for DC473 within the nucleus, suggesting it could also be present in the nucleolus, where it does not show fluorescence due to solvatochromatic effects in a more polar environment. The detection of the compound in the latter, and its polar characteristic, would not have been possible using either of the two techniques used here on their own, *i.e.*, a fluoRaman confocal imaging approach offers a clear advantage for studying this kind of reporter molecules. We anticipate that, by application of the types of chemical design principles we have employed herein, combined with the novel fluoRaman methods, this type of imaging has significant potential in a host of cellular and tissue imaging contexts. This approach can likely be readily applied to a range of other imaging probes and drugs, potentially providing new insights into the cellular behaviour of not only these compounds, but enabling new and unexplored molecular probing methods to be developed.

## Conflicts of interest

AW and CAA are shareholders and directors of LightOx Ltd, a company licensed to pursue commercial applications for the novel compound described in this manuscript.

## Acknowledgements

We thank Dr N. West for providing the SW480 cell line. J. G. P. thanks the EPSRC Award (1654179) and D. B. and S. D. E. the MRC grant (MR/M009084/1) for funding, S. D. E. also acknowledges support from NIHR (MIC-2016-004) and EPSRC (EP/P023266/1). The data used in the figures of this paper will be available at <https://doi.org/10.5518/465>. D. R. C thanks the EPSRC, BBSRC and High Force Research Ltd for doctoral funding. AKN thanks the Fulbright Foundation for a scholarship. TBM thanks the Julius-Maximilians-Universität Würzburg for support.

## Notes and references

- 1 S. Shashkova and M. C. Leake, *Biosci. Rep.*, 2017, **37**, BSR20170031.
- 2 A. S. Klymchenko, *Acc. Chem. Res.*, 2017, **50**, 366–375.
- 3 D. Zhang, P. Wang, M. N. Slipchenko and J.-X. Cheng, *Acc. Chem. Res.*, 2014, **47**, 2282–2290.



- 4 N. K. Das, Y. Dai, P. Liu, C. Hu, L. Tong, X. Chen and Z. J. Smith, *Sensors*, 2017, **17**, 1592.
- 5 J. Phillip Kennedy, P. Jeffrey Conn and C. W. Lindsley, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 3204–3208.
- 6 Z. Huang, H. Lui, D. I. McLean, M. Korbelik and H. Zeng, *Photochem. Photobiol.*, 2005, **81**, 1219.
- 7 N. Uzunbajakava and C. Otto, *Opt. Lett.*, 2003, **28**, 2073.
- 8 R. P. Carney, S. Hazari, M. Colquhoun, D. Tran, B. Hwang, M. S. Mulligan, J. D. Bryers, E. Girda, G. S. Leiserowitz, Z. J. Smith and K. S. Lam, *Anal. Chem.*, 2017, **89**, 5357–5363.
- 9 B. Kang, M. M. Afifi, L. A. Austin and M. A. El-Sayed, *ACS Nano*, 2013, **7**, 7420–7427.
- 10 S. Jeong, Y. Kim, H. Kang, G. Kim, M. G. Cha, H. Chang, K. O. Jung, Y.-H. Kim, B.-H. Jun, D. W. Hwang, Y.-S. Lee, H. Youn, Y.-S. Lee, K. W. Kang, D. S. Lee and D. H. Jeong, *Sci. Rep.*, 2015, **5**, 9455.
- 11 E.-O. Ganbold, J. Yoon, D. Kim and S.-W. Joo, *Phys. Chem. Chem. Phys.*, 2015, **17**, 3019–3023.
- 12 Y. Zeng, S. Zhao, H. Wei, M. P. Tucker, M. E. Himmel, N. S. Mosier, R. Meilan and S.-Y. Ding, *Biotechnol. Biofuels*, 2015, **8**, 126.
- 13 H.-J. van Manen and C. Otto, *Nano Lett.*, 2007, **7**, 1631–1636.
- 14 N. Uzunbajakava, A. Lenferink, Y. Kraan, E. Volokhina, G. Vrensen, J. Greve and C. Otto, *Biophys. J.*, 2003, **84**, 3968–3981.
- 15 X. Li, M. Jiang, J. W. Y. Lam, B. Z. Tang and J. Y. Qu, *J. Am. Chem. Soc.*, 2017, **139**, 17022–17030.
- 16 A. F. Palonpon, M. Sodeoka and K. Fujita, *Curr. Opin. Chem. Biol.*, 2013, **17**, 708–715.
- 17 S. Hong, T. Chen, Y. Zhu, A. Li, Y. Huang and X. Chen, *Angew. Chem., Int. Ed.*, 2014, **53**, 5827–5831.
- 18 Z. Chen, D. W. Paley, L. Wei, A. L. Weisman, R. A. Friesner, C. Nuckolls and W. Min, *J. Am. Chem. Soc.*, 2014, **136**, 8027–8033.
- 19 H. Yamakoshi, K. Dodo, M. Okada, J. Ando, A. Palonpon, K. Fujita, S. Kawata and M. Sodeoka, *J. Am. Chem. Soc.*, 2011, **133**, 6102–6105.
- 20 H. Yamakoshi, K. Dodo, A. Palonpon, J. Ando, K. Fujita, S. Kawata and M. Sodeoka, *J. Am. Chem. Soc.*, 2012, **134**, 20681–20689.
- 21 F. Hu, C. Zeng, R. Long, Y. Miao, L. Wei, Q. Xu and W. Min, *Nat. Methods*, 2018, **15**, 194–200.
- 22 J. Ferlay, I. Soerjomataram, R. Dikshit, S. Eser, C. Mathers, M. Rebelo, D. M. Parkin, D. Forman and F. Bray, *Int. J. Cancer*, 2015, **136**, E359–E386.
- 23 C. C. Applegate and M. A. Lane, *World J. Gastrointest. Oncol.*, 2015, **7**, 184.
- 24 R. Abdel-Samad, P. Aouad, H. Gali-Muhtasib, Z. Sweidan, R. Hmadi, H. Kadara, E. L. D'Andrea, A. Fucci, C. Pisano and N. Darwiche, *Am. J. Cancer Res.*, 2018, **8**, 39–55.
- 25 D. E. J. G. J. Dolmans, D. Fukumura and R. K. Jain, *Nat. Rev. Cancer*, 2003, **3**, 380–387.
- 26 W. Fan, P. Huang and X. Chen, *Chem. Soc. Rev.*, 2016, **45**, 6488–6519.
- 27 N. Shirasu, S. O. Nam and M. Kuroki, *Anticancer Res.*, 2013, **33**, 2823–2831.
- 28 C. Ambler, D. R. Chisholm and A. Whiting, Synthetic retinoid (in cell killing), University of Durham, *UK Patent*, GB1613712.7, 2016.
- 29 M. Koch, C. Suhr, B. Roth and M. Meinhardt-Wollweber, *J. Raman Spectrosc.*, 2017, **48**, 336–342.
- 30 M. Krämer, U. H. F. Bunz and A. Dreuw, *J. Phys. Chem. A*, 2017, **121**, 946–953.
- 31 C. Fenante, U. Kensity and B. Dick, *J. Phys. Chem.*, 1993, **97**, 13457–13463.
- 32 Y. Hirata, T. Okada and T. Nomoto, *Chem. Phys. Lett.*, 1997, **278**, 133–138.
- 33 M. Szyzkowska, I. Bylińska and W. Wiczak, *J. Photochem. Photobiol., A*, 2016, **326**, 76–88.
- 34 M. Biswas, P. Nguyen, T. B. Marder and L. R. Khundkar, *J. Phys. Chem. A*, 1997, **101**, 1689–1695.
- 35 J. C. Collings, A. C. Parsons, L. Porrès, A. Beeby, A. S. Batsanov, J. A. K. Howard, D. P. Lydon, P. J. Low, I. J. S. Fairlamb and T. B. Marder, *Chem. Commun.*, 2005, 2666–2668.
- 36 P. Nguyen, Z. Yuan, L. Agocs, G. Lesley and T. B. Marder, *Inorg. Chim. Acta*, 1994, **220**, 289–296.
- 37 J. H. Barnard, J. C. Collings, A. Whiting, S. A. Przyborski and T. B. Marder, *Chem. – Eur. J.*, 2009, **15**, 11430–11442.
- 38 G. Clemens, K. R. Flower, P. Gardner, A. P. Henderson, J. P. Knowles, T. B. Marder, A. Whiting and S. Przyborski, *Mol. Biosyst.*, 2013, **9**, 3124.
- 39 V. B. Christie, J. H. Barnard, A. S. Batsanov, C. E. Bridgens, E. B. Cartmell, J. C. Collings, D. J. Maltman, C. P. F. Redfern, T. B. Marder, S. Przyborski and A. Whiting, *Org. Biomol. Chem.*, 2008, **6**, 3497.
- 40 C. Dehu, F. Meyers and J. L. Bredas, *J. Am. Chem. Soc.*, 1993, **115**, 6198–6206.
- 41 P. Nguyen, G. Lesley, T. B. Marder, I. Ledoux and J. Zyss, *Chem. Mater.*, 1997, **9**, 406–408.
- 42 D. R. Chisholm, G. L. Zhou, E. Pohl, R. Valentine and A. Whiting, *Beilstein J. Org. Chem.*, 2016, **12**, 1851–1862.
- 43 D. Tsikritsis, S. Richmond, P. Stewart, A. Elfick and A. Downes, *Analyst*, 2015, **140**, 5162–5168.
- 44 S. A. Schønberg, A. G. Lundemo, T. Fladvad, K. Holmgren, H. Bremseth, A. Nilsen, O. Gederaas, K. E. Tvedt, K. W. Egeberg and H. E. Krokan, *FEBS J.*, 2006, **273**, 2749–2765.
- 45 M. Okada, N. I. Smith, A. F. Palonpon, H. Endo, S. Kawata, M. Sodeoka and K. Fujita, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 28–32.
- 46 J. Gala de Pablo, F. J. Armistead, S. A. Peyman, D. Bonthron, M. Lones, S. Smith and S. D. Evans, *J. Raman Spectrosc.*, 2018, 1–10.
- 47 L. Ashton and E. W. Blanch, *J. Mol. Struct.*, 2010, **974**, 132–138.
- 48 Z. Movasaghi, S. Rehman and I. U. Rehman, *Appl. Spectrosc. Rev.*, 2007, **42**, 493–541.
- 49 A. Barth and C. Zscherp, *Q. Rev. Biophys.*, 2002, **35**, S0033583502003815.
- 50 Z. Movasaghi, S. Rehman and D. I. ur Rehman, *Appl. Spectrosc. Rev.*, 2008, **43**, 134–179.
- 51 D. F. H. Wallach, S. P. Verma and J. Fookson, *Biochim. Biophys. Acta, Rev. Biomembr.*, 1979, **559**, 153–208.

