



Cite this: *Nanoscale*, 2022, **14**, 263

## Pulmonary delivery of mucosal nanovaccines

Wei Tang,<sup>\*a</sup> Yu Zhang<sup>b</sup> and Guizhi Zhu <sup>\*b</sup>

Mucosal vaccination can elicit both systemic and mucosal immunity, and therefore has the potential to not only treat mucosal immune diseases, prevent the pathogen infection at the mucosal entry sites, but also treat distant or systemic immune disorders. However, only a few mucosal vaccines have been approved for human use in the clinic. Effective mucosal immunization requires the delivery of immunogenic agents to appropriate mucosal surfaces, which remains significantly challenging due to the essential biological barriers presenting at mucosal tissues. In the past decade, remarkable progress has been made in the development of pulmonary mucosal nanovaccines. The nanovaccines leverage advanced nanoparticle-based pulmonary delivery technologies on the characteristics of large surface area and rich antigen presentation cell environment of the lungs for triggering robust immune protection against various mucosal diseases. Herein, we review current methods and formulations of pulmonary delivery, discuss the design strategies of mucosal nanovaccines for potent and long-lasting immune responses, and highlight recent advances in the application of lipid-based pulmonary nanovaccines against mucosal diseases. These advances promise to accelerate the development of novel mucosal nanovaccines for the prophylaxis and therapy of infectious diseases, and cancer, as well as autoimmune disorders at mucosal tissues.

Received 3rd October 2021,  
Accepted 29th November 2021

DOI: 10.1039/d1nr06512b

rs.c.li/nanoscale


<sup>a</sup>Department of Pharmacy and Department of Diagnostic Radiology, Faculty of Science and Yong Loo Lin School of Medicine, National University of Singapore, Singapore, 119077, Singapore. E-mail: wei.tang@nus.edu.sg

<sup>b</sup>Department of Pharmaceutics and Center for Pharmaceutical Engineering and Sciences, Institute for Structural Biology and Drug Discovery, School of Pharmacy, The Developmental Therapeutics Program, Massey Cancer Center, Richmond, VA 23298, USA. E-mail: gzhu2@vcu.edu

### 1. Introduction

The emergency use authorization (EUA) of mRNA vaccines (mRNA-1273 from Moderna and BNT162b from Pfizer) for coronavirus disease 2019 (COVID-19) marks a milestone for nanovaccines. By taking advantage of lipid nanoparticle-based delivery technologies, the mRNA vaccines successfully protect antigen-encoding nucleic acids from degradation, facilitate their cellular uptake and release, and generate systemic neutralizing antibodies.<sup>1–3</sup> However, the mRNA vaccines and other authorized COVID-19 vaccines are administered *via* an intramuscular injection. They are able to elicit systemic immune responses to block the development of COVID-19, but not to induce mucosal immunity at the entry ports of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) to prevent the viral transmission or infection.<sup>4</sup> To achieve robust mucosal immunization, a vaccine requires to be delivered to the appropriate mucosal surfaces,<sup>4–6</sup> for example, the nasal or lung tissue in the case of COVID-19. Nanovaccines that leverage nanotechnologies to overcome biological barriers to antigen delivery, representing an excellent opportunity for inducing strong immune protection at mucosal surfaces (Fig. 1).

Currently, mucosal nanovaccines are urgently needed for not only COVID-19, but also many other mucosal diseases caused by enteric pathogens, sexually transmitted agents, and mucosa-accessible oncogenic viruses (Fig. 2a).<sup>4,7,8</sup> Mucosal membranes of respiratory, digestive, and urogenital tracts cover the largest exposure surface areas of a human body.<sup>5,6</sup> These tissues form a highly specialized adaptive and innate




**Guizhi Zhu**

*Guizhi (Julian) Zhu is currently an Assistant Professor of Pharmaceutics at Virginia Commonwealth University (VCU). He was a National Institutes of Health (NIH) KL2 Scholar (2018–2021) and an NIH/NIGMS MIRA (R35) awardee (2021). He received a BS in Biotechnology from Nankai University, and PhD in Biomedical Sciences from the University of Florida, followed by a postdoc on immune drug phar-*

*macoimaging and delivery in NIH. His multidisciplinary group (<https://www.guizhizhu.org/>) studies the engineering and delivery of nucleic acid immunotherapeutics and vaccines for the treatment and/or prophylaxis of cancer, infectious diseases, and autoimmune diseases. He has published >90 peer-reviewed papers with >9000 citations.*





**Fig. 1** Schematic illustration of pulmonary delivery of mucosal nanovaccines. (a) Representative delivery vectors of mucosal nanovaccines. (b) Illustration of biological barriers at the lung tissue. (c) Immune responses stimulated upon the delivery of mucosal vaccines. Nanoparticles can be tailored to facilitate antigens and/or adjuvants/immunostimulators to overcome the delivery barriers and elicit robust protective mucosal immune responses.

mucosal immune system, which contains mucus as the first barrier against enteric pathogenic microorganisms and mucosa-associated lymphoid tissues (MALTs) for providing protective immune responses.<sup>5,6,9</sup> Mucosal vaccination has potential beneficials over conventional parenteral vaccination since it stimulates both systemic and mucosal immune defenses.<sup>10</sup> A key characteristic of mucosal adaptive immune response is the local production and secretory of immunoglobulin A (IgA) antibodies. Secretory IgA (sIgA) facilitates entrapment and clearance of pathogenic antigens or microorganisms in the mucus, thus effectively excluding pathogen invasion at the mucosal sites.<sup>5,6,11</sup> The other important characteristic is the generation of tissue-resident memory T cells ( $T_{RM}$ s), which offer rapid protection in the tissue by recruiting circulating memory T cells and producing inflammatory cytokines/chemokines.<sup>12,13</sup> Moreover, the crosstalk between mucosal compartments allows immunization at one mucosal surface to establish immunity at distant mucosal sites. Taken together, mucosal vaccination holds great promise in the control of mucosal diseases. Recently, a head-to-head comparison study was performed to investigate the protection activity of ChAd-SARS-CoV-2 vaccine, a chimpanzee adenoviral vaccine encoding stabilized spike protein (Fig. 2b).<sup>14</sup> It found that a single-

dose intranasal vaccination could evoke strong systemic and mucosal IgA and T cell responses in mice to almost completely prevent SARS-CoV-2 infections in both upper and lower respiratory tracts, whereas intramuscular immunization was only able to activate systemic immune responses that partially reduced lung infections.<sup>14</sup> This study clearly suggests that the route of immunization exerts a substantial impact on vaccine efficacy. However, a majority of vaccines in use today are administered by an intramuscular or subcutaneous route. There are only nine mucosal vaccines are approved for human use, including eight oral vaccines against cholera, rotavirus, poliovirus, and salmonella, and one intranasally administered live attenuated influenza vaccine.<sup>4</sup> However, oral vaccines need to survive harsh acidic environment in the stomach as well as to overcome variable pH conditions and the tight monolayer of endothelial cells lining throughout the gastrointestinal tract.<sup>15</sup> Intranasal delivery have risks of exposing therapeutics to the central nervous system (CNS),<sup>16,17</sup> which may cause adverse side effects. Therefore, continuous efforts are highly desired to develop safe and effective mucosal vaccines. Recently, pulmonary vaccine delivery has attracted increasing attention due to the large surface area and the rich antigen presentation cells (APCs) as desired vaccine target cells in the microenvironment





**Fig. 2** Mucosal vaccines are urgently needed for mucosal diseases and they have potential benefits over conventional parenteral vaccination. (a) Burden of mucosal diseases with unmet vaccine needs. In addition to their centrality in the pathogenesis of infectious disease, mucosal tissues are frequent sites of infectious disease and tumor development. CRC, colorectal cancer; ETEC, enterotoxigenic *Escherichia coli*; p.a., per annum; RSV, respiratory syncytial disease; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2. Reproduced with permission.<sup>4</sup> Copyright 2021, Nature Publishing Group. (b) Comparison of immunogenicity of single-dose administration of ChAd-SARS-CoV-2-S, a chimpanzee adenoviral vaccine encoding stabilized spike protein, by intramuscular and intranasal delivery routes. The mucosal immunity generated by intranasal inoculation of ChAd-SARS-CoV-2 likely controlled infections at both upper and lower respiratory tract, potentially protecting against SARS-CoV-2 infection and transmission. Reproduced with permission.<sup>14</sup> Copyright 2020, Elsevier Inc.

of lung tissue.<sup>18</sup> Pulmonary vaccination has demonstrated successful control of respiratory pathogen infections, lung cancer metastasis, and even sexually transmitted viral infections.<sup>13,19</sup>

In this review, we briefly overview the methods and formulations of pulmonary drug delivery. Then, we discuss the key factors that need to be considered in the design of delivery vectors, including the overcoming of pulmonary mucosal barriers, specific targeting of pulmonary epithelial and immune cells, as well as co-delivery of adjuvants and/or cytokines with antigens. We also highlight some significant achievements of lipid-based pulmonary mucosal nanovaccines in literature. Finally, we provide our perspectives on the future possibilities and challenges in pulmonary delivery technologies as well as in the development of mucosal nanovaccines.


## 2. Formulations and methods for pulmonary delivery

Unlike parenteral vaccination, pulmonary immunization does not need the use of a needle and syringe, thus offering improved safety, lowered cost per dose, reduced pain for patients, and increased feasibility of mass vaccinations. Formulations for pulmonary delivery can be (1) solutions/suspensions that can be administered by nebulizers or pressurized metered dose inhalers, or (2) dry powder formulations that administered by active or passive pressurized dry powder

inhalers (DPIs). The nebulizers and inhalers have their unique advantages. In general, nebulizers can consistently deliver high doses of drugs with deeper deposition, while inhalers are portable and less expensive. The most important requirement for the formulations is that antigens are present in a particulate form and the particles can be dispersed in fine aerosol with an aerodynamic particle size below 5  $\mu\text{m}$ .<sup>20</sup> Details of the pulmonary delivery technologies and devices have been reviewed elsewhere.<sup>20–22</sup>

For liquid preparations, stability is a major challenge due to the high molecular mobility, which requires the use of cold chains for storage and transportation. One solution to enhance the stability is to carefully adjust the pH and osmolarity of the vehicles, as well as the ionic strength of the buffering salts.<sup>20,23</sup> In most cases, the vaccines remain their maximum activity in the pH range of 5–8, and have a better stability at low osmolarity of the vehicles as well as relatively low ionic strength ( $I < 0.15$ ) of the buffering salts.<sup>24,25</sup> Another solution is to first freeze dry the vaccines and then reconstitute them before use. In this approach, both antigen and delivery vectors need to avoid intermolecular reactions and remain stable during the freezing and dehydration. Recently, the Anderson group synthesized hyperbranched poly(beta amino esters) (hPBAs), hC32-118 and hDD90-118 (Fig. 3a), and loaded them with firefly luciferase-encoding mRNA.<sup>26</sup> The resulting mRNA-hPBAs polyplexes (0.5 mg mL<sup>-1</sup> mRNA) were nebulized to mice by using a vibrating mesh nebulizer connected to a






**Fig. 3** Pulmonary delivery with different devices and formulations. (a–e) Delivery of liquid formulations with a nebulizer. (a) Synthesis scheme of hDD90-118 and hC32-118 hyperbranched poly(beta amino esters) (PBAEs). (b) A vibrating mesh nebulizer connected to a whole-body chamber was used to deliver *in vitro* transcribed (IVT)-mRNA encoding for firefly luciferase to mice. The nebulizer generates micrometer sized droplets optimal for lung deposition, containing nanoparticles for intracellular delivery. (c) Electron microscopy of hDD90-118 particles before (left) and after (right) nebulization, particles had an average size of 137 nm ( $\pm 21$ ) and 146 nm ( $\pm 40$ ), respectively ( $\pm$ SD,  $n = 13-15$ ). (d) Particles have narrow size distribution with polydispersity indices of 0.10 before (black) and 0.11 after (red, dashed) nebulization. (e) Bioluminescence 24 h after inhalation of polyplexes, hDD90-118 vectors produced significantly higher level of radiance ( $\pm$ SD,  $n = 4$ ) and luciferase production ( $***P < 0.001$ ,  $\pm$ SD,  $n = 5-6$ ) in the lungs, compared to hC32-118 and bPEI. Statistical analysis using one-way ANOVA with *post-hoc* Tukey test. Reproduced with permission.<sup>19</sup> Copyright 2019 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim. (f–i) Delivery of dry powder formulations with a dry powder inhaler (DPI). (f) Schematic illustration of the preparation, thin-film freeze-drying (TFFD), and pulmonary delivery of aerosolizable siRNA-encapsulated solid lipid nanoparticles (LNPs). (g) Representative scanning electron microscope (SEM) images of dry powders of LNPs prepared by TFFD (upper panel) or spray drying (lower panel). The scale bar indicates 2  $\mu$ m. (h) A comparison of siRNA-encapsulated LNPs particle size distribution by intensity. (i) Transmission electron microscopy (TEM) images of siRNA-encapsulated LNPs before (left panel) and after (right panel) they were subjected to TFFD and reconstitution. Reproduced with permission.<sup>32</sup> Copyright 2021, Elsevier B.V.

whole-body chamber (Fig. 3b). This nebulization process didn't change the average size and polydispersity index (PDI) of mRNA-hPBAE polyplexes (Fig. 3c and d). In control groups, naked firefly luciferase-encoding mRNA or branched 25 kDa polyethylenimine (bPEI)-packed mRNA were subjected to nebulized delivery to mice. After 24 hours, all mice were sacrificed and major organs were collected for bioluminescence imaging. As shown in Fig. 3e, positive signals were only localized in the lungs but not in the spleen and liver, which was significantly

different from the biodistribution after intravenous injection. The luminescent radiance in lungs was quantified to be  $4.8 \times 10^5$  p s<sup>-1</sup> cm<sup>-2</sup> sr<sup>-1</sup> and  $4.3 \times 10^4$  p s<sup>-1</sup> cm<sup>-2</sup> sr<sup>-1</sup> in the hDD90-118 and hC32-118 groups, respectively, which were much higher than that in the bPEI control group ( $2.9 \times 10^4$  p s<sup>-1</sup> cm<sup>-2</sup> sr<sup>-1</sup>) (Fig. 3e). In good agreement with the radiance values, mRNA-hPBAE polyplexes induced a more efficient luciferase protein production than mRNA-bPEI (Fig. 3e). This study demonstrated that hPBAE is an effective pulmonary







**Fig. 4** PEGylation strategy to facilitate overcoming of mucosal barriers to delivery vectors. (a) Representative TEM images of tetra(piperazino)fullerene epoxide (TPFE)-based delivery nanoparticles (scale bars = 100 nm). (b) Hydrodynamic diameter changes in a mucin solution at 37 °C over time ( $n = 3$ ). (c) Representative confocal images demonstrating distribution of transgene expression mediated by naked pZG, pZG/TPFE and pZG/TPFE-PEG in the lungs of wild-type C57BL/6 and *Scnn1b*-Tg mice after intratracheal administration. Scale bar = 100  $\mu$ m. (d) Percentages of cells with reporter transgene expression according to images in (c) ( $n = 3$ ) and overall level of transgene expression determined by homogenate-based luciferase assay ( $n = 5$ ) in the lungs of both wild-type C57BL/6 and *Scnn1b*-Tg mice. \*\*\*\* $P < 0.001$ .<sup>47</sup> Copyright 2021, WILEY-VCH GmbH.

expression in lung tissue sections of pZG/TPFE-PEG-treated wild-type C57BL/6 mice (>65% coverage). In contrast, negligible fluorescent signals were observed in both pZG and pZG/TPFE control groups (<2% coverage) (Fig. 4c and d). These results well agreed with the overall level of transgene expression that was measured by lung homogenate-based luciferase assays. Pulmonary delivery of pGL3/TPFE-PEG induced two orders of magnitude higher luciferase activity compared to naked pGL3 or pGL3/TPFE in lungs of wild-type C57BL/6 mice (Fig. 4d). Similar results were observed in a chronic lung disease transgenic *Scnn1b*-Tg mouse model, which featured mucus accumulation and stasis. Although encountering more challenging delivery barriers, the TPFE-PEG delivery system remained their mucosal penetration capacity and induced a widespread (~35% coverage) as well as high-level overall transgene expression in the lungs of *Scnn1b*-Tg mice (Fig. 4c and d). Collectively, PEG surface coating could endow nanoparticles with the capacity to overcome essential mucosal barriers in the lungs, even in pathological conditions characterized by the strengthened barrier properties.


However, success in mucosal barrier crossing doesn't necessarily guarantee potent mucosal immune protection. Without proper targeting, the cargo antigens may not induce

efficient cellular internalization and antigen cross-presentation. Undoubtedly, targeting APCs in the lungs is an attractive approach to address this problem. The target receptors that have been extensively explored in parenteral nanovaccines can apply to the design of mucosal nanovaccines, for example, the well-characterized dendritic cell (DC) receptor CD205<sup>50</sup> and mannose receptor.<sup>51,52</sup> In addition to APCs, pulmonary epithelial cells represent another important cell component for targeting mucosal nanovaccines. In particular, the neonatal Fc receptor (FcRn), which is overexpressed on some type I mucosal epithelial cells and involved in the bidirectionally transepithelial transport of IgG and proteins,<sup>53</sup> providing great opportunities for targeted delivery of protein-based or protein hitchhiking-mediated nanovaccines.<sup>13,54</sup> Moreover, M cell-targeted strategy can achieve highly effective mucosal immune responses.<sup>42,55,56</sup> The most widely investigated M cell-targeting ligand is *ulex europaeus* agglutinin 1 (UEA-1),<sup>56,57</sup> which specifically bind to  $\alpha$ -L-fucose residues on the surfaces of murine intestinal M cells. Nevertheless, the targeting ligands of pulmonary M cells were seldomly reported. To identify an effective pulmonary M-cell targeting strategy, the M cell characteristics should be taken into consideration as highlighted by Longet *et al.* in 2018.<sup>55</sup>



Despite effective delivery, mucosal nanovaccines with antigens alone may not induce sufficiently strong immune responses for long-lasting protection against viruses associated with mucosal diseases.<sup>58,59</sup> Co-delivery of adjuvants/immunostimulators with antigens could significantly potentiate mucosal immunization.<sup>54,60</sup> This strategy has been verified by using polymeric nanoparticle delivery systems, such as poly-(D, L-lactide-co-glycolide) (PLGA), poly(lactic acid) (PLA), and chitosan, which could simultaneously act as mucosal adjuvants and delivery vectors in nanovaccines.<sup>56,61,62</sup> In most cases, however, adjuvants need to be delivered separately.<sup>60</sup> For example, a phosphatidylserine (PS)-coated and stimulator of interferon genes (STING) agonist cyclic guanosine monophosphate-adenosine monophosphate (cGAMP)-loaded liposome (NP-cGAMP) mucosal nanovaccine was able to target pulmonary APCs and boost anticancer immunity against lung metastases (Fig. 5a).<sup>63</sup> Notably, in NP-cGAMP, the exposed PS functioned as the “eat-me” signal, which could be recognized and internalized by APCs through their PS receptors. The cGAMP was

expected to release in cytosol, bind to STING in APCs, stimulate STING signaling and type I interferons production, and finally promote pro-inflammatory tumor microenvironment for increased immune protection. The NP-cGAMP and its counterparts had an average diameter of around 120 nm (Fig. 5b and c). After inhalation, the aerosolized NP-cGAMP predominately resided in lung tissues at a relatively high concentration for at least 48 hours, which was attributed to the PS-mediated APC targeting (Fig. 5d). Furthermore, pulmonary delivery of NP-cGAMP synergized with fractionated radiation (IR) could successfully activate anticancer immunity in a B16F10-OVA tumor metastasis model. In animal studies, fractionated IR was only delivered to the right lung of mice, which resulted in successful tumor inhibition in the right lung but little therapeutic efficacy against metastatic foci in the non-irradiated left lung. The mice treated with inhalable NP-cGAMP caused a moderate reduction in the number of tumor foci in both lungs. Impressively, combination of NP-cGAMP and fractionated IR (right lung only) induced almost complete



**Fig. 5** APC-targeting and immunostimulator co-delivery strategies to improve protective immunity at mucosal sites. (a) Schematic illustration of the inhalable phosphatidylserine-coated and STING agonist cyclic guanosine monophosphate-adenosine monophosphate (cGAMP)-loaded liposomes (NP-cGAMP) for enhancing antitumor immunity against lung metastases. APC, antigen presenting cell. DLN, draining lymph node. IR, irradiation. TA, tumor antigen. (b) Size distribution, polydispersity index, and surface charge of NP-cGAMP, NP labeled with DIR (NP-DIR), and NP labeled with Rhob (NP-Rhob) by dynamic light scattering (DLS). (c) TEM images of NP-cGAMP. (d) Representative *ex vivo* fluorescence imaging of major organs dissected from 4T1-luc lung metastases-bearing mice at 1, 24, and 48 h post inhalation of DiR-labeled PS-coated NPs. Light signals were exclusively from both lungs and the lung signals were quantified ( $n = 3$  biologically independent mice/time;  $***P < 0.01$ , Student's *t* test). (e) NP-cGAMP inhalation synergized with radiotherapy by eliciting APC-mediated adaptive immunity in B16F10-OVA melanoma lung metastasis model. On day 5, after confirming development of multifocal metastases in both lungs, the mice were treated with fractionated radiation to the right lung (IR, 8 Gy  $\times$  3), inhalation of NP-cGAMP (24 h after each IR for three doses), or a combination of them. NP-CTR (2'5'-GpAp) served as a control of NP-cGAMP. To deplete pulmonary APCs, NP-clodronate was administered *via* inhalation 6 h before each of the three NP-cGAMP inhalations. To deplete CD4<sup>+</sup> or CD8<sup>+</sup> T cells, anti-CD4 Ab or anti-CD8 $\alpha$  Ab was injected intraperitoneal (i.p.) (400  $\mu$ g), respectively, one day before IR and repeated 7 days later. The mice ( $n = 6$ ) were sacrificed on day 18 and representative lungs from the treatment groups were displayed. (f) Lungs after treatments in (e) were examined under a dissecting microscope and a total of metastatic lung foci on each lung were counted. Data are shown as mean  $\pm$  SD of  $n = 6$  biologically independent samples.  $*P < 0.05$ ,  $**P < 0.01$ , and  $***P < 0.001$  by Student's *t* test. Reproduced with permission.<sup>63</sup> Copyright 2019, Nature Publishing Group.



tumor regression in both lungs (Fig. 5e and f). These data demonstrated that pulmonary delivery of immunostimulators, especially by using APC-targeting delivery vectors, could synergize with *in situ* tumor antigen release and enhance protective immunity against tumor development at the lung tissue. However, this study didn't take mucosal immune responses into careful considerations when exploring the mechanisms behind the synergistic therapeutic effects. A more comprehensive analysis of the protective immunity would be needed before applying the NP-cGAMP delivery strategy to other pulmonary nanovaccines. Although adjuvants could potentiate antigen-specific immune responses, an improper adjuvant may result in severe side effects. For example, an enterotoxin-adjuvanted nasal influenza vaccine caused a substantial number of Bell's palsy in a clinical trial, which was attributed to the high binding affinity between enterotoxin and GM1 ganglioside moieties of olfactory nerves.<sup>64–66</sup> Pollen grains have been reported as an adjuvant to elicit allergic immune response and promote DC maturation,<sup>67</sup> which therefore should be taken into consideration for the design of pulmonary vaccines. A pulmonary vaccine with too strong adjuvants may be at higher risks for triggering allergy, especially when it is administrated in pollen seasons.

#### 4. Pulmonary nanovaccines for protective immunity against mucosal diseases

As more extensively reviewed elsewhere, synthetic nanoparticles varying from LNPs, liposomes, polymeric nanoparticles, and hybrid nanoparticles, can be delivered *via* nasal, oral, sublingual, or colorectal routes for mucosal nanovaccines.<sup>42,68,69</sup> These nanoparticles may also find broad applications in pulmonary mucosal nanovaccines. For example, aerosolized PLGA and PLA nanoparticles were used for pulmonary delivery of hepatitis B vaccine, which elicited enhanced humoral, mucosal and cytokine responses in comparison to the delivery of free antigens.<sup>70</sup> In this section, inspired by the tremendous success of LNPs in parenteral nanovaccines for COVID-19, an infectious disease at pulmonary mucosal site, we will focus our discussion on lipid-based pulmonary nanovaccines.

Liposome represents a paradigm of delivery vectors in the clinical translation of pulmonary mucosal nanovaccines. Very recently, Ming *et al.* prepared a bionic virus-like nanovaccine that comprised viral “capsid”-mimicking pulmonary surfactant liposomes as delivery vectors, viral genetic material-mimicking poly(I:C) (polyinosinic-polycytidylic acid) as adjuvants, and “spike”-mimicking receptor binding domain (RBD) proteins on the liposome surfaces as antigens. This inhalable mucosal vaccine elicited high titer of sIgA in respiratory secretions, effectively neutralized intranasally delivered pseudovirus, and demonstrated a much stronger mucosal protective immune responses than subcutaneous vaccination.<sup>71</sup> It is worth noting that the lamellarity of liposomes can greatly impact the vaccine

immunogenicity.<sup>19</sup> Pulmonary immunization with interbilayer-crosslinked multilamellar liposome vesicles (ICMVs) were reported to activate high-frequency, long-lived, and antigen specific effector memory T cells ( $T_{EMs}$ ) at multiple mucosal sites.<sup>19</sup> The ICMVs were first reported by the Irvine group in 2011.<sup>72</sup> They were composed of phospholipids 1,2-dioleoyl-*sn*-glycero-3-phosphocholine (DOPC), anionic 1,2-di-(9Z-octadecenoyle)-snglycero-3-phospho-(1'-*rac*-glycerol) (DOPG), and anionic maleimide-headgroup lipid 1,2-dioleoyl-*sn*-glycero-3-phosphoethanolamine-*N*-[4-(*p*-maleimidophenyl) butyramide] (MPB) at a molar ratio of 4 : 1 : 5. To synthesize them, simple liposomes were first prepared and fused to multilamellar vesicles (MLVs), in which functionalized lipid headgroups of adjacent bilayers were crosslinked to generate the ICMVs (Fig. 6a). The ICMVs allowed stable entrapment of protein antigens in the core and lipid-based immunostimulators in their multi-layer walls under extracellular conditions and rapid release of the cargoes in cell endosomes, therefore promoting antigen cross-presentation and eliciting robust humoral and cellular immune responses upon subcutaneous injection.<sup>72</sup> In their following-up study, ICMVs were employed to pack antigen OVA and two TLR agonists, monophosphoryl lipid A (MPLA) and poly(I:C), for pulmonary vaccination. To investigate the distant mucosal immunity, luciferase-expressing OT-1 CD8<sup>+</sup> T cells (OT-1-luc) were intravenously transferred into mice twenty-four hours prior to intratracheal or subcutaneous administration of ICMV or soluble vaccines. According to bioluminescence imaging on day 3, T cell proliferation was only observed in draining lymph nodes (dLNs) in all vaccinated mice. In contrast, by day 5, higher T cell expansion was observed in the pulmonary ICMV group, showing bioluminescent signals in not only lymph nodes and spleen but also reproductive tracts and gut (Fig. 6b). These results indicated the unique mucosal homing property of T cell responses in mucosal immunization, which was attributed to the substantial expression of mucosal homing integrin  $\alpha_4\beta_7$  receptors in the tetramer<sup>+</sup> peripheral blood OT-1 cells after pulmonary vaccination (Fig. 6c). Moreover, pulmonary ICMV vaccination biased T cell responses toward effector memory phenotype at both local and distal mucosal tissues. When tested in subcutaneous OVA-expressing B16F10 mouse melanoma tumor model, ICMV OVA pulmonary vaccination led to complete tumor rejection and long-term survival (over 18 weeks) of all animals. In contrast, soluble OVA pulmonary vaccination delayed tumor growth but failed to improve ultimate survival of mice (Fig. 6d). Similar results were observed in prophylactic mucosal viral challenge models (Fig. 6e). Collectively, pulmonary ICMV nanovaccine could prime  $T_{EMs}$  that circulated to both local and distal mucosal tissues as well as systemic compartments, therefore providing robust protection in both therapeutic tumor and prophylactic viral infection settings. Further evaluation on the safety and efficacy of ICMV mucosal nanovaccines that carrying antigens of relevant mucosal disease, such as mucosal tumors and HIV, would provide significant guidance for their bench-to-bed translation.

In addition to liposome nanoparticles, lipid conjugates with “albumin hitching” property have demonstrated to effec-







**Fig. 6** Liposome nanoparticle-based pulmonary vaccine. (a) Schematic illustration of interbilayer-crosslinked multilamellar vesicles (ICMVs) synthesis and cryoelectron microscope images: (i) anionic, maleimide-functionalized liposomes are prepared from dried lipid films, (ii) divalent cations are added to induce fusion of liposomes and the formation of multilamellar vesicles (MLVs), (iii) membrane-permeable dithiols are added, which crosslink maleimide-lipids on apposed lipid bilayers in the vesicle walls, and (iv) the resulting lipid particles are PEGylated with thiol-terminated PEG. Cryo-EM images from each step of the synthesis show (i) initial liposomes, (ii) MLVs, and (iii) ICMVs with thick lipid walls. Scale bars = 100 nm. Right-hand image of (iii) shows a zoomed image of an ICMV wall, where stacked bilayers are resolved as electron-dense striations; scale bar = 20 nm. Reproduced with permission.<sup>49</sup> Copyright 2011, Nature Publishing Group. (b) OT-I-luc CD8<sup>+</sup> T cells were adoptively transferred into C57BL/6 mice (n = 5) 1 day before ICMV or soluble OVA vaccine immunization via intratracheal or subcutaneous routes. Trafficking and proliferation of OT-I-luc T cells were monitored by flow cytometry and bioluminescence imaging on days 3 and 5 after immunization. Flow cytometry histograms show representative CFSE dilutions in transferred T cells on day 3 after immunization at draining lymph nodes (dLNs) (mdLNs for intratracheal vaccines and ingLNs for subcutaneous vaccinations). Lungs (L) and gastrointestinal tracts (G) were dissected on day 5 and imaged to identify T cell localization. V, vaginal tract; PP, Peyer's patches. (c) Flow cytometry analyses of integrin  $\alpha_4\beta_7$  OT-I T cells in blood on day 5. (d) C57BL/6 mice (n = 5 to 6 per group) inoculated with  $5 \times 10^5$  B16F10-OVA cells subcutaneously in the flank on day 0 were treated on days 3 and 10 with intratracheal administration of ICMVs or soluble OVA vaccines. Survival of tumor-bearing mice was tracked for 18 weeks. \*P < 0.05, log-rank (Mantel-Cox) test. (e) C57BL/6 mice were immunized subcutaneously or intratracheally on days 0 and 28 with antigen AL11 (AAVKNWMTQTL) and the universal CD4<sup>+</sup> T cell helper epitope PADRE peptide ICMV or soluble vaccines, and then challenged by intratracheal administration of simian immunodeficiency virus (SIV) gag-expressing vaccinia virus ( $1 \times 10^6$  PFU) on day 42. Body weight changes were recorded over time. Dots show titers from individual animals. Data are means  $\pm$  SEM with n = 3 to 7 animals per group. \*P < 0.5, \*\*\*P < 0.01, \*\*\*\*P < 0.001, by one-way ANOVA (c) or two-way ANOVA (e). Reproduced with permission.<sup>19</sup> Copyright 2013, American Association for the Advancement of Science.

tively navigate amphiphilic phospholipid-coupling peptide antigens or molecular adjuvants (amph-vaccines) to dLNs.<sup>33,73,74</sup> In this strategy, the lipid tail of the amph-vaccines served as an albumin-binding domain. Upon injection, endogenous albumin was bound to and chaperoned amph-vaccines to dLNs, and then transferred the lipid tails from albumin to insert in cell membranes.<sup>33,74</sup> The Irvine group explored this amph-vaccine approach in pulmonary vaccination on the basis that (1) albumin is abundant in the bronchoalveolar lavage (BAL) fluid of naïve mice and its concentration can be rapidly increased from approximate 200  $\mu$ g

$\text{mL}^{-1}$  to a few  $\text{mg mL}^{-1}$  after pulmonary administration of the TLR9 agonist CpG; (2) FcRn is highly expressed on pulmonary epithelial cells, which can facilitate the transportation of albumin across the airway epithelium.<sup>13</sup> The amph-vaccines comprised a PEG spacer linked with a melanoma peptide antigen gp100<sub>20-39</sub> and 1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine (DSPE) phospholipid tail as well as a similar C18 lipid tail-coupled adjuvant CpG DNA (Fig. 7a). The amph-gp100 conjugates led to about 10 times higher accumulation in the lung parenchyma than free gp100 peptide 24 h after intratracheal administration, and remained easily detectable





**Fig. 7** Albumin-binding lipid-based pulmonary vaccine. (a) Structure of amph-vaccines. Upper: schematic of 1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine (DSPE) conjugated to cysteine(Cys)-terminated peptide via a polyethylene glycol (PEG) spacer. Lower: schematic of diacyl lipid tail conjugated to CpG. (b) Representative immunofluorescence images of the lungs and mediastinal lymph nodes (MLNs) harvested from C57BL/6 mice ( $n = 3$ ) at 48 hours after immunization with 20  $\mu\text{g}$  of fluorescein amidite (FAM)-labeled (green) gp100 peptide or an equimolar amount of labeled amph-gp100. Scale bars, 100  $\mu\text{m}$ . DAPI (blue) was used to stain nuclei. (c) Quantification of peptide signals from the lungs at 24 and 48 hours after the immunization in (b). (d) Quantification of peptide signals from the MLNs at 48 hours after the immunization in (b). (e) C57BL/6 mice ( $n = 5$ ) were vaccinated with 5  $\mu\text{g}$  of AL11 peptide combined with 8  $\mu\text{g}$  of CpG or equimolar doses of amph-AL11 and amph-CpG administered intratracheally or subcutaneously on days 0 and 28 and then challenged with a lethal dose of SIV-Gag Vaccinia virus ( $10^6$  PFU) intratracheally on day 150. (f) Animal weights over time after Vaccinia virus challenge in (e). (g) Animal survival after Vaccinia virus challenge in (e). (h) C57BL/6 mice ( $n = 10$ ) were left naïve or vaccinated with 10  $\mu\text{g}$  of gp100, 10  $\mu\text{g}$  of Trp1, and 10  $\mu\text{g}$  of Trp2 peptide antigens combined with 8  $\mu\text{g}$  of CpG or vaccinated with equimolar doses of amph-peptides and amph-CpG on days 0 and 14. Thirty days after boost, mice were challenged with  $4 \times 10^5$  B16F10 melanoma cells intravenously, and survival over time is plotted. In each cohort, the number of mice surviving at 80 days was indicated. Data are represented as means  $\pm$  SEM from three independent experiments (e–g) and two independent experiments (b–d and h); \* $P < 0.05$ , \*\* $P < 0.001$ , and \*\*\*\* $P < 0.0001$  as determined by one-way ANOVA. Survival data were analyzed using a log-rank test. Reproduced with permission.<sup>13</sup> Copyright 2021, American Association for the Advancement of Science.

in lungs and mediastinal lymph nodes (MLNs) at 48 h (Fig. 7b–d). Similar results were found in the amph-CpG conjugates. Importantly, pulmonary vaccination with amph-vaccines stimulated prolonged antigen presentation in MLNs for over two weeks, high levels of antigen-specific lung  $T_{\text{RM}}$ s, and enhanced protection against viral and tumor challenge. In a prophylactic viral infection model, mice were first intratracheally immunized with amph-AL11/amph-CpG vaccines, in which AL11 is a murine gag peptide. After five months, the mice were challenged with a lethal dose of gag epitope-expressing Vaccinia virus (Fig. 7e). All mice immunized with pulmonary amph-vaccines survived. In control groups, mice kept losing body weight and succumbed by day 7 (Fig. 7f and g). To test the protective capacity in tumor models, the antigens of

the amph-vaccines in Fig. 7e were replaced with a combination of three melanoma peptides (Trp1<sub>455–463</sub>, Trp2<sub>180–188</sub>, and gp100<sub>20–39</sub>) and mice were challenged 30 days after full vaccination by intravenously injecting B16F10 melanoma cells. According to survival curves in Fig. 7h, 80% mice in the pulmonary vaccine group survived at 100 days after the tumor inoculation, while only 40% mice were protected by subcutaneous vaccination. Taken together, pulmonary amph-vaccines primed long-lived antigen-specific  $T_{\text{RM}}$ s in the lungs, generated robust mucosal immunity, and effectively protected against viral or tumor challenge.

LNPs are another important class of lipid-based nanoparticles. LNPs are usually constructed by four lipid components: (1) ionizable/cationic lipids that protect the cargoes





SARS-CoV-2 RNA in lung tissue, mitigated influenza virus A or SARS-CoV-2 replication, and reduced disease symptoms in rodents.<sup>27</sup> Besides, intratracheal administration of multiplexing activity-based nanosensors achieved sensitive and specific lung cancer detection.<sup>84</sup> Combination of pulmonary nanovaccines with these applications may provide new solutions for the management of mucosal diseases. Taken together, pulmonary mucosal nanovaccines will provide powerful tools for preventing infection and transmission of mucosal pathogens, as well as for the treatment of pre-existing diseases such as infections and cancer at mucosal sites and other distant tissues.

## Conflicts of interest

G. Z. is a co-founder and CSO of AmpedRNA Biosciences, LLC, which develops novel vaccine technologies for cancer and other diseases.

## Acknowledgements

The work was supported by the National University of Singapore Start-Up funding (W. T.). G. Z. acknowledges support from NIH-NIGMS (R35GM143014), NIH-NINDS (R21NS114455), NIH-NCATS KL2 Scholarship (KL2TR002648) and Endowment Fund from VCU C. Kenneth and Dianne Wright Center for Clinical and Translational Research (UL1TR002649), and start-up fund from the Center for Pharmaceutical Engineering and Sciences-VCU School of Pharmacy.

## References

- 1 L. R. Baden, H. M. El Sahly, B. Essink, K. Kotloff, S. Frey, R. Novak, D. Diemert, S. A. Spector, N. Rouphael, C. B. Creech, J. McGettigan, S. Khetan, N. Segall, J. Solis, A. Brosz, C. Fierro, H. Schwartz, K. Neuzil, L. Corey, P. Gilbert, H. Janes, D. Follmann, M. Marovich, J. Mascola, L. Polakowski, J. Ledgerwood, B. S. Graham, H. Bennett, R. Pajon, C. Knightly, B. Leav, W. Deng, H. Zhou, S. Han, M. Ivarsson, J. Miller, T. Zaks and C. S. Group, *N. Engl. J. Med.*, 2021, **384**, 403.
- 2 F. P. Polack, S. J. Thomas, N. Kitchin, J. Absalon, A. Gurtman, S. Lockhart, J. L. Perez, G. P. Marc, E. D. Moreira, C. Zerbini, R. Bailey, K. A. Swanson, S. Roychoudhury, K. Koury, P. Li, W. V. Kalina, D. Cooper, R. W. Frencck Jr., L. L. Hammitt, O. Tureci, H. Nell, A. Schaefer, S. Unal, D. B. Tresnan, S. Mather, P. R. Dormitzer, U. Sahin, K. U. Jansen, W. C. Gruber and C. C. T. Group, *N. Engl. J. Med.*, 2020, **383**, 2603.
- 3 E. J. Anderson, N. G. Rouphael, A. T. Widge, L. A. Jackson, P. C. Roberts, M. Makhene, J. D. Chappell, M. R. Denison, L. J. Stevens, A. J. Pruijssers, A. B. McDermott, B. Flach, B. C. Lin, N. A. Doria-Rose, S. O'Dell, S. D. Schmidt, K. S. Corbett, P. A. Swanson, M. Padilla, K. M. Neuzil, H. Bennett, B. Leav, M. Makowski, J. Albert, K. Cross, V. V. Edara, K. Floyd, M. S. Suthar, D. R. Martinez, R. Baric, W. Buchanan, C. J. Luke, V. K. Phadke, C. A. Rostad, J. E. Ledgerwood, B. S. Graham and J. H. Beigel, *N. Engl. J. Med.*, 2020, **383**, 2427.
- 4 E. C. Lavelle and R. W. Ward, *Nat. Rev. Immunol.*, 2021, DOI: 10.1038/s41577-021-00583-2.
- 5 J. Holmgren and C. Czerkinsky, *Nat. Med.*, 2005, **11**, S45–S53.
- 6 M. R. Neutra and P. A. Kozlowski, *Nat. Rev. Immunol.*, 2006, **6**, 148.
- 7 C. J. Wei, M. C. Crank, J. Shiver, B. S. Graham, J. R. Mascola and G. J. Nabel, *Nat. Rev. Drug Discovery*, 2020, **19**, 239.
- 8 S. Nambulli, Y. Xiang, N. L. Tilston-Lunel, L. J. Rennick, Z. Sang, W. B. Klimstra, D. S. Reed, N. A. Crossland, Y. Shi and W. P. Duprex, *Sci. Adv.*, 2021, **7**, eabh0319.
- 9 J. A. Whittsett and T. Alenghat, *Nat. Immunol.*, 2015, **16**, 27.
- 10 Z. Zuo, Y. Zou, Q. Li, Y. Guo, T. Zhang, J. Wu, C. He and F. O. Eko, *Sci. Rep.*, 2021, **11**, 10389.
- 11 D. Sterlin, A. Mathian, M. Miyara, A. Mohr, F. Anna, L. Claer, P. Quentric, J. Fadlallah, H. Devilliers, P. Ghillani, C. Gunn, R. Hockett, S. Mudumba, A. Guihot, C. E. Luyt, J. Mayaux, A. Beurton, S. Fourati, T. Bruel, O. Schwartz, J. M. Lacorte, H. Yssel, C. Parizot, K. Dorgham, P. Charneau, Z. Amoura and G. Gorochoy, *Sci. Transl. Med.*, 2021, **13**, eabd2223.
- 12 G. Stary, A. Olive, A. F. Radovic-Moreno, D. Gondek, D. Alvarez, P. A. Basto, M. Perro, V. D. Vrbanac, A. M. Tager, J. Shi, J. A. Yethon, O. C. Farokhzad, R. Langer, M. N. Starnbach and U. H. Von Andrian, *Science*, 2015, **348**, aaa8205.
- 13 K. Rakhra, W. Abraham, C. Wang, K. D. Moynihan, N. Li, N. Donahue, A. D. Baldeon and D. J. Irvine, *Sci. Immunol.*, 2021, **6**, eabd8003.
- 14 A. O. Hassan, N. M. Kafai, I. P. Dmitriev, J. M. Fox, B. K. Smith, I. B. Harvey, R. E. Chen, E. S. Winkler, A. W. Wessel, J. B. Case, E. Kashentseva, B. T. McCune, A. L. Bailey, H. Zhao, L. A. VanBlargan, Y. N. Dai, M. Ma, L. J. Adams, S. Shrihari, J. E. Danis, L. E. Gralinski, Y. J. Hou, A. Schafer, A. S. Kim, S. P. Keeler, D. Weiskopf, R. S. Baric, M. J. Holtzman, D. H. Fremont, D. T. Curiel and M. S. Diamond, *Cell*, 2020, **183**, 169.
- 15 D. Hartl, R. Tirouvanziam, J. Laval, C. M. Greene, D. Habiell, L. Sharma, A. O. Yildirim, C. S. Dela Cruz and C. M. Hogaboam, *J. Innate Immun.*, 2018, **10**, 487.
- 16 S. U. Islam, A. Shehzad, M. B. Ahmed and Y. S. Lee, *Molecules*, 2020, **25**, 1929.
- 17 L. A. Keller, O. Merkel and A. Popp, *Drug Delivery Transl. Res.*, 2021, **11**, 1.
- 18 D. Hartl, R. Tirouvanziam, J. Laval, C. M. Greene, D. Habiell, L. Sharma, A. O. Yildirim, C. S. Dela Cruz and C. M. Hogaboam, *J. Innate Immun.*, 2018, **10**, 487.
- 19 A. V. Li, J. J. Moon, W. Abraham, H. Suh, J. Elkhader, M. A. Seidman, M. Yen, E. J. Im, M. H. Foley,



- D. H. Barouch and D. J. Irvine, *Sci. Transl. Med.*, 2013, **5**, 204ra130.
- 20 M. Hellfritzsch and R. Scherließ, *Pharmaceutics*, 2019, **11**, 375.
- 21 W. Liang, H. W. Pan, D. Vllasaliu and J. K. W. Lam, *Pharmaceutics*, 2020, **12**, 1025.
- 22 C. A. Ruge, J. Kirch and C. M. Lehr, *Lancet Respir. Med.*, 2013, **1**, 402.
- 23 T. E. Rajapaksa, K. M. Bennett, M. Hamer, C. Lytle, V. G. Rodgers and D. D. Lo, *J. Biol. Chem.*, 2010, **285**, 23739.
- 24 W. M. Colwell, D. G. Simmons, J. R. Harris, T. G. Fulp, J. H. Carrozza and T. A. Maag, *Avian Dis.*, 1975, **19**, 781.
- 25 D. T. Brandau, L. S. Jones, C. M. Wiethoff, J. Rexroad and C. R. Middaugh, *J. Pharm. Sci.*, 2003, **92**, 218.
- 26 A. K. Patel, J. C. Kaczmarek, S. Bose, K. J. Kauffman, F. Mir, M. W. Heartlein, F. DeRosa, R. Langer and D. G. Anderson, *Adv. Mater.*, 2019, **31**, e1805116.
- 27 E. L. Blanchard, D. Vanover, S. S. Bawage, P. M. Tiwari, L. Rotolo, J. Beyersdorf, H. E. Peck, N. C. Bruno, R. Hincapie, F. Michel, J. Murray, H. Sadhwani, B. Vanderheyden, M. G. Finn, M. A. Brinton, E. R. Lafontaine, R. J. Hogan, C. Zurla and P. J. Santangelo, *Nat. Biotechnol.*, 2021, **39**, 717.
- 28 Z. Li, Y. Wang, Y. Shen, C. Qian, D. Oupicky and M. Sun, *Sci. Adv.*, 2020, **6**, eaaz9240.
- 29 A. Minne, H. Boireau, M. J. Horta and R. Vanbever, *Eur. J. Pharm. Biopharm.*, 2008, **70**, 839.
- 30 J. Raula, F. Thielmann, M. Naderi, V. P. Lehto and E. I. Kauppinen, *Int. J. Pharm.*, 2010, **385**, 79.
- 31 C. Weiler, M. Egen, M. Trunk and P. Langguth, *J. Pharm. Sci.*, 2010, **99**, 303.
- 32 J. L. Wang, M. S. Hanafy, H. Xu, J. Leal, Y. Zhai, D. Ghosh, R. O. Williams III, H. D. C. Smyth and Z. Cui, *Int. J. Pharm.*, 2021, **596**, 120215.
- 33 L. Ma, T. Dichwalkar, J. Y. H. Chang, B. Cossette, D. Garafola, A. Q. Zhang, M. Fichter, C. Wang, S. Liang, M. Silva, S. Kumari, N. K. Mehta, W. Abraham, N. Thai, N. Li, K. D. Wittrup and D. J. Irvine, *Science*, 2019, **365**, 162.
- 34 G. Zhu, L. Mei, H. D. Vishwasrao, O. Jacobson, Z. Wang, Y. Liu, B. C. Yung, X. Fu, A. Jin, G. Niu, Q. Wang, F. Zhang and X. Chen, *Nat. Commun.*, 2017, **8**, 1482.
- 35 G. Zhu, G. M. Lynn, O. Jacobson, K. Chen, Y. Liu, H. Zhang, Y. Ma, F. Zhang, R. Tian, Q. Ni, S. Cheng, Z. Wang, N. Lu, B. C. Yung, Z. Wang, L. Lang, X. Fu, A. Jin, I. D. Weiss, H. Vishwasrao, G. Niu, H. Shroff, D. M. Klinman, R. A. Seder and X. Chen, *Nat. Commun.*, 2017, **8**, 1954.
- 36 M. E. Aikins, C. Xu and J. J. Moon, *Acc. Chem. Res.*, 2020, **53**, 2094.
- 37 S. Yan, Z. Luo, Z. Li, Y. Wang, J. Tao, C. Gong and X. Liu, *Angew. Chem., Int. Ed.*, 2020, **59**, 17332.
- 38 C. Zhang and K. Pu, *Chem. Soc. Rev.*, 2020, **49**, 4234.
- 39 W. Sang, Z. Zhang, Y. Dai and X. Chen, *Chem. Soc. Rev.*, 2019, **48**, 3771.
- 40 Y. Zhang, S. Lin, X. Y. Wang and G. Zhu, *Wiley Interdiscip. Rev.: Nanomed. Nanobiotechnol.*, 2019, **11**, 1559.
- 41 W. Tang, W. Fan, J. Lau, L. Deng, Z. Shen and X. Chen, *Chem. Soc. Rev.*, 2019, **48**, 2967.
- 42 K. A. Woodrow, K. M. Bennett and D. D. Lo, *Annu. Rev. Biomed. Eng.*, 2012, **14**, 17.
- 43 S. K. Lai, Y.-Y. Wang and J. Hanes, *Adv. Drug Delivery Rev.*, 2009, **61**, 158.
- 44 J. K. Sheehan, K. Oates and I. Carlstedt, *Biochem. J.*, 1986, **239**, 147.
- 45 J. S. Suk, Q. Xu, N. Kim, J. Hanes and L. M. Ensign, *Adv. Drug Delivery Rev.*, 2016, **99**, 28.
- 46 Q. Xu, L. M. Ensign, N. J. Boylan, A. Schön, X. Gong, J. C. Yang, N. W. Lamb, S. Cai, T. Yu, E. Freire and J. Hanes, *ACS Nano*, 2015, **9**, 9217.
- 47 D. Chen, S. Liu, D. Chen, J. Liu, J. Wu, H. Wang, Y. Su, G. Kwak, X. Zuo, D. Rao, H. Cui, C. Shu and J. S. Suk, *Angew. Chem., Int. Ed.*, 2021, **60**, 15225.
- 48 E. Nakamura and H. Isobe, *Chem. Rec.*, 2010, **10**, 260–270.
- 49 H. Isobe, S. Sugiyama, K. I. Fukui Ki, Y. Iwasawa and E. Nakamura, *Angew. Chem., Int. Ed.*, 2001, **40**, 3364.
- 50 C. Macri, C. Dumont, A. P. Johnston and J. D. Mintern, *Clin. Transl. Immunol.*, 2016, **5**, 66.
- 51 A. R. Poh and M. Ernst, *Front. Oncol.*, 2018, **8**, 49.
- 52 D. S. Wilson, S. Hirose, M. M. Racz, L. Bonilla-Ramirez, L. Jeanbart, R. Wang, M. Kwissa, J. F. Franetich, M. A. S. Broggi, G. Diaceri, X. Quaglia-Thermes, D. Mazier, M. A. Swartz and J. A. Hubbell, *Nat. Mater.*, 2019, **18**, 175.
- 53 M. Yoshida, S. M. Claypool, J. S. Wagner, E. Mizoguchi, A. Mizoguchi, D. C. Roopenian, W. I. Lencer and R. S. Blumberg, *Immunity*, 2004, **20**, 769.
- 54 S. Jangra, J. De Vrieze, A. Choi, R. Rathnasinghe, G. Laghali, A. Uvyn, S. Van Herck, L. Nuhn, K. Deswarte, Z. Zhong, N. Sanders, S. Lienenklaus, S. David, S. Strohmeier, F. Amanat, F. Krammer, H. Hammad, B. N. Lambrecht, L. Coughlan, A. Garcia-Sastre, B. de Geest and M. Schotsaert, *Angew. Chem., Int. Ed.*, 2021, **60**, 9467.
- 55 S. Longet, M. L. E. Lundahl and E. C. Lavelle, *Bioconjugate Chem.*, 2018, **29**, 613.
- 56 M. Li, Y. Wang, Y. Sun, H. Cui, S. J. Zhu and H.-J. Qiu, *Immunol. Lett.*, 2020, **217**, 116.
- 57 L. Du, Z. Yu, F. Pang, X. Xu, A. Mao, W. Yuan, K. He and B. Li, *Front. Cell. Infect. Microbiol.*, 2018, **8**, 7.
- 58 T. Azegami, Y. Yuki, R. Nakahashi, H. Itoh and H. Kiyono, *Mol. Immunol.*, 2018, **98**, 19.
- 59 F. C. Knight, P. Gilchuk, A. Kumar, K. W. Becker, S. Sevimli, M. E. Jacobson, N. Suryadevara, L. Wang-Bishop, K. L. Boyd, J. E. Crowe Jr., S. Joyce and J. T. Wilson, *ACS Nano*, 2019, **13**, 10939.
- 60 N. Lycke, *Nat. Rev. Immunol.*, 2012, **12**, 592.
- 61 A. Muralidharan, M. S. Russell, L. Larocque, C. Gravel, S. Sauvé, Z. Chen, C. Li, W. Chen, T. Cyr, M. Rosu-Myles, L. Wang and X. Li, *Vaccine*, 2019, **37**, 4031.
- 62 L. Feng, X. R. Qi, X. J. Zhou, Y. Maitani, S. C. Wang, Y. Jiang and T. Nagai, *J. Controlled Release*, 2006, **112**, 35.
- 63 Y. Liu, W. N. Crowe, L. Wang, Y. Lu, W. J. Petty, A. A. Habib and D. Zhao, *Nat. Commun.*, 2019, **10**, 5108.



- 64 M. Mutsch, W. Zhou, P. Rhodes, M. Bopp, R. T. Chen, T. Linder, C. Spyr and R. Steffen, *N. Engl. J. Med.*, 2004, **350**, 896.
- 65 F. W. van Ginkel, R. J. Jackson, N. Yoshino, Y. Hagiwara, D. J. Metzger, T. D. Connell, H. L. Vu, M. Martin, K. Fujihashi and J. R. McGhee, *Infect. Immun.*, 2005, **73**, 6892.
- 66 J. H. Rhee, S. E. Lee and S. Y. Kim, *Clin. Exp. Vaccine Res.*, 2012, **1**, 50.
- 67 Z. Allakhverdi, S. Bouguermouh, M. Rubio and G. Delespesse, *Allergy*, 2005, **60**, 1157.
- 68 A. Thakur and C. Foged, *Nanoengineered Biomaterials for Advanced Drug Delivery*, Elsevier, 2020, pp. 603–646, DOI: 10.1016/b978-0-08-102985-5.00025-5.
- 69 V. Bernasconi, K. Norling, M. Bally, F. Höök and N. Y. Lycke, *J. Immunol. Res.*, 2016, **2016**, 1.
- 70 C. Thomas, A. Rawat, L. Hope-Weeks and F. Ahsan, *Mol. Pharm.*, 2011, **8**, 405.
- 71 B. Zheng, W. Peng, M. Guo, M. Huang, Y. Gu, T. Wang, G. Ni and D. Ming, *Chem. Eng. J.*, 2021, **418**, 129392.
- 72 J. J. Moon, H. Suh, A. Bershteyn, M. T. Stephan, H. Liu, B. Huang, M. Sohail, S. Luo, S. H. Um, H. Khant, J. T. Goodwin, J. Ramos, W. Chiu and D. J. Irvine, *Nat. Mater.*, 2011, **10**, 243.
- 73 H. Liu, K. D. Moynihan, Y. Zheng, G. L. Szeto, A. V. Li, B. Huang, D. S. Van Egeren, C. Park and D. J. Irvine, *Nature*, 2014, **507**, 519.
- 74 K. D. Moynihan, R. L. Holden, N. K. Mehta, C. Wang, M. R. Karver, J. Dinter, S. Liang, W. Abraham, M. B. Melo, A. Q. Zhang, N. Li, S. L. Gall, B. L. Pentelute and D. J. Irvine, *Cancer Immunol. Res.*, 2018, **6**, 1025.
- 75 Q. Cheng, T. Wei, L. Farbiak, L. T. Johnson, S. A. Dilliard and D. J. Siegwart, *Nat. Nanotechnol.*, 2020, **15**, 313.
- 76 M. D. Buschmann, M. J. Carrasco, S. Alishetty, M. Paige, M. G. Alameh and D. Weissman, *Vaccines*, 2021, **9**, 65.
- 77 A. Akinc, A. Zumbuehl, M. Goldberg, E. S. Leshchiner, V. Busini, N. Hossain, S. A. Bacallado, D. N. Nguyen, J. Fuller, R. Alvarez, A. Borodovsky, T. Borland, R. Constien, A. de Fougerolles, J. R. Dorkin, K. Narayanannair Jayaprakash, M. Jayaraman, M. John, V. Koteliansky, M. Manoharan, L. Nechev, J. Qin, T. Racie, D. Raitcheva, K. G. Rajeev, D. W. Sah, J. Soutschek, I. Toudjarska, H. P. Vornlocher, T. S. Zimmermann, R. Langer and D. G. Anderson, *Nat. Biotechnol.*, 2008, **26**, 561.
- 78 S. C. Semple, A. Akinc, J. Chen, A. P. Sandhu, B. L. Mui, C. K. Cho, D. W. Sah, D. Stebbing, E. J. Crosley, E. Yaworski, I. M. Hafez, J. R. Dorkin, J. Qin, K. Lam, K. G. Rajeev, K. F. Wong, L. B. Jeffs, L. Nechev, M. L. Eisenhardt, M. Jayaraman, M. Kazem, M. A. Maier, M. Srinivasulu, M. J. Weinstein, Q. Chen, R. Alvarez, S. A. Barros, S. De, S. K. Klimuk, T. Borland, V. Kosovrasti, W. L. Cantley, Y. K. Tam, M. Manoharan, M. A. Ciufolini, M. A. Tracy, A. de Fougerolles, I. MacLachlan, P. R. Cullis, T. D. Madden and M. J. Hope, *Nat. Biotechnol.*, 2010, **28**, 172.
- 79 J. A. Kulkarni, D. Witzigmann, S. Chen, P. R. Cullis and R. van der Meel, *Acc. Chem. Res.*, 2019, **52**, 2435.
- 80 K. K. Sahu and R. S. Pandey, *Int. Immunopharmacol.*, 2016, **39**, 343.
- 81 K. K. Sahu, M. Kaurav and R. S. Pandey, *Int. Immunopharmacol.*, 2019, **66**, 317.
- 82 M. P. Lokugamage, D. Vanover, J. Beyersdorf, M. Z. C. Hatit, L. Rotolo, E. S. Echeverri, H. E. Peck, H. Ni, J.-K. Yoon, Y. Kim, P. J. Santangelo and J. E. Dahlman, *Nat. Biomed. Eng.*, 2021, **5**, 1059.
- 83 Z. Li, Z. Wang, P.-U. C. Dinh, D. Zhu, K. D. Popowski, H. Lutz, S. Hu, M. G. Lewis, A. Cook, H. Andersen, J. Greenhouse, L. Pessaint, L. J. Lobo and K. Cheng, *Nat. Nanotechnol.*, 2021, **16**, 942.
- 84 J. D. Kirkpatrick, A. D. Warren, A. P. Soleimany, P. M. K. Westcott, J. C. Voog, C. Martin-Alonso, H. E. Fleming, T. Tammela, T. Jacks and S. N. Bhatia, *Sci. Transl. Med.*, 2020, **12**, eaaw0262.

