

Energy Advances

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: E. Rossin, Y. Yang, M. Chirico, G. Rossi, P. Galloni and A. Sartorel, *Energy Adv.*, 2024, DOI: 10.1039/D4YA00273C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Data availability statements not applicable for a perspective article

[View Article Online](#)
DOI: 10.1039/D4YA00273C

Open Access Article. Published on 02/07/2024. Downloaded on 02/07/2024. This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.



PERSPECTIVE

Quinacridone dyes: versatile molecules and materials for photo- and photoelectrochemical processes

Elena Rossin,^{a†} Yunshuo Yang,^{a†} Martina Chirico,^a Greta Rossi,^a Pierluca Galloni,^{*b} Andrea Sartorel^{*a}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

The renaissance of photochemistry and the explosion of photo- and photoelectro-catalysis open new opportunities in organic photocatalyst design and applications towards solar fuels and sustainable organic reactivity. In this perspective, we discuss the relevant case of quinacridone (QA) dyes: these have long been known to the scientific community, but their application in photocatalysis is recent and still explored in a limited way. This is somehow surprising given that QA is a cheap and readily available organic pigment, and in front of the appealing properties of QA derivatives, including intense absorption in the visible region, balanced redox properties making them suitable for both oxidative and reductive photochemistry, and versatility to several operative conditions. We will discuss recent examples of photo- and photoelectrochemical processes taking advantage of QA dyes, from solution photocatalysis to photoactive materials and devices (nanoparticles, covalent organic frameworks, photoelectrodes); the target applications include water splitting, carbon dioxide reduction, and organic transformations. We aim to show the potential of organic photocatalyst design and implementation, and to inspire the readers with new opportunities in this field.

Introduction

In 1912, ahead of his time, the day-dreamer Italian photochemist Giacomo Ciamician envisioned solar light as an inexhaustible and clean source of energy to fuel the world, in contrast to the exploitation of fossil fuels.¹ Nowadays, considering the energy crisis and the climate change, this concept is more current than ever.

This led to the renaissance of photochemistry applications and to an explosion of reports dealing with photocatalysis starting from the beginning of XXI Century.^{2–8} Photocatalysis is often associated to redox chemistry for the production of solar fuels, with relevant examples dealing with water splitting^{9,10} and carbon dioxide reduction,^{11–13} and for the development of sustainable organic synthesis.^{14–16} The progress in new photocatalyst families design and the expansion of novel photochemical processes are therefore high-target objectives for the scientific community.

Among the necessary features of future investigations, we identify three important aspects.

First, photocatalysts should be affordable and cheap, avoiding the use of noble, rare, or expensive elements. Second, the photocatalyst should be modular through a chemical design, to afford the required photophysical and redox properties.^{4,17}

Third, the development of photocatalytic materials and devices

should be considered.¹⁸ As possible technologies for water splitting, Harry B. Gray identified photovoltaic modules coupled to electrolyzers, photoelectrochemical cells or mixed photocatalytic colloids,¹⁹ with this latter being the less developed but more scalable solution. Indeed, this include the recently assessed wireless, particulate materials based on aluminum-doped strontium titanate in a 100 m² prototype composed of an array of panel photoreactors, enabling hydrogen production for one year.²⁰ A record Solar-to-Hydrogen value of 9.2% has been achieved using a particulate photocatalysts based on indium gallium nitride.²¹ Other examples of photocatalytic particulate materials include dye-sensitised nanoparticles²² and covalent-organic frameworks, with some cases discussed in the following paragraphs of this perspective.

In order to respond to the requirements above, the use of organic photocatalysts is a promising road to follow.¹⁷ Organic photocatalysts can be either commercially available or synthesisable in large scale with few steps; many of them have low or null toxicity and are exploited in food or cosmetic industry. The molecular structure allows to reach tuneable chemical, photophysical and redox properties, upon suitable design and/or functionalisation of the parent scaffold.⁴ Finally, they can be implemented in heterogeneous systems exploiting chemical bonds or physisorption to surfaces.^{23,24}

In this perspective, we will discuss these aspects dealing with **quinacridone** (QA) dyes as a selected example (scheme 1). Quinacridone (pigment violet 19) belongs to industrially relevant pigments, it is non-toxic, commercially available and cheap, and is exploited in inks and paints.²⁵ Its pentacyclic aromatic structure provides an extended absorption in the visible region of the electromagnetic spectrum. The N-H and

^a Elena Rossin, Yunshuo Yang, Martina Chirico, Greta Rossi, Andrea Sartorel, Dept. Chemical Sciences, University of Padova, via Marzolo 1 35131 Padova (Italy). andrea.sartorel@unipd.it


^b Pierluca Galloni, Department of Chemical Science and Technologies, University of Rome "Tor Vergata" via della Ricerca Scientifica snc, 00133 Rome, Italy. galloni@scienze.uniroma2.it

† E.R. and Y.Y. equally contributed.



C=O groups induce the formation of an extended hydrogen bonding network, providing features as a semiconducting pigment.²⁶ Among its applications, it was considered in field-effect transistors²⁷ and in light emitting diodes,²⁸ while more recently it was the subject of several investigations towards photocatalysis.

Herein, we will focus on photo- and photoelectrochemical processes based on QA derivatives: we will discuss the versatility of this molecule towards several photocatalytic processes (including water splitting, carbon dioxide reduction, organic functionalisation) and concerning its application in several operative conditions including homogeneous systems, nanoparticles, covalent organic frameworks and photoelectrodes, scheme 1. Our aim is to show the potential of organic photocatalyst design and implementation, inspiring the readers with new opportunities in this field.



Scheme 1. Representation of the linear trans-isomer of quinacridone (QA), and of its versatility for application in photo- and photo-electrocatalysis, which is the subject of this perspective article. The numeration of the atoms in QA scaffold is also reported.²⁵

Photophysical and redox properties of QA derivatives

Photophysical properties.

The absorption properties of QA depend on the degree of aggregation of the molecules through intramolecular hydrogen bonding; it appears yellowish in diluted solutions, while it is intense red in aggregated forms and in the solid state.

Non-aggregated states. In a non-aggregated state in DMSO solution, QA displays an intense absorption in the visible region characterised by four vibronic components (450 – 550 nm, ϵ up to $1.4 \times 10^4 \text{ M}^{-1}\text{cm}^{-1}$), and due to the transition from the ground to the lowest electronic excited state, associated to a $\pi \rightarrow \pi^*$ molecular orbitals; few nm

bathochromic shifts are observed by increasing the polarity of the solvent.²⁹ The emission spectrum is specular to the absorption, and is characterised by an excited state lifetime of about 20 ns, a fluorescence quantum yield ϕ_F of 0.76,³⁰ low intersystem crossing yield, and an energy of the excited state $E_{0-0} = 2.34\text{--}2.36 \text{ eV}$,³¹ given by the interception of normalised absorption and emission spectra, Figure 1 (λ interception in the range 525–531 nm). N-alkylation of QA does not impact significantly on the optical properties,³² while N,N'-Boc functionalisation (Boc = tert-butyloxycarbonyl, as a transient protecting group) leads to a blue-shift of absorption and emission, with a significant increase of E_{0-0} up to 2.49 eV, Figure 1 (in the case of QA-Boc, a fluorescence quantum yield ϕ_F of 0.94 was reported in DMF solution).^{33,34} Insertion of both electron-donating and electron-withdrawing substituents in the aromatic scaffold leads to a redshift of the absorption and to a reduction of the E_{0-0} energy, due to a contraction of the HOMO/LUMO energy gap.^{35,36}

Aggregated states. Aggregated states of QA include nanoparticles and thin films, where QA can arrange in different crystalline or polymorphic states.³⁷ As a general trend, the aggregation through hydrogen bonding leads to a redshift of absorption and emission spectra, to a decrease of lifetime of the excited state and to an abatement of the fluorescence quantum yield.³⁸ The hydrogen bonding network aligns transition dipoles in a “head-to-tail” fashion, inducing the absorption redshift due to excitonic interactions.²⁹ For instance, 20 nm sized QA nanoparticles prepared through the reprecipitation method display absorption up to 600 nm (ϵ ca $5 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$ at 550 nm) and ϕ_F below 10^{-4} , with the excited state quantitatively relaxing in a 10–110 ps timescale.³⁰ In thin 40-nm thick films prepared by vacuum sublimation of QA onto fluorine doped tin oxide (FTO) slides, absorption peaks are ca 40 nm redshifted with respect to the solution state; the emission of the QA film is characterized by a maximum at 610 nm, and by a broad band centered at 690 nm with a lifetime of 1.7 ns, ascribable to differently aggregated forms of the pigment, Figure 1.³¹ The intersection of the normalized absorption and emission spectra allows to estimate an E_{0-0} energy of 2.10 eV for *QA within the film (λ interception 590 nm).




Figure 1. Normalised absorption and emission properties of QA derivatives (QA-Boc, QA-alkyl) and QA spectra are reported in DMSO solution; spectra of QA films prepared by vacuum sublimation over FTO are registered under air atmosphere. For the emission spectra, the major lifetime component is also indicated.

Redox properties.

The QA scaffold allows the occurrence of both oxidative and reductive redox processes. Due to its poor solubility, electrochemical characterisation of parent QA is typically measured in heterogeneous conditions,^{31,39} and in the presence of protic medium the redox processes are proton-coupled involving the C=O and N-H moieties (*vide infra*). N-functionalised QA are instead amenable of solution analysis.^{32,35} For instance, in DCM solution N,N'-dioctyl quinacridone (DOQA) shows reversible one-electron anodic waves at +0.68 and +1.15 V, and two reversible one-electron cathodic processes at -1.82 and -2.18 V (all potentials reported vs Fc^{+/0}), as shown in the differential pulse voltammograms in Figure 2.^{32,35} Notably, although the reversibility of the processes may depend on the solvent, the bis-oxidised and bis-reduced species are characterised by fully conjugated systems with an even number of electrons (Scheme 2). For QA-Boc, the waves are shifted to more positive potentials by 300/400 mV. More importantly, the redox properties of QA derivatives are tuneable and balanced,⁴⁰ allowing their application in both oxidative and reductive photocatalytic processes, as it will be presented in the next sections.





Figure 2. Differential pulse voltammograms registered for N,N'-dioctyl QA (DOQA) in DCM. Reproduced from ref. 32 with permission from the Royal Society of Chemistry, copyright 2017. DOI: 10.1039/D4YA00275E




Scheme 2. Representation of oxidised and reduced forms of DOQA.

Application of QA in photo- and photoelectrocatalysis

Homogeneous photocatalysis.

Functionalized QAs (on the ring or alkylating the N) show higher solubility in common organic solvents, such as DMF or chlorinated ones. The solubility is even better in fluorinated alcohols, in particular hexafluoroisopropanol (HFIP). This feature allows to use QA derivatives as photocatalysts in homogeneous conditions.

Moreover, considering the spectroscopic and electrochemical properties of QA derivatives, these compounds can act as photocatalysts both in reducing and oxidation processes. In fact, the potentials of the excited state, calculated combining the absorption spectra and the redox potential of the ground state,^{41,42} show suitable values in order to promote photoinduced oxidative or reductive quenching pathways (Scheme 3).



Scheme 3. Representation of possible reaction pathways of the *QA excited state through oxidative and reductive quenching, leading to the generation of oxidised QA⁺ and reduced QA⁻, respectively. A and A' represent electron acceptors, while D and D' represent electron donors. The two equations enable the estimation of *E_{ox} and *E_{red} of the *QA excited state.^{41,42}

N,N'-disubstituted QAs were probed as efficient photocatalysts in polymerization of methyl (MMA) or butyl methacrylate (BMA). In this system QA is functionalized at the nitrogen sites with Boc or a long alkyl chain (Figure 3). The resulting catalysts



are soluble in DMF and toluene, giving interesting results in polymerization reactions. In particular, the QA-Boc derivative showed the best results in term of polymer yield. The mechanism involves a halogenated initiator that is reduced by the photoexcited QA photocatalyst obtaining a radical initiator and a halide X^- , together with the oxidised photocatalyst (oxidative quenching, with the oxidised PC^+ and X^- forming an ion pair). This radical starts the chain reaction with the monomer (MMA or BMA) propagating the polymer, while the PC^+/X^- ion pair was responsible for halogenation of radical intermediates, that re-enter the catalytic cycle (Figure 3).




Figure 3. QA based derivatives employed for photocatalysed polymerization reaction of MMA, and proposed mechanism (Adapted from ref. 34 with permission from Elsevier, copyright 2019)

Recently, N-functionalized QA compounds have been used as photocatalysts in thioethers oxidation in homogenous reactions.⁴³ The N,N' di-butyl quinacridone derivative (DBQA) was obtained from the reaction of deprotonated QA and *n*-butyl bromide in a 65% yield, and was characterized in apolar DCM and polar, protic HFIP, showing large difference in terms of maximum absorption spectra, oxidation and reduction potentials, and consequently on the redox energy of the excited state.

In the reaction involving thioethers, the excited state of DBQA can oxidise directly the substrate (reductive quenching pathway, with $E(DBQA^+/DBQA^-)$ estimated +1.29 V vs SCE in HFIP; reduced $DBQA^-$ is then oxidised by O_2) or can be oxidised by molecular oxygen (oxidative quenching, with $E(DBQA^+/DBQA^*)$ estimated -1.28 V vs SCE in HFIP; oxidised $DBQA^+$ further reacts with the thioether), Figure 4. The overall result is a very efficient catalytic reaction, obtaining selectively the desired sulphoxide in excellent yields and very short reaction time. DBQA was also effective in the selective oxidation of thianthrene to thiantrene-5-oxide, which is a valuable reagent in organo-catalysis.⁴³




Figure 4. Photocatalytic oxidation of thioethers with QA derivatives, and proposed reaction mechanism on the basis of scavenging experiments. Reproduced from ref. 43 with permission from Wiley Inc., copyright 2023.

The possibility of QA to react with oxygen through the oxidative quenching pathway may be exploited for the generation of reactive oxygen species (ROS), that can be competent agents towards the degradation of organic contaminants. An example deals with a photocatalytic system exploiting a Ni-QA complex for the aerobic oxidation of acridine orange (AO) in water, Figure 5.⁴⁴ The complex Ni-QA can be suspended in a water solution of AO, and after irradiation with a Xe lamp the AO visible absorption is completely suppressed in 3 hours, whereas using a heterogeneous solution QA the degradation of AO is only around 50%. Although this system is actually heterogeneous, we report it in this section given the similar mechanism discussed previously for the oxidation of thioethers. Indeed, the Ni-QA complex acts as reducing agent of molecular oxygen (oxidative quenching) forming the superoxide radical anion that degrades the AO.




Figure 5. Ni-AQ complex (obtained as the nitrate salt) for the aerobic photodegradation of acridine orange (AO) in water with visible light.⁴⁴

The photoinduced reactivity of organic dyes towards oxygen can be exploited for the photochemical synthesis of hydrogen peroxide.⁴⁵ H_2O_2 is considered a green oxidant (47% oxygen efficiency, with water as the sole by-product), while it is industrially synthesized through the anthraquinone process,



that requires harsh and inconvenient reaction conditions (high temperatures, noble metal catalysts, high pressure of H₂, use of organic solvents). In the case of QA, photosynthesis of H₂O₂ was achieved with a water soluble 2,9-disulfonate derivative; in aqueous medium, when coupled with electron donors such as formate, glucose or oxalate, the disulfonate QA was able to photogenerate H₂O₂ in a mM concentration range, although with a limited turnover number of ca 5 (Figure 6).⁴⁶ The one electron oxygen reduction to superoxide radical anion is the first key step, with O₂^{•-} further converting to H₂O₂. We will discuss an example of H₂O₂ photosynthesis also in the photoelectrochemical cells section.




Figure 6. Water soluble QA disulfonate derivative, and photochemical hydrogen peroxide production coupled to oxidation of donors D. Adapted from ref. 46 with permission from Wiley Inc., copyright 2019.

QA sensitised nanoparticles, covalent organic frameworks, nanostructures.

The investigation of QA sensitised nanoparticles for photocatalysis applications was pioneered by Shan et al,⁴⁷ who reported the sol-gel deposition of a TiO₂/QA layer onto the inner pore wall of mesoporous SBA-15 silica. The resulting material was employed in photocatalytic decomposition of indigo carmine with visible light irradiation, showing enhanced activity with respect to anatase TiO₂ and Ti-modified SBA-15 materials.⁴⁷

Following a similar approach, Lin et al. prepared TiO₂/SBA-15 particles sensitised with 2,9-dichloro QA by impregnation method. The material was exploited for photocatalytic carbon dioxide reduction to methanol in aqueous solution, achieving a production of 0.44 μmol h⁻¹ per gram of photocatalyst, significantly higher than the productivity of P25 TiO₂ sensitised particles of 0.16 μmol h⁻¹ per gram of photocatalyst, ascribed to a higher quinacridone loading on the material.⁴⁸

A more recent report using QA composites as photocatalyst is related to a photochemical reduction of CO₂ using QA deposited onto reduced graphene oxide (rGO), Figure 7.⁴⁹ The hybrid photocatalyst was prepared by irradiation with a Xe lamp of a DMSO solution of QA and suspended GO, with the resulting QA/rGO composite that precipitates and can be recovered by filtration. As a reference material, the authors also prepared QA nanoparticles, and they showed the heterogeneous QA/rGO photocatalyst gave the best result in term of CO₂ reduction, showing a synergic effect of the QA and rGO components. In particular, excited QA can reduce directly the CO₂ or reduce the

rGO that then transfer the electron to the carbon dioxide. Using a sacrificial electron donor such as triethanolamine (TEOA) allowed to obtain photocatalytic production of CO and CH₄ with rates of 450 and 275 μmol h⁻¹ per gram of catalyst.⁴⁹




Figure 7. Reduction of CO₂ over QA/rGO photocatalyst, and schematic representation of CO₂ reduction to CO and CH₄, coupled with oxidation of triethanolamine (TEOA); the potential of the conduction and valence bands (CB and VB, respectively) are reported versus the reversible hydrogen electrode.⁴⁹

More recently, photocatalytic reduction of CO₂ to CO was achieved over a Z-scheme inspired Quinacridone/BiVO₄ nanocomposites.⁵⁰ The QA/nBiVO₄ hybrid materials were prepared in different compositions (n=0.2–20, n is mass ratio of BiVO₄ to QA) by hydrogen-bonding self-assembly of QNC over hydroxylated BiVO₄ nanosheets.

The QA/nBiVO₄ materials were proven to photocatalyse the reduction of CO₂ to CO as the main product (traces of methane and H₂ were also detected), when immobilised onto a glass sheet (5 mg of material onto a 9 cm² surface), immersed and irradiated in a 4:1 water:TEOA solution, saturated with CO₂. The optimal material was found to be QA/10BiVO₄, achieving 407 μmol h⁻¹ production of CO per gram of photocatalyst, 24 times higher than QA alone, while BiVO₄ nanosheets almost showed no activity (Figure 8; indeed, the conduction band of BiVO₄ does not reach the thermodynamic potential to drive reduction of CO₂). Interestingly, the proposed mechanism foresees a Z-scheme, with excitation of both QA and BiVO₄, and electron transfer from the conduction band of BiVO₄ to the previous HOMO of QA. The holes on BiVO₄ are exploited for oxidation of TEOA, while reducing equivalents conveyed to QA drive the CO₂ reduction. It is worth to mention that similarly to quinones,⁵¹ electrogenerated reduced forms of QA are capable of reversible carbon dioxide binding, leading to the formation of a QA-carbonate anion.⁵² The formation of such intermediates could be responsible for the observed photocatalytic activity of QA/nBiVO₄ composites.






Figure 8. Top panel: Production rates from CO_2 photoreduction (average over 10 h) on BiVO_4 , QA, and $\text{QA}/n\text{BiVO}_4$ for 10 h. n is the mass ratio between BiVO_4 and QA, impacting the photocatalytic performance. Bottom panel: Schematic representation of photochemical CO_2 reduction on QA/ BiVO_4 composites. Reproduced from ref. 50 with permission from Wiley Inc., copyright 2020.

The planar structure of QA makes it suitable for its incorporation in 2D covalent organic frameworks, as in the case of the report by Zhang, Chen and co-workers, Figure 9.⁵³ QA-TAPT-COF and QA-TPB-COF were prepared by solvothermal synthesis starting from the molecular components, necessary to form the imine linkers. The resulting COF are characterised by large pore size (> 4.7 nm).

The two COFs show a redshifted absorption with respect to the molecular QA monomers, due to the extended conjugated structure of the materials; the band gaps of 2.03 and 2.09 eV for QA-TAPT-COF and QA-TPB-COF, respectively, are in the range expected for aggregate states of QA (*vide supra*).

The QA-based 2D COFs were applied as photocatalysts for the Povarov reaction, targeting the condensation of maleimide and N,N -dimethylaniline derivatives under aerobic atmosphere to produce tetrahydroquinolines (Figure 9). Both COFs were effective in photocatalysing the reaction (quantitative yields were achieved in ca 40 minutes irradiation in DMF as the solvent), with a superior reactivity of QA-TAPT-COF with respect to QA-TPB-COF; QA-TAPT-COF was also more active than soluble QA derivatives alkylated at the nitrogen sites, and could also be recycled up to 10 times. Concerning the mechanism, the first step of the Povarov reaction typically involves oxidation by single electron transfer of the arylamine by the excited photocatalyst, with formation of the arylamine radical cation and of the reduced photocatalyst (reductive quenching), that is then re-oxidised by oxygen to form $\text{O}_2^{\cdot-}$, this latter being involved in an oxidation step leading to the final product.⁴⁰ As a

mechanistic proof, the authors were able to support $\text{O}_2^{\cdot-}$ formation by trapping it with 5,5-dimethyl-1-pyrroline N -oxide (DMPO) and characterising the adduct by electron paramagnetic resonance.⁵³




Figure 9. Incorporation of QA derivatives in 2D covalent organic frameworks and photochemical application towards the Povarov reaction. Reproduced from ref. 53 with permission from Elsevier, copyright 2022.

As a final selected example of QA based particulate materials for photocatalysis application, we report the development of supramolecular nanobelts.⁵⁴ In particular, these were designed through alkylation of both nitrogen atoms of QA with n -butyl or n -octyl chains, and by insertion of hydrophilic pyridine-2,6-dicarboxylic acid moieties at positions 3 and 10 of the pentacyclic core (Figure 10). The molecular system is characterised by donor-acceptor interactions between the electron rich QA donor and the electron-deficient pyridine ring acceptor. The monomers self-assemble in water through hydrogen bonding and π - π stacking interactions, forming nanobelts with a width of ca 50 nm and a length of hundreds of nm for SQNCP-C4, and a width of 100 nm and a length up to micrometers for SQNCP-C8. As expected and similarly to the cases discussed above, the formation of supramolecular aggregated systems leads to a substantial redshift of the nanobelts with respect to the isolated monomers, and the band gap of SQNCP-C4 and SQNCP-C8 was measured to be 1.76 eV and 1.94 eV, respectively. Both of them show emission peaks around 650 nm, with the emission intensity that was significantly lower in the case of SQNCP-C4; this was interpreted as a proof of intramolecular charge separation within the donor/acceptor system.

The materials were tested towards photocatalytic reduction of water to hydrogen, in aqueous solution in the presence of



ascorbate as electron donor. The SQNCP-C4 material exhibited the highest activity of ca. 656 $\mu\text{mol h}^{-1}$ per gram of photocatalyst; the H_2 production rate was increased by ca 3 times in the presence of 1% w/w Pt co-catalyst. The material maintained the activity over three runs, for a total of 15 h irradiation (Figure 10).




Figure 10. Formation of QA based nanobelts, and application in photocatalytic water reduction to hydrogen. Reproduced from ref. 54 with permission from the Royal Society of Chemistry, copyright 2020.

Photoelectrochemical cells.

Photoelectrochemical cells are considered a viable technology to conduct redox transformations; they are composed by an anode and a cathode, where at least one of them is activated by light. In photoelectrochemical cell, upon light absorption the electrons are thus conveyed through an external circuit, and thus it is not necessary to use sacrificial electron donors or acceptors, as it is often employed in solution photocatalysis. The first example of a photoelectrochemical cell was provided by Fujishima and Honda, that developed a TiO_2 based photoanode coupled to a Pt cathode for overall water splitting into hydrogen and oxygen.⁵⁵ Since then, many efforts have been conducted to design more efficient photoelectrodes, with one strategy being the integration of molecular components for more efficient light absorption (photosensitizers) and for accelerating the redox reactions (catalysts).^{56,57}

Exploiting the well-balanced redox properties of QA, its derivatives have been recently employed at both the photoanode and photocathode compartments of a photoelectrochemical cell for water splitting, capable to operate in the absence of external applied bias (Figure 11).⁵⁸ In particular, the QA was functionalised at 2 and 9

carbon positions with pyridine dicarboxylic acid pendants to anchor the dye to TiO_2 (n-type semiconductor) and to NiO (p-type semiconductor), to develop the photoanode and photocathode, respectively. QA was also functionalized at both nitrogen sites with long alkyl chain substituents (C4–C16), to prevent excessive adhesion of the dye to the hydrophilic surface of titania (likely mitigating unproductive charge recombination events).

The water oxidation catalyst was a single site Ru coordination complex, still bearing the pyridine dicarboxylic acid ligand; anchoring of the catalyst to the QA dye was achieved through Zr(IV) bridges, as pioneered by T.J. Meyer and co-workers.⁵⁹ The water reduction catalyst was instead a Cobalt cobaloxime,⁶⁰ directly anchored to the NiO through phosphonate pendants. When the two photoelectrodes were combined in a photoelectrochemical cell and both irradiated with visible light in aqueous solution (pH 7), water splitting was achieved in the absence of applied bias, reaching a photocurrent density of ca 110 $\mu\text{A cm}^{-2}$, a faradaic efficiency for hydrogen evolution of 89% and a solar to hydrogen efficiency of 0.11%.




Figure 11. QA based photoelectrochemical cell for water splitting into hydrogen and oxygen: representation of the energy levels (a) and molecular assembly of the cell (b). Reproduced from ref. 58 with permission from the Royal Society of Chemistry, copyright 2022.

An operative advantage of photoelectrochemical cells development is that the photoelectrodes can be designed and optimized independently, by coupling them with a suitable counter-electrode. Moreover, other processes with respect to water splitting can be targeted. As two final case studies, we will discuss the development of QA photocathodes for photoelectrochemical synthesis of hydrogen peroxide and of QA based photoanodes for photoelectrochemical oxidative C-H activation.

Epindolidione (EPI) and QA based photocathodes for hydrogen peroxide production were developed by Głowacki and co-workers, and constituted the first example of an organic semiconductor photocathode for H_2O_2 photosynthesis (Figure 12).³⁹ Polycrystalline films of QA (80 nm thickness) were prepared on gold electrode surface by vacuum sublimation of



the pigment; these films behave as organic semiconductors, displaying charge mobility in the range $0.1 - 1 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$.³⁹ A cathodic photocurrent response up to $10 \mu\text{A cm}^{-2}$ was observed in aqueous electrolyte in a wide pH range from 1 to 12 and attributed to oxygen reduction (indeed, the photocurrent response was 10-20 times lower when measured in N_2 degassed solutions). At pH 1, the photocurrent of the QA/Au photocathode was associated to H_2O_2 production with a $80 \pm 5\%$ faradaic yield; under these conditions and under illumination the photocathode was operative for 4 days, with photocurrent response in the range $1-4 \mu\text{A cm}^{-2}$. The initial photocathodic current drop was attributed to oxygen consumption in the cell, while the subsequent photocurrent increase was ascribed to a recrystallization of the QA dye, as shown in the scanning electron microscopy images taken before and after 20 hours of electrolysis, showing the conversion of rod-like crystallites into rectangular platelets (Figure 12). As a final remark, key features of the QA films for photoelectrochemically reducing oxygen to H_2O_2 are the high reducing level of the conduction band (or of the LUMO), and the two electron reductive chemistry couple to proton transfer in aqueous solution, similarly to quinones.^{61,62}




Figure 12. a-b: Epindolidione (EPI) and QA photocathodes for photoelectrochemical synthesis of H_2O_2 upon reduction of oxygen. Panel c represents the operating mechanism and the energy levels involved. Panel d shows the morphological evolution of QA films along the photoelectrocatalysis by scanning electron microscopy. Reproduced from ref. 39 with permission from Wiley Inc., copyright 2016.

The proton coupled electron transfer (PCET) reactivity of QA can be achieved also along its oxidative chemistry, and is the subject of the last example discussed in this perspective. Indeed, cyclic voltammograms of QA/carbon inks casted on glassy carbon registered in aqueous solution (pH between 4.5 and 8.3) show that the QA dye is characterized by an anodic wave with a potential shifting with the pH close to 58 mV / pH unit , typical of a Nernstian behavior associated to a PCET event. The oxidation potential can thus be conveniently reported as 1.23 V vs the reversible hydrogen electrode (RHE), and is associated to the oxidation of the QA core with release of H^+ from the nitrogen sites to the aqueous solution, and formation of a nitrogen centered radical (Figure 13).³¹ The oxidative PCET reactivity of QA was then exploited in photoanodes, sensitizing TiO_2 or SnO_2 mesoporous semiconductors through vacuum sublimation of QA; the photoelectrodes were characterized by an absorption typical of QA aggregate films (40 nm thickness) with an absorbance of 0.2 at 564 nm . Ultrafast spectroscopy confirmed electron injection from the QA dye to the semiconductor surface in a ps timescale (oxidative quenching of the QA dye), likely accompanied by proton transfer from the N-H group of the dye to the metal-oxide surface. This process is expected to generate a nitrogen centered radical on the QA dye, for which a Bond Dissociation Free Energy (BDFE) of $80.5 \pm 2.3 \text{ kcal mol}^{-1}$ can be estimated. Consistently, photoelectrochemical reactivity through hydrogen atom abstraction towards C-H bonds was observed. Indeed, photoelectrochemical oxidation of γ -terpinene to p-cymene was observed, occurring through the initial hydrogen atom transfer followed by a radical chain mechanism involving dissolved oxygen.

Expanding the scope to other C-H moieties, a photocurrent response was observed for substrates containing allylic or benzylic C-H bonds with a BDFE lower than the threshold of $80.5 \text{ kcal mol}^{-1}$ for the N-H bond in QA. As expected, more challenging substrates (containing C-H bonds with $\text{BDFE} > 80.5 \text{ kcal mol}^{-1}$) provided a null photocurrent enhancement since they could not be oxidized by hydrogen atom abstraction.

It is worth to mention that oxidative C-H bond activation is considered a Holy Grail in organic chemistry, while this is the first report where this reaction is conducted photoelectrochemically with a dye-sensitized electrode. Therefore, this proof-of-concept can open a novel way of conducting this challenging transformation.






Figure 13. QA based photoanode for oxidative C-H activation. The right figure shows the photoelectrochemical response (given by the ratio of the photocurrent in the presence and in the absence of the organic substrate) versus the BDFE of the C-H bond of the substrate. Reproduced from ref. 31 with permission from Wiley Inc., copyright 2023.

Conclusions and perspectives

In this perspective article, we have discussed examples of photocatalytic and photoelectrocatalytic processes taking advantage of QA derivatives. QA provides an example of a cheap and readily available organic pigment, showing intense absorption in the visible region of the electromagnetic spectrum and whose properties can be modulated by chemical design. Moreover, its versatility allows its use in several operative conditions, spanning from homogeneous photocatalysis to photoactive materials and devices. Finally, its balanced redox properties allow its application in both oxidative and reductive routines, embracing water splitting, carbon dioxide reduction, organic photocatalysis, as discussed in the previous sections.

We believe that QA represents an appropriate example (although investigated only in a limited manner) of the potential of organic photocatalyst design. This research field is in rapid evolution, and examples of emerging research lines in the recent literature landscape include reactivity in flow,^{63,64} multiphoton excitation,^{65,66} or the combination of light with electrochemistry to promote photochemistry of oxidised and reduced photocatalysts.⁶⁷ Targets for the future include the development of new processes and an improved sustainability of existing ones, together with a fundamental understanding of the reaction mechanism and identification of active species.⁶⁸

Author Contributions

All authors discussed the concepts and contributed to the first draft of the manuscript. P. G. and A. S. supervised and edited the writing, and were responsible for funding acquisition.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

A.S. acknowledges Dr. Eric Daniel Glowacki (Brno University of Technology) for having inspired his interest in QA pigments and for fruitful discussions. Financial support from the Italian MUR (PRIN 2022KPK8WM PROMETEO, to A.S. and P.G.), and by the European Union – Next Generation UE (PRIN2022 PNRR PHOTOCORE, P2022ZSPWF to A.S.) is gratefully acknowledged.

Abbreviations

AO=acridine orange
 BDFE=bond Dissociation Free Energy
 BMA= butyl methacrylate
 Boc = tert-butyloxycarbonyl
 COF=covalent organic framework
 DBQA=N,N' di-butyl quinacridone
 DCM=dichloromethane
 DMF=N,N'-dimethylformamide
 DMPO=5,5-dimethyl-1-pyrroldine N-oxide
 DMSO=dimethylsulfoxide
 DOQA=N,N'-dioctyl quinacridone
 EPI=Epindolidione
 FTO=fluorine doped tin oxide
 HFIP=hexafluoroisopropanol
 MMA=methyl methacrylate
 PCET=proton coupled electron transfer
 QA=quinacridone
 RHE=reversible hydrogen electrode
 rGO=reduced graphene oxide
 ROS=reactive oxygen species
 TEOA=triethanolamine

Notes and references

- G. Ciamician, *Science (80-.)*, 1912, **36**, 386–399.
- S. Wu, J. Kaur, T. A. Karl, X. Tian and J. P. Barham, *Angew. Chemie - Int. Ed.*, 2022, **61**, e202107811.
- J. P. Barham and B. König, *Angew. Chemie - Int. Ed.*, 2020, **59**, 11732–11747.
- A. Vega-Peñalosa, J. Mateos, X. Companyó, M. Escudero-Casao and L. Dell'Amico, *Angew. Chemie, Int. Ed.*, 2021, **60**, 1082–1097.
- G. E. M. Crisenza, D. Mazzarella and P. Melchiorre, *J. Am. Chem. Soc.*, 2020, **142**, 5461–5476.
- G. E. M. Crisenza and P. Melchiorre, *Nat. Commun.*, 2020, **11**, 8–11.
- N. A. Romero and D. A. Nicewicz, *Chem. Rev.*, 2016, **116**, 10075–10166.
- P. R. D. Murray, J. H. Cox, N. D. Chiappini, C. B. Roos, E. A. McLoughlin, B. G. Hejna, S. T. Nguyen, H. H. Ripberger, J. M. Ganley, E. Tsui, N. Y. Shin, B. Koronkiewicz, G. Qiu and R. R. Knowles, *Chem. Rev.*, 2022, **122**, 2017–2291.
- C. Bie, L. Wang and J. Yu, *Chem*, 2022, **8**, 1567–1574.
- B.-J. Ng, L. K. Putri, X. Y. Kong, Y. W. Teh, P. Pasbakhsh and S.-P. Chai, *Adv. Sci.*, 2020, **7**, art. number 1903171.
- W. Zhang, A. R. Mohamed and W. J. Ong, *Angew. Chemie - Int. Ed.*, 2020, **59**, 22894–22915.



- 12 V. Nikolaou, C. Govind, E. Balanikas, J. Bharti, S. Diring, E. Vauthey, M. Robert and F. Odobel, *Angew. Chemie - Int. Ed.*, 2024, **63**, e202318299.
- 13 C. Bizzarri, *European J. Org. Chem.*, 2022, e202200185.
- 14 M. A. Ischay, M. E. Anzovino, J. Du and T. P. Yoon, *J. Am. Chem. Soc.*, 2008, **130**, 12886–12887.
- 15 D. A. Nicewicz and D. W. C. MacMillan, *Science (80-.)*, 2008, **322**, 77–80.
- 16 D. M. Schultz and T. P. Yoon, *Science (80-.)*, 2014, **343**, 1239176.
- 17 T. Bortolato, S. Cuadros, G. Simionato and L. Dell'Amico, *Chem. Commun.*, 2022, **58**, 1263–1283.
- 18 D. Cappelletti, M. Barbieri, A. Aliprandi, M. Maggini and L. Đorđević, *Nanoscale*, 2024, Accepted article 10.1039/d4nr00383g.
- 19 J. R. McKone, N. S. Lewis and H. B. Gray, *Chem. Mater.*, 2014, **26**, 407–414.
- 20 H. Nishiyama, T. Yamada, M. Nakabayashi, Y. Maehara, M. Yamaguchi, Y. Kuromiya, Y. Nagatsuma, H. Tokudome, S. Akiyama, T. Watanabe, R. Narushima, S. Okunaka, N. Shibata, T. Takata, T. Hisatomi and K. Domen, *Nature*, 2021, **598**, 304–307.
- 21 P. Zhou, I. A. Navid, Y. Ma, Y. Xiao, P. Wang, Z. Ye, B. Zhou, K. Sun and Z. Mi, *Nature*, 2023, **613**, 66–70.
- 22 D. A. Kader and S. J. Mohammed, *RSC Adv.*, 2023, **13**, 26484–26508.
- 23 M. Bonchio, Z. Syrgiannis, M. Burian, N. Marino, E. Pizzolato, K. Dirian, F. Rigodanza, G. A. Volpato, G. La Ganga, N. Demitri, S. Berardi, H. Amenitsch, D. M. Guldi, S. Caramori, C. A. Bignozzi, A. Sartorel and M. Prato, *Nat. Chem.*, 2019, **11**, 146–153.
- 24 T. Gobatto, F. Rigodanza, E. Benazzi, P. Costa, M. Garrido, A. Sartorel, M. Prato and M. Bonchio, *J. Am. Chem. Soc.*, 2022, **144**, 14021–14025.
- 25 S. S. Labana and L. L. Labana, *Chem. Rev.*, 1967, **67**, 1–18.
- 26 M. Gsänger, D. Bialas, L. Huang, M. Stolte and F. Würthner, *Adv. Mater.*, 2016, **28**, 3615–3645.
- 27 E. D. Głowacki, M. Irimia-Vladu, M. Kaltenbrunner, J. Gasiorowski, M. S. White, U. Monkowius, G. Romanazzi, G. P. Suranna, M. Piero, T. Sekitani, S. Bauer, T. Someya, L. Torsi and N. S. Sariciftci, *Adv. Mater.*, 2013, **25**, 1563–1569.
- 28 E. D. Głowacki, R. R. Tangorra, H. Coskun, D. Farka, A. Operamolla, Y. Kanbur, F. Milano, L. Giotta, G. M. Farinola and N. S. Sariciftci, *J. Mater. Chem. C*, 2015, **3**, 6554–6564.
- 29 J. Mizuguchi and T. Senju, *J. Phys. Chem. B*, 2006, **110**, 19154–19161.
- 30 Y. Miyashita, H. Yokoyama, M. Tanabe, H. Kasai, H. Nakanishi and T. Miyashita, *J. Photochem. Photobiol. A Chem.*, 2009, **201**, 208–213.
- 31 Y. Yang, G. A. Volpato, E. Rossin, N. Peruffo, F. Tumbarello, C. Nicoletti, R. Bonetto, L. Paoloni, P. Umari, E. Colusso, L. Dell'Amico, S. Berardi, E. Collini, S. Caramori, S. Agnoli and A. Sartorel, *ChemSusChem*, 2023, e202201980.
- 32 P. Kurzep, Ł. Skórka, M. Zagorska, P. A. Guńka, M. Banasiewicz, B. Kozankiewicz and I. Kulszewicz-Bajer, *RSC Adv.*, 2017, **7**, 8627–8632.
- 33 M. Sytnyk, E. D. Głowacki, S. Yakunin, G. Voss, W. Schöfberger, D. Kriegner, J. Stangl, R. Trotta, C. Gollner, S. Tollabimazraehno, G. Romanazzi, Z. Bozkurt, M. Havlicek, N. S. Sariciftci and W. Heiss, *J. Am. Chem. Soc.*, 2014, **136**, 16522–16532.
- 34 T. Jia, S. Huang, H. Bohra and M. Wang, *Dye. Pigment.*, 2019, **165**, 223–230.
- 35 J. J. A. Chen, T. L. Chen, B. Kim, D. A. Poulsen, J. L. Mynar, J. M. J. Fréchet and B. Ma, *ACS Appl. Mater. Interfaces*, 2010, **2**, 2679–2686.
- 36 J. Jia, D. Feng, Y. Sha, C. Zhou, G. Liang and Y. She, *Tetrahedron*, 2020, **76**, 131057.
- 37 G. D. Potts, W. Jones, J. F. Bullock, S. J. Andrews and S. J. Maginn, *J. Chem. Soc. Chem. Commun.*, 1994, 2565–2566.
- 38 F. Würthner, C. R. Saha-Möllner, B. Fimmel, S. Ogi, P. Leowanawat and D. Schmidt, *Chem. Rev.*, 2016, **116**, 962–1052.
- 39 M. Jakešová, D. H. Apaydin, M. Sytnyk, K. Oppelt, W. Heiss, N. S. Sariciftci and E. D. Głowacki, *Adv. Funct. Mater.*, 2016, **26**, 5248–5254.
- 40 J. Mateos, F. Rigodanza, A. Vega-Peñalosa, A. Sartorel, M. Natali, T. Bortolato, G. Pelosi, X. Companyó, M. Bonchio and L. Dell'Amico, *Angew. Chemie - Int. Ed.*, 2020, **59**, 1302–1312.
- 41 F. Puntoriero, S. Serroni, G. La Ganga, A. Santoro, M. Galletta, F. Nastasi, E. La Mazza, A. M. Cancelliere and S. Campagna, *Eur. J. Inorg. Chem.*, 2018, 3887–3899.
- 42 V. Balzani, G. Bergamini and P. Ceroni, *Angew. Chemie - Int. Ed.*, 2015, **54**, 11320–11337.
- 43 M. Forchetta, V. Baia, F. Sabuzi, V. Conte and P. Galloni, *European J. Org. Chem.*, 2023, **26**, e202300523.
- 44 N. Mousavi, H. Hadadzadeh, A. A. Ensafi, H. Farrokhpour and K. Z. Mousaabadi, *Mater. Chem. Phys.*, 2023, **303**, 127823.
- 45 M. Gryszel, T. Schlossarek, F. Würthner, M. Natali and E. D. Głowacki, *ChemPhotoChem*, 2023, **7**, e202300070.
- 46 M. Gryszel, R. Rybakiewicz and E. D. Głowacki, *Adv. Sustain. Syst.*, 2019, **3**, 1900027.
- 47 H. Ding, H. Sun and Y. Shan, *J. Photochem. Photobiol. A Chem.*, 2005, **169**, 101–107.
- 48 C.-J. Lin, Y.-H. Liou, S.-Y. Chen and M.-C. Tsai, *Sustain. Environ. Res.*, 2012, **22**, 167–172.
- 49 P. Yang, S. Guo, X. Yu, F. Zhang, B. Yu, H. Zhang, Y. Zhao and Z. Liu, *Ind. Eng. Chem. Res.*, 2019, **58**, 9636–9643.
- 50 X. Yu, F. Wen, F. Zhang, P. Yang, Y. Zhao, Y. Wu, Y. Wang and Z. Liu, *ChemSusChem*, 2020, **13**, 5565–5570.
- 51 K. M. Diederichsen, R. Sharifian, J. S. Kang, Y. Liu, S. Kim, B. M. Gallant, D. Vermaas and T. A. Hatton, *Nat. Rev. Methods Prim.*, 2022, **2**, Article ID: 68.
- 52 D. H. Apaydin, E. D. Głowacki, E. Portenkirchner and N. S. Sariciftci, *Angew. Chemie - Int. Ed.*, 2014, **53**, 6819–6822.
- 53 Z. Liu, X. Yang, Z. Yang, X. Su, Z. Xie, W. Chen, W. Zhang and L. Chen, *Appl. Catal. B Environ.*, 2022, **312**, 121406.
- 54 M. Xu, K. Kong, H. Ding, Y. Chu, S. Zhang, F. Yu, H. Ye, Y. Hu and J. Hua, *J. Mater. Chem. C*, 2020, **8**, 930–934.
- 55 K. Fujishima, A., & Honda, *Nature*, 1972, **238**, 37–38.
- 56 L. Fei, L. Lei, T. J. Meyer and D. Wang, *Accounts Mater. Res.*, 2024, **5**, 124–135.



Journal Name

ARTICLE

- 57 T. Gobbato, G. A. Volpato, A. Sartorel and M. Bonchio, *Chem. Sci.*, 2023, **14**, 12402–12429.
- 58 L. Shen, S. Zhang, H. Ding, F. Niu, Y. Chu, W. Wu, Y. Hu, K. Hu and J. Hua, *Chem. Commun.*, 2021, **57**, 5634–5637.
- 59 D. Wang, M. V. Sheridan, B. Shan, B. H. Farnum, S. L. Marquard, B. D. Sherman, M. S. Eberhart, A. Nayak, C. J. Dares, A. K. Das, R. M. Bullock and T. J. Meyer, *J. Am. Chem. Soc.*, 2017, **139**, 14518–14525.
- 60 A. Fihri, V. Artero, M. Razavet, C. Baffert, W. Leibl and M. Fontecave, *Angew. Chemie - Int. Ed.*, 2008, **47**, 564–567.
- 61 M. T. Huynh, C. W. Anson, A. C. Cavell, S. S. Stahl and S. Hammes-Schiffer, *J. Am. Chem. Soc.*, 2016, **138**, 15903–15910.
- 62 E. J. Son, J. H. Kim, K. Kim and C. B. Park, *J. Mater. Chem. A*, 2016, **4**, 11179–11202.
- 63 G. Laudadio, Y. Deng, K. Van Der Wal, D. Ravelli, M. Nunõ, M. Fagnoni, D. Guthrie, Y. Sun and T. Noël, *Science (80-.)*, 2020, **369**, 92–96.
- 64 J. Schütte, D. Corsi, W. Haumer, S. Schmid, J. Žurauskas and J. P. Barham, *Green Chem.*, DOI:10.1039/d4gc00804a.
- 65 G. Han, G. Li, J. Huang, C. Han, C. Turro and Y. Sun, *Nat. Commun.*, 2022, **13**, art. number 2288.
- 66 B. Pfund, D. Gejsnæs-schaad, O. S. Wenger and B. Lazarevski, *Nat. Commun.*, DOI:10.1038/s41467-024-49006-5.
- 67 J. Žurauskas, S. Boháčová, S. Wu, V. Butera, S. Schmid, M. Domański, T. Slanina and J. P. Barham, *Angew. Chemie - Int. Ed.*, 2023, **62**, e202307550.
- 68 C. Wang, H. Li, T. H. Bürgin and O. S. Wenger, *Nat. Chem.*, DOI:10.1038/s41557-024-01482-4.

View Article Online
DOI: 10.1039/D4YA00273C

Open Access Article. Published on 02/07/2024. Downloaded on 2024/07/08 4:25:52.
This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.



Energy Advances Accepted Manuscript