


ChemComm

Chemical Communications

rsc.li/chemcomm



ISSN 1359-7345



COMMUNICATION

Takehiko Yoshimitsu *et al.*

Total synthesis of (–)-agelastatin A: an S_N2' radical azidation strategy



Cite this: *Chem. Commun.*, 2018, 54, 9893

Received 14th July 2018,
Accepted 1st August 2018

DOI: 10.1039/c8cc05697h

rsc.li/chemcomm

Total synthesis of (–)-agelastatin A: an S_H2' radical azidation strategy†

Izuru Tsuchimochi,^{ab} Yuta Kitamura,^b Hiroshi Aoyama,^b Shuji Akai,^b Keiyo Nakai^a and Takehiko Yoshimitsu^{*a}

A reagent generated from $TMSN_3/KMnO_4/BnEt_3NCl$ was found to promote an S_H2' radical azidation of a bromo silyl enol ether to furnish an azido silyl enol ether via olefin transposition. With the present azidation protocol, a new synthetic approach to agelastatin A, a potent antitumor marine alkaloid, has been established.

(–)-Agelastatin A (**1**), along with its congener agelastatin B (**2**), was first isolated as a cytotoxic constituent from the Coral Sea sponge *Agelas dendromorpha* by Pietra and co-workers in 1993 (Fig. 1).¹ Thereafter, Molinski and co-workers identified the Indian Ocean sponge *Cymbastela* sp. as another source that produces **1** along with agelastatins C (**3**) and D (**4**), two additional agelastatin members.² In 2010, Al-Mourabit and co-workers reported the isolation of agelastatins E (**5**) and F (**6**) from the New Caledonian sponge *A. dendromorpha*.³ Early biological assessments of agelastatins conducted by the aforementioned laboratories have revealed that compound **1** exhibits remarkable properties, including antitumor activity,^{1,3} brine shrimp toxicity,² and insecticidal activity.² In addition, Meijer and Pettit have found that agelastatin A (**1**) is a potent inhibitor of GSK-3 β , a pivotal serine/threonine kinase.⁴ Hale and El-Tanani have reported that agelastatin A (**1**) dramatically decreases β -catenin levels in cancer cells and inhibits cancer cell proliferation by arresting cell cycle at G2 phase.⁵

The biological significance of agelastatin A (**1**) has made it an attractive target for medicinal studies.^{6,7} For instance, Movassaghi's comparative cytotoxicity assay of all agelastatin members, i.e., A (**1**) to F (**6**), has successfully validated the relevance of agelastatin A (**1**)




Fig. 1 Agelastatin alkaloids.

as a promising anticancer agent.^{7a} In addition, structure–activity relationship (SAR) studies on agelastatin analogues have recently been disclosed by the groups of Molinski,⁸ Romo/Liu,⁹ and Movassaghi,¹⁰ boosting the applications of agelastatin particularly to blood cancer chemotherapy.

Our group has also been engaged in synthetic and medicinal studies on **1** and has demonstrated that agelastatin analogues potentially attenuate brain cancer.¹¹ Furthermore, our SAR study has revealed that structural modifications of the N1-substituent of the D-ring of **1** could retain the *in vitro* and *in vivo* therapeutic efficacies of agelastatin analogues.^{12,13} Movassaghi's group has further clarified that D-ring modifications expand the scope of derivatization of agelastatins to access potent analogues.¹⁰

In the present study, we have established a new route to agelastatin A (**1**) through an S_H2' radical azidation protocol using $TMSN_3/KMnO_4/BnEt_3NCl$ that enables the allylic transposition of a bromo silyl enol ether into an azido silyl enol ether, which serves as a useful D-ring precursor of the target natural product (Scheme 1).


The synthesis was commenced with Boc-protected amino-alcohol derivative **7** (>99% ee).¹⁴ The Boc group of **7** was removed with hydrochloric acid (HCl) generated *in situ* from $TMSCl$ in aq. MeCN to provide an ammonium salt (structure not shown). After evaporation of the solvents under reduced pressure, the resultant crude product was coupled with pyrrole-2-carboxylic acid using EDC, Et_3N , and DMAP in MeCN to furnish compound **8** in 84% yield. Then, compound **8** was hydrolyzed with LiOH in aq. EtOH to provide alcohol **9** in 88% yield. PDC oxidation of alcohol **9** in DMF delivered enone **10** in

^a Division of Pharmaceutical Sciences, Graduate School of Medicine, Dentistry, and Pharmaceutical Sciences, Okayama University, 1-1-1 Tsushima-naka, Kita-ku, Okayama 700-8530, Japan. E-mail: yoshimit@okayama-u.ac.jp;

Tel: +81-86-251-7930

^b Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 565-0871, Japan

† Electronic supplementary information (ESI) available: Experimental procedures, characterization of new compounds including NMR spectra. CCDC 1852628. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8cc05697h




Jiao and co-workers have reported that Mn(III) generated from MnBr₂ in the presence of molecular oxygen serves as an effective catalyst to generate an azide radical from TMSN₃.^{19a} We have examined Mn(OAc)₃·2H₂O in combination with TMSN₃ (6 equiv.) as a possible source of azido radical and found that desired material **14** could be similarly produced in 42% yield along with **17** (8%) (entry 9).^{19b} This result suggests that Mn(III) azide complex is likely responsible for the present radical azidation. Based on these observations, we currently assume that metastable Mn(V)

Table 1 Azidation of bromide **13** with various reagents

Entry	Reagents (equiv.)	Time	Yield ^a (%)			
			14	17	18	13 ^b
1	KMnO ₄ (0.3), BnEt ₃ NCl (0.3), TMSN ₃ (10), MeCN	40 min	50	21	3	Trace
2	KMnO ₄ (0.1), BnEt ₃ NCl (0.1), TMSN ₃ (10), MeCN	40 min	30	4	12	30
3	KMnO ₄ (0.1), BnEt ₃ NCl (0.1), TMSN ₃ (10), MeCN, O ₂	40 min	31	5	11	21
4	KMnO ₄ (0.6), BnEt ₃ NCl (0.6), TMSN ₃ (10), MeCN	40 min	43	11	4	9
5	NaN ₃ (1.1), DMF ^c	15 min	—	—	—	— ^d
6	BnEt ₃ NCl (0.3), TMSN ₃ (10), MeCN	75 min	—	—	—	90
7	TMSN ₃ (10), MeCN	70 min	—	—	—	89
8	PhIO (1.2), TMSN ₃ (2.4), CH ₂ Cl ₂ ^e	40 min	24	17	14	6
9	Mn(OAc) ₃ ·2H ₂ O (3), TMSN ₃ (6), MeCN ^c	11 h	42	8	—	—

^a Isolated yields after purification by column chromatography. ^b Recovered unreacted starting material. ^c The reaction was conducted at r.t.

^d Bromoketone (60%) was produced. ^e The reaction was conducted at −78 °C.



Scheme 2 Plausible generation of Mn(v) azide species.

species is generated from Mn(vii) with excess TMSN₃ and that Mn(v) provides 3 equiv. of azido radical to finally become Mn(ii), which no longer serves as a radical source. To elucidate the formation of the meta-stable Mn species, we carried out a comparison experiment: after stirring the reagents for 60 min, excess remaining TMSN₃ was completely removed under reduced pressure. Then, the residual solid that likely contains the Mn species was diluted with MeCN and mixed with substrate **13**. As a result, almost identical yields of products **14** (48%), **17** (22%), and **18** (6%) were obtained as in the case of entry 1, indicating that the Mn(v) azide complex is generated as a reactive meta-stable reagent.

The formation of compounds **17** and **18**, which provides an insight into the mechanism of the present azidation, also requires elaboration (Scheme 3). When azide **14** and isomeric azide **17** were separately subjected to the same reaction conditions for 1 h, only a trace amount of corresponding azide **17** and **14** was produced along with the unreacted starting azides, respectively. This indicates that both azides **14** and **17**, once produced, were hardly susceptible to the S_H2' azidation. In contrast, when isomeric bromide **18** was treated with the reagent, compounds **14** (34%), **17** (27%), and **18** (12%) were obtained similar to the case of **13**. Based on these results, we propose that the addition of an azide radical to bromide **13** generates a Br radical that undergoes rapid addition to substrate **13** to generate regioisomeric bromide **18**. Then, **18** is further converted into compound **17** via a radical azidation.

With azide **14** in possession, we further endeavored to accomplish the total synthesis. Thus, azide **14** was subjected to catalytic hydrogenation followed by one-pot urea formation with Batey's reagent²⁰

Scheme 3 Plausible mechanisms of the production of regioisomeric byproducts **17** and **18**.

and subsequent desilylative cyclization with CsF to afford tetracyclic compound **15** in 51% yield over three steps. It should be mentioned that no purification was required in the three-step sequence, allowing ease of experimental operations. Finally, the known bromination protocol was applied to compound **15** to furnish (−)-agelastatin A (**1**).

In conclusion, we have established a new approach to (−)-agelastatin A (**1**) by the strategic implementation of brominative olefin transposition and subsequent S_H2' radical azidation. The present approach features a late-stage construction of D-ring that would allow facile production of D-ring analogues. We believe that the present synthesis would facilitate further development of new agelastatin analogues.

This work was supported by a grant [KAKENHI #15K14977] generously provided by the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT), The NOVARTIS Foundation (Japan) for the Promotion of Science, and the Hoansha Foundation.

Conflicts of interest

The authors declare no conflicts of interest.

Notes and references

- (a) M. D'Ambrosio, A. Guerriero, C. Debitus, O. Ribes, J. Pusset, S. Leroy and F. J. Pietra, *J. Chem. Soc., Chem. Commun.*, 1993, 1305; (b) M. D'Ambrosio, A. Guerriero, G. Chiasera and F. Pietra,

- Helv. Chim. Acta*, 1994, **77**, 1895; (c) M. D'Ambrosio, A. Guerriero, M. Ripamonti, C. Debitus, J. Waikedre and F. Pietra, *Helv. Chim. Acta*, 1996, **79**, 727.
- 2 T. W. Hong, D. R. Jimenez and T. F. Molinski, *J. Nat. Prod.*, 1998, **61**, 158.
 - 3 S. Tilvi, C. Moriyou, M. Martin, J. Gallard, J. Sorres, K. Patel, S. Petek, C. Debitus, L. Ermolenko and A. Al-Mourabit, *J. Nat. Prod.*, 2010, **73**, 720.
 - 4 (a) L. Meijer, A. M. Thunnissen, A. W. White, M. Garnier, M. Nikolic, L. H. Tsai, J. Walter, K. E. Cleverley, P. C. Salinas, Y. Z. Wu, J. Biernat, E. M. Mandelkov, S. H. Kim and G. R. Pettit, *Chem. Biol.*, 2000, **7**, 51; (b) G. R. Pettit, S. Ducki, D. L. Herald, D. L. Doubek, J. M. Schmidt and J.-C. Chapuis, *J. Oncol. Res.*, 2005, **15**, 11.
 - 5 (a) C. K. Mason, S. McFarlane, P. G. Johnston, P. Crowe, P. J. Erwin, M. M. Domostoj, F. C. Campbell, S. Manaviazar, K. J. Hale and M. El-Tanani, *Mol. Cancer Ther.*, 2008, **7**, 548; (b) M. Harmata, *Strategies and Tactics in Organic Synthesis*, Elsevier Academic Press, London, 2005, ch. 11, vol. 6, pp. 352–394.
 - 6 For selected reviews on total synthesis of agelastatins, see: (a) G. Dong, *Pure Appl. Chem.*, 2010, **82**, 2231; (b) T. Yamaoka, Y. Ichikawa and H. Kotsuki, *J. Synth. Org. Chem., Jpn.*, 2012, **70**, 615.
 - 7 (a) D. Stien, G. T. Anderson, C. E. Chase, Y. Koh and S. M. Weinreb, *J. Am. Chem. Soc.*, 1999, **121**, 9574; (b) K. S. Feldman and J. C. Saunders, *J. Am. Chem. Soc.*, 2002, **124**, 9060; (c) K. S. Feldman, J. C. Saunders and M. L. Wroblewski, *J. Org. Chem.*, 2002, **67**, 7096; (d) K. J. Hale, M. M. Domostoj, D. A. Tocher, E. Irving and F. Scheinmann, *Org. Lett.*, 2003, **5**, 2927; (e) M. M. Domostoj, E. Irving, F. Scheinmann and K. J. Hale, *Org. Lett.*, 2004, **6**, 2615; (f) F. A. Davis and J. Deng, *Org. Lett.*, 2005, **7**, 621; (g) B. M. Trost and G. Dong, *J. Am. Chem. Soc.*, 2006, **128**, 6054; (h) Y. Ichikawa, T. Yamaoka, K. Nakano and H. Kotsuki, *Org. Lett.*, 2007, **9**, 2989; (i) D. P. Dickson and D. J. Wardrop, *Org. Lett.*, 2009, **11**, 13414; (j) N. Hama, T. Matsuda, T. Sato and N. Chida, *Org. Lett.*, 2009, **11**, 2687; (k) P. M. Wehn and J. Du Bois, *Angew. Chem., Int. Ed.*, 2009, **48**, 3802; (l) F. A. Davis, J. Zhang, Y. Zhang and H. Qiu, *Synth. Commun.*, 2009, **39**, 1914; (m) B. M. Trost and G. Dong, *Chem. – Eur. J.*, 2009, **15**, 6910; (n) M. Movassaghi, D. S. Siegel and S. Han, *Chem. Sci.*, 2010, **1**, 561; (o) Y. Menjo, A. Hamajima, N. Sasaki and Y. Hamada, *Org. Lett.*, 2011, **13**, 5744; (p) T. Kano, R. Sakamoto, M. Akakura and K. Maruoka, *J. Am. Chem. Soc.*, 2012, **134**, 7516; (q) J. C. P. Reyes and D. Romo, *Angew. Chem., Int. Ed.*, 2012, **51**, 6870; (r) P. A. Duspara and R. A. Batey, *Angew. Chem., Int. Ed.*, 2013, **52**, 10862; (s) A. H. Antropow, K. Xu, R. J. Buchsbaum and M. Movassaghi, *J. Org. Chem.*, 2017, **82**, 7720; (t) Y. Yao, X. Wang and G. Liang, *Tetrahedron*, 2017, **73**, 4538.
 - 8 E. P. Stout, M. Y. Choi, J. E. Castro and T. F. J. Molinski, *J. Med. Chem.*, 2014, **57**, 5085.
 - 9 (a) M. Jouanneau, B. McClary, J. C. P. Reyes, R. Chen, Y. Chen, W. Plunkett, X. Cheng, A. Z. Milinichik, E. F. Albone, J. O. Liu and D. Romo, *Bioorg. Med. Chem. Lett.*, 2016, **26**, 20927; (b) B. McClary, B. Zinshteyn, M. Meyer, M. Jouanneau, S. Pellegrino, G. Yusupova, A. Schuller, J. C. P. Reyes, J. Lu, Z. Gou, S. Ayinde, C. Luo, Y. Dang, D. Romo, M. Yusupov, R. Green and J. O. Liu, *Cell Chem. Biol.*, 2017, **24**, 605.
 - 10 A. H. Antropow, K. Xu, R. J. Buchsbaum and M. Movassaghi, *J. Org. Chem.*, 2017, **82**, 7720.
 - 11 (a) T. Yoshimitsu, T. Ino and T. Tanaka, *Org. Lett.*, 2008, **10**, 5457; (b) T. Yoshimitsu, T. Ino, N. Futamura, T. Kamon and T. Tanaka, *Org. Lett.*, 2009, **11**, 34025; (c) D. Shigeoka, T. Kamon and T. Yoshimitsu, *Beilstein J. Org. Chem.*, 2013, **9**, 860.
 - 12 (a) Z. Li, T. Kamon, D. A. Personett, T. Caulfield, J. A. Copland, T. Yoshimitsu and H. W. Tun, *Med. Chem. Commun.*, 2012, **3**, 233; (b) Z. Li, D. Shigeoka, T. R. Caulfield, T. Kawachi, Y. Qiu, T. Kamon, M. Arai, H. W. Tun and T. Yoshimitsu, *Med. Chem. Commun.*, 2013, **4**, 1093.
 - 13 H. W. Tun, T. Yoshimitsu, D. Shigeoka, T. Kamon, Z. Li, Y. Qiu and T. R. Caulfield, *US Pat.*, US9464093B2, 2016.
 - 14 (a) M. J. Mulvihill, J. I. Gage and M. J. J. Miller, *J. Org. Chem.*, 1998, **63**, 3357; (b) C. Cesario, L. P. Tardibono and M. J. J. Miller, *J. Org. Chem.*, 2009, **74**, 448.
 - 15 The optical purity was unambiguously confirmed by ¹H NMR analysis of Mosher esters derived from an alcohol that was prepared by reduction of ketone **11** with NaBH₄ (for the details, see the ESI†).
 - 16 (a) A. Deyine, G. Dujardin, M. Mammeri and J.-M. Poirier, *Synth. Commun.*, 1998, **28**, 1817; (b) K. Inanaga, Y. Ogawa, Y. Nagamoto, A. Daigaku, H. Tokuyama, Y. Takemoto and K. Takasu, *Beilstein J. Org. Chem.*, 2012, **8**, 658.
 - 17 P. Magnus, M. B. Roe and C. J. Hulme, *J. Chem. Soc., Chem. Commun.*, 1995, 263.
 - 18 The structure of regioisomeric bromide **18** was unambiguously confirmed by X-ray crystallographic analysis (CCDC 1852628)†.
 - 19 (a) X. Sun, X. Li, S. Song, Y. Zhu, Y.-F. Liang and N. Jiao, *J. Am. Chem. Soc.*, 2015, **137**, 6059; (b) Y. Zhao, Y. Hu, H. Wang, X. Li and B. Wan, *J. Org. Chem.*, 2016, **81**, 4412.
 - 20 P. A. Duspara, Md. S. Islam, A. J. Lough and R. A. Batey, *J. Org. Chem.*, 2012, **77**, 10362.