



Cite this: *Chem. Commun.*, 2018, 54, 9761

Received 26th July 2018,
Accepted 7th August 2018

DOI: 10.1039/c8cc06088f

rsc.li/chemcomm

Elucidation of the inverse *trans* influence in uranyl and its imido and carbene analogues *via* quantum chemical simulation†

Izaak Fryer-Kanssen and Andy Kerridge *

The inverse *trans* influence (ITI) is investigated in uranyl, UO_2^{2+} , and its isoelectronic imido ($\text{U}(\text{NH})_2^{2+}$) and carbene ($\text{U}(\text{CH}_2)_2^{2+}$) analogues at the density functional and complete active space self consistent field levels of theory. The quantum theory of atoms in molecules is employed to quantify, for the first time, the effect of the ITI on covalent bond character and its relationship to bond lengths and complex stability.

The inverse *trans* influence (ITI) is manifested in intermediate and high valence f-elements as a shortening of bond lengths of ligands lying *trans* to highly polarising species. The classic examples of this effect are manifested in $[\text{PPh}_4][\text{UOCl}_5]$ and $[\text{Et}_4\text{N}]_2[\text{PaOCl}_5]$, where the *trans* An–Cl bond is some 4.0 and 8.3% shorter than that of equivalent *cis* interactions, respectively.^{1,2} The effect has also been observed in a number of other U(v)/U(vi) mono-oxide and -imide complexes,^{3–8} as well as U(iv) and Ce(iv) bis-carbenes.⁹ The interested reader is directed to a comprehensive recent review of the ITI in actinide complexes.³

A qualitative explanation of the ITI, as well as detailed analysis of bonding in the actinyls, has been given by Denning^{10,11} in terms of the character of the metal polarisation by the ligand: dipolar in the case of transition metals, which therefore exhibit the *trans* influence, and quadrupolar in the f-elements, which therefore exhibit the ITI. Both Denning¹¹ and Hoffmann¹² have also highlighted the importance of the pseudocore electronic $6p_\sigma$ orbital of the actinide centre in the ITI, and the role of the $6p$ shell was successfully demonstrated by O'Grady and Kaltsoyannis using quantum chemical simulations, identifying key molecular orbitals associated with the phenomenon.¹³ A quantification of the impact of the ITI on electronic structure has, however, remained elusive. Recently, analysis methods based on the experimentally observable electron density have been shown to be capable of characterising a

number of chemical phenomena in the f-elements,¹⁴ and an appealing feature of this approach is that it can also be applied to experimentally derived densities.^{15,16} In this contribution, we employ a novel application of the quantum theory of atoms in molecules (QTAIM) to study the ITI in the prototypical U(vi) compound, uranyl (UO_2^{2+}), along with its isoelectronic imido and carbene analogues. The imido analogue has been reported by Boncella,¹⁷ who has also reported chalcogenido-substituted uranyl complexes of the form $[\text{OUE}]^{2+}$ ($\text{E} = \text{S}, \text{Se}$),¹⁸ whereas Gibson has shown that linear US_2^{2+} exists 171 kJ mol^{−1} higher in energy than the $\eta_2\text{-S}_2$ isomer in the gas phase.¹⁹ To date the carbene analogue is synthetically unrealised, although Liddle and coworkers have reported an $[\text{OUCR}_2]^{2+}$ uranyl analogue.²⁰

Since the analogues are expected to be more weakly binding ligands, so the ITI might be expected to be less pronounced. This is the first investigation of the ITI in uranyl *via* analysis of CASSCF-generated electron densities, although we note Pierloot's authoritative CASSCF study of uranyl bonding.²¹ This contribution also represents the first study of the electronic structure of $\text{U}(\text{CH}_2)_2^{2+}$. Whilst the ITI is expected to be substantial in uranyl, its effects are not experimentally accessible in their entirety, although Denning has characterised the role of the $6p_\sigma$ orbital *via* O K-edge X-ray absorption and emission (XAS/XES) spectroscopies.²² More recently, Cl K-edge XAS has been used to great effect to probe bond covalency in OUCl_5^- , as well as chloride complexes of both uranyl and its imido analogue,^{23,24} however the lack of equatorial ligands in bare uranyl analogues means that theory is able to provide a unique insight into the phenomenon in these complexes.

In this study, all complexes were assumed to have D_{2h} symmetry.† Density functional theory (DFT) calculations employed the non-empirical PBE0 hybrid-GGA exchange–correlation functional, which we have previously shown to give an accurate description of electronic structure in f-element complexes.^{25–29} Complete active space self-consistent field (CASSCF) calculations employed a minimal active space correlating 12 electrons in 12 orbitals, CASSCF(12,12), for uranyl and the imido analogue, and correlating eight electrons in eight orbitals, CASSCF(8,8)

Department of Chemistry, Lancaster University, Bailrigg, Lancaster, LA1 4YB, UK.
E-mail: a.kerridge@lancaster.ac.uk

† Electronic supplementary information (ESI) available: Full data tables, structural parameters and natural orbital visualisations. See DOI: 10.1039/c8cc06088f





Fig. 1 PBE0-generated density differences associated with freezing the 6p shell. Blue (white) regions indicate greater (lesser) charge density in the fully-relaxed electronic structure. CASSCF density differences exhibit qualitatively similar features. All isosurfaces rendered at a value of $\rho = 0.01$ a.u.

for the carbene analogue.[§] These active spaces included only those orbitals with bonding or antibonding character between the uranium centre and the coordinating species. All simulations employed the ANO-RCC basis sets of polarised triple-zeta quality and were carried out using version 8.0 of the MOLCAS code.³⁰

Following the approach developed by O'Grady and Kaltsoyannis,¹³ we have focussed on the role played by the pseudocore 6p shell. To achieve this, we have performed a set of simulations in which this shell, along with all those lying lower in energy, are restricted, or frozen, to have atomic character. These simulations are compared to a set in which the entire electronic structure is allowed to relax.

We first consider the variation in the total electron density distribution associated with freezing the 6p shell while maintaining the geometry associated with the fully relaxed electronic structure. These density differences (Fig. 1), obtained at the PBE0 level of theory, immediately demonstrate the validity of Denning's quadrupolar characterisation of the metal polarisation, with a central region of charge accumulation lying between two regions of charge depletion. The density differences of uranyl and its imido analogue exhibit cylindrical symmetry, whereas that of the carbene analogue reflects the sp^2 hybridisation of the carbon centres.

Whilst these density differences illustrate the qualitative characteristics of the ITI, they cannot be employed to ascertain its magnitude. To investigate the ITI further, we have therefore analysed the simulated electron densities using the QTAIM. We have focussed on two parameters in particular: the delocalisation index, $\delta(U,L)$, and the magnitude of the electron density at the

bond critical point, ρ_{BCP} , between the uranium and coordinating species. $\delta(U,L)$ quantifies the number of electrons shared between a pair of atoms, and can, in the absence of polarisation, be considered a measure of bond order, whereas ρ_{BCP} gives insight into the degree of charge accumulation in the bonding region. The high symmetry of the complexes under consideration here allowed us to employ a novel application of this analysis whereby gerade (g) and ungerade (u) contributions to ρ_{BCP} and $\delta(U,L)$ can be unambiguously identified and, additionally, contributions to $\delta(U,L)$ can be further decomposed into contributions from σ - and π -type interactions (note that only σ -type interactions contribute to ρ_{BCP}). This latter feature is of great value to the present study, since any $6p_\sigma$ contributions to metal–ligand interactions will necessarily be of σ_u -character.

To clarify the role of the ITI, we again consider the consequences of freezing the 6p shell when the geometry is held fixed at that obtained with the fully relaxed electronic structure. In Table 1, we report the effect on $\delta(U,L)$. In passing, we note that the absolute magnitude of $\delta(U,L)$ is largest for the imido analogue (see Table S1 of ESI†). In the case of uranyl, the delocalisation index increases significantly when the 6p shell is frozen. This appears counterintuitive at first, since the effect of freezing this shell is to render it unavailable for chemical interactions. It can, however, be rationalised in the context of the “pushing from below” description of bonding in uranyl, which recognises that the $6p_\sigma$ orbital interacts with antibonding character.³¹ Antibonding interactions serve to lower the delocalisation index and so removing 6p contributions therefore leads to the increase found here. The same effect is seen for the imido analogue, although it is reduced in magnitude, particularly when using the CASSCF methodology. In the carbene analogue it is almost entirely absent.

The decomposition of $\delta(U,L)$ further supports this interpretation, since the σ_u component, in which $6p_\sigma$ interactions would manifest themselves, accounts for the majority of the overall variation. This component is positive for uranyl and the imido analogue, but small and negative for the carbene analogue. This trend suggests that the pushing from below mechanism is reduced when the oxo ligand is substituted by a less polarising species and is, perhaps, reversed in the carbene analogue. There is also a small σ_g component which is relatively constant across the different complexes and effectively cancels the σ_u contribution in the carbene complex.

Whilst $\delta(U,L)$ characterises the electron sharing between the uranium centre and the coordinating species, it tells us little

Table 1 Differences in uranium–ligand delocalisation indices, as well as the electron density at the U–L BCP, calculated with a frozen or unfrozen 6p shell, obtained at geometries optimised using a fully relaxed electronic structure

Complex	Method	$\Delta\delta(U,L)$					$\Delta\rho_{BCP}$		
		Total	σ_u	σ_g	π_u	π_g	Total	u	g
UO_2^{2+}	DFT	0.097	0.065	0.005	0.012	0.014	0.009	0.003	0.006
	CASSCF	0.082	0.044	0.014	0.008	0.016	0.008	0.003	0.005
$U(NH_2)_2^{2+}$	DFT	0.081	0.036	0.018	0.010	0.016	0.002	−0.003	0.006
	CASSCF	0.042	0.014	0.016	0.002	0.010	0.005	0.000	0.006
$U(CH_2)_2^{2+}$	DFT	0.008	−0.015	0.015	0.000	0.009	−0.007	−0.011	0.004
	CASSCF	−0.005	−0.015	0.008	0.001	0.003	−0.007	−0.008	0.001



Complex	Method	$Q(U)$	$Q(L)$	$\lambda(U)$	$\lambda(L)$
UO_2^{2+}	DFT	+3.270(+3.101)	-0.635(-0.551)	86.446(86.498)	7.438(7.289)
	CASSCF	+3.328(+3.148)	-0.664(-0.574)	86.857(86.955)	7.724(7.590)
$U(NH)_2^{2+}$	DFT	+2.934(+2.753)	-0.467(-0.377)	86.634(86.726)	7.197(7.060)
	CASSCF	+3.004(+2.850)	-0.502(-0.425)	87.070(87.270)	7.574(7.430)
$U(CH_2)_2^{2+}$	DFT	+2.303(+2.171)	-0.152(-0.086)	87.517(87.638)	6.978(6.884)
	CASSCF	+2.354(+2.250)	-0.177(-0.125)	88.130(88.230)	7.352(7.310)

Complex	Method	r_{UL} (Å)	Δr_{UL} (Å)
UO ₂ ²⁺	DFT	1.684 (1.740)	0.056
	CASSCF	1.709 (1.775)	0.066
U(NH) ₂ ²⁺	DFT	1.746 (1.801)	0.055
	CASSCF	1.783 (1.874)	0.091
U(CH ₂) ₂ ²⁺	DFT	1.905 (1.954)	0.049
	CASSCF	1.970 (2.061)	0.091

Table S4 of ESI.† As expected, this reduces its magnitude although the obtained trends are very similar. Broadly speaking, the same trends can be found in the electronic structure data, as seen in Tables S1–S7 of ESI.†

In summary, we have, for the first time, characterised the ITI in uranyl and its analogues *via* analysis of electron densities calculated at the DFT and CASSCF levels of theory, finding that both approaches yield qualitatively similar results. Denning's assertion of the nature of uranium polarisation by strongly coordinating ligands is verified, and the increase in electron sharing and charge accumulation in the metal–ligand bonding region found when the 6p shell is forced to assume localised atomic character can be rationalised as due to an antibonding $6p_{\sigma}$ contribution to the metal–ligand interaction, commensurate with the pushing from below mechanism. Of the complexes considered here, the effect of the ITI on electronic structure is found to be most pronounced in uranyl, consistent with the strong polarising ability of the oxo ligands, and this is reflected in the CPSE energies, where the ITI is found to stabilise the *trans* U–O_{yl} bonding interactions by approximately 0.76 eV. Interestingly, the magnitude of the ITI does not correlate with the elongation of the bond found when the ITI is suppressed, and this is interpreted as being due to an interplay of increased covalent and decreased ionic interactions, as previously asserted by La Pierre and Meyer.³

The previously recognised key role of the $6p_{\sigma}$ interactions in the ITI is supported by this study, and the analysis approach taken here is applicable to any systems in which σ -type metal–ligand interactions can be identified *via* symmetry considerations. Current investigations are focussed on oxohalides of the form AnOX₅, for which structural and spectroscopic data are available.

Conflicts of interest

There are no conflicts to declare.

Notes and references

‡ Both uranyl and its imido analogue were found to be linear, whereas the D_{2h} -symmetry carbene analogue was found to lie just 3 meV higher in energy than the C_{∞} -symmetry isomer at the DFT level. This energy difference was sufficiently small that the D_{2h} -symmetry complex was deemed usable for ease of later analysis.

§ Test CASSCF(12,16) uranyl simulations, which included the four non-bonding 5f orbitals, gave structural parameters to within 0.001 Å of those calculated with CASSCF(12,12).

- 1 K. W. Bagnall, J. du Preez, J. Barry and J. H. Holloway, *Dalton Trans.*, 1975, 1963–1968.
- 2 B. D. Brown, C. T. Reynolds and P. T. Moseley, *Dalton Trans.*, 1972, 857–859.
- 3 H. S. La Pierre and K. Meyer, *Inorg. Chem.*, 2013, 52, 529–539.
- 4 H. S. La Pierre, M. Rosenzweig, B. Kosog, C. Hauser, F. W. Heinemann, S. T. Liddle and K. Meyer, *Chem. Commun.*, 2015, 51, 16671–16674.
- 5 O. P. Lam, S. M. Franke, H. Nakai, F. W. Heinemann, W. Hieringer and K. Meyer, *Inorg. Chem.*, 2012, 51, 6190–6199.
- 6 A. J. Lewis, K. C. Mullane, E. Nakamaru-Ogiso, P. J. Carroll and E. J. Schelter, *Inorg. Chem.*, 2014, 53, 6944–6953.
- 7 A. J. Lewis, P. J. Carroll and E. J. Schelter, *J. Am. Chem. Soc.*, 2013, 135, 13185–13192.
- 8 B. Kosog, H. S. La Pierre, F. W. Heinemann, S. T. Liddle and K. Meyer, *J. Am. Chem. Soc.*, 2012, 134, 5284–5289.
- 9 M. Gregson, E. Lu, D. Mills, F. Tuna, E. McInnes, C. Hennig, A. Scheinost, J. McMaster, W. Lewis, A. Blake, A. Kerridge and S. T. Liddle, *Nat. Commun.*, 2017, 8, 14137.
- 10 R. G. Denning, *Struct. Bonding*, 1992, 79, 215–276.
- 11 R. G. Denning, *J. Phys. Chem. A*, 2007, 111, 4125–4143.
- 12 K. Tatsumi and R. Hoffmann, *Inorg. Chem.*, 1980, 19, 2656–2658.
- 13 E. O'Grady and N. Kaltsoyannis, *J. Chem. Soc., Dalton Trans.*, 2002, 1233–1239.
- 14 A. Kerridge, *Chem. Commun.*, 2017, 53, 6685–6695.
- 15 V. V. Zhurov, E. A. Zhurova, A. I. Stash and A. A. Pinkerton, *J. Phys. Chem. A*, 2011, 115, 13016–13023.
- 16 C. G. Gianopoulos, V. V. Zhurov, S. G. Minasian, E. R. Batista, C. Jelsch and A. A. Pinkerton, *Inorg. Chem.*, 2017, 56, 1775–1778.
- 17 T. W. Hayton, J. M. Boncella, B. L. Scott, P. D. Palmer, E. R. Batista and P. J. Hay, *Science*, 2005, 310, 1941–1943.
- 18 J. L. Brown, S. Fortier, G. Wu, N. Kaltsoyannis and T. W. Hayton, *J. Am. Chem. Soc.*, 2013, 135, 5352–5355.
- 19 Y. Gong, C. L. Pereira, C. Michelini, J. Marc and J. K. Gibson, *Inorg. Chem.*, 2013, 52, 14162–14167.
- 20 D. P. Mills, O. J. Cooper, F. Tuna, E. J. L. McInnes, E. S. Davies, J. McMaster, F. Moro, W. Lewis, A. J. Blake and S. T. Liddle, *J. Am. Chem. Soc.*, 2012, 134, 10047–10054.
- 21 K. Pierloot and E. van Besien, *J. Chem. Phys.*, 2005, 123, 204309.
- 22 R. G. Denning, J. C. Green, T. E. Hutchings, C. Dallera, A. Tagliaferri, K. Giarda, N. B. Brookes and L. Braicovich, *J. Chem. Phys.*, 2002, 117, 8008–8020.
- 23 S. G. Minasian, J. M. Keith, E. R. Batista, K. S. Boland, D. L. Clark, S. D. Conradson, S. A. Kozimor, R. L. Martin, D. E. Schwarz, D. K. Shuh, G. L. Wagner, M. P. Wilkerson, L. E. Wolfsberg and P. Yang, *J. Am. Chem. Soc.*, 2012, 134, 5586–5597.
- 24 L. P. Spencer, P. Yang, S. G. Minasian, R. E. Jilek, E. R. Batista, K. S. Boland, J. M. Boncella, S. D. Conradson, D. L. Clark, T. W. Hayton, S. A. Kozimor, R. L. Martin, M. M. MacInnes, A. C. Olson, B. L. Scott, D. K. Shuh and M. P. Wilkerson, *J. Am. Chem. Soc.*, 2013, 135, 2279–2290.
- 25 A. Formanui, A.-M. Ariciu, F. Ortu, R. Beekmeyer, A. Kerridge, F. Tuna, E. J. L. McInnes and D. P. Mills, *Nat. Chem.*, 2017, 9, 578–583.
- 26 A. Formanui, F. Ortu, R. Beekmeyer, A. Kerridge, R. W. Adams and D. P. Mills, *Dalton Trans.*, 2016, 45, 2390–2393.
- 27 A. Formanui, F. Ortu, J. Liu, L. Nodaraki, F. Tuna, A. Kerridge and D. P. Mills, *Chem. – Eur. J.*, 2017, 23, 2290–2293.
- 28 A. C. Behrle, A. Kerridge and J. R. Walensky, *Inorg. Chem.*, 2015, 54, 11625–11636.
- 29 S. D. Woodall, A. N. Swinburne, A. Kerridge, P. Di Pietro, C. Adam, P. Kaden and L. S. Natrajan, *Chem. Commun.*, 2015, 51, 5402–5405.
- 30 G. Karlström, R. Lindh, P.-Å. Malmqvist, B. O. Roos, U. Ryde, V. Veryazov, P.-O. Widmark, M. Cossi, B. Schimmelpfennig, P. Neogrady and L. Seijo, *Comput. Mater. Sci.*, 2003, 28, 222–239.
- 31 C. K. Jorgensen and R. Reisfeld, *Struct. Bonding*, 1982, 50, 121.
- 32 P. Di Pietro and A. Kerridge, *Inorg. Chem.*, 2016, 55, 573–583.
- 33 G. Schreckenbach, P. J. Hay and R. L. Martin, *Inorg. Chem.*, 1998, 37, 4442–4451.

