Chemical Science



EDGE ARTICLE

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2019, 10, 3231

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 6th December 2018 Accepted 3rd February 2019

DOI: 10.1039/c8sc05445b

rsc.li/chemical-science

Nickel-catalyzed anti-Markovnikov hydroarylation of alkenes†

Julia Nguyen, Andrea Chong and Gojko Lalic **D**

We have developed a nickel-catalyzed hydroarylation of alkenes using aryl halides as coupling partners. Excellent anti-Markovnikov selectivity is achieved with aryl-substituted alkenes and enol ethers. We also show that hydroarylation occurs with alkyl substituted alkenes to yield linear products. Preliminary examination of the reaction mechanism suggests irreversible hydrometallation as the selectivity determining step of the hydroarylation.

Catalytic hydroarylation of alkenes has been targeted as an attractive synthetic route to alkylarenes, a common structural motif in organic chemistry. Traditionally, hydroarylation is accomplished either *via* classical Friedel–Crafts reaction¹ or through methods based on metal catalyzed C–H activation of arenes.² More recently, a third complementary strategy has been developed based on the reductive coupling of aryl halides and alkenes in the presence of a hydride donor. The key advance offered by reductive cross coupling relative to Friedel–Crafts and C–H activation methods is excellent site selectivity of the arene alkylation regardless of the substitution pattern of the starting arene.

The reductive cross coupling approach has been effectively used in several methods for the Markovnikov hydroarylation of alkenes. In 2016, Nakao *et al.* reported a Markovnikov hydroarylation of arylation of arylation of arylation and alkenes using a dual palladium/copper catalyst system.³ The same year, Buchwald *et al.* reported an enantioselective version of the reaction using a similar approach.⁴ Markovnikov hydroarylation of alkenes has also been achieved by Herzon *et al.*^{5,6} using metal hydride hydrogen atom transfer (HAT) to initiate radical alkylation of arenes. Concurrently, Shenvi *et al.*^{7,8} developed a distinct approach based on metal hydride HAT coupled with nickel-catalyzed cross-coupling.

Anti-Markovnikov selectivity in the reductive cross coupling of aryl halides and alkenes is relatively rare. At the time we were developing this project, it had been achieved only in reactions of simple terminal alkenes (Scheme 1a). Buchwald *et al.* have shown that a dual palladium/copper catalyst system promotes coupling of unactivated alkenes with aryl halides to generate linear hydroarylation products.^{9,10} Notably, the same catalyst system was previously shown to give Markovnikov selectivity in the hydroarylation of aryl alkenes.⁴ Recently, an alternative

 $\label{lem:prop:condition} Department of Chemistry, \ University of Washington, Seattle, \ Washington 98195, \ USA. \\ E-mail: lalic@chem.washington.edu$

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8sc05445b

approach based on reductive Heck reaction has been successfully used by Engle *et al.*¹¹ and later Hu *et al.*¹² to access the same products. Relative to the excellent scope of the Engle's reaction of terminal alkyl alkenes, the scope of the Hu's reaction with aryl alkenes was limited.

We were interested in developing a hydroarylation reaction that would complement existing methods and allow anti-Markovnikov hydroarylation of aryl alkenes and possibly other new substrate classes. Furthermore, we were hoping to achieve these transformations using a single metal complex as a catalyst. This catalyst would have to promote both the hydrometallation of alkenes and the subsequent cross coupling with aryl halides.

We focused on nickel complexes as particularly well suited for the task. Insertion of alkenes into nickel hydride complexes is well documented^{13,14} and alkyl nickel complexes have been implicated as intermediates in cross-coupling reactions with organohalides.¹⁵ Following an important precedent established by Liu *et al.*,¹⁶ the general feasibility of nickel-catalyzed hydroarylation was recently demonstrated by a reductive arylation reaction reported by Zhu *et al.* (Scheme 1b).^{17,18} This transformation involved extensive isomerization of the alkene and the exclusive formation of 1,1-diaryl products regardless of the original position of the alkene relative to the arene.¹⁹

In this work, we report the anti-Markovnikov hydroarylation of alkenes using a simple nickel catalyst. We are able to achieve this linear selectivity with a number of different substrate classes, including aryl alkenes, enol ethers and simple unactivated alkenes (Scheme 1c).

While exploring the reactivity of various nickel complexes in the catalytic hydroarylation of aryl alkenes, we discovered that NiCl₂(DME) provides the hydroarylation product with excellent anti-Markovnikov selectivity (Table 1, entry 1). Essential for the anti-Markovnikov selectivity and the overall success of the reaction was the absence of a ligand. A variety of ligands, including L1 used by Zhu *et al.*, resulted in no formation of the desired product (entries 2 and 3) (see ESI for additional data†).

a) Anti-Markovnikov hydroarylation of alkyl-substituted alkenes

b) Alkene isomerization in nickel-catalyzed hydroarylation

c) Nickel-catalyzed anti-Markovnikov hydroarylation of alkenes

Scheme 1 Reductive cross coupling of alkenes with aryl halides

Table 1 Reaction development^a

Chemical Science

Entry	Variations from above	Yield ^b (%)	$L: B^c$
1	None	89 (81)	55:1
2	L1 as a ligand	0	_
3	L2 as a ligand	0	_
4	IPr as a ligand	24	1.3:1
5	NiCl ₂ instead of NiCl ₂ (dme)	44	15:1
6	Ni(COD) ₂ instead of NiCl ₂ (dme)	39	12:1
7	Toluene instead of benzene	80	48:1
8	1,4-Dioxane	35	3:1
9	THF instead of benzene	8	2.5:1
10	(Me ₂ HSi) ₂ O instead of PMHS	57	29:1
11	LiOt-Bu instead of NaOt-Bu	30	7:1
12	KOt-Bu instead of NaOt-Bu	0	_
	$Ar \sim N \sim Ar$ IPr : $Ar = (2,6)$ - i Pr-Ph $PMHS = polymethylhydrosiloxane$	L1: R = Me L2: R = H	

^a Conditions: 1 (2 equiv.), 2 (1 equiv.), NaOt-Bu (4 equiv.), PMHS (4 equiv.). ^b Yields were determined by GC analysis of the crude reaction mixture using 1,3,5-trimethylbenzene as the internal standard. Number in parentheses are yields of isolated products. ^c L:B is the ratio of the linear product to the branched product as determined by GC analysis of the crude reaction mixture.

For those ligands that did generate product, such as N-heterocyclic carbine ligand **IPr**, regioselectivity and yield were low (entry 4). Using different nickel sources, such as NiCl₂ (entry 5), resulted in lower yields. Interestingly, Ni(COD)₂, a Ni(0)

source, proved to be competent in this reaction (entry 6). Aromatic solvents, such as benzene and toluene (entry 7), were essential for high yield of the hydroarylation product, while most other solvents were completely ineffective (see ESI for

Table 2 Scope of the hydroarylation reaction^a

able 2 scope of the flydrodryte		NiCl ₂ (DME) (10 mol%)			
	R + Ar-X	PMHS, NaO <i>t</i> -Bu benzene, 25 °C, 14 h	R Ar		
$Ar-X = 4-\text{MeOC}_{\theta}H_4-I$	Y = H 3, 81% (5 Ar t-Bu 4, 85% (4 MeO 5, 81% (4	6:1) Ar	Me	Ar	
	F 6 , 78% (2	7 , 80% (20:1)	8, 73% (66:1)	9 , 83% (34:1)	
Ar O	Ar	Ar	Ar	Ar Me	
10 , 90% (40:1)	11 , 94% (22:1)	12 , 84% (79:1)	13 , 86% (38:1)	14 , 95% (65:1)	
Ar N _{Me}	Ar N. Me	N Ar	Ar Me	Me	
15 , 78% (49:1)	16 , 95% (56:1)	17 , 66% (5:1)	18 , 72% (3:1)	19 , 56% (30:1)	
Ar-X = 4-MeOC ₆ H ₄ -I	Ar Y = MeO Bpin	20 , 74% (10:1) 21 , 61% (10:1)	Ar	Ar	
γ, ψ	F	22 , 74% (11:1)	23, 85% (10:1)	24 , 84% (7:1)	
Y = Et		Ar 7, 56% (10:1)	Ar 28, 61% (9:1)	Me 29, 78% (6:1)	
		1, 50% (10.1)	20, 0178 (0.1)	20, 7078 (0.1)	
Ph Y Y	= NMe ₂ 30, 85% (19:1) Cl 31, 46% (15:1) CO ₂ Me 32, 43% (24:1)	Ph 33, 86% (55:1)	Ph Me 34, 66% (71:1)	Me Ph 35, 78% (44:1)	
Ar-Br t-Bu	Ph	Рһ	S Ph F	Me N Boc	
36 , 81% (65:1)	37 , 99% (22:1)	38 , 70% (31:1)	39 , 73% (44:1)	40 , 81% (41:1)	
Ph	Ph	Ph	Ph Ne	Ph	
41 , 78% (13:1)	42 , 44% (21:1)	43 , 70% (11:1)	44 , 94% (36:1)	45 , 76% (41:1)	
Ph) N	Ph N N	Ph	N-Me	
46 , 93% (26:1)		47 , 70% (8:1)	48 , 69%	48, 69% (21:1)	

 $[^]a$ Conditions: aryl halide (1 equiv.), alkene (2 equiv.), PMHS (4 equiv.), NaOt-Bu (4 equiv.). Yields of isolated products are reported. The ratios of linear to branched products were determined by GC analysis of the crude reaction mixture and are given in parenthesis.

details†). Interestingly, ethereal solvents, such as 1,4-dioxane

Chemical Science

(entry 8) and THF (entry 9), provided significantly lower regioselectivity.

Polymethylhydrosiloxane (PMHS) was the best hydride source. Other silanes, including closely related (Me₂HSi)₂O (entry 10), provided lower yields and worse regioselectivity (see ESI† for details). Similarly, NaOt-Bu was essential for the reaction and even the closely related Li and K analogs (entries 11 and 12) provided significantly inferior results. Exclusion of any component of this reaction resulted in no formation of product 3.

Hydroarylation could be achieved with a range of aryl- and alkyl-substituted alkenes (Table 2). With aryl alkenes, ortho, meta and para substitution is tolerated (compounds 7, 8, 4). Aryl alkenes containing both electron-donating and electronwithdrawing substituents are viable substrates (5 and 6). Good selectivities and yields were also obtained in hydroarylation of aryl alkenes containing a wide range of heteroarenes, including benzofuran (13), indole (14, 15), pyrrole (16), benzopyrazole (17), and pyrrolopyridine (18). Finally, substitution on the α position of the alkene is also tolerated, as indicated by the formation of product 19 in the reaction with α -methylstyrene.

Nickel-catalyzed hydroarylation of various other classes of alkenes was also examined. Unactivated terminal alkenes were suitable coupling partners (20-23, 27-29), although the regioselectivity was somewhat lower. Surprisingly, a 1,3-diene substrate gave product 24 with high yield and good regioselectivity. Another unusual class of alkenes that were compatible with our reaction conditions are enol ethers. We were able to isolate the product of the reaction with ethyl vinyl ether (25) in good yield (78%) while the benzyl protected enol ether provided the expected product 26 in 63% yield. To our knowledge, these are the only examples of anti-Markovnikov reductive cross coupling reactions with 1,3-dienes or enol ethers.

We have also explored the performance of various arene coupling partners, and both aryl iodides and aryl bromides are viable substrates (Table 2). The electronic properties of aryl electrophiles seem to play a significant role as electron-neutral and electron-rich aryl halides generally gave high yields (30, 37, 46), while electron-poor aryl halides, though still competent, provided lower yields of the hydroarylation products (31 and 32). Again, we found that various heteroarenes are tolerated, including pyrrolopyridine (41), quinoline (42, 43), indole (44), benzofuran (45), pyrimidine (47), and benzopyrazole (48).

The unique aspect of this new hydroarylation reaction is the anti-Markovnikov selectivity obtained with a range of alkenes, including aryl alkenes and enol ethers. The observed selectivity is consistent with an addition of an aryl radical to the alkene, which would favour addition to the alkene terminus and formation of the more stable secondary alkyl radical. To test this mechanistic hypothesis, we performed experiments in the presence of a radical trap. The presence of 10 mol% of TEMPO did lower the yield of the desired product and a full equivalent completely blocked its formation (see ESI† for details). However, in all cases, we observed only trace amounts (<5% yield) of the aryl-TEMPO adduct, suggesting that TEMPO likely interacts with another intermediate20 and not with an aryl radical.21 In competition experiments (Scheme 2a), we found that simple styrene reacts faster than α -methylstyrene or α phenylstyrene, both of which should undergo radical addition faster than styrene. The results of these experiments are not consistent with a radical addition mechanism, leading us to explore other options.

A mechanism involving hydrometallation followed by arylation of the alkyl nickel intermediate was proposed by Zhu et al. for the related hydroarylation reaction (see Scheme 1b). We used the experiment shown in Scheme 2b to explore the intermediacy of the alkyl nickel complex in our hydroarylation reaction.

In the presence of 2 equiv. of EtOH, the hydroarylation product is formed in a lower yield than in the absence of EtOH (54% vs. 82%), while the alkane product (50) is formed in 36% yield. This result is consistent with competitive protonation of the alkyl nickel intermediate by EtOH. Based on these considerations, we propose that the hydroarylation reaction proceeds according to the mechanism shown in Scheme 3.

In the context of this mechanism, we were interested in the step that determines the regioselectivity of the reaction. In hydrofunctionalization reactions that proceed through hydrometallation, the product determining step varies depending on reversibility of the hydrometallation step. When

Scheme 2 Probing the reaction mechanism.

Edge Article

NaOt-Bu

NaOt-Bu

NiCl₂(DME) + R₃SiH +NaOt-Bu

R₃SiH

R₃SiOt-Bu

Ni-H

Scheme 3 Proposed mechanism of hydroarylation.

irreversible, hydrometallation is the regio-determining step. If hydrometallation is fast and reversible, there is a Curtin–Hammett scenario, and regioselectivity is determined by the relative rates of the irreversible reductive elimination at different sites and not by hydrometallation. Alkene insertion into nickel hydride complexes is often fast and reversible, ²² as documented by Zhu *et al.* in their reductive arylation of alkenes. ¹⁷ Because the C–C bond-forming reductive elimination from nickel complexes tends to be exothermic, ²³ the rates of reductive elimination are often based on the ground state stability of the alkyl metal intermediates. The result is that the reductive elimination from the most stable alkyl metal complex is favoured.

To explore the reversibility of the hydrometallation step in the hydroarylation reaction, we used deuterium-labelled styrene substrates (Scheme 2c and d). In the reaction with substrate 1- βD we observed no deuterium scrambling, while the reaction with 1- αD , resulted in minimal scrambling of the deuterium label.²⁴ In the context of the reaction mechanism shown in Scheme 3, the two deuterium scrambling experiments point to hydrometallation being irreversible and product determining step of the reaction. These results indicate that a strong kinetic preference for the formation of a homobenzylic alkyl nickel intermediate over the benzylic regioisomer is responsible for the regioselectivity of the hydroarylation.

The unusual selectivity in the hydrometallation of styrenes raises questions about the nature of the active catalyst. The absence of ligand in our hydroarylation reaction makes the formation of nanoparticles likely, and we observe an immediate formation of particles upon the addition of the alkoxide to the reaction mixture. Mixtures of alkoxides and reducing reagents have previously been used to generate highly reactive nickel nanoparticle catalysts, and sodium alkoxide has been proposed to stabilize "ligandless" nickel catalysts. 25,26 Furthermore, in the hydroarylation reaction, we observed a burst of initial activity (64% yield after 10 minutes), followed by a very slow reaction over the next 14 h when the maximum yield is achieved. Finally, we found that the addition of mercury significantly lowers the yield of the hydroarylation (56% vs. 89% GC yields) (Scheme 2e).27 These observations are consistent with a heterogeneous active catalyst.

Conclusions

We have developed a new method for the anti-Markovnikov hydroarylation of alkenes. The reaction involves reductive cross coupling of alkenes and aryl halides in the presence of a silane as a hydride donor. Anti-Markovnikov selectivity is obtained with a wide range of alkenes, including alkyl- and aryl-substituted alkenes, and enol ethers. Preliminary investigation of the reaction mechanism provides evidence for a mechanism that involves hydrometallation of the alkene, followed by coupling of the alkyl nickel intermediate with an aryl halide. The evidence points to highly selective hydrometallation of the alkene as the regioselectivity determining step of the reaction.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Financial support by NSF (NSF CAREER Award 1254636) and NIH (1R01GM125791-01A1) is acknowledged.

Notes and references

- 1 G. A. Olah, R. Krishnamurti and G. K. Surya Prakash, in *Comprehensive Organic Synthesis*, ed. B. M. Trost and I. Fleming, Pergamom Press, Oxford, 1991, vol. 3, pp. 293–339.
- 2 For a comprehensive review of this large research area, see: Z. Dong, Z. Ren, S. J. Thompson, Y. Xu and G. Dong, *Chem. Rev.*, 2017, 117, 9333.
- 3 K. Semba, K. Ariyama, H. Zheng, R. Kameyama, S. Sakaki and Y. Nakao, *Angew. Chem., Int. Ed.*, 2016, **55**, 6275–6279.
- 4 S. D. Friis, M. T. Pirnot and S. L. Buchwald, *J. Am. Chem. Soc.*, 2016, 138, 8372.
- 5 X. Ma and S. B. Herzon, J. Am. Chem. Soc., 2016, 138, 8718.
- 6 X. Ma, H. Dang, J. A. Rose, P. Rablen and S. B. Herzon, *J. Am. Chem. Soc.*, 2017, **139**, 5998.
- 7 S. A. Green, J. L. M. Matos, A. Yagi and R. A. Shenvi, *J. Am. Chem. Soc.*, 2016, **138**, 12779.
- 8 S. L. Shevick, C. Obradors and R. A. Shenvi, *J. Am. Chem. Soc.*, 2018, **140**, 12056–12068.
- 9 S. D. Friis, M. T. Pirnot, L. N. Dupuis and S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2017, 56, 7242.
- 10 For a recent example of anti-Markovnikov hydropyridylation of alkenes using photocatalysis, see: A. J. Boyington, M. L. Y. Riu and N. T. Jui, J. Am. Chem. Soc., 2017, 139, 6582.
- 11 J. A. Gurak Jr and K. M. Engle, ACS Catal., 2018, 8, 8987–8992.
- 12 L. Jin, J. Qian, N. Sun, B. Hu, Z. Shen and X. Hu, *Chem. Commun.*, 2018, 54, 5752.
- 13 N. A. Eberhardt and H. Guan, Chem. Rev., 2016, 116, 8373.
- 14 R. Cariou, T. W. Graham and D. W. Stephan, *Dalton Trans.*, 2013, 42, 4237; L. C. Liang, P. S. Chien and P. Y. Lee, *Organometallics*, 2008, 27, 3082; J. Breitenfeld, R. Scopelliti and X. Hu, *Organometallics*, 2012, 31, 2128.

15 T. J. Anderson, G. D. Jones and D. A. Vicic, J. Am. Chem. Soc., 2004, 126, 8100; J. Breitenfeld, J. Ruiz, M. D. Wodrich and X. Hu, J. Am. Chem. Soc., 2017, 139, 13929.

Chemical Science

- 16 Liu et al. have reported two examples of nickel-catalyzed hydroarylation reaction: X. Lu, B. Xiao, Z. Zhang, T. Gong, W. Su, J. Yi, Y. Fu and L. Liu, Nat. Commun., 2016, 7, 11129.
- 17 F. Chen, K. Chen, Y. Zhang, Y. He, Y. M. Wang and S. Zhu, *J. Am. Chem. Soc.*, 2017, **139**, 13929.
- 18 For examples of related reductive cross-coupling reactions, see: P. Shukla, A. Sharma, B. Pallavi and C. H. Cheng, *Tetrahedron*, 2015, 71, 2260; Q. Xurong, L. M. W. Yao and Z. J. Steve, *Angew. Chem., Int. Ed.*, 2017, 56, 12723.
- 19 Interestingly, Zhao *et al.* have also reported Markovnikov selectivity in a redox neutral nickel-catalysed hydroarylation of styrenes, see: L. J. Xiao, L. Cheng, W. M. Feng, M. L. Li, J. H. Xie and Q. L. Zhou, *Angew. Chem., Int. Ed.*, 2018, 57, 461.
- 20 TEMPO has been shown to react rapidly with metal hydrides. See: A. C. Albéniz, P. Espinet, R. López-Fernández and A. J. Sen, J. Am. Chem. Soc., 2002, 124, 11278TEMPO has also been used as an oxidant in nickel-catalyzed C-H

- activation reactions. See: X. Wu, Y. Zhao and H. Ge, Chem.-Eur. J., 2014, 20, 9530.
- 21 We cannot exclude the possibility that the Ar–TEMPO adduct reacts further under the reaction conditions.
- 22 C. A. Tolman, J. Am. Chem. Soc., 1972, 94, 2994; R. Cramer and R. V. Lindsey, J. Am. Chem. Soc., 1966, 88, 3534.
- 23 V. P. Ananikov, ACS Catal., 2015, 5, 1964; S. A. Macgregor,
 G. W. Neave and C. Smith, Faraday Discuss., 2003, 124, 111.
- 24 The deuterium scrambling is also observed in the recovered 1-αD and is higher at higher conversions. Furthermore, if the reaction with 1-αD is stopped after 10 minutes, product 51 is isolated in 49% yield and no deuterium scrambling is observed in either the product or the recovered starting material (see page S31 of the ESI†). These results suggest that the scrambling may be independent of the hydroarylation reaction.
- 25 I. Buslov, F. Song and X. Hu, Angew. Chem., Int. Ed., 2016, 55, 12295; A. G. Sergeey, J. D. Webb and J. F. Hartwig, J. Am. Chem. Soc., 2012, 134, 20226.
- 26 J. J. Brunet, D. Besozzi, A. Courtois and P. Caubere, *J. Am. Chem. Soc.*, 1982, **104**, 7130.
- 27 R. H. Crabtree, Chem. Rev., 2012, 112, 1536.