

ChemComm

This article is part of the

Glycochemistry & glycobiology web themed issue

Guest Editors are Professor Peter H. Seeberger and Dr Daniel B. Werz

Please visit the website to access the other articles in this issue:-

<http://www.rsc.org/chemcomm/glycochemistry>



Cite this: *Chem. Commun.*, 2011, **47**, 10260–10262

www.rsc.org/chemcomm

COMMUNICATION

Progress toward developing a carbohydrate-conjugate vaccine against *Clostridium difficile* ribotype 027: synthesis of the cell-surface polysaccharide PS-I repeating unit†‡Christopher E. Martin,^{ab} Markus W. Weishaupt^{ab} and Peter H. Seeberger^{*ab}

Received 17th June 2011, Accepted 21st July 2011

DOI: 10.1039/c1cc13614c


Clostridium difficile strain ribotype 027 is a hypervirulent pathogen that is responsible for recent, severe outbreaks of serious nosocomial infections. As a foundation for the development of a preventative carbohydrate-based vaccine, we have synthesized a pentasaccharide cell wall repeating unit from PS-I unique to this strain, by the linear assembly of four monosaccharide building blocks.

Clostridium difficile is a Gram-positive, spore-forming anaerobic bacterium that colonizes the intestinal tract of humans thus leading to *C. difficile* infections (CDI).¹ CDI have become the most commonly diagnosed cause of hospital-acquired diarrhea, particularly in risk groups including the elderly and immunodeficient patients, as well as those receiving antibiotic treatment. A steep rise in the incidence of CDI over the past decade is attributed to the emergence of the hypervirulent, and now predominant strain ribotype 027 that causes epidemic outbreaks with increased morbidity, mortality and high rates of relapse.² These infections significantly increase the treatment costs of patients, particularly in the case of recurring CDI.³ Prevention of CDI is therefore desirable, and vaccination of risk groups may be a useful and cost-efficient means to avoid future infections. Although models have shown that vaccination against *C. difficile* should be economically feasible,⁴ a vaccine has not yet been developed.

Carbohydrates exposed on the cell-surface of pathogens are often immunogenic and constitute potential candidates for vaccine development. When covalently connected to a carrier protein, a carbohydrate antigen can elicit long lasting, T-cell-dependent protection.⁵ The chemical structures of two *C. difficile* cell-surface polysaccharides, PS-I and PS-II were recently elucidated.⁶ Initially, the focus was directed towards the PS-II hexasaccharide repeating unit antigen⁷ because it is common to several *C. difficile* strains. PS-I consists of a

pentasaccharide phosphate repeating unit with the structure $[\rightarrow 4)\text{-}\alpha\text{-Rhap-(1} \rightarrow 3)\text{-}\beta\text{-Glc-(1} \rightarrow 4)\text{-}[\alpha\text{-Rhap-(1} \rightarrow 3)\text{-}\alpha\text{-Glc-(1} \rightarrow 2)\text{-}\alpha\text{-Glc-(1} \rightarrow \text{P}]$ and, so far, has been found to be expressed by *C. difficile* ribotype 027 only.⁶ The branched pentasaccharide repeating unit **1** of PS-I is shown in Fig. 1. Units A and B are 1 \rightarrow 2 linked α -Glc residues, wherein A is equipped with the aminopentyl linker at the reducing terminus, and B constitutes the branching point. B connects to α -Rha residue D at C-3 and β -Glc residue C at C-4. Finally, residue D' is the terminal α -Rha, linked to C-3 of C.

Here we report the first total synthesis of pentasaccharide **1**, the synthetic challenge was met by devising a linear strategy using a set of building blocks **2–7** (Fig. 1). This linear strategy allowed for the installation of the challenging 1,2-*cis* glycosidic linkages of both residues A and B early in the synthetic pathway by using building blocks **2** and **3**. Formation of the linkage between C and the AB fragment was investigated using three different glycosylating agents of C, namely thioglycoside


Fig. 1 Retrosynthesis of the PS-I pentasaccharide repeating unit **1**.

^a Max Planck Institute of Colloids and Interfaces, Department of Biomolecular Systems, Am Mühlenberg 1, 14476 Potsdam, Germany. E-mail: peter.seeberger@mpikg.mpg.de

^b Freie Universität Berlin, Institute for Chemistry and Biochemistry, Arnimallee 22, 14195 Berlin, Germany

† This article is part of the ChemComm 'Glycochemistry and glycobiology' web themed issue.

‡ Electronic supplementary information (ESI) available: Experimental details and full spectroscopic data for all new compounds. See DOI: 10.1039/c1cc13614c




Scheme 1 Synthesis of building block **2**. Reagents and conditions: (a) NaH, NAPBr, DMF, 0 °C to rt, 92%; (b) HO(CH₂)₅NBnCbz, NIS, TfOH, toluene/dioxane, −40 °C to −20 °C; (c) DDQ, DCM, H₂O, 35% over 2 steps.

4, *N*-phenyl trifluoroacetimidate **5** and glycosyl phosphate **6**. In order to minimize the number of deprotection and glycosylation steps, both Rha residues D and D' were added to the intermediate trisaccharide ABC in a single bis-glycosylation reaction with building block **7**.

The synthesis of building block **2** is depicted in Scheme 1. The latent amine at the reducing end was introduced by union of thioglucoside **9** and the linker.⁸ Thereby, 2-naphthylmethyl (NAP) served as a non-participating temporary protecting group at C-2 in order to achieve formation of the α -glycosidic linkage. Subsequent 3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) mediated cleavage of the NAP ether resulted in building block **2**.⁹

All four building blocks **3–6** originate from the key intermediate **13** (Scheme 2). This common precursor is synthesized




Scheme 2 Synthesis of monosaccharide building blocks **3–6**. Reagents and conditions: (a) 2-methyl-5-*tert*-butylthiophenol, BF₃·OEt₂, DCM, 85%; (b) NaOMe, MeOH, rt; (c) benzaldehyde dimethyl acetal, CSA, MeCN, 87% over 2 steps; (d) TBS-Cl, imidazole, DMF, 0 °C, 69%; (e) NaH, BnBr, DMF, 0 °C to rt; (f) 1 M TBAF in THF, 0 °C to rt, 93% over 2 steps; (g) Fmoc-Cl, pyridine, DCM, 95%; (h) TES, TfOH, DCM, 4 Å MS, −78 °C, 73%; (i) Lev₂O, pyridine, DCM, 3 days, 79%; (j) BzCl, DMAP, pyridine, 70 °C, 88%; (k) TBAF·3H₂O, AcOH, DMF, 35 °C, 91%; (l) Fmoc-Cl, pyridine, DCM, 96%; (m) NIS, AgOTf, TTBP, MeCN, H₂O; (n) CF₃C(NPh)Cl, Cs₂CO₃, DCM, 67% over 2 steps; (o) HOPO(OBu)₂, NIS/TfOH, DCM, 4 Å MS, 0 °C, 81%.

in four steps from glucose pentaacetate **10**.¹⁰ We found that use of the nontoxic and odorless 2-methyl-5-*tert*-butylthiophenol group¹¹ ensures exclusive β -anomer formation of thioglucosides and results in storage-stable monomer units. The adaptable thiol group is readily converted to phosphates or imidates, thus allowing creation of the series of glycosylating agents **3–6**. The protecting group pattern of **13** enables the orthogonal protection of all hydroxyl groups of the hexose ring. This feature was exploited in the synthesis of building block **3**, where a non-participating benzyl group was installed at C-2 of intermediate **14** to favor the formation of the α -glycosidic linkage between the A and B fragments. Subsequent placement of the 3-*O*-Fmoc-protection furnished compound **15**. Finally, regioselective opening of the 4,6-*O*-benzylidene acetal with TES-TfOH and protection of the free 4-hydroxyl gave orthogonally-protected building block **3**.

Preparation of differentially-protected thioglycoside **4** from **13** followed a similar route. In anticipation of the formation of a 1,2-*trans* linkage between the B and C saccharide fragments, a participating benzoyl group was installed at the C-2 hydroxyl of intermediate **17**. During TBAF-mediated desilylation of **17**, careful control of the TBAF:AcOH ratio was essential to prevent benzoyl-migration from C-2 to C-3. Fmoc-protected thioglycoside **4** was further diversified to glycosyl imidate **5** and glycosyl phosphate **6**. Building blocks **4–6** were individually evaluated for their reactivity towards the AB disaccharide fragment in assembly of the ABC trisaccharide. The terminal D and D' fragments of the pentasaccharide were provided by building block **7**. Synthesis of this rhamnosyl unit commenced with the bis-benzoylation of *p*-methoxyphenyl glycoside **19** (Scheme 3).¹²

CAN-mediated removal of the anomeric *p*-methoxyphenyl group yielded the free lactol that was immediately converted into rhamnosyl *N*-phenyl trifluoroacetimidate **7**.¹³

Assembly of the pentasaccharide target was achieved in seven linear steps by combining the monosaccharide building blocks in sequence (Scheme 4). The 1,2-*cis* glycosidic linkage between residues A and B was formed using nucleophile **2** and glycosylating agent **3**. Disaccharide **21** was obtained in good yield and stereoselectivity when NIS and TfOH in Et₂O was employed as promoter system. Selective cleavage of the levulinic ester with hydrazine hydrate in pyridine/AcOH, did not compromise the integrity of the Fmoc-group but cleanly produced disaccharide acceptor **22**. Installation of the next glycosidic linkage between C and the AB fragment to form trisaccharide **23** was attempted using the different building blocks **4–6** and the three reactions were screened for their



Scheme 3 Synthesis of rhamnosyl building block **7**. Reagents and conditions: (a) BzCl, DMAP, pyridine, DCM, 0 °C to rt, 97%; (b) CAN, MeCN, H₂O; (c) CF₃C(NPh)Cl, Cs₂CO₃, DCM, 74% over 2 steps.

