

Cite this: *Dalton Trans.*, 2016, **45**, 1953

Received 21st April 2015,

Accepted 3rd June 2015

DOI: 10.1039/c5dt01512j

www.rsc.org/dalton

Formation of the spirocyclic, Si-centered cage cations  $[\text{ClP}(\text{NSiMe}_3)_2\text{Si}(\text{NSiMe}_3)_2\text{P}_5]^+$  and  $[\text{P}_5(\text{NSiMe}_3)_2\text{Si}(\text{NSiMe}_3)_2\text{P}_5]^{2+}$ †

M. H. Holthausen and J. J. Weigand\*

On account of our interest in  $\text{P}_4$  activation by phosphonium ion insertion into P–P bonds we have developed synthetic routes to bicyclic N–P–Si-heterocycle **7** and probed its reactivity towards  $\text{GaCl}_3$  and  $\text{P}_4$ . A  $\text{GaCl}_3$ -induced rearrangement of **7** leads to the *in situ* formation of spirocyclic, Si-centered phosphonium ions. Their insertion into P–P bonds of one or two  $\text{P}_4$  tetrahedra yields polyphosphorus cages  $[\text{ClP}(\text{NSiMe}_3)_2\text{Si}(\text{NSiMe}_3)_2\text{P}_5]^+$  (**19**<sup>+</sup>) and  $[\text{P}_5(\text{NSiMe}_3)_2\text{Si}(\text{NSiMe}_3)_2\text{P}_5]^{2+}$  (**13**<sup>2+</sup>).

## Introduction

White phosphorus ( $\text{P}_4$ ) is an archetypal building block for the syntheses of polyphosphorus cations featuring a high P to substituent ratio.<sup>1</sup> In contrast to highly substituted cations  $\text{R}_n\text{P}_m$  ( $n > m$ ), which are obtained *via* synthetic methods based on *catena* or *cyclic* polyphosphanes,<sup>2</sup> the targeted preparation of P-rich cations  $\text{R}_n\text{P}_m$  ( $n < m$ ) is achieved by taking advantage of the tetrahedral  $\text{P}_4$  framework.<sup>1,3</sup> In a seminal paper, Krossing *et al.* reported that dicoordinated phosphonium ions, like other predominantly electrophilic ambiphiles,<sup>4</sup> are able to insert into a P–P bond of the  $\text{P}_4$  tetrahedron.<sup>5</sup> This was exploited for the preparation of a series of  $\text{P}_5\text{X}_2^+$ -cages (X = Cl, Br, I).<sup>5,6</sup> We expanded this methodology and prepared a series of symmetrically and unsymmetrically-substituted  $\text{R}_2\text{P}_5^+$ - and  $\text{RP}_5\text{Cl}^+$ -cations (R = aryl, alkyl,  $\text{R}_2\text{N}$ ).<sup>7</sup> The additional stability added by organo-substituents allowed for the stepwise insertion of up to three  $[\text{Ph}_2\text{P}]^+$  phosphonium ions into P–P bonds of one  $\text{P}_4$  molecule yielding mono- to tri-cationic  $[\text{Ph}_2\text{P}_5]^+$ ,  $[\text{Ph}_4\text{P}_6]^{2+}$ - and  $[\text{Ph}_6\text{P}_7]^{3+}$ -cages.<sup>8</sup> A complementary study exploited 1,3-dichloro-*cyclo*-1,3-diphosphadiazane  $\text{ClP}(\text{NDipp})_2\text{P}(\text{Cl})$  (Dipp = 2,6-diisopropylphenyl), as a twofold phosphonium ion source for the stepwise activation of two  $\text{P}_4$  tetrahedra which yielded the mono- and dicationic species  $[\text{ClP}(\text{NDipp})_2\text{P}_5]^+$  (**1**<sup>+</sup>) and  $[\text{P}_5(\text{NDipp})_2\text{P}_5]^{2+}$  (**2**<sup>2+</sup>, Fig. 1).<sup>9</sup> The related  $\text{P}_5^+$ -species **3** $[\text{GaCl}_4]$  was obtained by the reaction of  $\text{P}_4$  with the four-membered heterocycle  $\text{Cl}_2\text{Si}(\text{NSiMe}_3)_2\text{P}(\text{Cl})$  and




Fig. 1  $\text{P}_5^+$ -cages **1**<sup>+</sup>, **2**<sup>2+</sup>, **3**<sup>+</sup> and **4** featuring a fused four-membered heterocycle.

$\text{GaCl}_3$  while the isolobale Al-species **4** is the result of the reaction of phosphonium zwitterion  $\text{Cl}_2\text{Al}(\text{NSiMe}_3)_2\text{P}$  with  $\text{P}_4$ .<sup>10</sup>

Only recently, cationic polyphosphorus cages have emerged as valuable synthetic building blocks for three purposes. First, they can be selectively fragmented in reactions with carbenes which results in the formation of  $\text{P}_n$ -species (*e.g.* a  $\text{P}_5^+$ -cage cation yields a  $\text{P}_2^-$ - and cationic  $\text{P}_3^+$ -species).<sup>11</sup> Second, they can be oxidized with selenium or sulphur which allows for the targeted preparation of cationic phosphorus–chalcogen cages.<sup>7c</sup> Third, they can be used for the controlled release of  $\text{P}_4$  due to the reversibility of the phosphonium ion insertion.<sup>10b</sup>

With the intention to further expand the range of methods for the *in situ* generation of phosphonium ions for  $\text{P}_4$  activation, we investigated the reactivity of bicyclic phosphorus–nitrogen–silicon heterocycle **7** with  $\text{GaCl}_3$  and  $\text{P}_4$ . A  $\text{GaCl}_3$ -induced rearrangement reaction was observed which formally gives access to the spirocyclic, Si-centered compound  $\text{ClP}(\text{NSiMe}_3)_2\text{Si}(\text{NSiMe}_3)_2\text{P}(\text{Cl})$ . Chloride abstraction by  $\text{GaCl}_3$  and insertion of the respective phosphonium ion in P–P bonds of one or two  $\text{P}_4$  molecules yields the Si-centered polyphosphorus cages  $[\text{ClP}(\text{NSiMe}_3)_2\text{Si}(\text{NSiMe}_3)_2\text{P}_5]^+$  (**19**<sup>+</sup>) and  $[\text{P}_5(\text{NSiMe}_3)_2\text{Si}(\text{NSiMe}_3)_2\text{P}_5]^{2+}$  (**13**<sup>2+</sup>).

TU Dresden, Fachrichtung Chemie und Lebensmittelchemie, Professur für Anorganische Molekülchemie, 01062 Dresden, Germany.

E-mail: jan.weigand@tu-dresden.de

† Electronic supplementary information (ESI) available: Crystallographic data, <sup>31</sup>P NMR spectra of reaction mixtures and general experimental information. CCDC 1060461–1060464. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5dt01512j




## Results and discussion

### Synthetic routes towards and characterization of 7

As part of our ongoing interest in four-membered phosphorus–nitrogen–element heterocycles as precursors for phosphonium ions<sup>12</sup> that can insert into a P–P bond of  $P_4^{10a}$  we revisited the synthesis of heterocycle 6. This compound is synthesized by the reaction of iminophosphane 5 with  $SiCl_4$  according to a procedure reported by Niecke and co-workers (Scheme 1).<sup>13</sup> Compound 6 was obtained from the reaction mixture by distillation (40 °C,  $8 \times 10^{-2}$  mbar) and isolated in 41% yield in accordance with the literature report. Surprisingly, however, a second fraction was obtained at higher temperatures (105 °C,  $2 \times 10^{-3}$  mbar) and identified as bicyclic compound 7 (18% yield).<sup>14</sup> The  $^{31}P\{^1H\}$  NMR spectrum of 7 dissolved in  $C_6D_6$  shows a singlet resonance at  $\delta(P) = 211.8$  ppm which is in the typical chemical shift region of diphosphadiazanes featuring amino-groups on P.<sup>15</sup>

The molecular structure of 7 is  $C_{2v}$ -symmetric and features an distorted planar four-membered  $[NP]_2$ -ring (largest deviation from the plane 0.195 Å, Fig. 2). All four N atoms exhibit a trigonal planar arrangement (angular sums range from 358.8 (3)° to 360.0(3)°) whereas the P atoms are involved in pyramidal bonding environments (angular sums P1: 286(3)° and P2: 286.2(2)°). The P1–N1 and P2–N2 bonds are almost orthogonal to the  $[NP]_2$ -plane (N1–P1...P2: 95.84(4)°, N2–P2...P1: 95.84(4)°). The  $C_{2v}$ -symmetric arrangement of the  $N_2SiCl_2$ -moiety in 7 is rare<sup>16</sup> and contrasts other known diamido-cyclodiphosphazane compounds coordinating main group fragments.<sup>15</sup> Compounds of type  $(PNR)_2NR_2EX_n$  typically exhibit  $C_s$ -symmetric, *seco*-heterocube type structures in which the main group element fragment (e.g.  $EX_n = Mg, BPh, AlCl, GaCl, Ge, Sn, SnCl_2, SiCl_2, AsCl, SbCl, BiCl, R = t-Bu, Ph$ ) is coordinated by three N atoms and occupies the edge of a distorted cube.<sup>15</sup>


The molecular arrangement observed for 7 in the solid state also persists in solution. The observation of two sets of resonances for its chemically inequivalent  $Me_3Si$ -groups in the  $^1H$  NMR spectrum ( $\delta(H) = 0.21$  and 0.40 ppm) confirm the  $C_{2v}$ -symmetry. The  $^{13}C\{^1H\}$  NMR spectrum shows a triplet resonance which is assigned to the carbon atoms of the  $Me_3Si$ -groups attached to the  $[PN]_2$ -ring ( $\delta(C) = -0.1$  ppm,  $^3J(CP) = 3.6$  Hz). A *pseudo*-triplet is observed for the  $Me_3Si$ -groups adjacent to the  $N_2SiCl_2$ -moiety ( $\delta(C) = 2.3$  ppm,  $|^3J(CP) + ^5J(CP)| = 5.0$  Hz) and is a result of an  $AA'X_3X'_3$  spin system ( $A = ^{31}P$ ,



**Fig. 2** Molecular structure of 7 (hydrogen atoms are omitted for clarity and thermal ellipsoids are displayed at 50% probability); selected bond lengths [Å] and angles [°]: N1–P1 1.734(1), N3–P1 1.721(1), N4–P1 1.719(4), N2–P2 1.738(1), N3–P2 1.724(1), N4–P2 1.720(1), P1...P2 2.5235(6), N1–Si1 1.708(1), N1–Si2 1.773(1), N2–Si1 1.710(1), N2–Si3 1.774(1), N3–Si4 1.728(1), N4–Si5 1.740(1), Si1–Cl2 2.0554(6), Si2–Cl1 2.5235(6); P1–N4–P2 94.39(6), P1–N3–P2 94.19(7), N3–P1–N4 84.06(7), N3–P2–N4 83.96(7), N1–P1–N3 100.48(6), N1–P1–N4 102.10(6), N2–P2–N3 100.29(7), N2–P2–N4 101.91(6), N1–P1...P2 95.84(4), N2–P2...P1 95.58(4), Si1–N1–P1 117.14(7), Si1–N2–P2 117.18(8), N1–Si1–N2 114.19(6), Cl1–Si1–Cl2 104.35(2).


$X = ^{13}C$ ) with a comparatively large  $^2J(P_A P_A')$  coupling constant.<sup>17</sup> The same arguments account for the *pseudo*-triplet resonance in the  $^{29}Si\{^1H\}$  NMR spectrum of 7 ( $\delta(Si) = 7.7$  ppm) which is assigned to the  $Me_3Si$ -groups adjacent to the  $N_2SiCl_2$ -moiety ( $|^2J(SiP) + ^4J(SiP)| = 12.5$  Hz). The triplet resonance corresponding to the  $Me_3Si$ -groups attached to the  $[PN]_2$ -ring ( $\delta(Si) = 1.8$  ppm,  $^3J(SiP) = 11.7$  Hz) and the resonance of the  $SiCl_2$  moiety ( $\delta(Si) = -47.4$  ppm) are observed in the expected regions.<sup>18</sup>

Due to the low isolated yield of 7, it was of interest to develop an alternative synthetic approach. Compound 8 is the head to tail dimer of iminophosphane 5 and was obtained according to a literature known procedure.<sup>17</sup> The solvent free reaction of 8 with an excess of  $SiCl_4$  gave selectively and quantitatively compound 9 via  $Me_3SiCl$  elimination (Scheme 2).<sup>19</sup> Compound 9 is the *trans*-conformer of an unsymmetrically-substituted diphosphadiazane derivative and, thus, shows an AX spin system in its  $^{31}P\{^1H\}$  NMR spectrum ( $\delta(P_A) = 219.5$  ppm,  $\delta(P_X) = 232.3$  ppm,  $^2J(P_A P_X) = 12.0$  Hz). The  $^1H$  NMR spectrum of 9 shows singlet resonances for the  $SiMe_3$ -groups on the  $[NP]_2$ -ring and for both pointing towards the plane of the four-membered ring. The  $SiMe_3$  group pointing away from the ring, however, shows a relatively large  $^4J(HP)$  coupling constant ( $\delta(H) = 0.27$  ppm,  $^4J(P_A H) = 3.7$  Hz). For structurally related compounds, this was rationalized by the



**Scheme 1** Preparation of compounds 6 and 7; (a)  $+SiCl_4, -Me_3SiCl$ , neat, 80 °C, 3 d.






**Scheme 2** Synthesis of **9** via reaction of **8** and  $\text{SiCl}_4$ : (a)  $+\text{SiCl}_4$ ,  $-\text{Me}_3\text{SiCl}$ , rt, 12 h, 96%; and subsequent transformation to **7**: (b) neat, 185 °C, 10 min, 25%.

close proximity of the  $\text{CH}_3$ -groups to the lone pair of electrons on the adjacent P atom.<sup>20</sup> Due to the same reason, two relatively large  $^4J(\text{PH})$  coupling constants (3.4 and 3.7 Hz) were observed for the isomer of **9** in which the  $\text{SiCl}_3$ -group points towards the face of the four-membered ring (**9'**).<sup>19</sup> The chloro-substituted Si atom in **9** appears as doublet resonance in the typical range in the  $^{29}\text{Si}$  NMR spectrum ( $\delta(\text{Si}) = -27.3$  ppm)<sup>21</sup> with a remarkably large coupling constant of  $^2J(\text{SiP}_x) = 26.0$  Hz. A coupling constant of similar magnitude is observed for the  $\text{Me}_3\text{Si}$ -group pointing away from the four-membered ring ( $\delta(\text{Si}) = 7.8$  ppm,  $^2J(\text{SiP}_A) = 32.4$  Hz), whereas those pointing to the faces of the  $[\text{NP}]_2$ -ring exhibit rather small coupling constants ( $\delta(\text{Si}) = 4.9$  and  $0.2$  ppm;  $^2J(\text{SiP}) = 4.2$  and  $3.0$  Hz). Similarly the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum reveals rather small coupling constants for the latter  $\text{Me}_3\text{Si}$ -groups ( $^3J(\text{CP}) = 8.0$  and  $10.5$  Hz) and a large coupling constant for the  $\text{Me}_3\text{Si}$ -group pointing away from the  $[\text{NP}]_2$ -ring plane ( $^3J(\text{CP}_A) = 19.8$  Hz).

It is assumed, that **7** is obtained from **9** via an intermediate of type **10** which forms upon rotation<sup>22</sup> of the P–N bond involving the  $\text{SiCl}_3$ -substituted N atom and inversion<sup>23</sup> of one of the P centers in **9** (Scheme 2). The arrangement of the  $\text{SiCl}_3$ -group and one of the  $\text{SiMe}_3$ -groups in intermediate **10** allows for  $\text{Me}_3\text{SiCl}$  elimination yielding **7**. Indeed, heating **9** for ten minutes to 185 °C results in conversion to **7** in 25% yield.<sup>24</sup>

### Reactivity of **7** towards $\text{GaCl}_3$ and $\text{P}_4$


Diphosphadiazanes and related compounds bearing chloro- and  $\text{Me}_3\text{Si}$ -substituents are known to undergo a variety of distinct reactions with Lewis acids. Next to halide abstraction,<sup>10</sup> also elimination of  $\text{Me}_3\text{SiCl}$ ,<sup>25</sup> migration of  $\text{Me}_3\text{Si}$ -,<sup>26</sup> chloro- or methyl-substituents<sup>27</sup> and P–N bond cleavage reactions are reported.<sup>28</sup> Especially the latter reaction is of interest, since it is assumed to proceed via phosphonium ion intermediates. Thus, bicyclic compound **7** is a promising substrate for the generation of phosphonium ions and, subsequently, insertion of the latter into P–P bonds of  $\text{P}_4$ .



**Scheme 3** Formation of Lewis-acid/base adduct **11** via the reaction of **7** and  $\text{GaCl}_3$ : (a)  $+\text{GaCl}_3$ ,  $\text{C}_6\text{H}_5\text{F}$ , r.t., 1 h.

Thus, the reaction of **7** with the Lewis acid  $\text{GaCl}_3$  was probed (Scheme 3). The addition of one equivalent of  $\text{GaCl}_3$  to a solution of **7** in  $\text{C}_6\text{H}_5\text{F}$  gave a reddish colored reaction mixture which was investigated by means of  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy. The formation of the Lewis-acid/base adduct **11** is proposed on the basis of the observation of two broad resonances in an approximate ratio of 1 : 1 ( $\delta(\text{P}_A) = 115$  ppm,  $\Delta\nu_{1/2} = 1200$  Hz;  $\delta(\text{P}_X) = 179.6$  ppm,  $\Delta\nu_{1/2} = 600$  Hz). The coordination of  $\text{GaCl}_3$  to a P atom is very likely, since they have been identified as the most basic sites in related compounds.<sup>29</sup> The resonance at low field is tentatively assigned to the tri-coordinated P atom. Accordingly, the resonance at high field corresponds to the tetra-coordinated P atom and exhibits a stronger line broadening due to the coordination of  $\text{GaCl}_3$ .

Interestingly,  $\text{P}_4$  does not react with **11** which indicates that the latter is not a suitable phosphonium ion source. Adding two equivalents of  $\text{GaCl}_3$  to a solution of **7** in  $\text{C}_6\text{H}_5\text{F}$ , however, results in the rapid consumption of *in situ* generated **11**. This reaction yields a complex mixture of not identified products and bodes well for the generation of reactive intermediates that might be able to activate  $\text{P}_4$ . Indeed, mixtures of **7**,  $\text{P}_4$  and  $\text{GaCl}_3$  in 1 : 1 : 2 and 1 : 2 : 4 stoichiometries were bright red and the consumption of  $\text{P}_4$  accompanied by a color change to brown was observed. Subsequent investigation by means of  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy revealed the formation of  $\text{P}_5^+$  cage cation **12**<sup>+</sup> and bridged *bis*( $\text{P}_5^+$ )-cage dication **13**<sup>2+</sup> by their characteristic  $\text{A}_2\text{MOX}_2$  and  $\text{A}_2\text{MX}_2$  spin systems (Scheme 4).




**Scheme 4** Reaction of **7** with  $\text{P}_4$  and  $\text{GaCl}_3$  in various stoichiometries; (a)  $+\text{P}_4$ ,  $+2$ ,  $\text{GaCl}_3$ ,  $\text{C}_6\text{H}_5\text{F}$ , r.t., 24 h; (b)  $+2$ ,  $\text{P}_4$ ,  $+4$ ,  $\text{GaCl}_3$ , r.t.,  $\text{C}_6\text{H}_5\text{F}$ , 24 h; the equation depicts the major products and is not balanced; anions of the products are not depicted.



In addition, both spectra reveal significant amounts of  $P_4$  but neither remaining **7** nor the corresponding adduct **11** were observed which is attributed to unidentified side reactions. The products  $12^+$  and  $13^{2+}$  are formally obtained by the insertion of phosphonium ions based on  $ClP(NSiMe_3)_2Si(NSiMe_3)_2PCl$  which features a Si-centered spiro[3.3]heptane-motif, into P–P bonds of  $P_4$ . Interestingly, both reactions yield  $12^+$  and  $13^{2+}$  in comparable ratios (1 : 1.8 for (a) and 1 : 1.5 for (b), Scheme 4). This contrasts the anticipated increase of the amount of dication  $13^{2+}$  in the presence of excess  $P_4$  and  $GaCl_3$ . In addition, the formation of large quantities of  $13^{2+}$  in the reaction of 1 : 1 : 2 stoichiometry indicates that the reaction rate of a  $GaCl_3$ -induced rearrangement of **7** yielding the spiro-motif is slow compared to that of the subsequent phosphonium ion insertion.


A tentative mechanism for the formation of such a Si-centered spiro[3.3]heptane-type species from **7** is illustrated in Scheme 5. While the reaction of **7** with one equivalent  $GaCl_3$  gives the Lewis-acid/base adduct **11** (*vide supra*), an excess of  $GaCl_3$  is assumed to initiate a chloride abstraction from the  $SiCl_2$ -moiety. This is facilitated by the tendency of bicyclic compounds of type **7** to form a *seco*-heterocube-type structure.<sup>15</sup> Thus, compound **7** might be in equilibrium with the related derivative **7<sup>1</sup>**. Compound **7<sup>1</sup>** features a hypervalent penta-coordinated Si-moiety which might favor the sequestering of a chloride anion. Thus, it is assumed that the reaction with  $GaCl_3$  proceeds *via* chloride abstraction and formation of the cationic *seco*-heterocube **14<sup>+</sup>** featuring a tetra-coordinated



**Scheme 5** Suggested  $GaCl_3$ -induced rearrangement mechanism of **7** to phosphonium ion **18<sup>+</sup>** featuring a Si-centered spiro[3.3]heptane-type motif; R denotes  $Me_3Si$ -substituents; for reasons of simplification the reactions of intermediates or products possessing a di-coordinated P atom with  $P_4$  as well as retransfer of a chloride ion to a di-coordinated P atom are not considered.


N atom. The next two steps constitute P–N bond cleavage and formation reactions yielding intermediates **15<sup>+</sup>** and **16<sup>+</sup>** *via* formal retention of the *seco*-heterocube-type structure. Similar P–N bond ruptures were reported as decomposition pathways of diamino-*cyclo*-diphosphadiazanes.<sup>26</sup> Intermediate **16<sup>+</sup>** features a diphosphadiazane  $[NP]_2$ -ring which is assumed to react *via* a *cyclo*-reversion reaction to intermediate **17<sup>+</sup>**. Cation **17<sup>+</sup>** features an aminoiminophosphane moiety similar to **5** which is tethered to a four-membered  $SiN_2P$ -ring. Intramolecular nucleophilic attack of the imino-N atom on the chloro-substituted Si atom initiates a  $GaCl_3$ -mediated transfer of a chloride anion from the Si atom to a di-coordinated P atom to give formally the phosphonium ion **18<sup>+</sup>**. Cation **18<sup>+</sup>** features the Si-centered spiro[3.3]heptane-motif and, thus, is assumed to be accountable for the formation of cages cations **12<sup>+</sup>** and **13<sup>2+</sup>** *via* insertion into a P–P bond of  $P_4$ , chloride abstraction by  $GaCl_3$  and subsequent insertion into a P–P bond of a second  $P_4$  tetrahedron.

Attempts to isolate a gallate salt of **12<sup>+</sup>** from both reaction mixtures were unsuccessful, possibly due to the fluxional coordination behavior of the  $GaCl_3$ -molecule to the PCl-function. However, slow diffusion of *n*-hexane into the reaction mixture of 1 : 1 : 2 stoichiometry yielded compound **19** $[GaCl_4]$  in low yields (10%, Scheme 6). The  $^{31}P\{^1H\}$  NMR spectrum of **19** $[GaCl_4]$  dissolved in  $CD_2Cl_2$  shows an  $ABMX_2Y$  spin system which is in accordance with the  $C_s$  symmetry of the molecule (Fig. 3). The mirror plane is defined by the tetra-coordinated P atom and the two adjacent P atoms. The P–Cl unit is included in the plane and exhibits a spatial proximity to one of the two bridge-head P atoms. The chemical shifts and coupling constants involving the  $P_5^+$ -cage motif are similar to those observed for the cage cation **3<sup>+</sup>**.<sup>10a</sup> The chloro-substituted P atom in **19<sup>+</sup>** exhibits a singlet resonance ( $\delta(P_Y) = 166.8$  ppm) in the typical range of silyl-substituted diamino-chlorophosphanes.<sup>13,30</sup> A  $^4J(PP)$  coupling between this P atom and the tetra-coordinated P atom of the  $P_5^+$ -cage is not resolved. The  $^1H$  NMR spectrum of **19<sup>+</sup>** reveals three singlet resonances assigned to the chemically different  $Me_3Si$ -groups which integrate in a 2 : 1 : 1 ratio. The high-field resonance ( $\delta(H) = 0.31$  ppm) exhibits the highest intensity and is assigned to the two  $Me_3Si$ -groups bonded to the four-membered ring which incorporates the P–Cl moiety. The chemical shift is comparable to related four-membered ring compounds (**6**:  $\delta(H) = 0.18$  ppm,<sup>13</sup> **7**:  $\delta(H) = 0.21$  ppm). Both resonances assigned to the  $Me_3Si$ -groups bonded to the N atoms adjacent to the



**Scheme 6** Reaction of **19** $[GaCl_4]$  with  $GaCl_3$ ; (a)  $+GaCl_3$ ,  $CD_2Cl_2$ , r.t., 30 min.






**Fig. 3**  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of  $19[\text{GaCl}_4]$  ( $\text{CD}_2\text{Cl}_2$ , r.t.); insets show experimental (upwards) and fitted spectra (downwards);  $\text{A}_2\text{MO}_2\text{X}_2$  spin system of  $19^+$ :  $\delta(\text{P}_\text{A}) = -322.9$  ppm,  $\delta(\text{P}_\text{B}) = -320.2$  ppm,  $\delta(\text{P}_\text{M}) = 30.4$  ppm,  $\delta(\text{P}_\text{X}) = 165.5$  ppm,  $\delta(\text{P}_\text{Y}) = 166.8$  ppm,  $^1J(\text{P}_\text{A}\text{P}_\text{B}) = -189.8$  Hz,  $^1J(\text{P}_\text{A}\text{P}_\text{X}) = -143.4$  Hz,  $^1J(\text{P}_\text{B}\text{P}_\text{X}) = -147.7$  Hz,  $^1J(\text{P}_\text{M}\text{P}_\text{X}) = -245.1$  Hz,  $^2J(\text{P}_\text{A}\text{P}_\text{M}) = 18.8$  Hz,  $^2J(\text{P}_\text{B}\text{P}_\text{M}) = 18.0$  Hz.


$\text{P}_5^+$ -moiety exhibit a low field shift ( $\delta(\text{H}) = 0.67$  ppm and  $0.76$  ppm) and are comparable to the corresponding resonance of  $3^+$  ( $\delta(\text{H}) = 0.68$  ppm).<sup>10a</sup> The  $^{29}\text{Si}\{^1\text{H}\}$  NMR spectrum exhibits a resonance at  $\delta(\text{Si}) = -58.0$  ppm which is assigned to the Si spiro-atom. This resonance reveals a doublet of doublet splitting caused by  $^2J(\text{SiP})$ -couplings to the tetra-coordinated P atom ( $^2J(\text{SiP}_\text{M}) = 8.5$  Hz) and the chloro-substituted P atom ( $^2J(\text{SiP}_\text{Y}) = 18.5$  Hz).

The molecular structure of  $19^+$  is depicted in Fig. 4 and the P–P bond lengths and angles in the  $\text{P}_5^+$ -moiety are comparable to those of  $3^+$ .<sup>10a</sup> Both four-membered rings are almost planar (largest deviation from the planes N1:  $0.026$  Å and N3:  $0.022$  Å) and exhibit a perpendicular arrangement (angle between both planes:  $89.79(9)^\circ$ ). Due to the steric limitations of the four-membered heterocycles the spiro-Si atom exhibits a distorted tetrahedral arrangement with two rather small (N1–Si3–N2:  $85.4(1)^\circ$ , N3–Si3–N4:  $87.7(1)^\circ$ ) and two widened (N1–Si3–N3:  $122.9(1)^\circ$ , N2–Si3–N4:  $118.8(1)^\circ$ ) N–Si–N angles. Alternating bond lengths are observed within the two four-membered rings. The P–N bonds involving the tetra-coordinated P atom are shorter (N2–P1:  $1.656(3)$  Å, N1–P1:  $1.661(3)$  Å) than those involving the tri-coordinated P atom (N3–P6:  $1.716(3)$  Å, N4–P6:  $1.711(3)$  Å) and both magnitudes of bond lengths are also observed in the related cages  $3^+$  and  $2^+$ .<sup>9,10a</sup> The Si–N bonds in the  $[\text{SiN}_2\text{P}]$ -ring fused to the  $\text{P}_5^+$ -cage (N1–Si3:  $1.755(3)$  Å, N2–Si3:  $1.751(3)$  Å) are of similar lengths as observed for  $3^+$ .<sup>10a</sup> In contrast, the Si–N bonds in the second  $[\text{SiN}_2\text{P}]$ -ring are shorter (N3–Si3:  $1.711(3)$  Å, N4–Si3:  $1.704(3)$  Å).

The addition of  $\text{GaCl}_3$  to a solution of  $19[\text{GaCl}_4]$  in  $\text{CD}_2\text{Cl}_2$  yields the previously mentioned Lewis-acid base adduct  $12^+$  (Scheme 6). The  $\text{A}_2\text{MO}_2\text{X}_2$  spin system observed in the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of  $12^+$  suggests a time averaged  $\text{C}_{2v}$ -symmetry of the molecule in solution which can be explained by a fluxional behavior of the coordinated  $\text{GaCl}_3$  molecule (Fig. 5). The  $\text{A}_2\text{MX}_2$  part of the  $\text{A}_2\text{MO}_2\text{X}_2$  spin system corresponds to the  $\text{P}_5^+$ -cage moiety and reveals comparable chemical shifts and coupling constants as observed for  $3^+$ .<sup>10a</sup> The resonance corresponding to the O part of the spin system is very broad ( $\delta(\text{P}_\text{O}) =$



**Fig. 4** Molecular structure of  $19^+$  in compound  $19[\text{GaCl}_4]\cdot\text{C}_6\text{H}_5\text{F}$  (hydrogen atoms are omitted for clarity and thermal ellipsoids are displayed at 50% probability); selected bond lengths [Å] and angles [ $^\circ$ ]: N2–P1  $1.656(3)$ , N1–P1  $1.661(3)$ , N1–Si3  $1.755(3)$ , N2–Si3  $1.751(3)$ , N3–Si3  $1.711(3)$ , N4–Si3  $1.704(3)$ , N3–P6  $1.716(3)$ , N4–P6  $1.711(3)$ , Cl1–P6  $2.238(2)$ , P1...Si3  $2.445(1)$ , P6...Si3  $2.472(1)$ , P1–P2  $2.167(1)$ , P1–P3  $2.165(1)$ , P2–P4  $2.249(2)$ , P2–P5  $2.238(2)$ , P3–P4  $2.251(2)$ , P3–P5  $2.229(2)$ , P4–P5  $2.169(2)$ ; N1–P1–N2  $91.5(1)$ , P1–N1–Si3  $91.4(1)$ , P1–N2–Si3  $91.7(1)$ , N1–Si3–N2  $85.4(1)$ , N3–Si3–N4  $87.7(1)$ , N1–Si3–N3  $122.9(1)$ , N2–Si3–N4  $118.8(1)$ , N3–P6–N4  $87.2(1)$ , N3–P6–Cl1  $102.2(1)$ , N4–P6–Cl1  $101.7(1)$ , P3–P1–P2  $91.15(5)$ , P1–P2–P5  $83.95(5)$ , P1–P2–P4  $82.53(5)$ , P5–P2–P4  $57.82(5)$ , P5–P4–P2  $60.83(5)$ , P2–P5–P3  $87.67(5)$ .



**Fig. 5**  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of  $12[\text{GaCl}_4]$  ( $\text{CD}_2\text{Cl}_2$ , r.t.);  $\text{A}_2\text{MO}_2\text{X}_2$  spin system of  $12^+$ :  $\delta(\text{P}_\text{A}) = -318.3$  ppm,  $\delta(\text{P}_\text{M}) = 33.4$  ppm,  $\delta(\text{P}_\text{O}) = 144$  ppm ( $\Delta\nu_{1/2} = \sim 1200$  Hz),  $\delta(\text{P}_\text{X}) = 170.8$  ppm,  $^1J(\text{P}_\text{A}\text{P}_\text{X}) = -144.8$  Hz,  $^1J(\text{P}_\text{M}\text{P}_\text{X}) = -248.6$  Hz,  $^2J(\text{P}_\text{A}\text{P}_\text{M}) = 19.0$  Hz.

$144$  ppm,  $\Delta\nu_{1/2} = \sim 1200$  Hz) which is caused by the fluxional behavior of the coordinated  $\text{GaCl}_3$  molecule and its quadrupole moment.

Dication  $13^{2+}$  was isolated as a  $[\text{Ga}_2\text{Cl}_7]^-$  salt from the reaction of  $7$ ,  $\text{P}_4$  and  $\text{GaCl}_3$  in a  $1 : 2 : 4$  stoichiometry (Scheme 4) by the addition of  $n$ -hexane. This gave a brown oil which was isolated by decanting the supernatant and upon addition of small amounts of  $1,2\text{-C}_6\text{H}_4\text{F}_2$  yielded a suspension containing a yellow microcrystalline material of  $13[\text{Ga}_2\text{Cl}_7]_2$ . This compound was isolated by filtration in low yields (20%) in an approximate purity of 75%. Further purification by recrystallization from  $\text{CH}_2\text{Cl}_2/n$ -hexane leads to significantly decreased yields (7% yield in >90% purity). The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of  $13[\text{Ga}_2\text{Cl}_7]_2$  dissolved in  $\text{CD}_2\text{Cl}_2$  shows an  $\text{A}_2\text{MX}_2$ -spin system in accordance with the  $\text{C}_{2v}$ -symmetry of the molecule






Fig. 6  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of  $13[\text{Ga}_2\text{Cl}_7]_2$  ( $\text{CD}_2\text{Cl}_2$ , r.t.); unidentified side products are marked with asterisks;  $\text{A}_2\text{MX}_2$  spin system of  $13^{2+}$ :  $\delta(\text{P}_\text{A}) = -320.4$  ppm,  $\delta(\text{P}_\text{M}) = 32.2$  ppm,  $\delta(\text{P}_\text{X}) = 158.8$  ppm,  $^1J(\text{P}_\text{A}\text{P}_\text{X}) = -142.7$  Hz,  $^1J(\text{P}_\text{M}\text{P}_\text{X}) = -259.1$  Hz,  $^2J(\text{P}_\text{A}\text{P}_\text{M}) = 20.0$  Hz.

(Fig. 6). The  $C_2$ -axis includes both tetra-coordinated P atoms and the spiro-Si atom and the mirror planes are defined by the four-membered  $[\text{SiN}_2\text{P}]$ -rings. The resonances and coupling constants of  $13^{2+}$  are similar to those observed for the related compounds  $19^+$  and  $3^{+10a}$

Single crystals of compound  $13[\text{Ga}_2\text{Cl}_7]_2$  were obtained by diffusion of *n*-hexane in a  $\text{CH}_2\text{Cl}_2$  solution of  $13[\text{Ga}_2\text{Cl}_7]_2$  at  $-35$  °C. The compound crystallizes with two independent formula units in the asymmetric unit. Two of the four  $\text{Ga}_2\text{Cl}_7^-$  anions are highly disordered exhibiting unusually high thermal displacement parameters (see ESI† for details). Single crystals of  $13[\text{Ga}_2\text{Cl}_7][\text{GaCl}_4]$  were obtained by layering the supernatant solution of the reaction mixture of the synthesis of  $13[\text{Ga}_2\text{Cl}_7]_2$  with *n*-hexane at  $-35$  °C. The data obtained by X-ray single crystal structure determination was of higher quality and, thus, the molecular structure of  $13[\text{Ga}_2\text{Cl}_7][\text{GaCl}_4]$  is discussed (Fig. 7). The P–P bond lengths and angles in the  $\text{P}_5^+$ -moieties are comparable to those of related  $\text{P}_5^+$ -cage compounds.<sup>9,10a</sup> The P–N bonds in  $13^{2+}$  are rather short (*av.* P–N: 1.666(8) Å) which is a typical feature of P–N bonds involving phosphonium moieties. The Si–N bond lengths in the four-membered rings are nearly identical (*av.* Si–N: 1.732(8) Å) and are between the two types of bond lengths observed for  $19^+$  (*av.* Si–N: 1.753(6) Å and Si–N: 1.7007(6) Å).

## Conclusions

The bicyclic P–N–Si heterocycle **7** was targeted as source for the *in situ* generation of phosphonium cations for  $\text{P}_4$  activation. In this context, two distinct synthetic protocols for its preparation were thoroughly investigated and gave insights into the formation of the bicycle. The reaction of **7** with  $\text{GaCl}_3$  initially yields adduct **11**. This adduct is not stable and subsequently rearranges to give *in situ* spirocyclic, Si-centered compound  $\text{ClP}(\text{NSiMe}_3)_2\text{Si}(\text{NSiMe}_3)_2\text{P}\text{Cl}$ . The latter species gives access to polyphosphorus cage cations  $[\text{ClP}(\text{NSiMe}_3)_2\text{Si}(\text{NSiMe}_3)_2\text{P}_5]^+$  ( $19^+$ ) and  $[\text{P}_5(\text{NSiMe}_3)_2\text{Si}(\text{NSiMe}_3)_2\text{P}_5]^{2+}$  ( $13^{2+}$ ) in the presence of  $\text{GaCl}_3$  and  $\text{P}_4$ . We are continuing to investigate the Lewis-acid mediated generation of phosphonium ions




Fig. 7 Molecular structure of  $13^{2+}$  in compound  $13[\text{Ga}_2\text{Cl}_7][\text{GaCl}_4]$  (hydrogen atoms are omitted for clarity and thermal ellipsoids are displayed at 50% probability); selected bond lengths [Å] and angles [°]: N1–P1 1.667(2), N2–P1 1.671(2), N3–P6 1.663(2), N4–P6 1.665(2), Si1–N1 1.731(2), Si1–N2 1.732(2), S1–N3 1.736(2), Si1–N4 1.728(2), P1…Si1 2.4452(8), P6…Si1 2.4277(8), P1–P2 2.1540(8), P1–P3 2.1499(8), P2–P4 2.245(1), P2–P5 2.245(1), P3–P4 2.246(1), P3–P5 2.2464(9), P4–P5 2.168(1), P6–P7 2.1511(8), P6–P8 2.1572(2), P7–P9 2.240(1), P7–P10 2.252(1), P8–P9 2.2415(9), P8–P10 2.242(1), P9–P10 2.172(1); N1–P1–N2 90.81(9), N3–P6–N4 90.99(9), N1–Si1–N2 86.73(9), N3–Si1–N4 86.49(9), P1–N1–Si1 91.26(9), P1–N2–Si1 91.12(9), P6–N3–Si1 91.14(9), P6–N4–Si1 91.34(9), P3–P1–P2 92.59(3), P1–P2–P5 81.73(3), P1–P2–P4 83.19(3), P5–P2–P4 57.69(3), P5–P4–P2 61.25(3), P2–P5–P3 87.60(3), P7–P6–P8 92.30(3), P6–P7–P10 82.39(3), P6–P7–P9 82.89(3), P10–P7–P9 57.83(3), P10–P9–P7 61.36(3), P7–P10–P8 87.48(3).

for  $\text{P}_4$  activation from related phosphorus–nitrogen-element bicycles. Furthermore, studies directed towards the utilization of  $12^+$ ,  $19^+$  and  $13^{2+}$  as synthetic building blocks will be the target of future efforts.

## Experimental

### General


General information on materials and methods as well as  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra of reaction mixtures are given in the ESI.†

### Synthesis of $\text{Cl}_2\text{Si}(\text{NSiMe}_3)_2(\text{PNSiMe}_3)_2$ (**7**)

**Method A:** The literature reported synthesis of **5** was performed on a 20 mmol scale.<sup>13</sup> Compound **5** was removed by distillation from the reaction mixture (40 °C,  $8 \times 10^{-2}$  mbar). The remaining colorless, slushy residue was dissolved in  $\text{C}_6\text{H}_5\text{F}$  (5 mL) yielding a turbid suspension. The solvent was removed *in vacuo* yielding a sludgy residue which was redistilled employing a short Vigreux column (5 cm). Compound **7** was obtained as colorless oil (1.789 g, 3.51 mmol, 18%, 105 °C,  $2 \times 10^{-3}$  mbar) which solidified shortly after distillation. **Method B:** **9** (305 mg, 0.50 mmol, 1.0 eq.) was heated to 185 °C for 10 min. In the course of the reaction a colorless liquid is formed accompanied by the condensation of  $\text{Me}_3\text{SiCl}$  on




colder parts of the reaction vessel. After cooling to ambient temperature the reaction mixture remains a liquid. Isolation of 7 from this mixture proceeds as described in method A.



**m.p.:** 55.6–57.8 °C; **Raman (300 mW, [cm<sup>-1</sup>]):**  $\nu$  = 2959 (390), 2899 (100), 1410 (11), 690 (10), 645 (22), 613 (51), 562 (4), 349 (34), 185 (30), 141 (6), 75 (10); **IR (ATR, [cm<sup>-1</sup>]):**  $\nu$  = 2956 (w), 1408 (vw), 1249 (s), 1098 (vw), 973 (vw), 942 (w), 883 (w), 830 (vs), 776 (vw), 754 (vw), 713 (w), 682 (w), 643 (vw), 556 (s), 450 (m); **<sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, [ppm]):**  $\delta$  = 0.21 (18H, s, H1), 0.39 (18H, s, H2); **<sup>13</sup>C{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, [ppm]):**  $\delta$  = -0.08 (6C, t, C1, <sup>3</sup>J(CP) = 3.6 Hz), 2.7 (6C, pseudo-t, C2, <sup>3</sup>J(CP) = 5.0 Hz); **<sup>29</sup>Si{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, [ppm]):**  $\delta$  = -47.4 (1Si, s, Si3), 1.8 (2Si, t, Si1, <sup>2</sup>J(SiP) = 11.7 Hz), 7.7 (2Si, pseudo-t, Si2, <sup>2</sup>J(SiP) = 11.8 Hz); **<sup>15</sup>N NMR (C<sub>6</sub>D<sub>6</sub>, [ppm]):**  $\delta$  = -397 (t, N1, <sup>1</sup>J(NP) = 55 Hz), -374 (d, N2, <sup>1</sup>J(NP) = 75 Hz); **<sup>31</sup>P{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, [ppm]):**  $\delta$  = 211.8 (s); **elemental analysis for C<sub>12</sub>H<sub>36</sub>Cl<sub>2</sub>N<sub>4</sub>P<sub>2</sub>Si<sub>5</sub>:** calcd: C 28.3, H 7.1, N 11.0; found: C 28.5, H 7.3, N 10.6; **MS-ESI-EM:** 473.0943 [M-Cl<sup>-</sup>], calcd: for C<sub>12</sub>H<sub>36</sub>Cl<sub>1</sub>N<sub>4</sub>P<sub>2</sub>Si<sub>5</sub>: 473.0945.

### Synthesis of (SiMe<sub>3</sub>)<sub>2</sub>N(PNSiMe<sub>3</sub>)<sub>2</sub>N(SiMe<sub>3</sub>)(SiCl<sub>3</sub>) (9)

Compound 8 (1.114 g, 2.00 mmol, 1.0 eq.) was suspended in SiCl<sub>4</sub> (6.796 g, 40.0 mmol, 40.0 eq.) and stirred for 12 h at ambient temperature. After removal of all volatiles *in vacuo* 9 was isolated in quantitative yields as colorless solid (1.187 g, 1.92 mmol, 96%).




**m.p.:** 180.2–182.5 °C; **Raman (300 mW, [cm<sup>-1</sup>]):**  $\nu$  = 2957 (37), 2906 (100), 1410 (15), 686 (13), 651 (4), 641 (64), 586 (21), 488 (4), 436 (38), 351 (21), 206 (37), 107 (27), 78 (18); **IR (ATR, [cm<sup>-1</sup>]):**  $\nu$  = 3139 (vw), 3047 (m), 2958 (vw), 1406 (m), 1251 (s), 1060 (m), 923 (m), 834 (vs), 778 (vw), 755 (w), 712 (w), 677 (w), 572 (vw), 559 (m), 506 (w), 432 (vw); **<sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, [ppm]):**  $\delta$  = 0.17 (18H, s, H5), 0.27 (9H, d, H3, <sup>4</sup>J(HP) = 3.7 Hz), 0.52 (9H, s, H4), 0.64 (9H, s, H1); **<sup>13</sup>C{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, [ppm]):**  $\delta$  = 1.4 (6C, t, C5, <sup>3</sup>J(CP) = 2.5 Hz), 4.7 (3C, d, C1, <sup>3</sup>J(CP) = 10.5 Hz), 4.8 (3C, d, C3, <sup>3</sup>J(CP) = 19.8 Hz), 5.0 (3C, d, C4, <sup>3</sup>J(CP) = 8.0 Hz); **<sup>29</sup>Si{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, [ppm]):**  $\delta$  = -27.3 (1Si, d, Si2, <sup>2</sup>J(SiP) = 26.0 Hz), 0.2 (1Si, d, Si4, <sup>2</sup>J(SiP) = 3.0 Hz), 1.5 (2Si, t, Si5, <sup>2</sup>J(SiP) = 7.4 Hz), 4.5 (1Si, d, Si2, <sup>2</sup>J(SiP) = 4.2 Hz), 7.6 (1Si, d, Si3, <sup>2</sup>J(SiP) = 32.4 Hz); **<sup>15</sup>N NMR (C<sub>6</sub>D<sub>6</sub>, [ppm]):**  $\delta$  = 67.1 (t, N2, <sup>1</sup>J(NP) = 50 Hz), 93.8 (d, N1, <sup>1</sup>J(NP<sub>X</sub>) = 90 Hz), 113.9 (d, N3, <sup>1</sup>J(NP<sub>A</sub>) = 90 Hz); **<sup>31</sup>P{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, [ppm]):** AX spin system:  $\delta$ (P<sub>A</sub>) = 218.6 (d, P1,  $\Delta\nu_{1/2}$  = 42 Hz, <sup>2</sup>J(P<sub>A</sub>P<sub>X</sub>) = 12 Hz),  $\delta$ (P<sub>X</sub>) =

232.2 (d, P2,  $\Delta\nu_{1/2}$  = 37 Hz, <sup>2</sup>J(P<sub>A</sub>P<sub>X</sub>) = 12 Hz); **elemental analysis for C<sub>15</sub>H<sub>45</sub>Cl<sub>3</sub>N<sub>4</sub>P<sub>2</sub>Si<sub>6</sub>:** calcd: C 29.1, H 7.3, N 9.1; found: C 28.8, H 7.3, N 8.6.

### Reaction of 7, P<sub>4</sub> and GaCl<sub>3</sub> in 1 : 1 : 2 and 1 : 2 : 4 stoichiometries


**1 : 1 : 2:** Compound 7 (256 mg, 0.50 mmol, 1.0 eq.) was added to a suspension of P<sub>4</sub> (62 mg, 0.50 mmol, 1.0 eq.) in C<sub>6</sub>H<sub>5</sub>F (5 mL). A solution of GaCl<sub>3</sub> (176 mg, 1.00 mmol, 2.0 eq.) in C<sub>6</sub>H<sub>5</sub>F (2 mL) was added dropwise to the suspension giving a red colored reaction mixture which was stirred for 12 h at ambient temperature. In the course of the reaction the color of the reaction mixture changed to yellow and the dissolving of P<sub>4</sub> was observed. The reaction mixture was investigated by means of <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy (see ESI†). **1 : 2 : 4:** Compound 7 (256 mg, 0.50 mmol, 1.0 eq.) was added to a suspension of P<sub>4</sub> (124 mg, 1.00 mmol, 2.0 eq.) in C<sub>6</sub>H<sub>5</sub>F (5 mL). A solution of GaCl<sub>3</sub> (352 mg, 2.00 mmol, 4 eq.) in C<sub>6</sub>H<sub>5</sub>F (4 mL) was added dropwise to the suspension giving a red colored reaction mixture which was stirred for 12 h at ambient temperature. In the course of the reaction the color of the reaction mixture changed to brown and the dissolving of P<sub>4</sub> was observed. The reaction mixture was investigated by means of <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy (see ESI†). *n*-Hexane (2 mL) was added leading to the formation of a brown oil. The supernatant was removed, diluted with C<sub>6</sub>H<sub>5</sub>F (6 mL) and layered with *n*-hexane (3 mL) at -35 °C. Small amounts of crystalline material of 19[GaCl<sub>4</sub>] (41 mg, 10%), suitable for X-ray single crystal structure determination, were obtained within a few days. The remaining oil was washed with *n*-hexane (3 × 3 mL) transforming it into a brown sludge. All volatiles were removed *in vacuo* and the sludge was suspended in 1,2-C<sub>6</sub>H<sub>4</sub>F<sub>2</sub> (2 mL) leading to the formation of a yellow, microcrystalline solid. The supernatant was removed and the yellow powder was washed with 1,2-C<sub>6</sub>H<sub>4</sub>F<sub>2</sub>/*n*-hexane (1 : 1 mixture, 2 × 2 mL). The obtained yellow powder consisting of 13[Ga<sub>2</sub>Cl<sub>7</sub>] in an approximate purity of 75% (determined by <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy, 20% yield, 145 mg) was isolated by filtration and dried *in vacuo*. Recrystallization from a CH<sub>2</sub>Cl<sub>2</sub> solution by slow diffusion of *n*-hexane yielded crystalline material of 19 [Ga<sub>2</sub>Cl<sub>7</sub>]<sub>2</sub> (purity > 90%) which was suitable for single crystal structure determination. Isolation was conducted *via* filtration and removal of all volatiles *in vacuo* (7% yield, 55 mg). Single crystals of 13[Ga<sub>2</sub>Cl<sub>7</sub>]-[GaCl<sub>4</sub>] were obtained by layering the diluted supernatant of the reaction mixture with *n*-hexane at -35 °C.




**m.p.:** 115.0–116.9 °C; **Raman (300 mW, [cm<sup>-1</sup>]):**  $\nu$  = 2961 (37), 2998 (100), 1412 (24), 636 (37), 552 (63), 442 (22), 398 (24), 376 (14), 344 (43), 152 (44), 121 (13); **IR (ATR, [cm<sup>-1</sup>]):**



$\nu = 2956$  (vw), 2897 (vw), 1411 (vw), 1254 (m), 1123 (vw), 993 (s), 905 (s), 822 (vs), 756 (vw), 724 (s), 690 (w), 639 (w), 548 (vw), 509 (w), 462 (m);  $^1\text{H NMR}$  ( $\text{CD}_2\text{Cl}_2$ , [ppm]):  $\delta = 0.31$  (18H, s, H1), 0.67 (9H, s, H2), 0.76 (9H, s, H3);  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CD}_2\text{Cl}_2$ , [ppm]):  $\delta = 1.4$  (6C, d, C1,  $^3J(\text{CP}) = 3.1$  Hz), 2.2 (3C, m, C2), 3.1 (3C, m, C3);  $^{29}\text{Si}\{^1\text{H}\}$  NMR ( $\text{CD}_2\text{Cl}_2$ , [ppm]):  $\delta = -58.0$  (dd, Si4,  $^2J(\text{SiP}_Y) = 18.5$  Hz,  $^2J(\text{SiP}_M) = 8.5$  Hz), 7.4 (d, Si1,  $^2J(\text{SiP}) = 8.4$  Hz), 10.9 (s, Si2), 11.7 (s, Si3);  $^{15}\text{N}$  NMR ( $\text{CD}_2\text{Cl}_2$ , [ppm]):  $\delta = 96$  (s, N1), 115 (s, N2), 125 (s(br), N3);  $^{71}\text{Ga}\{^1\text{H}\}$  NMR ( $\text{CD}_2\text{Cl}_2$ , [ppm]):  $\delta = 249.6$  (s);  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{CD}_2\text{Cl}_2$ , [ppm]): ABMX<sub>2</sub>Y spin system:  $\delta(\text{P}_A) = -322.9$ ,  $\delta(\text{P}_B) = -320.2$ ,  $\delta(\text{P}_M) = 30.4$ ,  $\delta(\text{P}_X) = 165.5$ ,  $\delta(\text{P}_Y) = 166.8$ ,  $^1J(\text{P}_A\text{P}_B) = -189.8$  Hz,  $^1J(\text{P}_A\text{P}_X) = -143.4$  Hz,  $^1J(\text{P}_B\text{P}_X) = -147.7$  Hz,  $^1J(\text{P}_M\text{P}_X) = -245.1$  Hz,  $^2J(\text{P}_A\text{P}_M) = 18.8$  Hz,  $^2J(\text{P}_B\text{P}_M) = 18.0$  Hz; **elemental analysis for C<sub>12</sub>H<sub>36</sub>GaCl<sub>5</sub>P<sub>6</sub>N<sub>4</sub>Si<sub>5</sub>**: calcd: C 17.8, H 4.5, N 6.9; found: C 17.1, H 4.4, N 5.5.



$^1\text{H NMR}$  ( $\text{CD}_2\text{Cl}_2$ , [ppm]):  $\delta = 0.37$  (18H, s, H2), 0.75 (18H, s, H1);  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{CD}_2\text{Cl}_2$ , [ppm]): ABMOX<sub>2</sub> spin system:  $\delta(\text{P}_A) = -318.3$ ,  $\delta(\text{P}_M) = 33.4$ ,  $\delta(\text{P}_O) = 144$  ( $\Delta\nu_{1/2} = \sim 1200$  Hz),  $\delta(\text{P}_X) = 170.8$ ,  $^1J(\text{P}_A\text{P}_X) = -144.8$  Hz,  $^1J(\text{P}_M\text{P}_X) = -248.6$  Hz,  $^2J(\text{P}_A\text{P}_M) = 19.0$  Hz. Compound **12**[GaCl<sub>4</sub>] was independently synthesized by addition of GaCl<sub>3</sub> (18 mg, 0.10 mmol, 1.0 eq.) to a solution of **19**[GaCl<sub>4</sub>] (64 mg, 0.10 mmol, 1.0 eq.) in CD<sub>2</sub>Cl<sub>2</sub> (1 mL). The obtained colorless solution was stirred for 30 min at ambient temperature and subsequently investigated by  $^1\text{H}$  and  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy.



**m.p.:** 164.5–167.5 °C (decomposition); **Raman** (250 mW, [cm<sup>-1</sup>]):  $\nu = 2962$  (10), 2898 (17), 1095 (10), 633 (13), 549 (100), 441 (11), 397 (17), 384 (10), 354 (10), 138 (81) the Raman measurement was hampered by strong fluorescence effects; **IR** (ATR, [cm<sup>-1</sup>]):  $\nu = 2956$  (vw), 2898 (vw), 1409 (w), 1257 (s), 994 (vs), 899 (m), 814 (vs), 759 (vw), 729 (w), 692 (vw), 637 (w), 544 (w), 490 (w), 441 (vw), 409 (w);  $^1\text{H NMR}$  ( $\text{CD}_2\text{Cl}_2$ , [ppm]):  $\delta = 0.72$  (36H, s, CH<sub>3</sub>);  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CD}_2\text{Cl}_2$ , [ppm]):  $\delta = 2.7$  (12C, s, CH<sub>3</sub>);  $^{29}\text{Si}\{^1\text{H}\}$  NMR ( $\text{CD}_2\text{Cl}_2$ , [ppm]):  $\delta = 13.7$  (s, Si(CH<sub>3</sub>)<sub>3</sub>), the Si atom of the SiN<sub>4</sub>-moiety was not detected;  $^{15}\text{N}$  NMR ( $\text{CD}_2\text{Cl}_2$ , [ppm]):  $\delta = 115$  (s);  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{CD}_2\text{Cl}_2$ , [ppm]): A<sub>2</sub>MX<sub>2</sub> spin system:  $\delta(\text{P}_A) = -320.4$ ,  $\delta(\text{P}_M) = 32.2$ ,  $\delta(\text{P}_X) = 158.8$ ,  $^1J(\text{P}_A\text{P}_X) = -142.7$  Hz,  $^1J(\text{P}_M\text{P}_X) = -259.1$  Hz,  $^2J(\text{P}_A\text{P}_M) = 20.0$  Hz; **elemental analysis for C<sub>12</sub>H<sub>36</sub>Ga<sub>4</sub>Cl<sub>14</sub>P<sub>10</sub>N<sub>4</sub>Si<sub>5</sub>**: calcd: C 10.8, H 2.5, N 3.8; found: C 10.9, H 2.5, N 4.0.

## Acknowledgements

We gratefully acknowledge financial support from the Fonds der Chemischen Industrie (fellowship to M.H.H.) and the DFG (WE4621/2-1).

## Notes and references

- M. H. Holthausen and J. J. Weigand, *Chem. Soc. Rev.*, 2014, **43**, 6639.
- C. A. Dyker and N. Burford, *Chem. – Asian J.*, 2008, **3**, 28.
- P-rich cations R<sub>n</sub>P<sub>m</sub> ( $n < m$ ) were also prepared from P<sub>1</sub>-sources, however in most cases the reaction outcome is unpredictable, see: (a) M. Donath, E. Conrad, P. Jerabek, G. Frenking, R. Fröhlich, N. Burford and J. J. Weigand, *Angew. Chem., Int. Ed.*, 2012, **51**, 2964; (b) K.-O. Feldmann and J. J. Weigand, *Angew. Chem., Int. Ed.*, 2012, **51**, 7545; (c) M. Donath, M. Bodensteiner and J. J. Weigand, *Chem. – Eur. J.*, 2014, **20**, 17306.
- (a) Y. Peng, H. Fan, H. Zhu, H. W. Roesky, J. Magull and C. E. Hughes, *Angew. Chem., Int. Ed.*, 2004, **43**, 3443; (b) G. Prabusankar, A. Doddi, C. Gemel, M. Winter and R. A. Fischer, *Inorg. Chem.*, 2010, **49**, 7976; (c) W. Uhl and M. Benter, *Chem. Commun.*, 1999, 771; (d) Y. Xiong, S. Yao, M. Brym and M. Driess, *Angew. Chem., Int. Ed.*, 2007, **46**, 4511; (e) S. Khan, R. Michel, S. S. Sen, H.-W. Roesky and D. Stalke, *Angew. Chem., Int. Ed.*, 2011, **50**, 11786; (f) S. Khan, R. Michel, J. M. Dieterich, R. A. Mata, H. W. Roesky, J.-P. Demers, A. Lange and D. Stalke, *J. Am. Chem. Soc.*, 2011, **133**, 17889; (g) C. D. Martin, C. M. Weinstein, C. E. Moore, A. L. Rheingold and G. Bertrand, *Chem. Commun.*, 2013, **49**, 4486; For overviews of P<sub>4</sub> activation by nucleophiles, electrophiles and predominantly nucleophilic ambiphiles see: (h) M. Scheer, G. Balázs and A. Seitz, *Chem. Rev.*, 2010, **110**, 4236; (i) N. A. Giffin and J. D. Masuda, *Coord. Chem. Rev.*, 2011, **255**, 1342; (j) S. Khan, S. S. Sen and H. W. Roesky, *Chem. Commun.*, 2012, **48**, 2169.
- I. Krossing and I. Raabe, *Angew. Chem., Int. Ed.*, 2001, **40**, 4406; For insertion of a related NO<sup>+</sup> species into a P–P bond of P<sub>4</sub> see: (a) T. Köchner, S. Riedel, A. J. Lehner, H. Scherer, I. Raabe, T. A. Engesser, F. W. Scholz, U. Gellrich, P. Eiden, R. A. Paz Schmidt, D. A. Plattner and I. Krossing, *Angew. Chem., Int. Ed.*, 2010, **49**, 8139; (b) T. Köchner, T. A. Engesser, H. Scherer, D. A. Plattner, A. Steffani and I. Krossing, *Angew. Chem., Int. Ed.*, 2012, **51**, 6529.
- (a) M. Gonsior, I. Krossing, L. Müller, I. Raabe, M. Jansen and L. van Wüllen, *Chem. – Eur. J.*, 2002, **8**, 4475; (b) I. Krossing, *J. Chem. Soc., Dalton Trans.*, 2002, 500.
- (a) M. H. Holthausen, K.-O. Feldmann, S. Schulz, A. Hepp and J. J. Weigand, *Inorg. Chem.*, 2012, **51**, 3374; (b) M. H. Holthausen and J. J. Weigand, *Z. Anorg. Allg. Chem.*, 2012, **638**, 1103; (c) M. H. Holthausen, A. Hepp and J. J. Weigand, *Chem. – Eur. J.*, 2013, **19**, 9895.





- 8 J. J. Weigand, M. H. Holthausen and R. Fröhlich, *Angew. Chem., Int. Ed.*, 2009, **48**, 295.
- 9 M. H. Holthausen and J. J. Weigand, *J. Am. Chem. Soc.*, 2009, **131**, 14210.
- 10 (a) M. H. Holthausen, C. Richter, A. Hepp and J. J. Weigand, *Chem. Commun.*, 2010, **46**, 6921; (b) J. W. Dube, C. M. E. Graham, C. L. B. Macdonald, Z. D. Brown, P. P. Power and P. J. Ragogna, *Chem. – Eur. J.*, 2014, **20**, 6739.
- 11 M. H. Holthausen, S. K. Surmiak, P. Jerabek, G. Frenking and J. J. Weigand, *Angew. Chem., Int. Ed.*, 2013, **52**, 11078.
- 12 E. Niecke and R. Kröher, *Angew. Chem., Int. Ed. Engl.*, 1976, **15**, 692.
- 13 E. Niecke and W. Bitter, *Chem. Ber.*, 1976, **109**, 415; performing the reaction in 2 : 1 stoichiometry of **5** and SiCl<sub>4</sub> gave **6** and **7** in similar ratios.
- 14 Ref. 13 reports that after removal of **6** the residue consists mainly of polymeric material of unknown constitution and minor amounts of spirocyclic compound ClP(NSiMe<sub>3</sub>)<sub>2</sub>Si(NSiMe<sub>3</sub>)<sub>2</sub>P. Investigation of the residue by <sup>31</sup>P{<sup>1</sup>H} NMR (C<sub>6</sub>H<sub>5</sub>F solvent), however, revealed **7** is the main compound (~60%) next to several minor species, see ESI† for more information.
- 15 L. Stahl, *Coord. Chem. Rev.*, 2000, **210**, 203.
- 16 (a) N. Burford and D. J. LeBlanc, *Inorg. Chem.*, 1999, **38**, 2248; (b) I. Schranz, L. P. Grocholl and L. Stahl, *Inorg. Chem.*, 2000, **39**, 3037.
- 17 E. Niecke, W. Flick and S. Pohl, *Angew. Chem., Int. Ed. Engl.*, 1976, **15**, 309.
- 18 J. Heinicke, S. Mantey, A. Oprea, M. K. Kindermann and P. G. Jones, *Heteroat. Chem.*, 1999, **10**, 605.
- 19 Next to main species **7**, also minor amounts of compounds **8**, **9** and **9'** were observed in the <sup>1</sup>H and <sup>31</sup>P{<sup>1</sup>H} NMR spectra of the residue described in ref. 14, see ESI† for more information.
- 20 P. Paetzgold, C. von Plotho, E. Niecke and R. Rieger, *Chem. Ber.*, 1983, **116**, 1678.
- 21 M. L. Filleus-Blanchard, *Org. Mag. Res.*, 1979, **12**, 12.
- 22 This rotation is not observed at ambient temperature due to the high barrier of rotation reported for exocyclic P–N bonds in amino-substituted diphosphadiazanes (e.g. **8**: Δ*G* > 27 kcal mol<sup>-1</sup>, ref. 17).
- 23 Several isomerization pathways are conceivable: (a) L. Horner and H. Winkler, *Tetrahedron Lett.*, 1964, 461; (b) A. Rauk, L. C. Allen and K. Miskow, *Angew. Chem., Int. Ed. Engl.*, 1970, **9**, 400; (c) J. Silaghi-Dumitrescu, F. Lara-Ochoa and I. Haiduc, *Main Group Chem.*, 1998, **2**, 309; (d) I. Schranz, D. F. Moser, L. Stahl and R. J. Staples, *Inorg. Chem.*, 1999, **38**, 5814; (e) U. Wieringa, H. Voelker, H. W. Roesky, Y. Shermolovich, L. Markovski, I. Uson, M. Noltemeyer and H.-G. Schmidt, *J. Chem. Soc., Dalton Trans.*, 1995, 1951; (f) H.-J. Chen, R. C. Haltiwanger, T. G. Hill, M. L. Thompson, D. E. Coons and A. D. Norman, *Inorg. Chem.*, 1985, **24**, 4725.
- 24 Next to **7** (25%), compounds **5** (33%) and **6** (42%) are main products of this reaction. It is assumed, that **5** and Me<sub>3</sub>SiN=PN(SiCl<sub>3</sub>)(SiMe<sub>3</sub>) form upon *cyclo*-reversion of **9**. Compound Me<sub>3</sub>SiN=PN(SiCl<sub>3</sub>)(SiMe<sub>3</sub>) is assumed to eliminate Me<sub>3</sub>SiCl yielding four-membered heterocycle **6**.
- 25 A. Villinger, P. Mayer and A. Schulz, *Chem. Commun.*, 2006, 1236.
- 26 A. Villinger, A. Westenkirchner, R. Wustrack and A. Schulz, *Inorg. Chem.*, 2008, **47**, 9140.
- 27 (a) B. Luo, V. G. Young and W. L. Gladfelter, *J. Organomet. Chem.*, 2002, 268; (b) H. Schmidtbauer and W. Findeiss, *Angew. Chem., Int. Ed. Engl.*, 1964, **3**, 696.
- 28 M. Rastätter, P. W. Roesky, D. Gudat, G. B. Deacon and P. C. Junk, *Chem. – Eur. J.*, 2007, **13**, 7410.
- 29 (a) R. W. Shaw, *Phosphorus, Sulfur Silicon Relat. Elem.*, 1978, **4**, 101; (b) S. S. Krishnamurthy, *Phosphorus, Sulfur Silicon Relat. Elem.*, 1994, **87**, 101.
- 30 A. H. Cowley, M. Lattman and J. C. Wilburn, *Inorg. Chem.*, 1981, **20**, 2916.

