RSC Advances



PAPER View Article Online



Cite this: RSC Adv., 2019, 9, 11865

Aggregation-induced emission enhancement (AIEE)-active tetraphenylethene (TPE)-based chemosensor for Hg²⁺ with solvatochromism and cell imaging characteristics†

Aling Tang, Zhao Chen, **Diandian Deng, Gang Liu, **D Yayi Tu**D and Shouzhi Pu**

An aggregation-induced emission enhancement (AIEE)-active fluorescent sensor based on a tetraphenylethene (TPE) unit has been successfully designed and synthesized. Interestingly, the luminogen could detect Hg^{2+} with high selectivity in an acetonitrile solution without interference from other competitive metal ions, and the detection limit was 7.46×10^{-6} mol L⁻¹. Furthermore, the luminogen also showed interesting solvatochromic behavior and superior cell imaging performance.

Received 19th March 2019 Accepted 11th April 2019

DOI: 10.1039/c9ra02119a

rsc.li/rsc-advances


The design and synthesis of molecular sensors for the detection of metal ions, especially transition-metal ions, has attracted much attention. Among all transition-metal ions, mercury (Hg²⁺) is identified as one of the most dangerous and ubiquitous heavy metals. Indeed, it is not biodegradable, and can cause extreme injury to the environment as well as human health. Additionally, it can be accumulated through the food chain in the human body, consequently giving rise to several deleterious effects such as central nervous system defects, kidney damage, endocrine system disease and so on. Although many governments around the world have adopted strict regulations to limit Hg²⁺ emission, the global Hg²⁺ pollution caused by human activities is still serious. Therefore, it is highly desirable to develop a new method for the detection of Hg²⁺ with high selectivity and sensitivity.

Most traditional fluorescent sensors suffer from a detrimental phenomenon called aggregation-caused quenching (ACQ), which usually results in a poor solid-state emission efficiency. Fortunately, in 2001, Tang *et al.* reported a fluorescent molecule named 1-methyl-1,2,3,4,5-pentaphenylsilole. Interestingly, the fluorescence emission of this luminogen was induced by aggregation, a phenomenon referred to as aggregation-induced emission (AIE). Subsequently, in 2002, Park *et al.* reported an interesting phenomenon named aggregation-induced emission enhancement (AIEE). In fact, both AIE and AIEE can achieve highly efficient emission in the solid state or aggregated state. In the past years, AIE (or AIEE) phenomenon has attracted considerable research interest owing to the potential applications

Jiangxi Key Laboratory of Organic Chemistry, Jiangxi Science and Technology Normal University, Nanchang 330013, PR China. E-mail: chenzhao666@126.com; pushouzhi@tsinghua.org.cn

† Electronic supplementary information (ESI) available: Experimental section, NMR spectra, mass spectrum, and characterization data mentioned in the paper. See DOI: 10.1039/c9ra02119a

in a lot of fields, including bioimaging, fluorescence sensors, organic lighting emitting diode (OLED) devices and organic lasers.²⁸⁻³³ Meanwhile, many stimuli-responsive materials have been reported, including photochromism, mechanochromism, and solvatochromism.34-46 At present, the solvatochromism materials have been generally used in chemical and biological systems.47 To date, many fluorescent chemosensors for Hg2+ have been reported. In contrast, the corresponding chemosensors with AIE or AIEE effect are rare, not to mention solvatochromic AIE or AIEE-active fluorescent sensors for Hg2+ with good cell imaging behavior. Indeed, preparing such multifunctional sensors is challenging and significative. In this paper, we reported an AIEE-active tetraphenylethene (TPE)-based fluorescent sensor (Scheme 1) for the detection of Hg²⁺. Furthermore, the luminogen also showed remarkable solvatochromism and good cell imaging characteristics.



Scheme 1 The synthetic route of luminogen 1.

RSC Advances Paper

To investigate the AIEE phenomenon of luminogen 1, we initially studied the UV-visible absorption spectra and photoluminescence (PL) spectra in acetonitrile-water mixtures with different water fractions (f_w) . The results indicated that the absorption spectra exhibited level-off tails in the long wavelength region as the water content increased (Fig. S1†). It is wellknown that such tails are usually associated with the Mie scattering effect, which is the key signal of nano-aggregate formation. 48,49 As presented in Fig. 1, luminogen 1 showed weak fluorescence and the luminescence quantum yield (Φ) was as low as 1.35%. Interestingly, when the f_w in the acetonitrile solution was increased to 80%, an obvious emission band was formed, and a vellow-green fluorescence was observed. As the water content was increased to 90%, the emission intensity was further increased, and a bright yellow-green luminescence (Φ = 27.81%) with a λ_{max} at 545 nm could be observed.

Clearly, water is a poor solvent of luminogen 1. As a result, the generation of the yellow-green emission can be attributed to the aggregate formation. Moreover, as shown in Fig. 2, the nano-aggregates obtained were verified by dynamic light scatterings (DLS). Therefore, 1 is a typical AIEE-active fluorescent molecule, and its AIEE behavior is caused by the restricted intramolecular rotation. As shown in Fig. S2,† solid-state compound 1 showed a strong green emission ($\Phi = 14.50\%$) with a $\lambda_{\rm max}$ at 497 nm, and the corresponding fluorescence lifetime is 2.66 ns (Fig. S3†).

On the other hand, the luminogen 1 also displayed interesting solvatochromism effect. As presented in Fig. 3, the absorption spectra and fluorescence spectra of luminogen 1 in different solvents were investigated, respectively. Obviously, the absorption spectra was barely affected by the polarity of solvents. However, the photoluminescence peaks were gradually red-shifted from 515 nm to 625 nm, and thus 1 exhibited remarkable solvatochromism behavior. Obviously, the




Fig. 1 (a) Fluorescence spectra of the dilute solutions of 1 (2.0 \times 10^{-5} mol L $^{-1}$) in acetonitrile–water mixtures with different water content (0–90%). Excitation wavelength = 380 nm. (b) Changes the emission intensity of 1 at 545 nm in acetonitrile–water mixtures with different volume fractions of water (0–90%). (c) PL images of 1 (2.0 \times 10^{-5} mol L $^{-1}$) in acetonitrile–water mixtures with different $f_{\rm W}$ values under 365 nm UV illumination.




Fig. 2 Size distribution curve of luminogen 1 (2.0 \times 10⁻⁵ mol L⁻¹) in acetonitrile—water mixtures with $f_{\rm w}=90\%$.

molecular structure of luminogen 1 is distorted due to the presence of TPE unit, and the conjugation degree of molecule 1 is different in various solvents, and the intramolecular charge transfer (ICT) effect is possibly responsible for the solvatochromism behavior of 1.

Subsequently, the changes in the fluorescence of luminogen 1 induced by Hg^{2+} were investigated in acetonitrile (2.0 \times 10^{-5} mol L^{-1}) at room temperature. In the fluorometric titration experiments, as shown in Fig. 4, the emission intensity significantly decreased when the Hg^{2+} concentration increased from 0 to 7.0 equivalents in acetonitrile. Meanwhile, the fluorescent color changed from orange-red to colorless, and followed by a plateau upon further titration (Fig. 5). Remarkably, the emission intensity of luminogen 1 was almost quenched completely. Furthermore, based on the titration experiments, the detection limit of luminogen 1 for Hg^{2+} on the basis of LOD = $3 \times \sigma/\mathrm{B}$ (where σ is the standard deviation of blank sample and B is the slope between the fluorescence intensity *versus* Hg^{2+} concentration) was 7.46×10^{-6} M (Fig. S4†). Moreover,




Fig. 3 (a) UV-Vis absorption spectra and (b) normalized fluorescence spectra (Excitation wavelength =380 nm) of luminogen 1 in different solvents (2.0 \times 10^{-5} mol L $^{-1}$). (c) Photographs of 1 under 365 nm UV illumination in different solvents: tol (toluene); THF (tetrahydrofuran); EA (ethyl acetate); BT (acetone); MeCN (acetonitrile); DMSO (dimethyl sulfoxide).

Paper

90 75 0 eq. Hg²⁺ 7 eq. 500 550 600 650 700 750 Wavelength (nm)

Fig. 4 Fluorescence titration spectra of luminogen 1 (2.0 \times 10^{-5} mol L⁻¹) induced by Hg²⁺ (0–7.0 equiv.) in an acetonitrile solution. Excitation wavelength = 380 nm.




Fig. 5 The emission intensity changes of luminogen $\bf 1$ at 625 nm with different equivalents of Ha^{2+} .

a good linear relationship could be obtained (R = -0.9933) and the quenching constant of luminogen 1 with Hg^{2+} was 1.9×10^4 M⁻¹ (Fig. S5†). On the other hand, the binding ratio of luminogen 1 for Hg2+ was established through Job's plot and the results showed 1:1 stoichiometric complexation (Fig. S6†). Next, the binding interactions between 1 and Hg²⁺ were further investigated by NMR in acetonitrile- d_3 . As presented in Fig. 6, the signal of Ha from 9.42 ppm shifted to 9.62 ppm and the Hb changed from 8.43 ppm to 9.24 ppm. Besides, the Hc or Hd was slightly shifted for 0.04 ppm or 0.06 ppm, respectively. These consequences revealed that the N on the pyridine and the N on the pyrazine (near the N on the pyridine) are the most probable binding with Hg²⁺. Mass spectra were utilized to further demonstrate the binding mode of luminogen 1 toward Hg²⁺. The peak located at m/z = 820.0 was coincided well with the ensemble $[1 + Hg^{2+} + 2NO_3^- + Cl^-]^-$, confirming the binding ratio of luminogen 1 for Hg^{2+} with 1: 1 stoichiometry (Fig. S7†).

Next, in order to study the selectivity behavior of luminogen 1 as a fluorescent sensor for Hg^{2+} , other metal ions, such as Zn^{2+} , Cd^{2+} , Ba^{2+} , Sr^{2+} , Mn^{2+} , Mg^{2+} , Ca^{2+} , Pb^{2+} , Ni^{2+} , Co^{2+} , Cu^{2+} , Al^{3+} , Fe^{3+} , Cr^{3+} , Ag^+ and K^+ were also measured in acetonitrile




Fig. 6 1 H NMR (acetonitrile- d_{3} , 400 MHz) spectra changes of luminogen 1 in the presence of Hg²⁺.

under the same experimental conditions. The corresponding UV-Vis absorption spectra were shown in Fig. S8.† Furthermore, as showed in Fig. 7, when these cations were added separately into the solution containing luminogen 1, no obvious fluorescence changes were observed. Indeed, as noticed in Fig. S9,† no obvious interference was observed when Hg^{2+} (7.0 equiv.) was added with other ions (7.0 equiv.). These results indicated that luminogen 1 could be served as a highly selective fluorescent sensor for detection of Hg^{2+} .

Fluorescent probe is a powerful tool for optical imaging, which allows direct visualization of biological analytes. 53 Luminogen 1 was AIEE-active due to the restriction of intramolecular rotation in the aggregate state, which is beneficial for cell imaging. Indeed, the viability of HeLa cells incubated with luminogen 1 was evaluated by the standard MTT method (Fig. 8), and the result indicated that compound 1 exhibited low cytotoxicity. Next, cell imaging behavior of luminogen 1 was investigated using a confocal laser scanning microscopy (CLSM). HeLa cells were incubated with luminogen 1 (20 μM) for 30 min at 37 $^{\circ} C$ and the fluorescence images were obtained by CLSM.

As presented in Fig. 9, an intense yellow-green fluorescence, which was consistent with the fluorescence of luminogen 1 in




Fig. 7 (a) Fluorescence spectra of luminogen 1 (2.0×10^{-5} mol L $^{-1}$) towards various cations including Zn $^{2+}$, Cd $^{2+}$, Hg $^{2+}$, Ba $^{2+}$, Sr $^{2+}$, Mn $^{2+}$, Mg $^{2+}$, Ca $^{2+}$, Pb $^{2+}$, Ni $^{2+}$, Co $^{2+}$, Cu $^{2+}$, Al $^{3+}$, Fe $^{3+}$, Cr $^{3+}$, Ag $^{+}$ and K $^{+}$. Excitation wavelength = 380 nm. (b) Fluorescence photographs of luminogen 1 after addition of various metal ions (7.0 equiv.) under 365 nm light.

RSC Advances Paper




Fig. 8 The MTT assay of compound 1 for measuring cell viability.




Fig. 9 Fluorescence images of HeLa cells incubated with luminogen 1 (20 μM) for 30 min at 37 °C: (a) bright field image; (b) fluorescence image; (c) merge image (a and b).

acetonitrile-water mixture with high water content (80% or 90%), was displayed inside the cells. This result indicated that luminogen 1 tended to aggregate inside the cells, and thus the bright yellow-green luminescence was clearly observed. Furthermore, the merged picture c demonstrated that picture a and picture b overlapped very well. These results indicated that luminogen 1 showed superior cell imaging performance.

In summary, a TPE-based fluorescent molecule was designed and synthesized. The luminogen exhibited obvious AIEE phenomenon. Moreover, luminogen 1 could be used as a highly selective fluorescence turn-off chemosensor for Hg²⁺, and the detection limit was 7.46×10^{-6} mol L⁻¹. Furthermore, 1 also displayed interesting solvatochromic behavior and superior cell imaging performance. Further studies on the design of new AIE or AIEE-active fluorescent chemosensors are in progress.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors are grateful for the financial support from the National Natural Science Foundation of China (21702079 and 41867053), the "5511" Science and Technology Innovation Talent Project of Jiangxi Province (20165BCB18015), the key project of Natural Science Foundation of Jiangxi Province (20171ACB20025), the youth project of Natural Science

Foundation of Jiangxi Province (20171BAB213004), the Young Talents Project of Jiangxi Science and Technology Normal University (2017QNBJRC005), and the Project of the Science Fund of Jiangxi Education Office (GJJ180593).

Notes and references

- 1 C. Zhang, S. Pu, Z. Sun, C. Fan and G. Liu, J. Phys. Chem. B, 2015, 119, 4673-4682.
- 2 H. N. Kim, W. X. Ren, J. S. Kim and J. Yoon, Chem. Soc. Rev., 2012, 41, 3210-3244.
- 3 S. Xia, G. Liu and S. Pu, J. Mater. Chem. C, 2015, 3, 4023-4029.
- 4 J. D. Demers, J. D. Blum, S. C. Brooks, P. M. Donovan, A. L. Riscassi, C. L. Miller, W. Zheng and B. Gu, Environ. Sci.: Processes Impacts, 2018, 20, 686-707.
- 5 M. F. Wolfe, S. Schwarzbach and R. A. Sulalman, Environ. Toxicol. Chem., 1998, 17, 146-160.
- 6 X. Liu, X. Liu, M. Tao and W. Zhang, J. Mater. Chem. A, 2015, 3, 13254-13262.
- 7 W. Luo, H. Jiang, K. Zhang, W. Liu, X. Tang, W. Dou, Z. Ju, Z. Li and W. Liu, J. Mater. Chem. B, 2015, 3, 3459-3464.
- 8 S. Gupta and M. D. Milton, New J. Chem., 2018, 42, 2838-2849.
- 9 W. Fang, G. Zhang, J. Chen, L. Kong, L. Yang, H. Bi and J. Yang, Sens. Actuators, B, 2016, 229, 338-346.
- 10 C. Li and S. Liu, J. Mater. Chem., 2010, 20, 10716-10723.
- 11 O. Abollino, A. Giacomino, M. Malandrino, G. Piscionieri and E. Mentasti, Electroanalysis, 2008, 20, 75-83.
- 12 L. Yang, Y. Su, Y. Geng, H. Xiong, J. Han, Q. Fang and X. Song, Org. Biomol. Chem., 2018, 16, 5036-5042.
- 13 Z. Ruan, Y. Shan, Y. Gong, C. Wang, F. Ye, Y. Qiu, Z. Liang and Z. Li, J. Mater. Chem. C, 2018, 6, 773-780.
- 14 K. Wang, J. Li, S. Ji, L. Li, Z. Qiu, C. Pan, J. Zhang and Y. Huo, New J. Chem., 2018, 42, 13836-13846.
- 15 S.-L. Pan, K. Li, L.-L. Li, M.-Y. Li, L. Shi, Y.-H. Liu and X.-Q. Yu, Chem. Commun., 2018, 54, 4955-4958.
- 16 S. Chen, W. Wang, M. Yan, Q. Tu, S.-W. Chen, T. Li, M.-S. Yuan and J. Wang, Sens. Actuators, B, 2018, 255, 2086-2094.
- 17 Y. Chen, W. Zhang, Y. Cai, R. T. K. Kwok, Y. Hu, J. W. Y. Lam, X. Gu, Z. He, Z. Zhao, X. Zheng, B. Chen, C. Gui and B. Z. Tang, Chem. Sci., 2017, 8, 2047-2055.
- 18 J. Luo, Z. Xie, J. W. Y. Lam, L. Cheng, H. Chen, C. Qiu, H. S. Kwok, X. Zhan, Y. Liu, D. Zhu and B. Z. Tang, Chem. Commun., 2001, 1740-1741.
- 19 B.-K. An, S.-K. Kwon, S.-D. Jung and S. Y. Park, J. Am. Chem. Soc., 2002, 124, 14410-14415.
- 20 Z. Chen, J. Zhang, M. Song, J. Yin, G.-A. Yu and S. H. Liu, Chem. Commun., 2015, 51, 326-329.
- 21 Z. Chen, Z. Li, F. Hu, G.-A. Yu, J. Yin and S. H. Liu, Dyes Pigm., 2016, 125, 169-178.
- 22 F. Zhao, Z. Chen, G. Liu, C. Fan and S. Pu, Tetrahedron Lett., 2018, 59, 836-840.
- 23 Y. Yin, F. Zhao, Z. Chen, G. Liu and S. Pu, Tetrahedron Lett., 2018, 59, 4416-4419.
- 24 Z. Chen, G. Liu, S. Pu and S. H. Liu, Dyes Pigm., 2017, 143, 409-415.

Paper

25 L. Yu, Z. Wu, G. Xie, C. Zhong, Z. Zhu, D. Ma and C. Yang,

26 X. Ma, L. Hu, X. Han and J. Yin, *Chin. Chem. Lett.*, 2018, 29, 1489–1492.

Chem. Commun., 2018, 54, 1379-1382.

- 27 X. Han, Y. Liu, G. Liu, J. Luo, S. H. Liu, W. Zhao and J. Yin, *Chem.-Asian J.*, 2019, **14**, 890–895.
- 28 F. Hua and B. Liu, Org. Biomol. Chem., 2016, 14, 9931-9944.
- 29 Z. Chen, J. Liang, Y. Nie, X. Xu, G.-A. Yu, J. Yin and S. H. Liu, *Dalton Trans.*, 2015, **44**, 17473–17477.
- 30 X. Sun, A. Zebibula, X. Dong, G. Zhang, D. Zhang, J. Qian and S. He, *ACS Appl. Mater. Interfaces*, 2018, **10**, 25037–25046.
- 31 D. Wu, A. C. Sedgwick, T. Gunnlaugsson, E. U. Akkaya, J. Yoon and T. D. James, *Chem. Soc. Rev.*, 2017, 46, 7105–7123.
- 32 J. Huang, N. Sun, P. Chen, R. Tang, Q. Li, D. Ma and Z. Li, *Chem. Commun.*, 2014, **50**, 2136–2138.
- 33 F. Liu, J. Tu, X. Wang, J. Wang, Y. Gong, M. Han, X. Dang, Q. Liao, Q. Peng, Q. Li and Z. Li, *Chem. Commun.*, 2018, 54, 5598–5601.
- 34 S. Guo, C. Fan, G. Liu and S. Pu, RSC Adv., 2018, 8, 39854-39864.
- 35 B. Wang and C. Wei, RSC Adv., 2018, 8, 22806-22812.
- 36 Z. Chi, X. Zhang, B. Xu, X. Zhou, C. Ma, Y. Zhang, S. Liu and J. Xu, *Chem. Soc. Rev.*, 2012, 41, 3878–3896.
- 37 F. Chen, W. Zhang, Z. Liu, L. Meng, B. Bai, H. Wang and M. Li, *RSC Adv.*, 2019, **9**, 1–10.
- 38 J. Liang, Z. Chen, L. Xu, J. Wang, J. Yin, G.-A. Yu, Z.-N. Chen and S. H. Liu, *J. Mater. Chem. C*, 2014, 2, 2243–2250.
- 39 Z. Chen, Z. Li, L. Yang, J. Liang, J. Yin, G.-A. Yu and S. H. Liu, Dyes Pigm., 2015, 121, 170–177.

- 40 J. Liang, Z. Chen, J. Yin, G.-A. Yu and S. H. Liu, *Chem. Commun.*, 2013, 49, 3567–3569.
- 41 C. Gstrein, B. Zhang, M. A. Abdel-Rahman, O. Bertran, C. Alemán, G. Wegner and A. D. Schlüter, *Chem. Sci.*, 2016, 7, 4644–4652.
- 42 R. Tan, S. Wang, H. Lan and S. Xiao, Curr. Org. Chem., 2017, 21, 236–248.
- 43 R. Wang, Y. Sun, F. Zhang, M. Song, D. Tian and H. Li, *Angew. Chem., Int. Ed.*, 2017, **56**, 5294–5298.
- 44 W. Chen, Y. Pan, J. Chen, F. Ye, S. H. Liu and J. Yin, *Chin. Chem. Lett.*, 2018, **29**, 1429–1435.
- 45 J. Chen, D. Li, W. Chi, G. Liu, S. H. Liu, X. Liu, C. Zhang and J. Yin, *Chem.–Eur. J.*, 2018, 24, 3671–3676.
- 46 J. Chen, F. Ye, Y. Lin, Z. Chen, S. Liu and J. Yin, *Sci. China: Chem.*, 2019, **62**, 440–450.
- 47 C. Karthik, V. Manjuladevi, R. K. Gupta and S. Kumar, RSC Adv., 2015, 5, 84592–84600.
- 48 C. W. T. Leung, Y. Hong, S. Chen, E. Zhao, J. W. Y. Lam and B. Z. Tang, *J. Am. Chem. Soc.*, 2013, **135**, 62–65.
- 49 Z. Chen, D. Wu, X. Han, J. Liang, J. Yin, G.-A. Yu and S. H. Liu, *Chem. Commun.*, 2014, **50**, 11033–11035.
- 50 Z. Chen, X. Han, J. Zhang, D. Wu, G.-A. Yu, J. Yin and S. H. Liu, RSC Adv., 2015, 5, 15341–15349.
- 51 Z. Chen, G. Liu, S. Pu and S. H. Liu, *Dyes Pigm.*, 2018, **152**, 54–59.
- 52 J. Mei, N. L. C. Leung, R. T. K. Kwok, J. W. Y. Lam and B. Z. Tang, *Chem. Rev.*, 2015, **115**, 11718–11940.
- 53 Z. Xu, X. Huang, X. Han, D. Wu, B. Zhang, Y. Tan, M. Cao, S. H. Liu, J. Yin and J. Yoon, *Chem*, 2018, 4, 1609–1628.