Chemical Science



EDGE ARTICLE

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2022, 13, 13403

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 24th August 2022 Accepted 20th October 2022

DOI: 10.1039/d2sc04727f

rsc.li/chemical-science

Total synthesis of the antibacterial polyketide natural product thailandamide lactone†

Stereoselective total synthesis of the structurally intriguing polyketide natural product thailandamide lactone was accomplished, and done so using a convergent approach for the first time to the best of our knowledge. The key features of this synthesis included use of a Crimmins acetate aldol reaction, Evans methylation, Urpi acetal aldol reaction, Sharpless asymmetric epoxidation and subsequent γ -lactonization for the installation of six asymmetric centers and the use of the Negishi reaction, Julia-Kocienski olefination, cross metathesis, HWE olefination and intermolecular Heck coupling for construction of a variety of unsaturated linkages. Pd(i)-based Heck coupling was introduced, for the first time to the best of our knowledge, quite efficiently to couple the major eastern and sensitive western segments of the molecule. The antibacterial activity of thailandamide lactone was also evaluated.

Introduction

During the mining of the genome of Burkholderia thailandensis, a bacterium isolated from rice paddies in central and northeastern Thailand, Hertweck and co-workers1 in 2008 first observed the labile polyene polyketide thailandamide A (1, Fig. 1), albeit in minute quantities. To better understand thailandamide biosynthesis, the silent tha PKS-NRPS gene cluster of B. thailandensis was activated by a research group in 2010 through manipulation of a quorum sensing (QS) regulatory system that produced a mutant with a dramatically altered metabolic profile.2 This process resulted in the isolation of the structurally challenging new polyketide thailandamide lactone (2, Fig. 1), which was not detected in wild type broth initially. Moreover, the production of thailandamide A (1) was significantly greater here than when the wild type was used.² Later, the same group developed an elegant biosynthetic route yielding the first total synthesis of thailandamide A; this development enabled them to find another unstable metabolite, namely thailandamide B (3, Fig. 1), a geometrical isomer of thailandamide A.3 Broad biological screening of thailandamide A revealed its selective and potential inhibitory activity against various pathogenic Gram-positive and Gram-negative bacteria with a specific mode of action.4 However, the antibacterial activity of thailandamide lactone and thailandamide B remained undisclosed. The highly challenging architectural features and natural scarcity of thailandamide lactone and lack of a synthetic route to this lactone—together with our continual interest in natural products chemistry⁵—encouraged us to seek out its total synthesis. Structurally, thailandamide lactone² is a linear polyene polyketide where a tetraene conjugated with a γ -butyrolactone is fused with a conjugated triene through an enolized dione moiety. It consists of six asymmetric centers including a quaternary center at one terminus of the molecule and a phenolic moiety at the other terminus. Herein, we report a convergent and highly modular route for the first total synthesis of thailandamide lactone and report its antibacterial activity against various pathogenic and non-pathogenic bacterial strains.

Results and discussion

A retrosynthetic analysis of thailandamide lactone (2) is shown in Scheme 1. We envisioned that the target molecule could be constructed from vinyl iodide 4 and 1,3-dione-containing

Fig. 1 Chemical structures of thailandamide family natural products.

[&]quot;School of Chemical Sciences, Indian Association for the Cultivation of Science, Jadavpur, Kolkata-700032, India. E-mail: ocrkg@iacs.res.in

bSchool of Biological Sciences, Indian Association for the Cultivation of Science, Jadavpur, Kolkata-700032, India. E-mail: ritesh.pal@iacs.res.in

[†] Electronic supplementary information (ESI) available. CCDC 2184015 and 2184016. For ESI and crystallographic data in CIF or other electronic format see DOI: https://doi.org/10.1039/d2sc04727f

Scheme 1 Retrosynthetic analysis of thailandamide lactone (2).

polyene 5 adopting intermolecular Heck coupling as the key step. Vinyl iodide 4 could further be made from compounds 6 and 7 by performing amide coupling. Compound 6 could be assembled using the Crimmins acetate aldol reaction, Julia-Kocienski olefination and Evans methylation as the key steps, whereas compound 7 could be accessed utilizing cross olefin metathesis or Julia-Kocienski olefination and the Negishi reaction as the pivotal steps. On the other hand, keto-alkene 5 could be prepared from intermediates 8 and 9 using Heck coupling, whereas compound 9 could be synthesized using the Negishi reaction, Julia-Kocienski olefination, HWE olefination, Urpi acetal aldol reaction, Sharpless asymmetric epoxidation and subsequent γ -lactonization as the salient steps.

The synthesis of intermediate **6** was commenced with the known compound **11** (Scheme 2) prepared from commercially available 4-hydroxy benzaldehyde (**10**) following a literature procedure; compound **11** was subjected to a Crimmins acetate aldol reaction using the known auxiliary 12^{7b} in the presence of TiCl₄/DIPEA to obtain compound **13** as the major product in 80% yield along with its minor counterpart (dr = 4:1). The major aldol product was separated from other components using silica gel column chromatography and its structure was confirmed unambiguously using X-ray crystallographic analysis. Next, compound **13** was treated with TBSOTf/2,6-lutidine followed by NaBH₄ to access compound **14**, which was subjected to the Mitsunobu reaction using 1-phenyl-5-thiotetrazole (15) in the presence of DIAD/PPh₃ and oxidized further using

 $({\rm NH_4})_6{\rm Mo_7O_{24}}\cdot 4{\rm H_2O/H_2O_2}$ in ethanol^{5c} to achieve sulfone **16** in very good overall yield. Next, sulfone **16** was subjected to Julia-Kocienski olefination^{5c,9} with the known aldehyde **17** ¹⁰using KHMDS to obtain the corresponding *E*-coupled isomer as the major product along with its minor *Z*-isomer (dr = 4:1). The purified major isomer was subsequently treated with PPTS to obtain alcohol **18**, which finally was oxidized to acid **6** using Swern oxidation followed by Pinnick oxidation.¹¹

The synthesis of intermediates 7 and 8 is depicted in Scheme 3. The known vinyl iodide 19,12 prepared from propargylic alcohol using the Negishi reaction as the key step, was subjected to the Mitsunobu reaction using 1-phenyl-5-thiotetrazole (15) and the resultant sulfide was oxidized using (NH₄)₆-Mo₇O₂₄·4H₂O/H₂O₂ in dioxane¹³ to access sulfone 20. Notably, the production of sulfone 20 from its corresponding sulfide was found to be much more efficient in dioxane than in the commonly used ethanol. Sulfone 20 was then reacted with the known aldehyde 21 14 following the Julia-Kocienski olefination protocol.^{9,15} Several conditions were screened for synthesizing compound 7 (Table 1) and the use of KHMDS in DME (entry-4) was found to be the best (E/Z = 3:1). In parallel, the cross metathesis16 between the known alkenes 22 17 and 23 18 was also investigated and it was observed that HG-II produced compound 7 in 32% yield with much better selectivity (E: Z =10:1) compared to Julia-Kocienski olefination whereas G-II and HG-I functioned ineffectively, leaving a trace amount of the desired product. However, the geometrical isomers remained inseparable at this stage. On the other hand, commercially available Weinreb amide 24 was transformed to the known compound 25 following a literature procedure19 and subjected further to a reaction with vinyl magnesium bromide to access intermediate 8 in very good overall yield.

The synthesis of aldehyde **34** is described in Scheme 4. The known aldehyde **27** prepared from prenol (**26**) following a literature method²⁰ was converted to the corresponding acetal using (MeO)₃CH/CSA, which was subjected further to the Urpi acetal aldol reaction^{5b,21} in the presence of TiCl₄/DIPEA/SnCl₄ to access compound **29** with excellent selectivity (dr = 20:1). The purified compound was then treated with LiOH·H₂O/H₂O₂ followed by NaOMe/MeOH to obtain compound **30** in 72% yield. The stereochemistry of asymmetric centers newly generated using Urpi acetal aldol reaction was confirmed further from an X-ray

Scheme 2 Synthesis of intermediate 6.

Edge Article Chemical Science

Scheme 3 Synthesis of intermediates 7 and 8

Table 1 Efforts to optimize Julia-Kocienki olefination

Entry	Conditions	Yield $(E:Z)$
1	NaHMDS, THF, −78 °C	66% (1:1.2)
2	KHMDS, THF, −78 °C	70% (1:1)
3	LiHMDS, DME, −60 °C	62% (1.5:1)
4	KHMDS, DME, -60 °C	68% (3:1)

crystallographic analysis of compound 31, which was synthesized from compound 30 by performing tritylation. Next, compound 30 was reacted with BnBr/K2CO3 to obtain benzyl ester 32, which was subjected to Sharpless asymmetric epoxidation²² followed by hydrogenation to produce the corresponding epoxy acid. The stage was set for γ-lactonization.²³ The corresponding epoxy acid was treated with CSA/CH₂Cl₂ to access the 5-exo cyclized product 33 exclusively. The characteristic NOESY correlation of C₄-Me with C₂-H and C₃-H confirmed its structure unambiguously. We did not observe the formation of any other possible γ -lactone originating via 6-endo cyclization followed by concomitant acyl migration.23 Our exhaustive efforts for achieving oxidative cleavage of the diol moiety of compound 33 using either NaIO₄ or NaIO₄/NaHCO₃

Scheme 5 Synthesis of intermediate 9

did not produce aldehyde 34 in an isolable yield due to its rapid decomposition. Delightfully, silica-supported NaIO₄²⁴ provided considerable relief here, where the required aldehyde was obtained quantitatively.

The construction of compound 9 is shown in Scheme 5. Alcohol 19 was oxidized using the Swern condition and subjected to Julia-Kocienski olefination9 with the known sulfone 35^{25} to access compound 36 as a major product (dr = 5:1). The purified major isomer was reacted further with aldehyde 34 in the presence of NaHMDS/THF following the HWE olefination protocol25 to achieve intermediate 9 exclusively. The initially encountered isomerization problem with the α-methyl center was overcome by performing a controlled addition of NaHMDS and also by reducing the reaction time (see the optimization in Table S1 in ESI†).

The synthesis of major coupling partners (4 and 5) of thailandamide lactone is described in Scheme 6. Compound 7 was treated with 10% TFA/CH₂Cl₂ and the resultant Boc-deprotected amine was coupled with acid 6 to access the western segment 4 in 80% yield (over two steps). On the other hand, vinyl ketone 8 was subjected to intermolecular Heck coupling with compound 9 in the presence of Pd(OAc)₂/Bu₄NCl/Et₃N in DMF²⁶ to obtain the corresponding coupled product in complete regioselectivity, and this coupled species was further treated with CSA to obtain the corresponding β-hydroxy ketone in 77% yield after two steps. Substantial trials have been conducted to optimize its conversion to the eastern segment 5. Most of the oxidizing agents including DMP/NaHCO3 did not function properly as

Scheme 4 Synthesis of aldehyde 34.

Chemical Science Edge Article

Scheme 6 Synthesis of coupling partners 4 and 5.

their use ended up with complete decomposition of the product. However, DMP without NaHCO3 produced the required product 5 in 73% yield. The appearance of a signal at a δ of 15.6 ppm in the ¹H NMR spectrum of compound 5 and two carbonyl carbons at δ 183.0, 183.9 ppm in its ¹³C NMR spectrum clearly ascertained its existence as keto-enol tautomeric mixtures.

The completion of the total synthesis of thailandamide lactone is depicted in Scheme 7 where the stage was set for the crucial coupling between the western (4) and eastern (5) segments. Extensive efforts were made to optimize the Heck coupling (Table 2). Trials with PdCl₂(MeCN)₂/Et₃N/HCO₂H in MeCN (entry-1)²⁷ ended up with complete decomposition of staring materials, whereas those with Pd(PPh₃)₄/Et₃N/Bu₄NCl in

Scheme 7 Completion of the synthesis of thailandamide lactone (2).

DMF (entry-2)27 provided the coupled product 37 in a trace amount. Use of Pd(PPh₃)₂Cl₂/K₂CO₃/Bu₄NCl in DMF (entry-3), Pd(OAc)₂/Et₃N/Bu₄NCl in DMF (entry-4)²⁷ and Pd(OAc)₂/K₃PO₄ in DMF (entry-5)^{27c} resulted in the required product in 10%, 45% and 40% yields, respectively. A mixture of some unidentified compounds was formed along with the required compound 37 in most of the cases. Having moderate success in the transformation of compound 37 using either Pd(0) or Pd(II), we then turned our attention towards Pd(1)-catalyzed Heck coupling as it has provided excellent results in some cases.28 Thus, $[Pd(\mu-I)(Pt-Bu_3)]_2$ (entry-6), prepared from PdI_2/P^tBu_3 following a literature report,28a was then screened in the presence of DIPEA/toluene to furnish the coupled product in an improved yield (58%). Later, an equimolar mixture of Pd(OAc)₂ and Pd(PPh₃)₄ in the presence of K₃PO₄/DMF (entry-7)^{28e} was tested. Delightfully, this reaction was found to proceed in a considerably cleaner manner than those with all the other tested conditions, and the coupled product 37 was obtained in 77% yield. All the reactions were performed at room temperature to reduce the rate of decomposition. A detailed NMR study unambiguously confirmed the identity of compound 37 (see the 2D spectra in ESI†). Notably the attempted synthesis of the corresponding compound requisite for an alternative Heck coupling with compound 9 was not successful-because the corresponding β-hydroxy ketone obtained from the Heck coupling between compounds 4 and 8 followed by subsequent TES ether deprotection was found to be very sensitive to various oxidizing agents including DMP. Next, compound 37 was subjected to global deprotection using HF Py to access compound 229 in 89% yield. ¹H and ¹³C NMR data (see comparison Table S2 in ESI†), optical rotation results {observed $[\alpha]_D^{28} = -43.20$ (c 0.24, methanol); reported $[\alpha]_D = -45.76$, and HRMS, FT-IR and UV-visible spectra (see ESI†) of synthesized compound 2 were found to be in good agreement with reported data of the isolated thailandamide lactone, which unambiguously confirmed its first total synthesis.

Having thailandamide lactone in hand, we then screened its antibacterial activity against different non-pathogenic and pathogenic Gram-positive bacteria such as Bacillus subtilis (PY79), Bacillus megaterium (2G), Staphylococcus aureus as well as Gram-negative bacteria such as Vibrio cholerae (N16961), Enteropathogenic Escherichia coli (EPEC e2348/69), and Escherichia coli (MC1061)]. This screening revealed its moderate to potent antibacterial activity (Table 3). The efficacies of

Table 2	Optimization	of final	Heck	coupling
---------	--------------	----------	------	----------

Entry	[Pd] (mol%)	Condition	Yield (%)
1	PdCl ₂ (MeCN) ₂ (10)	Et ₃ N, HCO ₂ H, MeCN, rt, 3 h	Decomposition
2	$Pd(PPh_3)_4$ (5)	Et ₃ N, Bu ₄ NCl, DMF, rt, 6 h	Trace
3	$PdCl_2(PPh_3)_2$ (10)	K ₂ CO ₃ , Bu ₄ NCl, DMF, rt, 12 h	10
4	$Pd(Oac)_2$ (5)	Et ₃ N, Bu ₄ NCl, DMF, rt, 4 h	45'
5	$Pd(Oac)_2$ (5)	K ₃ PO ₄ , DMF, rt, 12 h	40
6	$[Pd(\mu-I)(P^tBu_3)]_2$ (7.5)	DIPEA, toluene, rt, 9 h	58
7	Pd(Oac) ₂ , (10) Pd(PPh ₃) ₄ (10)	K ₃ PO ₄ , DMF, rt, 18 h	77

Table 3 Antibacterial activities of thailandamide lactone

Staining type	Strains	MIC (μg ml ⁻¹)
Gram negative	V. cholerae (N16961) (pathogenic)	71.3
	EPEC (e2348/69) (pathogenic)	71.3
	E. coli (MC1061)	53.5
Gram positive	B. subtilis (PY79)	57.0
•	B. megaterium (2G)	53.5

thailandamide lactone even against Gram-negative strains were found to be promising.

Conclusions

Edge Article

In summary, we have developed a convergent route for the first total synthesis of a structurally challenging and labile polyketide natural product, namely thailandamide lactone, starting from the known compound 26 in 17 LLS with 8.5% overall yield. Our synthesis includes several coupling operations including two intermolecular Heck reactions. Notably, Pd(i)-based Heck coupling has been introduced for the first time, to the best of our knowledge, in the total synthesis of natural product. The possible site to couple the highly sensitive eastern and western polyene segments of thailandamide lactone was determined. The antibacterial activities of thailandamide lactone against different bacterial strains have been disclosed. Importantly, our modular strategy is expected to be amenable to thailandamide A, another member of this family as well as to structurally simplified designed analogues for further antibacterial study.

Data availability

Please see the ESI† for the data related to the manuscript.

Author contributions

R. K. G. conceived the idea, designed the hypothesis and managed overall manuscript preparation. R. R. P. planned and supervised the biological testing and assisted with manuscript preparation. H. S. and J. M. completed the total synthesis, A. K. G. was responsible for biological testing.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

H. S., J. M. and A. K. G. thank the Council of Scientific and Industrial Research (CSIR), New Delhi and the Indian Association for the Cultivation of Science, respectively, for their research fellowship. The financial support from the Science and Engineering Research Board (Project no-CRG/2019/001664) of India to carry out this work is gratefully acknowledged.

Notes and references

- 1 T. Nguyen, K. Ishida, H. Jenke-Kodama, E. Dittmann, C. Gurgui, T. Hochmuth, S. Taudien, M. Platzer, C. Hertweck and J. Piel, *Nat. Biotechnol.*, 2008, **26**, 225–233.
- 2 K. Ishida, T. Lincke, S. Behnken and C. Hertweck, *J. Am. Chem. Soc.*, 2010, **132**, 13966–13968.
- 3 K. Ishida, T. Lincke and C. Hertweck, *Angew. Chem., Int. Ed.*, 2012, **51**, 5470–5474.
- 4 Y. Wu and M. R. Seyedsayamdost, *Biochemistry*, 2018, 57, 4247–4251.
- 5 (a) S. Saha, D. Paul and R. K. Goswami, Chem. Sci., 2020, 11, 11259-11265; (b) T. K. Kuilya and R. K. Goswami, Org. Lett., 2017, 19, 2366-2369; (c) S. Das, D. Paul and R. K. Goswami, Org. Lett., 2016, 18, 1908-1911; (d) S. Guchhait, S. Chatterjee, R. S. Ampapathi and R. K. Goswami, J. Org. Chem., 2017, 82, 2414-2435; (e) D. Saha, G. H. Mandal and R. K. Goswami, J. Org. Chem., 2021, 86, 10006-10022.
- 6 A. Ruiz-Olalla, M. A. Wurdemann, M. J. Wanner, S. Ingemann, J. H. van Maarseveen and H. Hiemstra, *J. Org. Chem.*, 2015, 80, 5125–5132.
- 7 (a) M. T. Crimmins, B. W. King, E. A. Tabet and K. Chaudhary, *J. Org. Chem.*, 2001, 66, 894–902; (b) M. T. Crimmins, *Org. Lett.*, 2007, 9, 149–152.
- 8 O. Mitsunobu, Synthesis, 1981, 1, 1-28.
- 9 P. Liu and E. N. Jacobsen, J. Am. Chem. Soc., 2001, 123, 10772–10773.
- 10 A.-F. Salit, C. Meyer and J. Cossy, Synlett, 2007, 6, 934-938.
- 11 P. Li, J. Li, F. Arikan, W. Ahlbrecht, M. Dieckmann and D. Menche, *J. Am. Chem. Soc.*, 2009, **131**, 11678–11679.
- 12 C. L. Rand, D. E. Van Horn, M. W. Moore and E. Negishi, J. Org. Chem., 1981, 46, 4093–4096.
- 13 T. P. Ahern, H. O. Fong, R. F. Langler and P. M. Mason, *Can. J. Chem.*, 1980, 58, 878–883.
- 14 H. Yao and K. Yamamoto, *Chem.-Asian J.*, 2012, 7, 1542–1545.
- 15 T. J. Harrison, S. Ho and J. L. Leighton, *J. Am. Chem. Soc.*, 2011, **133**, 7308–7311.
- 16 (a) D. Saha, S. Guchhait and R. K. Goswami, Org. Lett., 2020,
 22, 745-749; (b) K. C. Nicolaou, P. G. Bulger and D. Sarlah,
 Angew. Chem., Int. Ed., 2005, 44, 4442-4489; (c)
 S. B. Garber, J. S. Kingsbury, B. L. Gray and A. H. Hoveyda,
 J. Am. Chem. Soc., 2000, 122, 8168-8179; (d) Y. Zhang,
 M. Dlugosch, M. Jubermann, M. G. Banwell and S. J. Ward,
 J. Org. Chem., 2015, 80, 4828-4833.
- 17 J. Ryan, M. Saiuciulis, A. Gomm, B. Maciá, E. O'Reilly and V. Caprio, J. Am. Chem. Soc., 2016, 138, 15798–15800.
- 18 S. Ma and E.-i. Negishi, J. Org. Chem., 1997, 62, 784-785.
- 19 J. L. Tyler, A. Noble and V. K. Aggarwal, *Angew. Chem., Int. Ed.*, 2021, **60**, 11824–11829.
- 20 J. A. González-Delgado, J. F. Arteaga, M. M. Herrador and A. F. Barrero, *Org. Biomol. Chem.*, 2013, 11, 5404–5408.
- 21 A. Cosp, P. Romea, P. Talavera, F. Urpí, J. Vilarrasa, M. Font-Bardia and X. Solans, *Org. Lett.*, 2001, 3, 615–617.
- 22 T. Katsuki and K. B. Sharpless, *J. Am. Chem. Soc.*, 1980, **102**, 5974–5976.

23 J. S. Clark, C. A. Baxter, A. G. Dossetter, S. Poigny, J. L. Castro and W. G. Whittingham, *J. Org. Chem.*, 2008, 73, 1040–1055.

Chemical Science

- 24 Y. L. Zhong and T. K. M. Shing, *J. Org. Chem.*, 1997, **62**, 2622–2624.
- 25 N. Cichowicz and P. Nagorny, *Org. Lett.*, 2012, **14**, 1058–1061. 26 R. F. Heck, *Org. React.*, 1982, 27, 345–390.
- 27 (a) I. P. Beletskaya and A. V. Cheprakov, Chem. Rev., 2000,
 100, 3009–3066; (b) A. de Meijere and F. E. Meyer, Angew. Chem., Int. Ed. Engl., 1994, 33, 2379–2411; (c) Q. Yao,
 E. P. Kinney and Z. Yang, J. Org. Chem., 2003, 68, 7528–7531.
- 28 (a) R. Vilar, D. M. P. Mingos and C. J. Cardin, J. Chem. Soc., Dalton Trans., 1996, 23, 4313–4314; (b) F. Schoenebeck,
 T. Sperger and C. Stirner, Synthesis, 2016, 49, 115–120; (c)
 M. Aufiero, T. Sperger, A. S. Tsang and F. Schoenebeck,
- Angew. Chem., Int. Ed., 2015, 54, 10322–10326; (d) F. Proutiere, M. Aufiero and F. Schoenebeck, J. Am. Chem. Soc., 2012, 134, 606–612; (e) G. Magnin, J. Clifton and F. Schoenebeck, Angew. Chem., Int. Ed., 2019, 58, 10179–10183; (f) C. Fricke, T. Sperger, M. Mendel and F. Schoenebeck, Angew. Chem., Int. Ed., 2021, 60, 3355–3366.
- 29 Compound 2 was found highly sensitive to light as well as quite unstable. It decomposed to a mixture of unidentified compounds. The colour changed from yellow to dark red. It was observed that use of Silica gel for purification and glass vessel for storing was not suitable as it underwent faster decomposition. Notably, the major part of compound stuck to reverse phase C18 column (Xbridge RP18) during purification.