
#### Volume 14 Number 31 21 August 2023 Pages 8225-8422

# Chemical Science



ISSN 2041-6539



#### **EDGE ARTICLE**

Bina Fu, Xueming Yang, Kaijun Yuan *et al.* Vacuum ultraviolet photodissociation of sulfur dioxide and its implications for oxygen production in the early Earth's atmosphere

### Chemical Science



#### **EDGE ARTICLE**

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2023, 14, 8255

dll publication charges for this article have been paid for by the Royal Society of Chemistry

Received 30th June 2023 Accepted 25th July 2023

DOI: 10.1039/d3sc03328g

rsc.li/chemical-science

## Vacuum ultraviolet photodissociation of sulfur dioxide and its implications for oxygen production in the early Earth's atmosphere†

Yao Chang, [b] ‡<sup>a</sup> Yanlin Fu,‡<sup>a</sup> Zhichao Chen,‡<sup>a</sup> Zijie Luo,<sup>ab</sup> Yarui Zhao,<sup>a</sup> Zhenxing Li,<sup>a</sup> Weiqing Zhang,<sup>a</sup> Guorong Wu, [b] <sup>a</sup> Bina Fu, [b] \*\*ade Dong H. Zhang, [b] <sup>aef</sup> Michael N. R. Ashfold, [b] <sup>c</sup> Xueming Yang [b] \*\*aef and Kaijun Yuan [b] \*\*ade

The emergence of molecular oxygen  $(O_2)$  in the Earth's primitive atmosphere is an issue of major interest. Although the biological processes leading to its accumulation in the Earth's atmosphere are well understood, its abiotic source is still not fully established. Here, we report a new direct dissociation channel yielding  $S(^1D) + O_2(a^1\Delta_g/X^3\Sigma_g^{-})$  products from vacuum ultraviolet (VUV) photodissociation of  $SO_2$  in the wavelength range between 120 and 160 nm. Experimental results show  $O_2$  production to be an important channel from  $SO_2$  VUV photodissociation, with a branching ratio of  $30 \pm 5\%$  at the H Lyman- $\alpha$  wavelength (121.6 nm). The relatively large amounts of  $SO_2$  emitted from volcanic eruptions in the Earth's late Archaean eon imply that VUV photodissociation of  $SO_2$  could have provided a crucial additional source term in the  $O_2$  budget in the Earth's primitive atmosphere. The results could also have implications for abiotic oxygen formation on other planets with atmospheres rich in volcanically outgassed  $SO_2$ .

#### Introduction

The provenance of oxygen  $(O_2)$  remains a crucial topic in the history of the Earth's evolution. The Earth's present atmosphere is notable for its remarkably high concentration of  $O_2$  ( $\sim$ 21% by volume), whereas geological and geochemical constraints suggest that free oxygen was anything but plentiful during the first half of the Earth's 4.5 billion year history. A permanent rise to appreciable concentrations of  $O_2$  in the atmosphere, known as the "Great Oxidation Event (GOE)", As is estimated to have occurred  $\sim$ 2.4 billion years ago (2.4 Ga). The GOE could have been a consequence of the emergence of oxygenic photosynthesis. Alternatively,  $O_2$  biogenesis may be much older; several pieces of evidence point to the first emergence of oxygenic photosynthesis long before the GOE (as early as  $\sim$ 3.8 Ga). At the emergence of the GOE could have been

a consequence of an abiotic shift in the balance of oxidants and reductants at the Earth's surface, *i.e.*, at early times, the biogenically produced  $\rm O_2$  was effectively consumed via reaction with reduced compounds, thereby suppressing  $\rm O_2$  levels ( $<10^{-6}$  of the present atmospheric level (PAL))<sup>15-17</sup> but, eventually, this source-sink balance shifted in favour of  $\rm O_2$  accumulation. Recent trace metal studies, *e.g.* of molybdenum and rhenium enrichment in the crust, <sup>10,18</sup> suggested "whiffs" of  $\rm O_2$  in the late Archaean (2.5–2.7 Ga), *i.e.*, intermittent periods before the GOE when the  $\rm O_2$ -sources overwhelmed the sinks. Such whiffs of oxygen have been suggested as a possible trigger for the GOE, but their duration, magnitude, and sources remain unknown.

Apart from biogenic processes leading to O<sub>2</sub> production, the widely accepted abiotic route to forming O2 is the three-body recombination reaction O + O + M → O<sub>2</sub> + M, involving O atoms produced by photolysis of CO2 or other oxygencontaining molecules. 19-22 Direct O2 production via vacuum ultraviolet (VUV) photodissociation of CO2 has also been identified, but deduced to be a very minor process compared with the indirect three-body recombination.<sup>23</sup> Dissociative electron attachment to CO2 has also been shown to lead to direct O2 production.24 Haqq-Misra et al.25 have suggested another abiotic O<sub>2</sub> production pathway, in which H<sub>2</sub>O<sub>2</sub> was first produced from the by-products of H<sub>2</sub>O photolysis, and then converted to O<sub>2</sub> via disproportionation reactions. Modelled O<sub>2</sub> concentrations up to 10<sup>-7</sup> PAL were reported. These findings have all informed knowledge of the history of the Earth's atmosphere and our understanding of planetary atmospheres and of interstellar photochemical processes.

<sup>&</sup>quot;State Key Laboratory of Molecular Reaction Dynamics, Dalian Coherent Light Source, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, 457 Zhongshan Road, Dalian, 116023, China. E-mail: kjyuan@dicp.ac.cn; bina@dicp.ac.cn; xmyang@dicp.ac.cn

<sup>&</sup>lt;sup>b</sup>Marine Engineering College, Dalian Maritime University, Liaoning, 116026, China <sup>c</sup>School of Chemistry, University of Bristol, Bristol, BS8 1TS, UK

<sup>&</sup>lt;sup>d</sup>University of Chinese Academy of Sciences, Beijing 100049, P. R. China

eHefei National Laboratory, Hefei 230088, China

Department of Chemistry, Center for Advanced Light Source Research, College of Science, Southern University of Science and Technology, Shenzhen 518055, China

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI https://doi.org/10.1039/d3sc03328g

<sup>‡</sup> These authors contributed equally.

Chemical Science Edge Article

SO<sub>2</sub> has also been considered as a possible source of oxygen. Astronomical observations have identified high SO<sub>2</sub> concentrations in the atmosphere of terrestrial exoplanets like Venus<sup>26,27</sup> and Io,<sup>28,29</sup> which are largely attributable to outgassing from volcanoes. In the Earth's late Archaean, prior to the GOE, subaerial volcanic degassing became important, yielding gases much richer in sulfur and dominated by SO<sub>2</sub>.<sup>30,31</sup> The photochemistry of volcanic SO<sub>2</sub> has been studied extensively in recent decades and has commonly been linked to the origin of the sulfur mass independent fractionation (S-MIF) in ancient rock samples.<sup>32–34</sup> SO<sub>2</sub> should be the major component from volcano eruptions, but the role of SO<sub>2</sub> photochemistry in the formation of molecular oxygen in the Earth's early atmosphere has hitherto been largely ignored.

SO<sub>2</sub> has two strong absorption bands in the UV region (see Section S2 and Fig. S2 in the ESI†). The more intense band, assigned to excitation from the ground  $\tilde{X}^1A_1(1^1A')$  in  $C_s$ ) state to the  $\tilde{C}^1B_2(2^1A')$  state, spans the wavelength range 185 <  $\lambda$  < 235 nm. <sup>35</sup> The other strong band at longer wavelengths,  $\lambda \sim 240$ – 350 nm, is associated with transition to the  $B^1B_1(2^1A'')$  state. The B<sup>1</sup>B<sub>1</sub>(2<sup>1</sup>A") state potential energy surface displays a conical intersection with that of the first excited singlet state (the  $A^1A_2(1^1A'')$  state). 33,36,37 Weak absorption at yet longer wavelengths is attributed to the spin-forbidden transition to the  $\tilde{a}^3B_1$  state. Photoexcitation of  $SO_2$  at wavelengths  $\lambda \leq 219$  nm results in predissociation, primarily to SO and O fragments, driven by nonadiabatic couplings to the lower-lying dissociative singlet and triplet states.<sup>38-41</sup> As Fig. S2† shows, the SO<sub>2</sub> absorption spectrum at shorter (VUV) wavelengths displays several intense diffuse absorption bands assigned to transitions to Rydberg states.

Early investigations of the VUV photolysis of SO<sub>2</sub> found indirect experimental evidence for the S + O2 channel through detection of OH fluorescence from SO<sub>2</sub>/H<sub>2</sub> mixtures.<sup>42</sup> More recently, Rosch et al.43 reported the first direct evidence for the  $S(^{3}P) + O_{2}$  channel from  $SO_{2}$  photolysis at 193 nm but, in the absence of quantitative measurements, the assessment of the importance (or otherwise) of O2 production from VUV photodissociation of SO<sub>2</sub> in the Earth's primitive atmosphere was not possible. The recent development of the intense VUV free electron laser (FEL) at the Dalian Coherent Light Source (DCLS) provides a unique tool for studying molecular photofragmentation dynamics across the entire VUV range. 22,44,45 Here, we present careful experimental studies of the  $S(^{1}D) + O_{2}$  product channel following SO2 photolysis at various wavelengths in the range  $120 < \lambda < 160$  nm using the VUV-pump and VUV-probe time-sliced velocity-map imaging (TS-VMI) technique, along with complementary electronic structure calculations. The quantitative assessment of this channel suggests that the VUV photochemistry of SO<sub>2</sub> could have been an important additional source of O<sub>2</sub> in the Earth's atmosphere in the late Archaean.

#### Results and discussion

#### The verification of the $S(^{1}D) + O_{2}$ product channel

In this study, the photodissociation dynamics of  $SO_2$  have been investigated using the recently constructed VUV pump and VUV probe TS-VMI apparatus, which is equipped with two

independently tunable VUV laser radiation sources (see Section S1 and Fig. S1 in the ESI†). A pulsed supersonic molecular beam generated from a gas mixture of about 1% SO $_2$  in Ar was irradiated with two counter-propagating VUV beams. The VUV FEL output was used to excite SO $_2$  molecules to different Rydberg states at wavelengths in the range  $120 < \lambda < 160$  nm. The S( $^1$ D) photofragments were then resonantly ionized with  $\lambda = 130.092$  nm photons, which were generated using a table-top VUV source and a difference frequency four-wave mixing (FWM) scheme, involving two 212.556 nm photons and one 580.654 nm photon that were overlapped in a Kr gas cell. Postionization, the S( $^1$ D) photoproducts were detected by the high resolution VMI detector.

Fig. 1 shows time-sliced ion images of the  $S(^1D)$  photofragments recorded following photolysis of  $SO_2$  at VUV wavelengths  $\lambda=121.6,\ 133.1,\ 140.0$  and 150.0 nm, respectively. Additional images, taken at  $\lambda=125.1,\ 130.1,\ 144.1$  and 154.1 nm, are displayed in Fig. S3 of the ESI.† The double headed arrow in Fig. 1 shows the direction of the polarization vector of the photolysis laser. Well-resolved, concentric rings with different intensities are clearly observable in the displayed images. These structures can be readily assigned to the population of different vibrational levels of the  $O_2$  co-product in its ground  $(X^3\Sigma_g^-)$  or first excited  $(a^1\Delta_g)$  electronic state arising via the photodissociation channel (1),

$$SO_2 + hv \rightarrow S(^1D) + O_2.$$
 (1)

The triple dissociation channel yielding  $S(^1D) + O + O$  products has a threshold energy of  $\sim 12.3 \text{ eV}^{46}$  and is thus not accessible in




Fig. 1 Time-sliced images of the S( $^1$ D) products from SO $_2$  photodissociation. The images were recorded at photolysis wavelengths  $\lambda=$  (A) 121.6 nm, (B) 133.1 nm, (C) 140.0 nm and (D) 150.0 nm. The double headed red arrow indicates the polarization direction of the photodissociation laser,  $\varepsilon$ . The concentric ring features reflect the population of different vibrational levels of the coincident  $O_2(X^3\Sigma_{\alpha}^{-}/a^1\Delta_{\alpha}, \nu)$  products.

the present experiment. The use of an off-axis biconvex LiF lens as the exit window for the four-wave mixing cell ensured that the 212.556 nm and 580.654 nm laser beams were dispersed from the photodissociation/photoionization region, thereby eliminating the possibility of unintended secondary dissociation of any primary SO fragments (from the rival SO + O dissociation channel) by absorption of another UV or IR photon. In addition, both the VUV FEL beam and the 130.092 nm probe beam were kept defocused to minimize any two-photon excitation effects. These steps ensured that no processes other than channel (1) yielded S(¹D) fragments under the prevailing experimental conditions.

The thermochemical threshold for process (1), i.e.  $D_0[SO_2(\tilde{X}, \nu)]$ = 0)  $\rightarrow$  S(<sup>1</sup>D) + O<sub>2</sub>(X,  $\nu$  = 0)], is  $\sim$ 7.1 eV (corresponding to an excitation wavelength,  $\lambda \sim 175$  nm).<sup>46</sup> Any energy provided by the VUV photon  $(E_{hv})$  in excess of this threshold energy will be deposited into the fragment kinetic energy and/or into the internal energy  $(E_{int})$  of the  $O_2$  products. The radii of the well resolved ring structures in the TS-VMI images can be used to determine the velocity distribution of the S(1D) products. These velocities can then be converted to a total kinetic energy release  $P(E_{\rm T})$  spectrum of the  $S(^{1}D) + O_{2}$  products using linear momentum conservation arguments. Fig. 2 and S4 (in the ESI†) display the  $P(E_{\rm T})$  spectra of the  $S(^{1}D) + O_{2}$  products obtained by integrating signals over all angles in the respective images. The internal energy distributions of the  $O_2$  co-products,  $E_{int}[O_2]$  formed at each wavelength can then be obtained from the corresponding  $P(E_T)$ spectrum using the law of energy conservation (eqn (2)).

$$E_{hv} - D_0 = E_{int}[O_2] + E_T[S(^1D) + O_2],$$
 (2)




Fig. 3 The relative population of the  $O_2(a^1\Delta_g)$  and  $O_2(X^3\Sigma_g^{-})$  products as a function of photolysis wavelength.

Each panel in Fig. 2 reveals two vibrational progressions for the  $O_2$  co-products, which can be assigned to the population of different vibrational levels of the ground  $(X^3\Sigma_g^-)$  and first excited electronic  $(a^1\Delta_g)$  state (the zero-point level of which lies 7882 cm<sup>-1</sup> above that of the ground state<sup>46</sup>). The onset of the strong progression in each  $P(E_T)$  spectrum accords well with the threshold of the  $S(^1D) + O_2(a^1\Delta_g)$  product channel, providing unambiguous evidence for the formation of molecular  $O_2$  in the VUV photodissociation of  $SO_2$ . Fig. 3 shows the relative populations of the  $O_2$  products formed in the two electronic states




Fig. 2 The product total kinetic energy ( $P(E_T)$ ) spectra for  $S(^1D) + O_2$  products following photolysis at  $\lambda = (A)$  121.6 nm, (B) 133.1 nm, (C) 140.0 nm and (D) 150.0 nm, derived from the images shown in Fig. 1, in red, along with the best-fit simulation of the spectra, in cyan dashed lines and orange dotted lines. The superposed combs indicate the  $E_T$  values associated with the formation of the various vibrational levels of  $O_2(X^3\Sigma_g^{-}/a^1\Delta_g, \nu)$ .

Chemical Science Edge Article

following  $SO_2$  photolysis at each wavelength studied, obtained from simulation of the  $P(E_T)$  spectra (Fig. 2 and  $S4\dagger$ ). Clearly, the  $S(^1D) + O_2(X^3\Sigma_g^-)$  channel is more important at the longest wavelengths investigated, but the  $S(^1D) + O_2(a^1\Delta_g)$  channel becomes increasingly dominant upon tuning to shorter wavelengths. The  $O_2(a^1\Delta_g)$  and  $O_2(X^3\Sigma_g^-)$  fragments formed at all but the very longest wavelengths both display inverted vibrational state population distributions, spanning a wide range of vibrational levels (Fig. S5 and S6 in the ESI $\dagger$ ). The finite bandwidth of the FEL source precludes detailed discussion of the rotational energy disposal in the  $O_2$  products from the observed vibrational peak profiles.

Potential energy surface (PES) calculations were also performed to gain insights into possible dissociation mechanisms for SO<sub>2</sub> molecules excited to low lying Rydberg states. Fig. 4 depicts two possible dissociation pathways. The left half of Fig. 4 illustrates a triplet-state pathway, in which SO<sub>2</sub> is initially excited to the 53A Rydberg state, and then undergoes nonadiabatic coupling to the 4<sup>3</sup>A" PES. The SO<sub>2</sub> molecule evolves on the 4<sup>3</sup>A" state PES, and one O-atom roams away from the SO partner and then returns to abstract the other O-atom. This mechanism involves an S-O-O intermediate and eventual dissociation to  $S(^{1}D) + O_{2}(X^{3}\Sigma_{g}^{-})$  products on the  $5^{3}A''$  PES. The right half of Fig. 4 illustrates a dissociation pathway via the singlet-state manifold. Initial photoexcitation in this case is to the 5<sup>1</sup>A" state, which is followed by non-adiabatic coupling to the  $3^1A''$  PES and dissociation to form  $S(^1D) + O_2(a^1\Delta_g)$  products. In this pathway, the interbond angle reduces, the two oxygen atoms approach towards a cyclic-SO2 intermediate and the O2 molecule is ejected. The identification of these pathways supports the experimental observation of  $O_2(X^3\Sigma_g^-)$  or  $O_2(a^1\Delta_g)$ products following VUV photodissociation of SO2. More details of the theoretical calculations are provided in Section S5 and Fig. S9 of the ESI.†

#### The quantum yield of $S(^{1}D) + O_{2}$ products

To quantify the importance of O<sub>2</sub> production from SO<sub>2</sub> photolysis in the Earth's primitive atmosphere, we have sought to determine the branching ratios of all active dissociation channels. Since the H Lyman-α wavelength (121.6 nm) is most abundant in stellar VUV radiation and about 80% of all SO2 photodissociation events occur around the Lyman-α line, 47 we have taken this as the representative wavelength to measure the branching ratios. Three fragment channels  $O(^{1}D) + SO$ ,  $O(^{1}S) +$ SO and  $S(^{1}D) + O_{2}$  formed by 121.6 nm photolysis have been detected using the VUV pump and VUV probe technique (Section S4 and Fig. S7 in the ESI†). Attempts to detect O(<sup>3</sup>P), S(<sup>3</sup>P) and S(<sup>1</sup>S) products under similar experimental conditions yielded no observable signals, suggesting that branching into each of the  $O(^3P)$  + SO,  $S(^3P)$  +  $O_2$  and  $S(^1S)$  +  $O_2$  channels is negligible at this wavelength. By calibrating the detection efficiencies for the O(<sup>1</sup>D), O(<sup>1</sup>S) and S(<sup>1</sup>D) fragments (Section S4 and Fig. S8 in the ESI $\dagger$ ), the branching ratio of the S( $^{1}$ D) + O<sub>2</sub> product channel was determined to be  $\sim 30 \pm 5\%$  at  $\lambda =$ 121.6 nm (as a fraction of all the detectable channels).

#### Insights into the O2 budget in the ancient atmosphere

The present experimental results imply that  $O_2$  production is a significant process following VUV excitation of  $SO_2$ . This finding could have profound implications for understanding the evolution of  $O_2$  in the Earth's primitive atmosphere. Current scenarios assume that biogenic  $O_2$  production started long before the GOE,  $^{10-14}$  but that any potential accumulation of  $O_2$  was offset by its consumption in reactions with reduced compounds emanating from the Earth's interior, *i.e.*  $O_2$  consumption balanced its production at these early times. This source-sink balance then shifted in favour of  $O_2$  accumulation, and finally triggered the GOE. This critical shift, relying on appreciable  $O_2$  production, is poorly understood. Geological




Fig. 4 Dissociation pathways of  $SO_2$  leading to  $O_2(X^3\Sigma_g^-) + S(^1D)$  products *via* the triplet state manifold (left) and to  $O_2(a^1\Delta_g) + S(^1D)$  products *via* the singlet state manifold. The schematics of the simplified PECs for the ground and the first five excited states of  $^3A''$  symmetry (left) and the first five states of  $^1A''$  symmetry (right) support different paths leading to  $S + O_2$  product formation. Dashed arrows indicate possible non-adiabatic pathways and the geometries and energies (in eV) of key intermediate structures along the dissociation pathways are also indicated. The horizontal axis represents the reaction coordinate, and the vertical axis is the potential energy (in eV).

studies suggest that the composition of the primitive atmosphere was probably determined by outgassing, through volcanism, and that the redox state of volcanic gases differs markedly between subaerial and submarine eruptions.  $^{30,31,48}$  Volcanic gases that erupted subaerially have generally equilibrated at high temperatures and low pressures with magmas close to the fayalite–magnetite–quartz buffer. Consequently, subaerial volcanism emitted  $SO_2$ -dominated gas into the primitive atmosphere. Volcanic  $SO_2$  gas could reach the stratosphere during eruptions, whereupon photodissociation would contribute an additional  $O_2$  source. Such a scenario would introduce additional  $O_2$  sources that, temporarily, would exceed the available sinks.

Here we attempt to estimate the total  $O_2$  production from the VUV photochemistry of volcanic  $SO_2$  in the late Archaean. We start by assuming that the average  $SO_2$  emission per year from volcanoes in the late Archaean was the same as the modern volcanic outgassing rate on Earth ( $\sim 2.3 \times 10^{13}$  gram per year (g per year)). When the volcanism subsided,  $SO_2$  was rapidly removed from the atmosphere by continued photolysis, gas-phase reactions and rain-out. Given that  $O_2$  production from  $SO_2$  photochemistry has here been shown to be a prominent pathway, we attempt to quantify the possible contribution of  $SO_2$  by assuming that VUV photodissociation coverts 1% of the total emitted  $SO_2$  directly into  $O_2$ . The accumulated amount of  $O_2$  in the atmosphere from VUV photodissociation of  $SO_2$  during the late Archaean eon ( $\sim 200$  million years) can then be estimated as follows,

$$W_{\rm O2} = 2.3 \times 10^{13} \text{ g per year} \times 200$$
  
  $\times 10^6 \text{ year} \times 1\% = 4.6 \times 10^{19} \text{ g}$  (3)

The total mass of  $O_2$  in the current Earth's atmosphere is  $\sim 1.07 \times 10^{21}$  g. Based on the above assumptions, the summed  $O_2$  from  $SO_2$  photolysis (eqn (3)) could be  $\sim 4.3\%$  of the present level of atmospheric  $O_2$  and this estimate should probably be viewed as a lower limit as it is likely that  $SO_2$  emissions and the UV flux were both higher in the late Archaean. Such arguments imply that  $O_2$  production from volcanic  $SO_2$  photochemistry could have provided substantial (and probably sporadic) contributions to the atmospheric  $O_2$  budget, given prevailing assumptions that most of the  $O_2$  sinks were already balanced by biogenic  $O_2$  production.  $^{10-14}$ 

We note that volcanic  $SO_2$  clouds and  $SO_2$ -based aerosols typically only survive in the atmosphere for a short time (several weeks to a few years) before sinking, and that most incident VUV excitation of  $SO_2$  would likely occur in the stratosphere during the volcanic eruptions. (The incipient  $O_3$  column in the late Archean is transparent to VUV radiation and the  $CO_2$  column is also transparent to VUV photons around 121.6 nm.<sup>50</sup>) Within the average lifetime of  $SO_2$  in the atmosphere, the transient accumulation of  $O_2$  from  $SO_2$  photolysis can be estimated as  $\sim 10^{18}$  to  $10^{19}$  molecules per cm<sup>2</sup> or  $\sim 10^{-5}$  to  $10^{-6}$  PAL. (This range was estimated assuming (i) a photon flux at  $\lambda \sim 121.6$  nm of  $\sim 2 \times 10^{12}$  photons per cm<sup>2</sup> per s,<sup>47</sup> (ii) that all VUV photons can be absorbed by  $SO_2$  clouds during the volcanic eruption and (iii) atmospheric lifetimes of  $SO_2$  between one month and one

year). Such events could have led to short-lived "oxygen oases", *i.e.*, localized or regional areas with significantly elevated O<sub>2</sub> during the volcanic eruption. The photochemical activity of SO<sub>2</sub> gas at 2.5–2.7 Ga has been linked with the abrupt rise of S-MIF signatures in sedimentary rocks around this time. <sup>1,51,52</sup> Spikes in Mo, Se, and Re concentrations at 2.5 to 2.66 Ga, and their isotopic excursions, have also recently been interpreted in terms of transient sources, or "whiffs", of O<sub>2</sub>. <sup>10,18,53,54</sup> The present findings are consistent with these scenarios. We propose that the VUV photochemistry of volcanic SO<sub>2</sub> can lead to efficient production of molecular O<sub>2</sub> and could have led to transient and localised accumulations of O<sub>2</sub> in the atmosphere before the GOE. Such transient elevations of O<sub>2</sub> from SO<sub>2</sub> photochemistry merit further consideration as a possible trigger for the GOE.

Furthermore, the elemental sulfur produced via channel (1) from  $SO_2$  photochemistry might polymerize into  $S_2$ ,  $S_3$ ,  $S_4$ , etc., and end up contributing to insoluble  $S_8$ -containing aerosols,  $^{55}$  the recycling of which could be responsible for the S-MIF signature in sedimentary rocks. If so, the tectonic reorganization, the abrupt rise of S-MIF signatures in sedimentary rocks and the "whiffs" of  $O_2$  in the late Archaean might be far from coincidental. Tectonics control volcanic eruptions, and the photochemistry of volcanic  $SO_2$  contributes to S-MIF and boosts the  $O_2$  budget.

 $O_2$  production from  $SO_2$  photochemistry could also be relevant in the contemporary atmospheres of other planets. For example, a layer of volcanic  $SO_2$  has been observed in the Venusian atmosphere, spanning heights from 48–65 km above the surface and with a measured maximal abundance of  $\sim 130$  ppm. <sup>56</sup> More recently,  $O_2$  molecules in the  $a^1\Delta_g$  state have also been detected (through the  $O_2(a^1\Delta_g)$  dayglow) in the Venusian atmosphere. <sup>57</sup> The present study provides unambiguous evidence for  $O_2(a^1\Delta_g)$  formation via  $SO_2$  photolysis at wavelengths around the H Lyman- $\alpha$  transition. Clearly, this is an  $O_2$  production mechanism that merits further scrutiny and, if necessary, incorporation into photochemical models for all planets with rich volcanically outgassed  $SO_2$ .

#### Data availability

The data supporting this study are available within the main text and the ESI.†

#### Author contributions

K. Y. conceived the research. K. Y. and X. Y. designed the experiments and supervised the research. Y. C., Z. C., Z. L., Y. Z. and Z. L. performed the experiments. W. Q. Z., G. R. W., and X. M. Y. operated the FEL facility. Y. F., B. F., and D. Z. performed the theoretical calculations. Y. C., B. F., K. Y., M. N. R. A. and X. Y. wrote the manuscript. All authors discussed the results and commented on the manuscript.

#### Conflicts of interest

There are no conflicts to declare.

#### Acknowledgements

The experimental work was supported by the National Natural Science Foundation of China (Grant No. 22241304, 22225303, and 22203093), the National Natural Science Foundation of China (NSFC Center for Chemical Dynamics (Grant No. 22288201)), the Scientific Instrument Developing Project of the Chinese Academy of Sciences (Grant No. GJJSTD20220001), the Innovation Program for Quantum Science and Technology (2021ZD0303304), and the Innovation Fund Project of Dalian Institute of Chemical Physics (DICP I202112). The theoretical work was supported by the National Natural Science Foundation of China (Grant No. 22173099) and the Innovation Program for Quantum Science and Technology (2021ZD0303305). X. Yang also thanks the Guangdong Science and Technology Program (Grant No. 2019ZT08L455 and 2019JC01X091) and the Shenzhen Science and Technology Program (Grant No. ZDSYS20200421111001787). Y. Chang thanks the Special Research Assistant Funding Project of Chinese Academy of Sciences and the China Postdoctoral Science Foundation (Grant No. 2021M693118).

#### References

- 1 J. Farquhar, H. Bao and M. Thiemens, *Science*, 2000, 289, 756–758.
- 2 H. D. Holland, *Geochim. Cosmochim. Acta*, 2009, **73**, 5241–5255.
- 3 D. E. Canfield, Annu. Rev. Earth Planet. Sci., 2005, 33, 1–36.
- 4 H. D. Holland, Geochim. Cosmochim. Acta, 2002, 66, 3811-3826.
- 5 H. D. Holland, Philos. Trans. R. Soc. London, Ser. B, 2006, 361, 903–915.
- 6 A. Bekker, H. D. Holland, P. L. Wang, D. Rumble 3rd, H. J. Stein, J. L. Hannah, L. L. Coetzee and N. J. Beukes, *Nature*, 2004, 427, 117–120.
- 7 G. Luo, S. Ono, N. J. Beukes, D. T. Wang, S. Xie and R. E. Summons, *Sci. Adv.*, 2016, 2, e1600134.
- 8 G. Luo, X. Zhu, S. Wang, S. Zhang and C. Jiao, *Sci. China: Earth Sci.*, 2022, **65**, 1646–1672.
- 9 J. J. Brocks, G. A. Logan, R. Buick and R. E. Summons, Science, 1999, 285, 1033–1036.
- 10 A. D. Anbar, Y. Duan, T. W. Lyons, G. L. Arnold, B. Kendall, R. A. Creaser, A. J. Kaufman, G. W. Gordon, C. Scott, J. Garvin and R. Buick, *Science*, 2007, 317, 1903–1906.
- 11 M. T. Rosing and R. Frei, Earth Planet. Sci. Lett., 2004, 217, 237–244.
- 12 K. L. French, C. Hallmann, J. M. Hope, R. Buick, J. J. Brocks and R. E. Summons, *Mineral. Mag.*, 2013, 77, 1110.
- 13 B. Rasmussen, I. R. Fletcher, J. J. Brocks and M. R. Kilburn, *Nature*, 2008, **455**, 1101–1104.
- 14 T. W. Lyons, C. T. Reinhard and N. J. Planavsky, *Nature*, 2014, **506**, 307–315.
- 15 A. A. Pavlov and J. F. Kasting, Astrobiology, 2002, 2, 27-41.
- 16 K. Zahnle, M. Claire and D. Catling, *Geobiology*, 2006, 4, 271–283.
- 17 D. C. Catling and K. J. Zahnle, Sci. Adv., 2020, 6, eaax1420.

- 18 J. Meixnerova, J. D. Blum, M. W. Johnson, E. E. Stueken, M. A. Kipp, A. D. Anbar and R. Buick, *Proc. Natl. Acad. Sci.* U. S. A., 2021, 118, e2107511118.
- 19 J. F. Kasting, S. C. Liu and T. M. Donahue, *J. Geophys. Res.: Oceans*, 1979, **84**, 3097–3107.
- 20 Y. Chang, Y. Yu, F. An, Z. Luo, D. Quan, X. Zhang, X. Hu, Q. Li, J. Yang, Z. Chen, L. Che, W. Zhang, G. Wu, D. Xie, M. N. R. Ashfold, K. Yuan and X. Yang, *Nat. Commun.*, 2021, 12, 2476.
- 21 J. F. Kasting and D. Catling, Annu. Rev. Astron. Astrophys., 2003, 41, 429-463.
- 22 Y. Chang, F. An, Z. Chen, Z. Luo, Y. Zhao, X. Hu, J. Yang, W. Zhang, G. Wu, D. Xie, K. Yuan and X. Yang, *Nat. Commun.*, 2021, 12, 6303.
- 23 Z. Lu, Y. C. Chang, Q. Z. Yin, C. Y. Ng and W. M. Jackson, *Science*, 2014, **346**, 61–64.
- 24 X. D. Wang, X. F. Gao, C. J. Xuan and S. X. Tian, *Nat. Chem.*, 2016, **8**, 258–263.
- 25 J. Haqq-Misra, J. F. Kasting and S. Lee, *Astrobiology*, 2011, **11**, 293–302.
- 26 X. Zhang, M.-C. Liang, F. Montmessin, J.-L. Bertaux, C. Parkinson and Y. L. Yung, *Nat. Geosci.*, 2010, 3, 834–837.
- 27 A. I. Stewart, D. E. Anderson, Jr., L. W. Esposito and C. A. Barth, *Science*, 1979, 203, 777–779.
- 28 C. T. Russell and M. G. Kivelson, *Science*, 2000, 287, 1998–1999.
- 29 R. Hu, S. Seager and W. Bains, Astrophys. J., 2013, 769, 6.
- 30 F. Gaillard, B. Scaillet and N. T. Arndt, *Nature*, 2011, 478, 229–232.
- 31 L. R. Kump and M. E. Barley, Nature, 2007, 448, 1033-1036.
- 32 S. O. Danielache, C. Eskebjerg, M. S. Johnson, Y. Ueno and N. Yoshida, *J. Geophys. Res.*, 2008, **113**, D17314.
- 33 C. Xie, X. Hu, L. Zhou, D. Xie and H. Guo, *J. Chem. Phys.*, 2013, **139**, 014305.
- 34 S. Ono, Annu. Rev. Earth Planet. Sci., 2017, 45, 301-329.
- 35 C. Leveque, A. Komainda, R. Taieb and H. Koppel, *J. Chem. Phys.*, 2013, 138, 044320.
- 36 A. R. Hoy and J. C. D. Brand, Mol. Phys., 1978, 36, 1409-1420.
- 37 J. Heicklen, N. Kelly and K. Partymiller, *Rev. Chem. Intermed.*, 1980, 3, 315–404.
- 38 H. Katagiri, T. Sako, A. Hishikawa, T. Yazaki, K. Onda, K. Yamanouchi and K. Yoshino, *J. Mol. Struct.*, 1997, 413–414, 589–614.
- 39 P. C. Ray, M. F. Arendt and L. J. Butler, J. Chem. Phys., 1998, 109, 5221–5230.
- 40 A. Okazaki, T. Ebata and N. Mikami, *J. Chem. Phys.*, 1997, 107, 8752–8758.
- 41 S. Becker, C. Braatz, J. Lindner and E. Tiemann, *Chem. Phys.*, 1995, **196**, 275–291.
- 42 C. Lalo and C. Vermeil, J. Photochem., 1975, 3, 441-454.
- 43 D. Rosch, R. Almeida, B. Sztaray and D. L. Osborn, *J. Phys. Chem. A*, 2022, **126**, 1761–1774.
- 44 Y. Chang, Y. Yu, H. Wang, X. Hu, Q. Li, J. Yang, S. Su, Z. He, Z. Chen, L. Che, X. Wang, W. Zhang, G. Wu, D. Xie, M. N. R. Ashfold, K. Yuan and X. Yang, *Nat. Commun.*, 2019, 10, 1250.

**Edge Article Chemical Science** 

- 45 J. Zhou, Y. Zhao, C. S. Hansen, J. Yang, Y. Chang, Y. Yu, G. Cheng, Z. Chen, Z. He, S. Yu, H. Ding, W. Zhang, G. Wu, D. Dai, C. M. Western, M. N. R. Ashfold, K. Yuan and X. Yang, Nat. Commun., 2020, 11, 1547.
- 46 On the basis of thermodynamic calculations with the data available from the thermochemical network (https:// atct.anl.gov) and references therein.
- 47 M. W. Claire, J. Sheets, M. Cohen, I. Ribas, V. S. Meadows and D. C. Catling, Astrophys. J., 2012, 757, 95.
- 48 Z.-X. Anser Li and C.-T. Aeolus Lee, Earth Planet. Sci. Lett., 2004, 228, 483-493.
- 49 S. A. Carn, V. E. Fioletov, C. A. McLinden, C. Li and N. A. Krotkov, Sci. Rep., 2017, 7, 44095.
- 50 D. L. Huestis and J. Berkowitz, Adv. Geosci., 2010, 25, 229-

- 51 J. Farquhar, M. Peters, D. T. Johnston, H. Strauss, A. Masterson, U. Wiechert and A. J. Kaufman, Nature, 2007, 449, 706-709.
- 52 D. E. Canfield, K. S. Habicht and B. Thamdrup, Science, 2000, 288, 658-661.
- 53 A. J. Kaufman, D. T. Johnston, J. Farguhar, A. L. Masterson, T. W. Lyons, S. Bates, A. D. Anbar, G. L. Arnold, J. Garvin and R. Buick, Science, 2007, 317, 1900-1903.
- 54 E. E. Stüeken, R. Buick and A. D. Anbar, Geology, 2015, 43, 259-262.
- 55 D. Babikov, Proc. Natl. Acad. Sci. U. S. A., 2017, 114, 3062-
- 56 E. Marcq, B. Bézard, P. Drossart, G. Piccioni, J. M. Reess and F. Henry, J. Geophys. Res., 2008, 113, E00B07.
- 57 J. C. Gérard, S. W. Bougher, M. A. López-Valverde, M. Pätzold, P. Drossart and G. Piccioni, Space Sci. Rev., 2017, 212, 1617-1683.