# Dalton Transactions



PAPER

View Article Online

View Journal | View Issue



**Cite this:** *Dalton Trans.*, 2025, **54**, 12463

# High-spin face-capped deltahedra in divanadadicarbaboranes are very different from the singlet structures of chromium analogues†

Horaţiu Casian, a Alexandru Lupan (10 \* a and R. Bruce King (10 \* b)

Using density functional theory (DFT), divanadadicarbaboranes  $Cp_2V_2C_2B_{n-4}H_{n-2}$  are found to have very different low-energy structures than the corresponding dichromadicarbaboranes  $Cp_2Cr_2C_2B_{n-4}H_{n-2}$ . Thus, the low-energy divanadadicarbaborane structures with n vertices have triplet or quintet states rather than singlet spin states, frequently based on an (n-1)-vertex  $VC_2B_{n-4}$  deltahedron having a face capped by the second vanadium atom bearing most of the spin density. Such structures are analogous to the low-energy structures of dimanganaboranes  $Cp_2Mn_2B_{n-2}H_{n-2}$ , even though the vanadium and manganese systems are not isoelectronic with each other. Most of the low-energy 8-vertex  $Cp_2V_2C_2B_4H_6$  structures are based on the hexagonal bipyramid, whereas most of the low-energy 9- and 10-vertex structures are based on the 9-vertex isocloso deltahedron. The bicapped square antiprism capped by a high-spin vanadium vertex is characteristic of the low-energy 11-vertex  $Cp_2V_2C_2B_7H_9$  structures. Similarly, an 11-vertex closo deltahedron capped by a high-spin vanadium vertex is the lowest energy closo closo deltahedron capped by a high-spin vanadium vertex is the lowest energy closo closo deltahedron capped by a substantial margin of closo clo

Received 27th June 2025, Accepted 16th July 2025 DOI: 10.1039/d5dt01520k

rsc li/dalton

#### 1. Introduction

The preferred structures of deltahedral boranes and isoelectronic carboranes are based on the most spherical closo deltahedra in which all of the faces are triangles and the vertices are as nearly similar as possible (Fig. 1).1-3 Each of the boron and carbon vertices in such structures uses one valence orbital for bonding to an external atom or group R such as hydrogen, a halogen, or a simple alkyl or aryl group. This leaves the remaining three vertex atom valence orbitals available to participate in skeletal bonding. Much of the early development of this chemistry was based on structures with an external hydrogen atom bonded to each boron or carbon vertex. In the electron bookkeeping for such systems, each BH vertex thus contributes two skeletal electrons and each CH vertex contributes three skeletal electrons. The preferred *n*-vertex deltahedral species were of the  $B_n H_n^{2-}$ ,  $CB_{n-1} H_n^{-}$ , and  $C_2 B_{n-2} H_n$  types with 2n + 2 skeletal electrons as discussed by Wade and Mingos<sup>4-6</sup> in early papers and

The research groups of Hawthorne<sup>9</sup> and Grimes<sup>10</sup> were the pioneers in synthesizing the first deltahedral boranes having transition metal vertices. Vertices of the CpM type (Cp = η-C<sub>5</sub>H<sub>5</sub>) were preferred owing to the robust nature of the cyclopentadienyl-metal bond. This allows a variety of chemical transformations to be performed on the metallaborane structure with the CpM vertices remaining intact. Cobalt vertices of the CpCo type were used in the early metallaborane and metallacarbaborane research. For electron bookkeeping purposes, a CpCo vertex is isoelectronic with a BH vertex as a contributor of two skeletal electrons. Thus, among the nine valence orbitals of the sp<sup>3</sup>d<sup>5</sup> manifold of the cobalt atom, three are used for external bonding to the Cp anion and three are occupied by lone pairs. This leaves three orbitals and two electrons available for skeletal bonding, corresponding to the favored 18-electron configuration for the cobalt atom. In the early work, Hawthorne's group<sup>9</sup> focused on metallaboranes obtainable using B<sub>10</sub>H<sub>14</sub> as the boron hydride source, whereas Grimes' group<sup>10</sup> focused on metallaboranes obtainable from B<sub>5</sub>H<sub>9</sub>. Because of the U.S. government's interest in the 1960s in the use of boron hydrides as possible rocket fuels, these key binary boron hydride starting materials were much more readily available then than they are now in modern times.

The pioneering work by Hawthorne and Grimes on metallaboranes was followed by work in the laboratory of Kennedy on


justified by chemical bonding models based on graph theory/topology<sup>7</sup> and tensor surface harmonics.<sup>8</sup>

<sup>&</sup>lt;sup>a</sup>Faculty of Chemistry and Chemical Engineering, Babeş-Bolyai University, Cluj-Napoca, Romania. E-mail: alexandru.lupan@ubbcluj.ro

<sup>&</sup>lt;sup>b</sup>Department of Chemistry and Center for Computational Quantum Chemistry, The University of Georgia, Athens, GA, USA. E-mail: rbking@uga.edu

<sup>†</sup> Electronic supplementary information (ESI) available: Initial structures, distance and energy ranking tables, orbital energies and HOMO/LUMO gaps, complete Gaussian09 reference (.pdf file), and the concatenated .xyz file containing the optimized structures that can be visualized using free software such as the Mercury program. See DOI: https://doi.org/10.1039/d5dt01520k

Paper






Fig. 1 The closo, isocloso, and oblatocloso deltahedra having 8 to 12 vertices. Vertices of degrees 4, 5, 6, and 7 are indicated in red, black, green, and blue, respectively.

systems using platinum group metals of the 4d and 5d transition series as vertex atoms. The Kennedy group identified alternative deltahedra known as isocloso<sup>14</sup> or hypercloso<sup>15–17</sup> deltahedra for the 9- and 10-vertex systems, particularly the latter (Fig. 1). 18-21 The isocloso 9- and 10-vertex deltahedra have a single degree 6 vertex, whereas the corresponding 9and 10-vertex closo deltahedra have exclusively degree 4 and 5 vertices (Fig. 1). The favored skeletal electron count for *n*-vertex iso*closo* metallaboranes was 2n, so they were hypoelectronic relative to the 2n + 2 skeletal electron count for the corresponding n-vertex closo metallaboranes. However, O'Neill and Wade<sup>22</sup> showed that the 8- and 9-vertex closo deltahedra have non-degenerate molecular orbitals in the frontier regions so that either 2n or 2n + 2 skeletal electrons could be favored for these systems. This is reflected experimentally in the longknown<sup>23,24</sup> deltahedral boron chlorides  $B_nCl_n$  (n = 8, 9) as well as the dirhodaboranes  $Cp_2^*Rh_2B_{n-2}H_{n-2}$  ( $Cp_3^* = \eta_1Me_5C_5$ ) synthesized by Ghosh and co-workers in recent years. 25,26

A seminal advance in the chemistry of metallaboranes was the discovery of dirhenaboranes  $Cp_2^*Re_2B_{n-2}H_{n-2}$  (n = 8, 9, 10, 11, 12) by Ghosh, Fehlner, and their coworkers. <sup>27</sup> X-ray crystallography showed that these dirhenaboranes have a non-spheri-

cal deltahedral structure approximating an oblate (flattened) ellipsoid in which one of the three primary axes is significantly shorter than the other two (Fig. 1). Because of this characteristic, deltahedra of this type can conveniently be designated as oblatocloso deltahedra.28 The only one of these five oblatocloso deltahedra that is clearly recognizable is the hexagonal bipyramid. In these oblatocloso deltahedra, the rhenium atoms are degree 6 or 7 vertices located at relatively flat points on the deltahedral surface, whereas the boron atoms are degree 4 or 5 vertices located at relatively high curvature points on the deltahedral surface. Electron bookkeeping using the Wade-Mingos assumption that the Cp\*Re vertices use three of the rhenium valence orbitals for skeletal bonding classifies them as more severely hypoelectronic systems with 2n - 4 Wadean skeletal electrons. This hypoelectronicity is illusory, however, since a more reasonable chemical bonding scheme considers the rhenium atoms at degree 6 and 7 vertex sites of low local curvature to use five rather than three valence orbitals for skeletal bonding, thereby making them effectively 2n + 4 skeletal electron systems.

Several years ago, we gratifyingly showed by density functional theory (DFT) calculations that these unusual experi-

Dalton Transactions Paper

mentally known oblato*closo*  $\operatorname{Cp^*}_2\operatorname{Re}_2B_{n-2}H_{n-2}$  (n=8,9,10,11,12) structures (Fig. 1) with the rhenium atoms in approximately antipodal positions are the lowest energy isomers. However, higher energy  $\operatorname{Cp^*}_2\operatorname{Re}_2B_{n-2}H_{n-2}$  isomers were found exhibiting a new structural paradigm, namely that of a *closo* deltahedron having adjacent rhenium atoms with a short rhenium–rhenium distance suggesting a multiple bond. Upon replacement of the two  $\operatorname{Cp^*Re}$  vertices with a single  $\operatorname{PnRe}_2$  unit ( $\operatorname{Pn} = \operatorname{pentalenyl} \{\eta^{5,5} - \operatorname{C}_8H_6\}$ ), thereby forcing the two rhenium atoms into adjacent vertices of the  $\operatorname{Re}_2B_{n-2}$  deltahedron, such *closo* deltahedral structures were found by DFT to be the lowest energy structures.

The synthetic methods used to prepare these dirhenaboranes of considerable structural interest not only involve the rare metal rhenium but also require several steps to prepare the Cp\*ReCl<sub>4</sub> starting material from typical rhenium sources such as perrhenate. Therefore, our subsequent theoretical studies were directed to explore the viability of structures similar to the dirhenaboranes but using the more abundant first-row transition metals. Our first work in this area studied dichromadicarbaboranes of the Cp<sub>2</sub>Cr<sub>2</sub>C<sub>2</sub>B<sub>n-4</sub>H<sub>n-2</sub> isoelectronic type with  $Cp_2^*Re_2B_{n-2}H_{n-2}$  systems.<sup>31</sup> In addition, Stone and co-workers<sup>32</sup> reported a species formulated as Cp2Cr2C2B8H10 shown by X-ray crystallography to have a central Cr<sub>2</sub>C<sub>2</sub>B<sub>8</sub> icosahedron with adjacent chromium vertices with a short chromium-chromium distance of 2.272 Å. Initially we thought that this short chromium-chromium distance might correspond to the formal quadruple bond required to provide enough skeletal electrons from the chromium atoms to give 26 (=2n + 2 for n = 12) skeletal electrons for a closo icosahedral system. However, our conclusion from this study based on the comparison of predicted chromium-chromium distances for various structures with the experimental chromium-chromium distance determined by X-ray crystallography suggested that the species reported by Stone and co-workers has a Cr=Cr triple bond bridged by two hydrogen atoms, not revealed in their X-ray crystallographic study dating back to the 1980s. In general, this study revealed low-energy oblatocloso structures  $Cp_2Cr_2C_2B_{n-4}H_{n-2}$  similar to those of their isoelectronic rhenium  $Cp_2^*Re_2B_{n-2}H_{n-2}$  analogues as well as closo and isocloso structures with short chromium-chromium distances, suggesting formal quadruple and triple bonds, respectively.

More recently, we used DFT to study  $Cp_2Mn_2B_{n-2}H_{n-2}$  systems that are the direct analogues of  $Cp_2Re_2B_{n-2}H_{n-2}$  systems for which the oblato*closo* structures are preferred.<sup>33</sup> However, low-energy oblato*closo* structures were not found for dimanganese systems. Instead, the energetically preferred structures for the  $Cp_2Mn_2B_{n-2}H_{n-2}$  systems were found to be higher spin state triplet and quintet structures, apparently a consequence of the lower ligand field strength in manganese complexes relative to analogous rhenium complexes. In general, the lowest energy  $Cp_2Mn_2B_{n-2}H_{n-2}$  structures were typically found to have a central  $MnB_{n-1}$  *closo* deltahedron with one face capped by the second CpMn unit.

Our studies on the  $Cp_2Cr_2C_2B_{n-4}H_{n-2}$  systems found a variety of interesting singlet oblatocloso, isocloso, and closo

structures as low-energy structures. We now report a DFT investigation of the corresponding divanadadicarbaboranes  $Cp_2V_2C_2B_{n-4}H_{n-2}$ . Unexpectedly, we found that the low-energy vanadadicarbaborane structures are of a totally different type from those of the dichromadicarbaboranes. Thus, the low-energy  $Cp_2V_2C_2B_{n-4}H_{n-2}$  structures are seen to resemble those of the dimanganaboranes  $Cp_2Mn_2B_{n-2}H_{n-2}$  in exhibiting higher spin state triplet and quintet structures with central  $VC_2B_{n-3}$  deltahedra capped by a high-spin CpV unit, even though the  $Cp_2V_2C_2B_{n-4}H_{n-2}$  and  $Cp_2Mn_2B_{n-2}H_{n-2}$  systems are not isoelectronic. Our results thus show that the two skeletal electron difference resulting in the substitution of vanadium for chromium in the dimetalladicarbaboranes  $Cp_2M_2C_2B_{n-4}H_{n-2}$  has a profound effect on the energetically preferred structure types.

#### 2. Theoretical methods

The initial  $Cp_2V_2C_2B_{n-4}H_{n-2}$  structures were constructed by systematic substitution of two boron vertices in  $B_nH_n^{2-}$  with two CpV units in various n-vertex polyhedral geometries, followed by further substitution of two of the remaining BH vertices by CH vertices. Thus, 337 structures of the 8-vertex  $Cp_2V_2C_2B_4H_6$ clusters, 347 structures of the 9-vertex structures of the 10-vertex  $Cp_2V_2C_2B_5H_7$ clusters, 538  $Cp_2V_2C_2B_6H_8$ clusters, 407 structures of the 11-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>7</sub>H<sub>9</sub> clusters, and 223 structures of the 12-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>8</sub>H<sub>10</sub> clusters were chosen as starting points for the optimizations (see the ESI†).

Full geometry optimizations were carried out on the  $Cp_2V_2C_2B_{n-4}H_{n-2}$  systems (n=8–12) at the B3LYP/6-31G(d) level of theory with all of them optimized, in turn, as neutral singlets, triplets, and quintets. For the singlet structures, the broken symmetry approach was also considered. The lowest-energy structures were then reoptimized at a higher level of theory, namely PBE0/Def2TZVP, and these are the structures presented in the manuscript.<sup>34</sup> The nature of the stationary points after the optimization was checked by calculations of harmonic vibrational frequencies. If significant imaginary frequencies were found, the optimization was continued by following the normal modes corresponding to the imaginary frequencies to ensure that genuine minima were obtained.

All calculations were performed using the Gaussian 09 package<sup>35</sup> with the default settings for the SCF cycles and geometry optimization. The Wiberg bond indices (WBIs) for the V–V interactions in the optimized  $Cp_2V_2C_2B_{n-4}H_{n-2}$  structures were obtained from the NBO analysis automatically provided in the Gaussian output.<sup>36</sup> All of the structures reported in this paper have appreciable HOMO–LUMO gaps of 2.59–4.40 eV (see Table S7 in the ESI†).

The  $Cp_2V_2C_2B_{n-4}H_{n-2}$  (n=8 to 12) structures are numbered as B(n-4)C2V2-xY where n is the total number of polyhedral vertices, x is the relative order of the structure on the energy scale (PBE0/Def2TZVP including zero-point corrections) and Y is the spin state designating singlets, triplets, and quintets as

Paper

B4C2V2-1T (C<sub>1</sub>)
0.0 kcal/mol

B4C2V2-2T (C<sub>s</sub>)
0.5 kcal/mol

3.1 kcal/mol

B4C2V2-4Q (C<sub>2h</sub>)

B4C2V2-5T (C<sub>2v</sub>)

B4C2V2-6T (C<sub>s</sub>)

Fig. 2 The lowest energy 8-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>4</sub>H<sub>6</sub> structures.

5.1 kcal/mol

S, T and Q. The lowest energy optimized structures discussed in this paper are depicted in Fig. 2 through 6. Only the lowest energy and thus potentially chemically significant structures are considered in detail in this paper. More comprehensive lists of structures, including higher energy structures, are given in the ESI.†

8.6 kcal/mol

11.1 kcal/mol

#### 3. Results and discussion

#### 3.1. Eight-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>4</sub>H<sub>6</sub> clusters

Six 8-vertex  $Cp_2V_2C_2B_4H_6$  structures were found within 11 kcal  $mol^{-1}$  of the lowest energy structure **B4C2V2-1T** (Fig. 2 and Table 1). The triplet structure **B4C2V2-1T** has a central  $VC_2B_4$  pentagonal bipyramid in which one face is capped by a CpV moiety with two hydrogen bridges to this vanadium atom from adjacent boron atoms. The spin density on the capping vanadium atom of 3.22 corresponds to the three unpaired electrons of high-spin vanadium(II) in a  $CpV^+$  moiety. This is analogous to the localization of five unpaired electrons in the capping  $CpMn^+$  moieties in many of the previously studied dimanganaboranes.<sup>33</sup> The opposing spin density of -1.17 on the other vanadium atom located at one of the apices of the

pentagonal bipyramid leads to a net vanadium spin of 2.05, consistent with the triplet spin state of B4C2V2-1T.

The other five low-energy Cp2V2C2B4H6 structures are all hexagonal bipyramids having the vanadium atoms in the axial positions with V-V distances of ~3 Å, suggesting some vanadium-vanadium interaction through the center of the hexagonal bipyramid (Fig. 2 and Table 1). These five structures can thus be considered as 8-vertex oblatocloso structures (Fig. 1). The B4C2V2-2T, B4C2V2-5T, and B4C2V2-6T structures lying 0.5, 8.6, and 11.1 kcal mol<sup>-1</sup>, respectively, above B4C2V2-1T, are triplet structures with the two carbon atoms located in meta (non-adjacent/non-antipodal), para (antipodal), and ortho (adjacent) positions in the equatorial hexagon, respectively. The quintet Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>4</sub>H<sub>6</sub> structures, **B4C2V2-3Q** and **B4C2V2-4Q**, lying 3.1 and 5.1 kcal mol<sup>-1</sup>, respectively, in energy above B4C2V2-1T, are quite similar to the carbon atoms located in the meta and para positions in the equatorial hexagon, respectively.

#### 3.2. Nine-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>5</sub>H<sub>7</sub> clusters

The potential energy surface for the 9-vertex  $Cp_2V_2C_2B_5H_7$  is more complicated than that of  $Cp_2V_2C_2B_4H_6$  with 10 structures lying within 11 kcal  $mol^{-1}$  of the lowest energy structure B5C2V2-1Q (Fig. 3 and Table 2). This quintet structure B5C2V2-1Q has a central  $VC_2B_4$  pentagonal bipyramid with the fifth boron atom capping a  $VB_2$  face. The remaining CpV moiety then caps the  $B_3$  face including this capping boron atom with B-H-V bridges from all three boron atoms in the capped face. This degree 3 capping vanadium atom has a spin density of 3.11 corresponding to a high-spin vanadium( $\pi$ )  $CpV^{\dagger}$  similar to that in the  $Cp_2V_2C_2B_4H_6$  structure B4C2V2-1T (Fig. 2 and Table 1).

Three of the 10 lowest energy  $Cp_2V_2C_2B_5H_7$  structures, namely the quintets **B5C2V2-2Q**, **B5C2V2-4Q**, and **B5C2V2-5Q** lying 0.9, 3.3, and 6.4 kcal  $mol^{-1}$  above **B5C2V2-1Q**, have central  $V_2C_2B_5$  9-vertex iso*closo* deltahedra (Fig. 3 and Table 2). The somewhat higher energy triplet 9-vertex iso*closo* deltahedral structures **B5C2V2-7T**, **B5C2V2-9T**, and **B5C2V2-10T** lie at 7.1, 9.8, and 10.3 kcal  $mol^{-1}$ , respectively, above **B5C2V2-1Q**. All six of these  $Cp_2V_2C_2B_5H_7$  structures have one of the vanadium atoms located at the unique degree 6 vertex of the 9-vertex iso*closo* deltahedron. Except for **B5C2V2-4Q**, the second vanadium vertex is not adjacent to the degree 6 vertices, leading to long V···V distances of ~3.7 Å with WBIs of

 Table 1
 The lowest energy  $Cp_2V_2C_2B_4H_6$  structures up to 11.5 kcal mol<sup>-1</sup>. Energies are given in kcal mol<sup>-1</sup> and distances are given in Å

Structure (symmetry)	$\Delta E$	Vanadium vertices				Net V	$V \cdots V$	$v \cdot \cdot \cdot v$	
		Degree	Spin	Degree	Spin	Spin	Distance	WBI	Polyhedron
B4C2V2-1T (C <sub>1</sub> )	0.0	3(2µ-H)	3.22	6	-1.17	2.05	3.24	0.72	Cap pent bipy
B4C2V2-2T $(C_s)$	0.5	6	2.91	6	-0.85	2.06	2.95	0.53	Hex bipy
$B4C2V2-3Q(C_s)$	3.1	6	2.45	6	2.42	4.87	3.04	0.25	Hex bipy
$B4C2V2-4Q(C_{2h})$	5.1	6	2.33	6	2.33	4.66	3.07	0.22	Hex bipy
B4C2V2-5T $(C_{2y})$	8.6	6	2.07	6	0.54	2.61	2.86	0.39	Hex bipy
<b>B4C2V2-6T</b> $(C_s)$	11.1	6	2.65	6	-0.24	2.41	3.26	0.40	Hex bipy

B5C2V2-1Q (C<sub>s</sub>) B5C2V2-2Q (C<sub>1</sub>) B5C2V2-3Q (C<sub>1</sub>) 2.7 kcal/mol 2.7 kcal/mol 3.3 kcal/mol 6.4 kcal/mol




Fig. 3 The lowest energy 9-vertex  $Cp_2V_2C_2B_5H_7$  structures.

B5C2V2-9T (C1)

9.8 kcal/mol

 $\sim$ 0.13 for the quintet structures and  $\sim$ 3.5 Å with somewhat higher WBIs for the triplet structures. The V–V distance in **B5C2V2-4Q** with adjacent vanadium atoms at a distance of 2.78 Å corresponds to a significantly higher WBI of 0.25.

B5C2V2-10T (C1)

10.3 kcal/mol

The three remaining  $Cp_2V_2C_2B_5H_7$  structures have other central polyhedra (Fig. 3 and Table 2). The quintet structure B5C2V2-3Q, lying 2.7 kcal  $mol^{-1}$  above B5C2V2-1Q, has a central  $VC_2B_5$  bisdisphenoid (the 8-vertex *closo* deltahedron—Fig. 1) with one of its faces capped by a high-spin  $CpV^+$  moiety with a spin density of 3.16 and one B-H-V bridge. The quintet structure B5C2V2-6Q, lying 6.8 kcal  $mol^{-1}$  above B5C2V2-1Q, has a tricapped trigonal prismatic structure (the 9-vertex *closo* deltahedron—Fig. 1) with one vanadium atom at a degree 5 vertex and the other vanadium atom at a degree 4 vertex. The degree 4 vanadium vertex in B5C2V2-6Q forms two V-H-B bridges with adjacent boron atoms. Finally, the  $V_2C_2B_5$  polyhedron in B5C2V2-8Q, lying 9.5 kcal  $mol^{-1}$  above B5C2V2-1Q in energy, is also a tricapped trigonal prism with the two carbon atoms and a boron atom as degree 4 vertices capping the underlying  $V_2B_4$  trigonal prism.

#### 3.3. Ten-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>6</sub>H<sub>8</sub> clusters

Five 10-vertex  $Cp_2V_2C_2B_6H_8$  structures were found within 13 kcal mol<sup>-1</sup> of the lowest energy structure **B6C2V2-1T** (Fig. 4 and Table 3). The central polyhedron in the triplet structure **B6C2V2-1T** is a 10-vertex iso*closo* polyhedron with one of the




Fig. 4 The lowest energy 10-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>6</sub>H<sub>8</sub> structures.

 Table 2
 The lowest energy  $Cp_2V_2C_2B_5H_7$  structures up to 11 kcal mol<sup>-1</sup>. Energies are given in kcal mol<sup>-1</sup> and distances are given in Å

	$\Delta E$	Vanadium vertices				Net V	$V \cdots V$	$v \cdots v$	
Structure (symmetry)		Degree	Spin	Degree	Spin	Spin	Distance	WBI	Polyhedron
B5C2V2-1Q (C <sub>s</sub> )	0.0	3(3µ-H)	3.11	6	1.69	4.80	3.99	0.03	Central pent bipy
$B5C2V2-2Q(C_1)$	0.9	6	2.28	5	2.32	4.60	3.69	0.14	9-Vertex isocloso
<b>B5C2V2-3Q</b> $(C_1)$	2.7	$3(1\mu-H)$	3.16	6	1.29	4.45	3.07	0.21	Cap bisdisphen
<b>B5C2V2-4Q</b> $(C_1)$	3.3	4(1μ-H)	2.50	6	2.24	4.74	2.78	0.25	9-Vertex iso <i>closo</i>
<b>B5C2V2-5Q</b> $(C_1)$	6.4	5(1μ-H)	2.37	6	2.29	4.66	3.67	0.12	9-Vertex isocloso
$B5C2V2-6Q(C_s)$	6.8	4(2μ-H)	2.62	5	2.50	5.12	3.91	0.06	Tricap trig prism
<b>B5C2V2-7T</b> $(C_1)$	7.1	5(1μ-H)	2.37	6	-0.08	2.29	3.50	0.18	9-Vertex iso <i>closo</i>
<b>B5C2V2-8Q</b> $(C_2)$	9.5	5	2.36	5	2.35	4.70	3.84	0.10	Tricap trig prism
B5C2V2-9T $(C_1)$	9.8	5	2.40	6	-0.09	2.31	3.54	0.15	9-Vertex iso <i>closo</i>
<b>B5C2V2-10T</b> $(C_1)$	10.3	5	2.40	6	-0.07	2.33	3.44	0.19	9-Vertex iso <i>closo</i>

**Table 3** The lowest energy  $Cp_2V_2C_2B_6H_8$  structures up to 13 kcal mol<sup>-1</sup>. Energies are given in kcal mol<sup>-1</sup> and distances are given in Å

Structure (symmetry)	$\Delta E$	Vanadium vertices				Net V	$V \cdots V$	$V \cdots V$	
		Degree	Spin	Degree	Spin	Spin	Distance	WBI	Polyhedron
B6C2V2-1T (C <sub>s</sub> )	0.0	4	3.09	6	-1.23	1.86	2.53	0.56	10v-isocloso
<b>B6C2V2-2Q</b> $(C_1)$	0.8	3(1µ-H)	3.15	7	1.32	4.47	3.31	0.10	Cap 9v isocloso
B6C2V2-3T $(C_s)$	1.1	3(3µ-H)	3.17	7	-1.25	1.92	3.36	0.11	Cap 9v iso <i>closo</i>
$B6C2V2-4Q(C_s)$	2.4	3(1μ-H)	3.17	7	1.29	4.46	4.01	0.18	Cap 9v iso <i>closo</i>
$\mathbf{B6C2V2-5Q} (C_{\mathbf{s}})$	3.3	3(3μ-H)	3.15	6	1.32	4.47	4.64	0.02	Cap 9v iso <i>closo</i>

vanadium atoms at the degree 6 vertex. The short V=V distance of 2.53 Å with a relatively high WBI of 0.56 in B6C2V2-1T suggests some type of multiple bond. The second vanadium atom in B6C2V2-1T is a degree 4 vertex with a high spin density of 3.09 corresponding to a high-spin vanadium( $\pi$ ) CpV<sup>+</sup> unit similar to that found in a number of the other low-energy Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>n-4</sub>H<sub>n-2</sub> structures.

The remaining four low-energy  $Cp_2V_2C_2B_6H_8$  structures, namely **B6C2V2-2Q**, **B6C2V2-3T**, **B6C2V2-4Q**, and **B6C2V2-5Q**, are closely spaced in energy, lying 0.8, 1.1, 2.4, and 3.3 kcal mol<sup>-1</sup>, respectively, above **B6C2V2-1T**. All four structures have a central 9-vertex iso*closo*  $VC_2B_6$  deltahedron (Fig. 1), in which one of the faces is capped by a high-spin vanadium(II) atom in a  $CpV^+$  unit with a spin density of ~3.15 (Fig. 4 and Table 3). These four nearly isoenergetic structures differ in their spin state and in which of the faces of the 9-vertex iso*closo* deltahedron is capped by the second vanadium atom.

#### 3.4. Eleven-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>7</sub>H<sub>9</sub> clusters

The relevant potential energy surfaces of the 11-vertex  $Cp_2V_2C_2B_7H_9$  system (Fig. 5 and Table 4) and the 12-vertex  $Cp_2V_2C_2B_8H_{10}$  system are simple compared with the systems with fewer vertices discussed above. Thus the 11-vertex  $Cp_2V_2C_2B_7H_9$  system has three structures within 5 kcal  $mol^{-1}$  of the lowest energy structure **B7C2V2-1Q**. However, any other




Fig. 5 The lowest energy 11-vertex  $Cp_2V_2C_2B_7H_9$  structures.

isomeric structures lie at least 18 kcal  $\mathrm{mol}^{-1}$  above B7C2V2-1Q (Fig. 5). All three of these structures have a central  $VC_2B_7$  bicapped square antiprism (the 10-vertex *closo* deltahedron—Fig. 1), in which one of the faces is capped by a high-spin  $\mathrm{CpV}^+$  unit with a spin density of  $\sim 3.15$ . The capping vanadium atoms form one to three V-H-B bridges to adjacent boron atoms (Table 4). In all three  $\mathrm{Cp}_2V_2C_2B_7H_9$  structures, the capped face includes one of the degree 4 vertices of the underlying deltahedron.

#### 3.5. Twelve-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>8</sub>H<sub>10</sub> clusters

The potential energy surface of the 12-vertex  $Cp_2V_2C_2B_8H_{10}$  system is even simpler since the lowest energy structure **B8C2V2-1Q** (Fig. 6) lies ~18 kcal mol<sup>-1</sup> below the next lowest energy structure, which has a similar geometry but with a triplet rather than quintet spin state. Structure **B8C2V2-1Q** is a capped 11-vertex *closo* deltahedron with one of the vanadium atoms at a degree 6 vertex. The other vanadium atom is the degree 3 capping atom with a spin density of 3.18 corresponding to a high-spin  $CpV^+$  unit. This capping vanadium atom forms three V-H-B bridges with adjacent boron atoms.



Fig. 6 The lowest energy  $Cp_2V_2C_2B_8H_{10}$  structure B8C2V2-1Q lying  $\sim \!\! 18$  kcal mol $^{-1}$  in energy below the next lowest energy structure.

Table 4 The lowest energy  $Cp_2V_2C_2B_7H_9$  structures up to 18 kcal mol<sup>-1</sup>. Energies are given in kcal mol<sup>-1</sup> and distances are given in Å

		Vanadium vertices				Net V	V···V	VV	
Structure (symmetry)	$\Delta E$	Degree	Spin	Degree	Spin	Spin	Distance	WBI	Polyhedron
$\overline{\mathbf{B7C2V2-1Q}\left(C_{\mathrm{s}}\right)}$	0.0	3(2µ-H)	3.15	6	1.55	4.70	4.19	0.02	Cap 10v closo
B7C2V2-2Q $(C_1)$	4.0	3(1μ-H)	3.14	6	1.55	4.69	3.32	0.10	Cap 10v closo
B7C2V2-3T $(C_s)$	4.6	3(3µ-H)	3.17	6	-1.35	1.82	3.35	0.11	Cap 10v closo

**Dalton Transactions** Paper

#### Conclusion

In a previous study,<sup>31</sup> the low-energy structures of the dichromadicarbaboranes  $Cp_2Cr_2C_2B_{n-4}H_{n-2}$  (n = 8 to 12) were found to be singlet spin state structures with flattened oblatocloso  $Cr_2C_2B_{n-4}$  deltahedra having degree 6 and 7 chromium vertices similar to the lowest energy<sup>29</sup> and experimentally known<sup>27</sup> isoelectronic dirhenaboranes the  $Cp_2Re_2B_{n-4}H_{n-2}$ . In addition, singlet spherical closo deltahedral structures having adjacent chromium vertices with surface chromium-chromium triple or quadruple bonds are found for the dichromadicarbaboranes at accessible energies.

Simple substitution of two vanadium atoms for the two chromium atoms in the dichromadicarbaboranes to give the divanadadicarbaboranes  $Cp_2V_2C_2B_{n-4}H_{n-2}$  (n = 8 to 12) leads to very different potential energy surfaces with all of the lowest energy structures being higher spin state triplet and quintet structures rather than singlet structures. A conspicuous feature in many of the low-energy structures is a degree 3 high-spin vanadium(II)  $CpV^{+}$  moiety capping a face of an (n-1)-vertex deltahedron with one to three B-H-V bridges to this vanadium atom. In this respect, the divanadadicarbaboranes resemble the dimanganaboranes<sup>33</sup>  $Cp_2Mn_2B_{n-2}H_{n-2}$ , even though the systems are not isoelectronic. Thus the low-energy dimanganaborane structures are all higher spin triplet and quintet spin state structures with the frequent feature of a similar high-spin  $CpMn^{+}$  moiety capping a face of an (n-1)-vertex deltahedron with one to three B-H-Mn bridges to the capping manganese

Most of the low-energy 8-vertex divanadadicarbaboranes Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>4</sub>H<sub>6</sub> exhibit oblatocloso structures having a V<sub>2</sub>C<sub>2</sub>B<sub>4</sub> hexagonal bipyramid with the vanadium atoms in antipodal positions at the low-curvature degree 6 vertices. The 9-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>5</sub>H<sub>7</sub> potential energy surface is rather complicated but the 9-vertex isocloso deltahedron (Fig. 1) is a key feature of many of the low-energy structures. Most of the low-energy structures for the 10-vertex Cp2V2C2B6H8 system are also based on a central 9-vertex VC<sub>2</sub>B<sub>6</sub> isocloso deltahedron but with one of the faces capped by the second vanadium atom in a high-spin vanadium(II) CpV<sup>+</sup> moiety with one to three B-H-V hydrogen bridges to this vanadium atom. The three lowest-energy structures of the 11-vertex Cp<sub>2</sub>V<sub>2</sub>C<sub>2</sub>B<sub>7</sub>H<sub>9</sub> system by a wide margin all have a central 10-vertex VC<sub>2</sub>B<sub>7</sub> closo deltahedron, namely the bicapped square antiprism (Fig. 1), in which one of the faces is capped by the second vanadium atom, likewise in a high-spin CpV<sup>+</sup> moiety bridged by one to three hydrogen atoms to the adjacent boron atoms. The lowest energy Cp2V2C2B8H10 structure by a substantial margin of ~18 kcal mol<sup>-1</sup> has a central 11-vertex closo deltahedron capped by a high-spin vanadium vertex with three V-H-B bridges.

#### Conflicts of interest

The authors declare no competing financial interest.

### Data availability

The data are available from the authors upon request.

## Acknowledgements

The computational facilities were provided by the Babes-Bolyai University under project POC/398/1/1/124155 - co-financed by the European Regional Development Fund (ERDF) through the Competitiveness Operational Program for Romania 2014-2020.

#### References

- 1 A. S. Weller, Carboranes including their metal complexes, in Organometallic Chemistry, ed. M. Green, RSC Publishing, 2002, vol. 30.
- 2 Boron Science New Technologies and Applications, ed. N. S. Hosmane, Taylor and Francis, 2011.
- 3 Handbook of Boron Science With Applications Organometallics, Catalysis, Materials and Medicine, ed. N. S. Hosmane and R. Eagling, World Scientific, 2018.
- 4 K. Wade, The structural significance of the number of skeletal bonding electron-pairs in carboranes, the higher boranes and borane anions, and various transition-metal carbonyl cluster compounds, J. Chem. Soc. D: Chem. Commun., 1971, 792-793.
- 5 D. M. P. Mingos, A general theory for cluster and ring compounds of the main group and transition elements, Nature Phys. Sci., 1972, 236, 99-102.
- 6 D. M. P. Mingos, Polyhedral skeletal electron pair approach, Acc. Chem. Res., 1984, 17, 311-319.
- 7 R. B. King and D. H. Rouvray, A graph-theoretical interpretation of the bonding topology in polyhedral boranes, carboranes, and metal clusters, J. Am. Chem. Soc., 1977, 99, 7834-7840.
- 8 A. J. Stone and M. J. Alderson, A new model of structure and bonding in the boron hydrides, Inorg. Chem., 1982, 21, 2297-2302.
- 9 K. P. Callahan and M. F. Hawthorne, Ten years of metallocarboranes, Adv. Organomet. Chem., 1976, 14, 145-186.
- 10 R. N. Grimes, The role of metals in borane clusters, Acc. Chem. Res., 1983, 16, 22-26.
- 11 S. C. Rasmussen, The 18-electron rule and electron counting in transition metal compounds: theory and application, ChemTexts, 2015, 1, 10.
- 12 C. A. Tolman, The 16 and 18 electron rule in organometallic chemistry and homogeneous catalysis, Chem. Soc. Rev., 1972, 1, 337-353.
- 13 D. P. Craig and G. Doggett, Theoretical basis of the "raregas rule", J. Chem. Soc., 1963, 4189.
- 14 J. D. Kennedy, Structure and bonding in recently isolated metallaboranes, Inorg. Chem., 1986, 25, 111-112.
- 15 R. T. Baker, Hyper-closo metallaboranes, Inorg. Chem., 1986, 25, 109-111.

Paper

16 R. L. Johnston and D. M. P. Mingos, Molecular orbital calculations relevant to the hypercloso vs. iso-closo controversy in metallaboranes, *Inorg. Chem.*, 1986, **25**, 3321–3323.

- 17 R. L. Johnston, D. M. P. Mingos and P. Sherwood, Bonding and electron counting in hyper *closo* metalloboranes and metallocarboranes, *New J. Chem.*, 1991, **15**, 831–841.
- 18 J. Bould, J. D. Kennedy and M. Thornton-Pett, Ten-vertex metallaborane chemistry. Aspects of the iridadecaborane *closo*→isonido→iso*closo* structural continuum, *J. Chem. Soc., Dalton Trans.*, 1992, 563–576.
- 19 J. D. Kennedy and B. Štibr, in *Current Topics in the Chemistry of Boron*, ed. G. W. Kabalka, Royal Society of Chemistry, Cambridge, 1994, pp. 285–292.
- 20 J. D. Kennedy, in *The Borane-Carborane-Carbocation Continuum*, ed. J. Casanova, Wiley, New York, 1998, ch. 3, pp. 85–116.
- 21 B. Štibr, J. D. Kennedy, E. Drdáková and M. Thornton-Pett, Nine-vertex polyhedral iridamonocarbaborane chemistry. Products of thermolysis of [(CO)(PPh<sub>3</sub>)<sub>2</sub>IrCB<sub>7</sub>H<sub>8</sub>] and emerging alternative cluster-geometry patterns, *J. Chem. Soc., Dalton Trans.*, 1994, 229–236.
- 22 M. E. O'Neill and K. Wade, *Closo* clusters with unusual electron numbers: The significance of frontier orbital degeneracies and bonding characteristics, *J. Mol. Struct.:* THEOCHEM, 1983, 103, 259–268.
- 23 J. A. Morrison, Chemistry of the polyhedral boron halides and the diboron tetrahalides, *Chem. Rev.*, 1991, **91**, 35–48.
- 24 C. Corminboeuf, M. W. Wodrich, R. B. King and P. v. R. Schleyer, Effects of halogen substitution on the properties of eight- and nine-vertex *closo*-boranes, *Dalton Trans.*, 2008, 1745–1751.
- 25 R. Borthakur, B. Mondal, P. Nandi and S. Ghosh, Hypoelectronic isomeric diiridaboranes [ $(Cp*Ir)_2B_6H_6$ ]: the "Rule-Breakers"( $Cp*=\eta 5\text{-}C_5Me_5$ ), *Chem. Commun.*, 2016, 52, 3199–3202.
- 26 D. K. Roy, R. Borthakur, R. Prakash, S. Bhattacharya, R. Jagan and S. Ghosh, Hypoelectronic 8–11-Vertex iridaand rhodaboranes, *Inorg. Chem.*, 2016, 55, 4764–4770.
- 27 For a review of much of the relevant chemistry from Fehlner's group, see: T. P. Fehlner, in *Group 13 Chemistry:* From Fundamentals to Applications, ed. P. J. Shapiro and

- D. A. Atwood, American Chemical Society, Washington, D. C., 2002, pp. 49–67.
- 28 R. B. King, The oblate deltahedra in dimetallaboranes: geometry and chemical bonding, *Inorg. Chem.*, 2006, 45, 8211–8216.
- 29 A. Lupan and R. B. King, Hypoelectronic dirhenaboranes having eight to twelve vertices: internal versus surface rhenium-rhenium bonding, *Inorg. Chem.*, 2012, 51, 7609– 7616.
- 30 A. Lupan and R. B. King, Dimetallaboranes with polyhedral surface metal-metal multiple bonds: deltahedral dirhenaboranes with pentalenedirhenium vertices, *Organometallics*, 2013, 32, 4002–4008.
- 31 S. Jakó, A. Lupan, A.-Z. Kun and R. B. King, Spherical *closo* deltahedra with surface metal-metal multiple bonding versus oblate deltahedra with internal metal-metal bonding in dichromadicarbaborane structures: The nature of Stone's icosahedral dichromadicarbaborane, *Inorg. Chem.*, 2019, **58**, 3825–3837.
- 32 G. K. Barker, M. R. Godfrey, M. Green, H. E. Parge, F. G. A. Stone and A. J. Welch, Synthesis and molecular structures of nido,-[9-((-C<sub>5</sub>H<sub>5</sub>)-7,8,9-C<sub>2</sub>NiB<sub>8</sub>H<sub>11</sub>], *nido*-[9-((-C<sub>5</sub>H<sub>5</sub>-μ<sub>10,11</sub>-(Ph<sub>3</sub>PAu)-7.8.9-C<sub>2</sub>NiB<sub>8</sub>H<sub>10</sub>], and *closo*-[1,3-((-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>-1,2,3,4-CrCCrCB<sub>8</sub>H<sub>10</sub>]: Evidence for a multiple metal-metal bond in a dimetallacarbaborane, *J. Chem. Soc., Chem. Commun.*, 1983, 277–279.
- 33 A. Găină-Gardiuta, A. Lupan and R. B. King, Triplet capped deltahedral structures rather than singlet oblatocloso, structures as energetically favored dimanganaborane structures: A new structural paradigm in hypoelectronic metallaborane chemistry, *Inorg. Chem.*, 2022, 61, 20793–20803.
- 34 C. Adamo and V. Barone, Toward reliable density functional methods without adjustable parameters: The PBE0 model, *J. Chem. Phys.*, 1999, **110**, 6158–6170.
- 35 Gaussian 09 (Revision A.02), Gaussian, Inc., Wallingford, CT, 2009. The complete reference is given in the ESI.†
- 36 F. Weinhold and C. R. Landis, *Valency and Bonding: A Natural Bond Order Donor-Acceptor Perspective*, Cambridge University Press, Cambridge, England, U. K., 2005, pp. 32–36.