



Cite this: *Chem. Sci.*, 2025, **16**, 18167

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 9th July 2025
Accepted 29th August 2025

DOI: 10.1039/d5sc05077d
rsc.li/chemical-science

Introduction

Enantioenriched aziridines serve as valuable scaffolds for various biologically active natural products, such as Mitomycins^{1–3} and Azinomycins.^{4–6} These structures are known for their antitumor, antibiotic, antimicrobial, neoplasm inhibiting, and glycosidase inhibitory properties.^{7–9} Due to the nature of their strained rings, aziridines serve as synthetic intermediates in ring-opening reactions^{10,11} to produce amine-derived products.¹² They are also useful in ring-expansion reactions to form larger heterocycles,¹³ such as β -lactams,¹⁴ γ -lactams,¹⁵ pyrrolidines,¹⁶ and piperidines,¹⁷ which are essential in both organic synthesis and medicinal chemistry.

Classic methods for the stereoselective synthesis of aziridines^{18–20} often involve the addition of nitrenes to olefins,^{21–23} transfer of carbenes to imines,^{24–27} and the intramolecular cyclization of chiral 1,2-vicinal haloamines^{28–30} or amino alcohols.³¹ Many of these approaches use transition metal catalysts due to their efficiency and broad applicability. However, concerns regarding the high costs and sustainability of these catalysts have driven a shift toward more environmentally friendly methods. To address this, researchers have increasingly turned to organocatalysis^{32,33} as a promising alternative for aziridine synthesis.


In this regard, the groups of Cordova,^{34,35} Hamada,³⁶ and Albrecht³⁷ have reported organocatalytic enantioselective aziridination reactions utilizing an aza-Michael-initiated ring-closing approach (Scheme 1a), which leads to good to

Sulfenate anion catalyzed enantio- and diastereoselective aziridination

Youse Pu, ^a Anthony M. Smaldone, ^a Javier Adrio ^{ab} and Patrick J. Walsh ^{ab}

The synthesis of enantioenriched aziridines is important for drug development due to their prevalence in bioactive molecules. Previous methods often use expensive catalysts, activated substrates, or show poor stereoselectivity. Herein, we report a novel organocatalytic approach using enantioenriched [2.2] paracyclophane (PCP)-based sulfenate anion catalysts, enabling the synthesis of 18 cyclopropanated aziridines from unactivated imines and commercially available benzyl chlorides in 50–99% yields with 73–99% ee and >20:1 dr. This approach fills a gap in the existing methods for aziridine synthesis, facilitating the generation of cyclopropyl-substituted aziridines with high stereoselectivity under mild and transition metal-free reaction conditions.

excellent yields and stereoselectivities. However, these reactions, are typically limited to electronically activated substrates, such as α,β -unsaturated carbonyl compounds.



Scheme 1 Recent advances in organocatalytic aziridination.

^aRoy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, 231 South 34th Street, Philadelphia, PA, USA. E-mail: pwalsh@sas.upenn.edu

^bDepartment of Organic Chemistry, Institute for Advance Research in Chemical Sciences (IAdChem), Universidad Autónoma de Madrid, Cantoblanco, 28049-Madrid, Spain. E-mail: javier.adrio@uam.es



To expand the scope of this process, Kürti and coworkers developed a ketone catalyzed aziridination of unactivated olefins (Scheme 1b).³⁸ This approach exploits an *in situ*-generated oxaziridine intermediate, enabling selective nitrogen transfer to unactivated carbon–carbon double bonds with excellent regio- and diastereoselectivity. This method has yet to be made enantioselective. Chein and coworkers³⁹ developed an organocatalytic method for asymmetric aziridination with benzyl bromides and imines *via* the imino Corey–Chaykovsky reaction, using a tetrahydrothiophene-based chiral sulfide as the catalyst (Scheme 1c). This method achieved aziridination of *N*-phosphonate-activated benzaldimines with excellent enantioselectivities. Although these methods expand the range of accessible aziridines, the diastereoselectivities were moderate ($dr = 54/46$ to $83/17$). Thus, complementary strategies for achieving high levels of enantio- and diastereoselectivity with unactivated substrates remain in demand.

Our group has been exploring sulfur-based organocatalysts and has successfully employed the sulfenate anion (RSO^\ominus), the conjugate base of sulfenic acids, in various catalytic reactions, including the synthesis of *trans*-stilbenes,^{40,41} stilbene-based polymers⁴² and a one-pot method to form all three bonds of diaryl alkynes.⁴³ The nucleophilic nature of the sulfenate anion allows it to effectively attack electrophiles such as benzyl chlorides, while its ability to function as a leaving group facilitates the closure of the catalytic cycle. The sulfur changes oxidation state in the catalytic cycle^{44–49} and the intermediate sulfoxide activates the α -hydrogens toward deprotonation. Higher-valent sulfur species such as S(iv)⁵⁰ and S(vi)^{26,51–53} have also been reported, illustrating the range of accessible oxidation states. Expanding on these advances, we have recently developed a diastereoselective method for the synthesis of racemic *trans*-aziridines from imines and benzylic or alkyl halides using sulfenate anion ($PhSO^\ominus$) catalysts⁵⁴ (Scheme 1d). While this method affords good yields and high diastereoselectivities (*trans* : *cis* > 20 : 1), the catalyst is achiral, and the products are racemic.

To develop an enantioselective aziridination to couple the two electrophilic partners, we envisioned introducing an enantioenriched sulfenate anion catalyst. Herein, we report the first example of an asymmetric sulfenate anion catalyzed enantio- and diastereoselective aziridine formation (Scheme 1e) utilizing an enantioenriched [2.2]paracyclophane-substituted sulfoxide precatalyst. To our knowledge, this is the first example of asymmetric catalysis using an enantioenriched sulfenate anion.

Results and discussion

Proposed mechanism

Prior to discussing the specifics of catalyst design, we first outline the working mechanism (Fig. 1), as it informs the design process. The catalytic cycle begins with the generation of the sulfenate anion A.⁵⁴ As a strong nucleophile, the sulfenate anion readily reacts with alkyl halides (B) to generate sulfoxide C with a change in oxidation state at sulfur. The resulting sulfoxide, in its more oxidized form, has activated α -hydrogens ($pK_a \sim 27.2$ in




Fig. 1 Proposed mechanism of enantio- and diastereoselective synthesis of aziridines.

$DMSO$)⁵⁵ that can be deprotonated by moderately strong bases to yield the deprotonated intermediate D. Intermediate D is also a strong nucleophile but preferentially reacts with the imine E, leading to the formation of F. The basic nitrogen in F can then act as a nucleophile, displacing the sulfenate anion and closing the catalytic cycle with formation of the aziridine G.

The challenge in developing asymmetric sulfenate anion catalysts for aziridine synthesis is that there are two steps that form three new stereocenters in the catalytic reaction, as depicted in the proposed mechanism in Fig. 1. When the sulfenate anion R^*SO^\ominus (A) reacts with benzyl chloride, the sulfur lone pairs—previously enantiotopic in the achiral catalyst $ArSO^\ominus$ —become diastereotopic in enantioenriched R^*SO^\ominus . To generate a single diastereomer of the catalyst at the sulfoxide intermediate (C) the benzylation reaction must proceed with complete diastereoselectivity. We hypothesize that the configuration at sulfur in catalytic intermediate C will impact the formation of the two carbon-stereocenters in the addition adduct, D. Selecting an appropriate R^* group presents a significant challenge, as the chirality of R^*SO^\ominus plays a role in the establishment of three contiguous stereocenters during the formation of intermediate F (one at sulfur and two at the carbons that will form the aziridine backbone).

Reaction development

We envisioned catalysts R^*SO^\ominus , where R^* is a planar chiral *para*-cyclophane (PCP). Fortunately, beautiful work by the Perri group on the *stoichiometric benzylation* of the *rac*-*para*-cyclophane, $PCP-SO^\ominus$, had been reported in 2008.⁵⁶ This team demonstrated that under their conditions the planar chirality of the PCP group completely controlled the central chirality at sulfur during the S_N2 reaction with benzyl bromide *via* the depicted conformation with the stereoselectivity shown (Table 1, entry 1–3). In this conformation (A0), one sulfur lone



Table 1 Diastereoselective sulfenate salt alkylation

Entry	RX	Base	Solvent	Temp	Yield ^c
1 ^a	BnBr	^t BuOK	THF	-78 °C	82%
2 ^a	BnBr	^t BuOK	THF	-40 °C	80%
3 ^a	BnBr	^t BuOK	THF	0 °C	77%
4 ^b	BnCl	LiN(SiMe ₃) ₂	CPME	80 °C	78%


^a Reactions performed by Perrio and coworkers. ^b Reaction performed using 2 equiv. of BnCl, 2 equiv. of LiN(SiMe₃)₂ with CPME (0.1 M). ^c Isolated yield.

pair extends outward from the ring, while the other is buried between the two aromatic rings. This spatial arrangement directs nucleophilic attack to occur predominantly from the more exposed lone pair, ensuring high stereoselectivity. However, our conditions for the asymmetric aziridination are expected to differ. For example, we previously demonstrated that silyl amide bases were far more effective in racemic aziridine formation than the *tert*-butoxide base used by the Perrio group in the benzylation.⁵⁴ Furthermore, the achiral sulfenate anion catalyst employed a Li⁺ counterion, rather than K⁺ and was performed in a different solvent. Given that main group counterions and solvents are well known to significantly impact reactivity and diastereoselectivity in organic reactions,^{57,58} these differences are likely to have significant implications in the benzylolation at sulfur.

With these considerations in mind, we chose conditions similar to our previous racemic aziridination studies,⁵⁴ using LiN(SiMe₃)₂ to generate *rac*-PCP-SO[⊖] at 80 °C in the presence of benzyl chloride. We were pleased to observe the formation of a single diastereomer of PCP-S(=O)CH₂Ph (**B1**) in 78% yield (Table 1, entry 4). This result supports the formation of a single sulfoxide intermediate, maintaining the stereochemical integrity of the catalyst and preventing the generation of mixed diastereomers that could compromise the enantio- and diastereoselectivity in the aziridine forming steps.

For proof-of-principle studies, we resolved racemic **B1** into its enantiomers using preparative chiral phase HPLC on a small scale. We then employed the enantioenriched **B1** (20 mol%) as catalyst in the presence of 1 equiv. (E)-N,1-Diphenylmethanimine (**1b**), benzyl chloride (**2a**) and 2 equiv. LiN(SiMe₃)₂ at 80 °C with the goal of preparing enantioenriched triphenyl aziridine (Scheme 2a). Despite achieving 92% yield of the desired product **4**, it was found to be racemic.

The observation of 0% ee in asymmetric catalysis is quite informative.⁵⁹ It usually indicates either that the enantioenriched “catalyst” is not actually catalyzing the reaction or there is a path for rapid racemization of the enantioenriched product. To investigate the origin of the observed 0% ee, we



Scheme 2 Enantioselective aziridination and studies on reversible aziridine ring-opening.

considered the possibility of aziridine racemization *via* thermal ring-opening. Aziridines are known to undergo thermally induced ring-opening to form ylide intermediates under certain conditions (Scheme 2b).^{60,61} To assess whether such a process contributes to racemization in our system, we examined the reactivity of fluorinated triphenyl aziridine **5** with dipolarophile **6** (5 equiv.) in CPME (0.1 M) at 45 °C in the presence of an internal standard (C₆F₆ in C₆D₆). The reaction was monitored by ¹⁹F{¹H} NMR spectroscopy (Scheme 2c). After 20 h at 45 °C, full conversion of the aziridine to the corresponding cycloadduct was observed (99% assay yield, determined by ¹⁹F{¹H} NMR). These results suggest that under the aziridine-forming reaction conditions in Scheme 2a, the aziridine product undergoes reversible ring-opening to form the higher energy achiral azomethine ylide, leading to racemization.

We hypothesized that an alkyl substituent on the imine would destabilize the azomethine ylide, thereby increasing the energy barrier for the ring opening/racemization process. To test this hypothesis, we examined the reactivity of cyclopropanated aziridine **3a** with **6** at slightly lower temperature (CPME, 60 °C; Scheme 2d). Notably, only trace amounts of the corresponding 3-pyrroline product were observed, while 96% of the starting aziridine was recovered. This experiment indicates that alkyl-substituted azomethine ylides are significantly less prone to racemization through the azomethine ylide than their aryl-substituted counterparts.

Encouraged by these findings, we next examined the reaction of imine **1b** and benzyl chloride using enantioenriched **B1** as the catalyst (20 mol%, Scheme 2e). Notably, the desired product was obtained in 80% yield with 87% ee, demonstrating that the presence of an alkyl group is critical for suppressing the reversible ring-opening process and preserving enantioselectivity.



Pre-catalyst design and synthesis


Next, we prepared sulfoxide pre-catalysts that enable direct entry into the catalytic cycle *via* the sulfenate anion.⁴¹ Based on our prior experience, we hypothesized that both pre-catalysts **A1** and **A2** (Scheme 3) would be synthetically accessible and capable of generating the PCP-substituted sulfenate anion *via* base-promoted elimination of isobutene or styrene, respectively.

Both **A1** (ref. 62 and 63) and **A2** (ref. 64 and 65) were synthesized from *rac*-PCP-Br, followed by lithiation, and addition to enantioenriched sulfinyl reagents, yielding enantioenriched diastereomers that were separated by chromatography (see the SI for details). The ee of both catalysts were determined to be up to 99% by chiral stationary phase supercritical fluid chromatography (SFC).

Reaction optimization


We then employed both pre-catalysts **A1** and **A2** in reaction optimization studies (Table 2). We began by optimizing the reaction conditions on a 0.1 mmol scale using model substrates: (*E*)-1-cyclopropyl-*N*-(4-methoxyphenyl)methanimine (**1a**) and benzyl chloride (**2a**), with a catalyst loading of 10 mol% (unless otherwise stated). **A1** was selected for initial screening due to its more straightforward and efficient synthetic preparation. In a base screening with $\text{LiN}(\text{SiMe}_3)_2$, $\text{NaN}(\text{SiMe}_3)_2$, and $\text{KN}(\text{SiMe}_3)_2$ (entries 1–3), only $\text{LiN}(\text{SiMe}_3)_2$ showed high diastereoselectivity, yielding exclusively the *trans* product with an assay yield (AY) of 55% (determined by ^1H NMR using dibromomethane as an internal standard, Table 2). $\text{NaN}(\text{SiMe}_3)_2$, and $\text{KN}(\text{SiMe}_3)_2$ both produced a significant amount of the *cis* product, and no enantioselectivity was observed.

Recognizing that higher temperatures might be required to facilitate the elimination of isobutene from **A1** and generate the sulfenate anion,⁴¹ we then started the reaction by pre-stirring the precatalyst, solvent, and base at 110 °C for 30 min. The reaction mixture was cooled to 80 °C, 60 °C, or 45 °C (entries 4–6), at which point the remaining reagents were added and the reaction was allowed to proceed for an additional 24 h. This pre-activation step significantly improved the yield from 55% (entry 1) to 99% (entries 4–5). Notably, entry 5 provided a higher ee of 80%. Further lowering the temperature did not



Scheme 3 Pre-catalysts for the generation of sulfenate anions through base-promoted elimination.

Table 2 Optimization of the aziridination reaction with (*E*)-1-cyclopropyl-*N*-(4-methoxyphenyl)methanimine (**1a**) and benzyl chloride (**2a**)



Entry	Pre-cat.	Solvent	Temp.	Yield ^{a,b}	Ee ^c , dr ^b
1	A1	1,4-Dioxane	80 °C	55%	80%, >20:1
2 ^d	A1	1,4-Dioxane	80 °C	41%	0%, 3:2
3 ^e	A1	1,4-Dioxane	80 °C	76%	0%, 1:1
4 ^f	A1	1,4-Dioxane	110 °C 30 min, then 80 °C	99%	76%, >20:1
5 ^f	A1	1,4-Dioxane	110 °C 30 min, then 60 °C	99%	80%, >20:1
6 ^f	A1	1,4-Dioxane	110 °C 30 min, then 45 °C	66%	83%, >20:1
7 ^f	A1	CPME	110 °C 30 min, then 60 °C	28%	81%, >20:1
8 ^f	A1	Toluene	110 °C 30 min, then 60 °C	Trace	N/A
9 ^f	A1	ⁿ Bu ₂ O	110 °C 30 min, then 60 °C	16%	N/A
10	A2	1,4-Dioxane	60 °C	70%	62%, >20:1
11	A2	CPME	60 °C	66%	88%, >20:1
12	A2	CPME	45 °C	70%	96%, >20:1
13	A2	CPME	25 °C	18%	94%, >20:1
14 ^g	A2	CPME	45 °C	84%	97%, >20:1
15 ^h	A2	CPME	45 °C	92%	97%, >20:1
16 ⁱ	A2	CPME	45 °C	76%	94%, >20:1

^a Reaction conditions: **1a** (0.10 mmol, 1.0 equiv.), **2a** (0.20 mmol, 2.0 equiv.), solvent (1.0 mL, 0.1 M), 24 h. ^b Assay yield and dr's were determined by ^1H NMR using CH_2Br_2 as the internal standard. ^c ee was determined by SFC using column Chiralcel OJ3 with 5% MeOH and 95% CO_2 . ^d $\text{NaN}(\text{SiMe}_3)_2$ was used as base. ^e $\text{KN}(\text{SiMe}_3)_2$ was used as base. ^f Pre-catalyst **A1**, base, and solvent were preheated at 110 °C for 30 min. ^g 0.2 M in CPME. ^h 0.4 M in CPME; 2 equiv. of **1a** and 1 equiv. of **2a** were used and the same yield and ee were observed. ⁱ 0.4 M in CPME and 5 mol% **A2** was used.



significantly increase the ee but instead resulted in a substantial yield reduction (entry 6). Next, we evaluated a series of solvents, including CPME, toluene, and ⁷Bu₂O (entries 7–9). While the reaction in CPME provided an enantioselectivity comparable to that achieved with dioxane, the yield decreased significantly. Toluene produced only trace amounts of the product (entry 8), and ⁷Bu₂O resulted in a 16% yield (entry 9), both too low to obtain reliable ee and dr data. Integrating the results from base, temperature, and solvent screenings, the optimal conditions for precatalyst **A1** were determined to be 10 mol% catalyst loading, 2 equiv. of LiN(SiMe₃)₂, and 1,4-dioxane (0.1 M), with pre-stirring at 110 °C for 30 min, followed by 60 °C for 24 h. Under these conditions, the final product **3a** was obtained in 99% yield with 80% ee (entry 5). Attempts to further enhance the enantioselectivity, including the addition of various additives were unsuccessful (see the SI for details).

Given that precatalyst **A2** was expected to generate the sulfenate anion more efficiently and under milder conditions, due to the greater acidity of its β -hydrogens and the stabilization of the styrene elimination product, it was subsequently used in the reaction optimization. Using the optimized conditions for pre-catalyst **A1**, we evaluated the two top-performing solvents with **A2** as the pre-catalyst. While 1,4-dioxane achieved a slightly higher yield of 70% (entry 10), CPME provided a comparable yield of 66% (entry 11) but with a significantly improved enantiomeric excess (88% vs. 62% for 1,4-dioxane). Based on these results, CPME was selected as the solvent for further optimization. Considering that lower temperatures might enhance the enantioselectivity by favoring the pathway with the lowest activation energy, we decreased the temperature from 60 °C to 45 °C (entry 12) and then to 25 °C (entry 13). We were pleased to observe enantioselectivities of 96% and 94%, respectively. Since 45 °C maintained the yield (70%) while 25 °C resulted in only 18% yield, we selected 45 °C as the reaction temperature for further optimization. Increasing the solution concentration from 0.1 M to 0.2 M and 0.4 M resulted in a notable yield improvement (from 70% to 84% and 92%, entries 14 and 15) while holding the ee at 97%. Reducing the precatalyst loading from 10 mol% to 5 mol% led to a decrease in yield from 92% to 76% (entry 16). Adjusting the ratio of imine to benzyl chloride from 1:2 to 2:1 yielded identical results in terms of yield and ee (entry 15). However, the 2:1 ratio provided higher yields for other substrates, likely due to reduced byproduct formation from coupling of benzyl chlorides to form *trans*-stilbenes.⁴⁰ As a result, the optimized reaction conditions for the aziridination employed 2 equiv. LiN(SiMe₃)₂ in CPME (0.4 M) with the pre-catalyst loading of 10 mol% at 60 °C and with a ratio of 2:1 imine to benzyl chloride (entry 15).

Substrate scope

With the optimized conditions in hand, we investigated the substrate scope with commercially available benzyl chlorides. As shown in Scheme 4, benzyl chlorides bearing electron-donating aryl groups such as 4-Me (**2b**), 4-^tBu (**2c**), 4-SMe (**2d**), 4-C₂H₄Ph (**2e**), and 1,3,5-Me₃ (**2f**) provided *trans*-

aziridines in 81–99% yield and 87–98% ee. Benzyl chlorides bearing electron-withdrawing substituents with 3-OPh (**2g**), 3-CF₃ (**2h**) or 4-OCF₃ (**2l**) cleanly afforded *trans*-aziridines **3g** (98% yield, 97% ee), **3h** (81% yield, 84% ee) and **3i** (81% yield, 97% ee).

We were particularly interested in halogen-substituted aziridines due to their potential for further functionalization *via* transition metal-catalyzed cross-coupling reactions. Accordingly, using benzyl chlorides substituted with 3-Br (**2j**) and 2-Br (**2k**) groups led to the formation of *trans*-aziridine **3j** in 88% yield with 90% ee and *trans*-aziridine **3k** in 73% yield with 73% ee. The lower yield of **3k** is attributed to the steric hindrance caused by the 2-bromo substituent. For benzyl chlorides bearing 4-F or 2-F, the enantioenriched *trans*-aziridine **3i** and **3m** were obtained in 96% with >99% ee and 94% yield with 95% ee, respectively. Similarly, 4-chlorobenzyl chloride was efficiently converted to *trans*-aziridine **3n** in 90% yield and 95% ee.

We tested 1-(chloromethyl)naphthalene, with extended conjugation, which successfully yielded *trans*-aziridine **3o** in 70% yield and 92% ee. Carbon–carbon double bonds are also of interest due to their potential for further elaboration, such as hydrofunctionalization,⁶⁶ oxidation,⁶⁷ or cross-coupling reactions.⁶⁸ Accordingly, 4-vinylbenzyl chloride (**2p**) was used to deliver *trans*-aziridine **3p** in 93% yield with 96% ee. Aziridines containing heterocycles are frequently used in drug molecules. As shown in Scheme 4, a pyrimidine-containing aziridine **3q** was obtained in 50% yield with 73% ee.


We were interested to determine if electrophiles other than benzyl chlorides were suitable. Allyl chloride was attractive because the product would be a vinyl aziridine. In the event, the use of allyl chloride (**2r**) furnished the vinyl-substituted aziridine **3r** in 58% yield with 89% ee, highlighting the potential of this method to make highly functionalized building blocks.^{69–71}

Other alkyl aldehydes were evaluated for imine formation, including pivaldehyde and cyclohexanecarbaldehyde, to generate (*E*)-*N*-(4-methoxyphenyl)-2,2-dimethylpropan-1-imine (**2s**), (*E*)-1-cyclohexyl-*N*-(4-methoxyphenyl)methanimine (**2t**), and (*E*)-1-phenyl-*N*-(3-(trifluoromethyl)phenyl)methanimine (**2u**), separately. However, imine **2s** did not yield any product likely due to increased steric hindrance. For imines with α -C-H's, like **2t**, tautomerization to the enamine occurred at room temperature, resulting in no formation of the desired product **3t**. Unfortunately, substrates that readily form enamines are not viable under our reaction conditions. In the case of **2u**, we attempted to install an electron-withdrawing 3-C₆H₄-CF₃ group on the imine nitrogen to reduce the stability of the possible ring-opening azomethine ylide intermediate (Scheme 2b). However, the resulting aziridine **3u** was obtained in 92% yield with 0% ee, indicating that racemization is still occurring under our reaction conditions due to the ring-opening process.


Gram-scale and X-ray structure determination

To illustrate the scalability of this aziridine synthesis, compound **3l** was prepared on a 6 mmol scale, yielding a 95% isolated yield (1.62 g) with 99% ee and >20:1 dr (Scheme 5). To





Scheme 4 Substrate scope (dr's > 20 : 1). ^aReaction was conducted for 36 hours. ^bReaction was conducted with 4 equiv. of 2r.



Scheme 5 Gram-scale synthesis of aziridine 3l.

determine the absolute configuration of the aziridines, a single crystal of the aziridine 3l was obtained through cooling 3l in a solution in hexanes from 40 °C to -16 °C. The X-ray crystal structure (Fig. 2, CCDC: 2422017) confirmed that the aziridine was the (2S,3S) isomer (see the SI for crystallographic data),




Fig. 2 X-ray structure of (2S,3S) aziridine 3l (CCDC: 2422017).

which was obtained by using the precatalyst **A2** with a R_p configuration.

Conclusions

In conclusion, we have developed the first enantioselective sulfenate anion-catalyzed method, which has been showcased in the highly enantioselective synthesis of *trans*-aziridines from simple benzyl chlorides and imines. The reaction proceeds with good to excellent yields and enantioselectivities. The [2.2]paracyclophane (PCP) scaffold is instrumental in providing excellent stereocontrol, not only in the formation of the sulfur stereocenter, but also in the generation of the two stereocenters in the aziridine backbone. Although the current scope is somewhat restricted due to ring opening of certain aziridines to achiral azomethane ylides, it provides proof-of-concept that this strategy is viable and enables the synthesis of enantioenriched cyclopropyl aziridines, a class of compounds not previously accessible with high levels of enantioselectivity. Further studies to develop more efficient oxidation state altering main group catalysts are under way in our laboratories.

Author contributions

P. J. W. conceived the project and supervised the research. J. A. conducted preliminary experiments and Y. P. conducted the experimental work and data analysis with A. S. P. J. W and Y. P. wrote the original draft of the manuscript. All authors contributed to the discussion and revision of the manuscript.

Conflicts of interest

The authors declare no competing financial interest.

Data availability

CCDC 2422017 contains the supplementary crystallographic data for this paper.⁷²

All data for this manuscript is included in the SI. Supporting information: The experimental procedures, characterization data and crystallographic data. Deposition Number 2422017 contains the supplementary crystallographic data for this paper. See DOI: <https://doi.org/10.1039/d5sc05077d>.

Acknowledgements

P. J. W. thanks the NSF CHE-2154593. AMS acknowledges the NSF REU program CHE-1851640. J. A. is grateful for financial support from the Ministerio de Educación, Cultura y Deporte, subprograma estatal de movilidad and Fulbright grant visiting scholar.

Notes and references

- 1 T. Hata, T. Hoshi, K. Kanamori, A. Matsumae, Y. Sano, T. Shima and R. Sugawara, Mitomycin, a new antibiotic from Streptomyces. I, *J. Antibiot. Ser. A*, 1956, **9**, 141–146.
- 2 T. Hata and R. Sugawara, Mitomycin, a new antibiotic from Streptomyces. II. Description of the strain, *J. Antibiot. Ser. A*, 1956, **9**, 147–151.
- 3 S. Wakaki, H. Marumo, K. Tomioka, G. Shimizu, E. Kato, H. Kamada, S. Kudo and Y. Fujimoto, Isolation of new fractions of antitumor mitomycins, *Antibiot. Chemother.*, 1958, **8**, 228–240.
- 4 T. J. Hodgkinson and M. Shipman, Chemical synthesis and mode of action of the azinomycins, *Tetrahedron*, 2001, **57**, 4467–4488.
- 5 G. T. Kelly, C. Liu, R. Smith, R. S. Coleman and C. M. H. Watanabe, Cellular Effects Induced by the Antitumor Agent Azinomycin B, *Chem. Biol.*, 2006, **13**, 485–492.
- 6 J. Foulke-Abel, H. Agbo, H. Zhang, S. Mori and C. M. H. Watanabe, Mode of action and biosynthesis of the azabicycle-containing natural products azinomycin and ficeleomycin, *Nat. Prod. Rep.*, 2011, **28**, 693–704.
- 7 P. A. S. Lowden, Aziridine Natural Products – Discovery, Biological Activity and Biosynthesis, in *Aziridines and Epoxides in Organic Synthesis*, ed. A. K. Yudin, 2006, pp. 399–442, DOI: [10.1002/3527607862.ch11](https://doi.org/10.1002/3527607862.ch11).
- 8 C. Henderson, Recent Advances in the Usage of Mitomycin. Proceedings of a symposium. Hawaii, March 21–24, 1991, *Oncology*, 1993, **50**(suppl. 1), 1–83.
- 9 M. Casely-Hayford and M. Searcey, The Azinomycins. Discovery, Synthesis, and DNA-Binding Studies, in *DNA and RNA Binders*, ed. M. Demeunynck, C. Bailly, W. D. Wilson, 2002, pp. 676–696, DOI: [10.1002/3527601783.ch24](https://doi.org/10.1002/3527601783.ch24).
- 10 S. Sabir, G. Kumar, V. P. Verma and J. L. Jat, Aziridine Ring Opening: An Overview of Sustainable Methods, *ChemistrySelect*, 2018, **3**, 3702–3711.
- 11 P. S. Bera, Y. K. Mirza, T. Sachdeva and M. Bera, Recent Advances in Transition Metal-Catalyzed Ring-Opening Reaction of Aziridine, *Compounds*, 2024, **4**, 626–649.
- 12 X. E. Hu, Nucleophilic ring opening of aziridines, *Tetrahedron*, 2004, **60**, 2701–2743.
- 13 N. Srivastava and H.-J. Ha, Regioselective ring opening of aziridine for synthesizing azaheterocycle, *Front. Chem.*, 2023, **11**, 1280633.
- 14 N. Piens and M. D'Hooghe, Carbonylation of Aziridines as a Powerful Tool for the Synthesis of Functionalized β -Lactams, *Eur. J. Org. Chem.*, 2017, **2017**, 5943–5960.
- 15 T. B. Bisol, A. J. Bortoluzzi and M. M. Sá, Nucleophilic Ring-Opening of Epoxide and Aziridine Acetates for the Stereodivergent Synthesis of β -Hydroxy and β -Amino γ -Lactams, *J. Org. Chem.*, 2011, **76**, 948–962.
- 16 T.-Y. Lin, H.-H. Wu, J.-J. Feng and J. Zhang, Divergent Access to Functionalized Pyrrolidines and Pyrrolines via Iridium-Catalyzed Domino-Ring-Opening Cyclization of Vinyl Aziridines with β -Ketocarbonyls, *Org. Lett.*, 2017, **19**, 6526–6529.
- 17 J. Eshon, K. A. Nicastri, S. C. Schmid, W. T. Raskopf, I. A. Guzei, I. Fernández and J. M. Schomaker, Intermolecular [3+3] ring expansion of aziridines to dehydropiperidines through the intermediacy of aziridinium ylides, *Nat. Commun.*, 2020, **11**, 1273.



18 L. Degennaro, P. Trinchera and R. Luisi, Recent Advances in the Stereoselective Synthesis of Aziridines, *Chem. Rev.*, 2014, **114**, 7881–7929.

19 D. Tanner, Chiral aziridines—their synthesis and use in stereoselective transformations, *Angew. Chem. Int. Ed. Engl.*, 1994, **33**, 599–619.

20 H. M. I. Osborn and J. Sweeney, The asymmetric synthesis of aziridines, *Tetrahedron Asymmetry*, 1997, **8**, 1693–1715.

21 L. Maestre, W. M. C. Sameera, M. M. Díaz-Requejo, F. Maseras and P. J. Pérez, A General Mechanism for the Copper- and Silver-Catalyzed Olefin Aziridination Reactions: Concomitant Involvement of the Singlet and Triplet Pathways, *J. Am. Chem. Soc.*, 2013, **135**, 1338–1348.

22 Y. Li, J. He, V. Khankhoje, E. Herdtweck, K. Köhler, O. Storcheva, M. Cokoja and F. E. Kühn, Copper(II) complexes incorporating poly/perfluorinated alkoxyaluminate-type weakly coordinating anions: syntheses, characterization and catalytic application in stereoselective olefin aziridination, *Dalton Trans.*, 2011, **40**, 5746–5754.

23 D. A. Evans, M. T. Bilodeau and M. M. Faul, Development of the Copper-Catalyzed Olefin Aziridination Reaction, *J. Am. Chem. Soc.*, 1994, **116**, 2742–2753.

24 K. B. Hansen, N. S. Finney and E. N. Jacobsen, Carbenoid Transfer to Imines: A New Asymmetric Catalytic Synthesis of Aziridines, *Angew. Chem. Int. Ed. Engl.*, 1995, **34**, 676–678.

25 J. R. Krumper, M. Gerisch, J. M. Suh, R. G. Bergman and T. D. Tilley, Monomeric Rhodium(II) Catalysts for the Preparation of Aziridines and Enantioselective Formation of Cyclopropanes from Ethyl Diazoacetate at Room Temperature, *J. Org. Chem.*, 2003, **68**, 9705–9710.

26 X. Zeng, X. Zeng, Z. Xu, M. Lu and G. Zhong, Highly Efficient Asymmetric Trans-Selective Aziridination of Diazoacetamides and N-Boc-imines Catalyzed by Chiral Brønsted Acids, *Org. Lett.*, 2009, **11**, 3036–3039.

27 A.-H. Li, L.-X. Dai and V. K. Aggarwal, Asymmetric Ylide Reactions: Epoxidation, Cyclopropanation, Aziridination, Olefination, and Rearrangement, *Chem. Rev.*, 1997, **97**, 2341–2372.

28 D. Guijarro, Ó. Pablo and M. Yus, Synthesis of γ -, δ -, and ε -Lactams by Asymmetric Transfer Hydrogenation of N-(tert-Butylsulfinyl)iminoesters, *J. Org. Chem.*, 2013, **78**, 3647–3654.

29 Y. Hayashi, T. Urushima, D. Sakamoto, K. Torii and H. Ishikawa, One-Pot Synthesis of Chiral Aziridines by a Domino Reaction by Using Desulfonylative Formation on the N-Tosyl Imine of Chloroacetaldehyde with an Asymmetric Mannich Reaction as a Key Step, *Chem.-Eur. J.*, 2011, **17**, 11715–11718.

30 J. Wei, Z. Chen, Y. Gao, P. Zhang, C. Wang, P. Zhao, Y. Wang and X. Shi, A Rapid and Simple Method for Quantitative Aziridination from Aminobrominated Derivatives of Olefins under Solvent-free and Mild Conditions, *Chin. J. Chem.*, 2012, **30**, 391–399.

31 S. K. Kim and E. N. Jacobsen, General Catalytic Synthesis of Highly Enantiomerically Enriched Terminal Aziridines from Racemic Epoxides, *Angew. Chem., Int. Ed.*, 2004, **43**, 3952–3954.

32 E. Roma, E. Tosi, M. Miceli and T. Gasperi, Asymmetric Organocatalytic Aziridination: Recent Advances, *Asian J. Org. Chem.*, 2018, **7**, 2357–2367.

33 Y. Zhu, Q. Wang, R. G. Cornwall and Y. Shi, Organocatalytic Asymmetric Epoxidation and Aziridination of Olefins and Their Synthetic Applications, *Chem. Rev.*, 2014, **114**, 8199–8256.

34 L. Deiana, G.-L. Zhao, S. Lin, P. Dziedzic, Q. Zhang, H. Leijonmarck and A. Córdova, Organocatalytic Enantioselective Aziridination of α -Substituted α,β -Unsaturated Aldehydes: Asymmetric Synthesis of Terminal Aziridines, *Adv. Synth. Catal.*, 2010, **352**, 3201–3207.

35 J. Vesely, I. Ibrahim, G.-L. Zhao, R. Rios and A. Córdova, Organocatalytic Enantioselective Aziridination of α,β -Unsaturated Aldehydes, *Angew. Chem., Int. Ed.*, 2007, **46**, 778–781.

36 H. Arai, N. Sugaya, N. Sasaki, K. Makino, S. Lectard and Y. Hamada, Enantioselective aziridination reaction of α,β -unsaturated aldehydes using an organocatalyst and tert-butyl N-arenesulfonyloxy carbamates, *Tetrahedron Lett.*, 2009, **50**, 3329–3332.

37 S. Frankowski, J. Bojanowski, M. Saktura, M. Romaniszyn, P. Drelich and Ł. Albrecht, Organocatalytic Synthesis of cis-2,3-Aziridine Aldehydes by a Postreaction Isomerization, *Org. Lett.*, 2017, **19**, 5000–5003.

38 Q.-Q. Cheng, Z. Zhou, H. Jiang, J. H. Siitonens, D. H. Ess, X. Zhang and L. Kürti, Organocatalytic nitrogen transfer to unactivated olefins via transient oxaziridines, *Nat. Catal.*, 2020, **3**, 386–392.

39 M.-T. Huang, H.-Y. Wu and R.-J. Chein, Enantioselective synthesis of diaryl aziridines using tetrahydrothiophene-based chiral sulfides as organocatalysts, *Chem. Commun.*, 2014, **50**, 1101–1103.

40 M. Zhang, T. Jia, H. Yin, P. J. Carroll, E. J. Schelter and P. J. Walsh, A New Class of Organocatalysts: Sulfenate Anions, *Angew. Chem., Int. Ed.*, 2014, **53**, 10755–10758.

41 M. Zhang, T. Jia, I. K. Sagamanova, M. A. Pericás and P. J. Walsh, tert-Butyl Phenyl Sulfoxide: A Traceless Sulfenate Anion Precatalyst, *Org. Lett.*, 2015, **17**, 1164–1167.

42 M. Li, S. Berritt, C. Wang, X. Yang, Y. Liu, S.-C. Sha, B. Wang, R. Wang, X. Gao, Z. Li, X. Fan, Y. Tao and P. J. Walsh, Sulfenate anions as organocatalysts for benzylic chloromethyl coupling polymerization via C=C bond formation, *Nat. Commun.*, 2018, **9**, 1754.

43 M. Zhang, T. Jia, C. Y. Wang and P. J. Walsh, Organocatalytic Synthesis of Alkynes, *J. Am. Chem. Soc.*, 2015, **137**, 10346–10350.

44 D. M. Freudendahl, S. Santoro, S. A. Shahzad, C. Santi and T. Wirth, Green Chemistry with Selenium Reagents: Development of Efficient Catalytic Reactions, *Angew. Chem., Int. Ed.*, 2009, **48**, 8409–8411.

45 A. Breder and S. Ortgies, Recent developments in sulfur- and selenium-catalyzed oxidative and isohypsic functionalization reactions of alkenes, *Tetrahedron Lett.*, 2015, **56**, 2843–2852.



46 S. Ortgies and A. Breder, Oxidative Alkene Functionalizations via Selenium- π -Acid Catalysis, *ACS Catal.*, 2017, **7**, 5828–5840.

47 L. Shao, Y. Li, J. Lu and X. Jiang, Recent progress in selenium-catalyzed organic reactions, *Org. Chem. Front.*, 2019, **6**, 2999–3041.

48 F. V. Singh and T. Wirth, Selenium reagents as catalysts, *Catal. Sci. Technol.*, 2019, **9**, 1073–1091.

49 A. Matviitsuk, J. L. Panger and S. E. Denmark, Catalytic, Enantioselective Sulfenofunctionalization of Alkenes: Development and Recent Advances, *Angew. Chem., Int. Ed.*, 2020, **59**, 19796–19819.

50 X. Zhang, R. Sun, D. Zeng, M. Wang and X. Jiang, Cu-catalyzed Bilateral Linkage of Sulfenamides for the Construction of Benzothiophene-based Cyclic Sulfilimines, *Eur. J. Org. Chem.*, 2025, **28**, e202401467.

51 M. Wang, S. Chen and X. Jiang, Construction of Functionalized Annulated Sulfone via SO₂/I Exchange of Cyclic Diaryliodonium Salts, *Org. Lett.*, 2017, **19**, 4916–4919.

52 D. Zeng, Y. Ma, W.-P. Deng, M. Wang and X. Jiang, Divergent sulfur(VI) fluoride exchange linkage of sulfonimidoyl fluorides and alkynes, *Nat. Synth.*, 2022, **1**, 455–463.

53 S. Zhao, D. Zeng, M. Wang and X. Jiang, C-SuFEx linkage of sulfonimidoyl fluorides and organotrifluoroborates, *Nat. Commun.*, 2024, **15**, 727.

54 Z. Zheng, Y. Pu, J. Adrio and P. J. Walsh, Sulfenate Anion Catalyzed Diastereoselective Synthesis of Aziridines, *Angew. Chem., Int. Ed.*, 2023, **62**, e202303069.

55 F. G. Bordwell, Equilibrium acidities in dimethyl sulfoxide solution, *Acc. Chem. Res.*, 1988, **21**, 456–463.

56 J.-F. Lohier, F. Foucoin, P.-A. Jaffrè, J. I. Garcia, J. Sopková-de Oliveira Santos, S. Perrio and P. Metzner, An Efficient and Straightforward Access to Sulfur Substituted [2.2]Paracyclophanes: Application to Stereoselective Sulfenate Salt Alkylation, *Org. Lett.*, 2008, **10**, 1271–1274.

57 T. X. Gentner and R. E. Mulvey, Alkali-Metal Mediation: Diversity of Applications in Main-Group Organometallic Chemistry, *Angew. Chem., Int. Ed.*, 2021, **60**, 9247–9262.

58 R. Sreedharan and T. Gandhi, Masters of Mediation: MN(SiMe₃)₂ in Functionalization of C(sp)–H Latent Nucleophiles, *Chem.-Eur. J.*, 2024, **30**, e202400435.

59 P. J. Walsh and M. C. Kozlowski, *Fundamentals of Asymmetric Catalysis*, University Science Books, 2009.

60 I. Coldham and R. Hutton, Intramolecular Dipolar Cycloaddition Reactions of Azomethine Ylides, *Chem. Rev.*, 2005, **105**, 2765–2810.

61 H. W. Heine and R. Peavy, Aziridines X.I. Reaction of 1,2,3-triphenylaziridine with diethylacetylene dicarboxylate and maleic anhydride, *Tetrahedron Lett.*, 1965, **6**, 3123–3126.

62 G. J. Rowlands and R. J. Seacome, Enantiospecific synthesis of [2.2]paracyclophane-4-thiol and derivatives, *Beilstein J. Org. Chem.*, 2009, **5**, 9.

63 D. J. Weix and J. A. Ellman, Improved Synthesis of tert-Butanesulfonamide Suitable for Large-Scale Production, *Org. Lett.*, 2003, **5**, 1317–1320.

64 P. B. Hitchcock, G. J. Rowlands and R. Parmar, The synthesis of enantiomerically pure 4-substituted [2.2]paracyclophane derivatives by sulfoxide–metal exchange, *Chem. Commun.*, 2005, 4219–4221, DOI: [10.1039/B507394D](https://doi.org/10.1039/B507394D).

65 D. Jishkariani, B. T. Diroll, M. Cargnello, D. R. Klein, L. A. Hough, C. B. Murray and B. Donnio, Dendron-Mediated Engineering of Interparticle Separation and Self-Assembly in Dendronized Gold Nanoparticles Superlattices, *J. Am. Chem. Soc.*, 2015, **137**, 10728–10734.

66 X. Yin, S. Li, K. Guo, L. Song and X. Wang, Palladium-Catalyzed Enantioselective Hydrofunctionalization of Alkenes: Recent Advances, *Eur. J. Org. Chem.*, 2023, **26**, e202300783.

67 C. Bonini and G. Righi, A critical outlook and comparison of enantioselective oxidation methodologies of olefins, *Tetrahedron*, 2002, **58**, 4981–5021.

68 L. J. Oxtoby, J. A. Gurak Jr, S. R. Wisniewski, M. D. Eastgate and K. M. Engle, Palladium-Catalyzed Reductive Heck Coupling of Alkenes, *Trends Chem.*, 2019, **1**, 572–587.

69 D. Kang, T. Kim, H. Lee and S. Hong, Regiodivergent Ring-Opening Cross-Coupling of Vinyl Aziridines with Phosphorus Nucleophiles: Access to Phosphorus-Containing Amino Acid Derivatives, *Org. Lett.*, 2018, **20**, 7571–7575.

70 D. C. D. Butler, G. A. Inman and H. Alper, Room Temperature Ring-Opening Cyclization Reactions of 2-Vinylaziridines with Isocyanates, Carbodiimides, and Isothiocyanates Catalyzed by [Pd(OAc)₂]/PPh₃, *J. Org. Chem.*, 2000, **65**, 5887–5890.

71 T. Zhang, S. Wang, D. Zuo, J. Zhao, W. Luo, C. Wang and P. Li, Palladium-Catalyzed Carbonylative [5+1] Cycloaddition of N-Tosyl Vinylaziridines: Solvent-Controlled Divergent Synthesis of α,β - and β,γ -Unsaturated δ -Lactams, *J. Org. Chem.*, 2022, **87**, 10408–10415.

72 Y. Pu, A. M. Smaldone, J. Adrio, P. Walsh, CCDC, Experimental Crystal Structure Determination, 2025, DOI: [10.5517/ccdc.csd.cc2m99iv](https://doi.org/10.5517/ccdc.csd.cc2m99iv).

