



**Controlling uranyl oxo group interactions to Group 14 elements using polypyrrolic Schiff-base macrocyclic ligands**

Journal:	<i>Dalton Transactions</i>
Manuscript ID	DT-ART-05-2016-001948.R1
Article Type:	Paper
Date Submitted by the Author:	20-Jun-2016
Complete List of Authors:	Love, Jason; University of Edinburgh, School of Chemistry Arnold, Polly; University of Edinburgh, EaStCHEM School of Chemistry Bell, Nicola; University of Edinburgh, Chemistry

## Controlling uranyl oxo group interactions to Group 14 elements using polypyrrolic Schiff-base macrocyclic ligands

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/


Nicola L. Bell,<sup>3</sup> Polly L. Arnold\*<sup>3a</sup> and Jason B. Love\*<sup>a</sup>

Heterodinuclear uranyl/group 14 complexes of the aryl- and anthracenyl-linked Schiff-base macrocyclic ligands L<sup>Me</sup> and L<sup>A</sup> were synthesised by reaction of UO<sub>2</sub>(H<sub>2</sub>L) with M{N(SiMe<sub>3</sub>)<sub>2</sub>}<sub>2</sub> (M = Ge, Sn, Pb). For complexes of the anthracenyl-linked ligand (L<sup>A</sup>) the group 14 metal sits out of the N<sub>4</sub>-donor plane by up to 0.7 Å resulting in relatively short M...OUO distances which decrease down the group; however, the solid state structures and IR spectroscopic analyses suggest little interaction occurs between the oxo and Group 14 metal. In contrast, the smaller aryl-linked ligand (L<sup>Me</sup>) enforces greater interaction between the metals; only the Pb<sup>II</sup> complex was cleanly accessible although this complex was relatively unstable in the presence of HN(SiMe<sub>3</sub>)<sub>2</sub> and some organic oxidants. In this case, the equatorial coordination of pyridine-N-oxide causes a 0.8 Å elongation of the endo UO bond and a clear interaction of the uranyl ion with the Pb(II) cation in the second donor compartment.

### Introduction

Uranyl [U<sup>VI</sup>O<sub>2</sub>]<sup>2+</sup> is the most stable and prevalent form of uranium in the environment.<sup>1,2</sup> The redox properties of uranyl are of interest in order to establish chemical routes to separate and immobilise actinide radioactive wastes.<sup>3</sup> Reduced [U<sup>V</sup>O<sub>2</sub>]<sup>+</sup> species are inherently unstable to disproportionation under aqueous conditions forming [U<sup>VI</sup>] and [U<sup>IV</sup>] products, with the latter being insoluble in aqueous waste streams and therefore immobilised by this process.<sup>4</sup> Oxidised U<sup>VI</sup>O<sub>3</sub> is also poorly soluble in water and T-shaped U-tri-oxo species have recently been proposed, alongside cisoid-UO<sub>2</sub> complexes, to be important in oxo-transfer processes in aqueous media.<sup>5</sup> As such, the synthesis of these motifs in molecular species bound within defined ligand environments can improve our understanding of actinide bonding and help us to predict uranyl speciation.

We have studied extensively the reduction of uranyl incorporated within a Schiff-base macrocyclic ligand with two aryl-linked coordination pockets (Scheme 1(a)).<sup>2,6,7,8</sup> Incorporation of two uranyl units into this ligand environment results in reduction to U<sup>V</sup> and oxo group rearrangement to form the homodinuclear 'butterfly' complex (Me<sub>3</sub>SiOU<sup>V</sup>)<sub>2</sub>{μ-(O)}<sub>2</sub>L<sup>Me</sup> which contains bridging oxo groups and acute OUO angles.<sup>2</sup>



**Scheme 1:** Metalation chemistry of the uranyl complexes of aryl (L<sup>Me</sup>) and anthracenyl (L<sup>A</sup>) linked ligands. (N<sup>+</sup> = N(SiMe<sub>3</sub>)<sub>2</sub>, Sol = THF, pyridine)

We have seen similar reduction during the formation of mixed-metal complexes from U<sup>VI</sup>O<sub>2</sub>(H<sub>2</sub>L<sup>Me</sup>). Coordination of an electropositive metal within the vacant compartment of the macrocycle was shown to activate the uranyl towards reduction so forming a range of stable mixed-metal, [U<sup>V</sup>O<sub>2</sub>]<sup>+</sup> complexes.<sup>7,8</sup> In contrast, incorporation of a less reducing


<sup>a</sup> EaStCHEM School of chemistry, Joseph Black Building, The King's Buildings, The University of Edinburgh, West Mains Road, Edinburgh EH9 3FJ

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

metal (e.g. Fe(II), Mn(II)) did not promote reduction and the corresponding uranyl(VI)  $U^{VI}O_2M(L^{Me})$  complexes were isolated.<sup>9</sup> In this ligand, the proximity of the uranyl endo-oxygen to the lower macrocyclic pocket facilitates interaction with the second metal. More recently we have developed the chemistry of an anthracenyl-linked analogue of this ligand and synthesised mono- and dinuclear uranyl complexes (Scheme 1(b)).<sup>10,11</sup> The anthracenyl linker enforces both a greater distance between the  $N_4$ -coordination compartments and a greater degree of coplanarity.

We have now explored and compared the reactivity of the uranyl complexes of these two ligands,  $UO_2(H_2L)$  ( $L = L^{Me}, L^A$ ) towards group 14 metal silylamides ( $M\{N(SiMe_3)_2\}_2$ ,  $M = Ge, Sn, Pb$ ). These latter metals have been used in order to target macrocyclic complexes of  $U^{VI}O_3$ , with the chalcophilic group 14 metal potentially charge balancing through oxidation from  $M^{II}$  to  $M^{IV}$ . The interaction between the two metal ions in the resulting complexes and any resulting activation of the  $UO_2$  bonds has been assessed spectroscopically, as has the reactivity of these complexes towards oxo-transfer reagents.



Scheme 2: Synthesis of complexes 1-4 using ligands  $L^A$  (top) and  $L^{Me}$  (bottom).  $N'' = N(SiMe_3)_2$

## Results and discussion

### Complexes of the anthracenyl-hinged macrocycle $L^A$

The reaction of a dark green solution of  $UO_2(H_2L^A)$  in THF or pyridine with  $M\{N(SiMe_3)_2\}_2$  ( $M = Sn, Pb$ ) results in the immediate formation of deep red solutions and the appearance of new sets of resonances in the  $^1H$  NMR spectra which are consistent with the formation of **1(Pb)** and **1(Sn)** (Scheme 2). In contrast, reaction of  $UO_2(H_2L^A)$  with  $Ge\{N(SiMe_3)_2\}_2$  in pyridine takes 6 h at room temperature to reach full conversion, forming **1(Ge)-py**, and in THF elevated temperatures for ca. 8 h are required to form **1(Ge)-THF** cleanly. Removal of solvent and workup yielded **1(M)-sol** for  $M = Sn, Ge$ , and  $Pb$ .


X-ray quality crystals of the THF solvate of each complex were grown by either slow cooling concentrated THF/ $C_6D_6$  solutions (**1(Pb)-THF**) or diffusion of hexanes into THF (**1(Ge)-THF** and **1(Sn)-THF**) (Figure 1(a): **1(Pb)-THF**; Figure S1: **1(Ge)-THF** and **1(Sn)-THF**). While the three structures are very similar there are some subtle differences (Table 1). Intriguingly all of the group 14 metal ions sit above the  $N_4$ -donor plane within the molecular cleft, so moving closer to the uranyl endo-oxygen. We have previously demonstrated that the flexibility of the  $N_4$ -donor pocket in  $L^{Me}$  allows the coordinated metal to

sit either above or below the plane (by up to ca. 0.6 Å) to accommodate the steric requirements of various co-ligands.<sup>12</sup> The structures show that descending group 14 causes the metal to move further out of the plane and closer to the endo-oxygen. This promotes a  $Pb \cdots OUO$  distance of as little as 3.06 Å, well within the range for bridging  $M-O-Pb$  bonds of 2.22–3.33 Å, although slightly longer than the  $M=O \cdots Pb$  interactions that have been characterised (all ca. 2.6 Å).<sup>13</sup> However, the question of whether the metal interacts with, and induces any weakening of the  $UO_2$  bonding is less clear. The X-ray crystal structures show only a very slight lengthening of the  $U=O_{endo}$  bond on descending the group (ca. 0.01 Å) and no significant change in the  $U=O_{exo}$  bonds. Similarly, the X-ray crystal structure of the uranyl(VI) complex  $UO_2Li(HL^{Me})$  (1.794(3) Å)<sup>7</sup> also does not show significant lengthening of the  $U=O$  bonds relative to its precursor  $UO_2(H_2L^{Me})$  (1.790(4) Å)<sup>14</sup> despite there being a significant interaction between the lithium and the endo-oxygen as evidenced in the solid state IR spectra of these compounds (899 vs. 908  $cm^{-1}$  respectively). The solid state IR spectra (Charts S1 and S3) of the three THF adducts of **1(Sn)**, **1(Ge)** and **1(Pb)** are consistent with a very slight lengthening of the  $UO_2$  bond down the group; in contrast, the solution state IR spectra (Chart S6) are inconclusive with multiple bands appearing in the  $[U^{VI}O_2]$  region (ca. 890–930  $cm^{-1}$ ). These data may suggest the proximity of the group 14 metal to the uranyl oxo group is simply a product of crystal packing effects with

the metal moving out of the N<sub>4</sub>-donor plane due to the increasing ionic radius of the metal down the group.

The <sup>119</sup>Sn NMR spectrum of **1(Sn)-THF** shows a resonance at -459 ppm which is deshielded from that of Sn<sub>2</sub>(L<sup>A</sup>) at -527 ppm; however, with the large anthracenyl-linked molecular cleft we have previously seen endo-solvent coordination in

homodinuclear complexes which may shield the metal.<sup>15</sup> Endo-coordination of solvent would not be possible in **1(Sn)-THF** due to the presence of the linear UO<sub>2</sub> group meaning the tin cation would appear relatively deshielded in the mixed metal complex.



**Figure 1:** Solid-state structures of (a) **1(Pb)-THF**; (b) **2(Pb)-THF**; (c) **3(Pb)-py**; (d) **3(Pb)-THF**. For clarity, all hydrogen atoms and solvent of crystallisation are omitted (displacement ellipsoids are drawn at 50% probability). Atom colours: green = uranium; blue = nitrogen; red = oxygen; light grey = Group 14 element; dark grey = carbon.

**Table 1:** Selected geometric parameters for X-ray crystal structures and IR spectroscopy data.

Crystal	M...O (Å)	M...N <sub>4</sub> -plane	U=O endo (Å)	U=O exo (Å)	IR (Nujol, cm <sup>-1</sup> ) <sup>1</sup>	IR (C <sub>6</sub> H <sub>6</sub> , cm <sup>-1</sup> ) <sup>1</sup>	M-sol (Å)	U-O-py (Å)
UO <sub>2</sub> (H <sub>2</sub> L <sup>A</sup> )(THF)	-	-	1.774(5)	1.776(4)	916	919/910	-	-
<b>1(Ge)-THF</b>	3.36	0.41	1.757(2)	1.762(2)	925	928/910	2.140(2)	-
<b>1(Sn)-THF</b>	3.23	0.52	1.781(3)	1.782(3)	921	927/911	2.325(3)	-
<b>1(Pb)-THF</b> <sup>2</sup>	3.13/ 3.06	0.55	1.767(4)/ 1.780(4)	1.764(4)/ 1.761(4)	916	905	2.487(4)/ 2.469(4)	-
<b>2(Pb)-THF</b>	3.0	0.59	1.787(7)	1.779(7)	-	-	2.455(7)	-
<b>2(Pb)-py</b>	3.0	0.70	1.779(4)	1.776(4)	902	-	2.480(5)	2.313(3)
<b>3(Pb)-THF</b>	2.5	-0.62	1.817(9)	1.78(1)	896	898	-	-
<b>3(Pb)-py</b>	2.91(1)	-0.70	1.77(1)	1.77(1)	908	895	2.58(1)	-
<b>4(Pb)-py</b>	2.612(8)	-0.40	1.853(8)	1.759(7)	893	-	2.73(1)	2.354(6)

<sup>1</sup> resolution = 2 cm<sup>-1</sup>; <sup>2</sup> two molecules in the asymmetric unit

Reaction of **1(Pb)** with one equivalent of pyridine N-oxide yields a mixture of compounds in both THF and pyridine solvent which does not change upon heating. However, addition of an excess of pyridine N-oxide (>8 eq.) allowed conversion to one new product by <sup>1</sup>H NMR spectroscopy which was assigned as **2(Pb)**. No free pyridine was observed in the <sup>1</sup>H NMR spectrum suggesting no oxidation of the Pb<sup>II</sup> centre has occurred and instead a broadened doublet of doublets at 6.26 ppm indicates coordination of one pyridine N-oxide molecule.

Single crystals of **2(Pb)** were grown by slow diffusion of hexane into a pyridine or THF solution containing excess pyridine N-oxide. The refined structures (Figure 1(b) for **2(Pb)-THF** and Figure S2 for **2(Pb)-py**) show coordination of the pyridine N-oxide to the fifth equatorial position of uranium while THF or pyridine solvent coordinates to the Pb centre in the exo- position. While the UO<sub>2</sub> bonds in both solvates are not significantly elongated relative to **1(Pb)-THF**, the IR spectrum demonstrates a significant effect of the strong

equatorial donor with the UO<sub>2</sub> stretching frequency decreasing (902 cm<sup>-1</sup> for **2(Pb)-py** cf. 916 cm<sup>-1</sup> for **1(Pb)-THF**; Chart S7).

Despite the lower oxidation potential of Sn<sup>II</sup> (-0.13 V cf. Pb<sup>II</sup> -1.8 V)<sup>16</sup> the reaction of **1(Sn)** with pyridine N-oxide similarly results in formation of the uranyl(VI) adduct **2(Sn)** which was found to be in equilibrium with solvent coordinated **1(Sn)** in the <sup>1</sup>H NMR spectrum. The <sup>119</sup>Sn NMR spectrum of **2(Sn)** shows a single resonance at -480 ppm which is very slightly shielded relative to **1(Sn)**.

Ge<sup>II</sup> is relatively reducing compared with Sn<sup>II</sup> and Pb<sup>II</sup> but treatment of **1(Ge)** with pyridine N-oxide yields a mixture of compounds in the <sup>1</sup>H NMR spectrum. This may suggest that the relatively electropositive Ge centre may coordinate pyridine-N-oxide in a similar manner to uranium giving a competitive equilibrium between the two metals or even one that displaces a Ge<sup>II</sup> cation from the macrocycle, in solution in the presence of excess oxidant.


Complexes of the aryl-hinged macrocycle  $L^{Me}$ 

Reaction of the uranyl complex of the smaller aryl-hinged ligand  $UO_2(H_2L^{Me})$  with  $Pb\{N(SiMe_3)_2\}_2$  in pyridine yields a red solution of **3(Pb)** immediately whereas in THF solvent the reaction takes several hours. The  $^1H$  NMR spectrum shows the formation of a new set of resonances consistent with **3(Pb)** and the loss of the NH signal whilst the  $^{29}Si$  NMR spectrum shows the formation of  $HN(SiMe_3)_2$ . However, degradation of this complex occurs upon removal of the solvent from the reaction mixture under reduced pressure and the residue was identified as consisting of  $UO_2(H_2L^{Me})$ , **3(Pb)** (from pyridine solution only) and minor impurities. Pure **3(Pb)** is accessed by adding an anti-solvent (hexane) to the reaction mixture and once isolated as a solid **3(Pb)** is stable under reduced pressure. Treatment of crystalline **3(Pb)-py** with  $D\{N(SiMe_3)_2\}$  followed by removal of the solvent under reduced pressure yielded a deep red residue which was redissolved in  $d_5$ -pyridine. The  $^1H$  NMR spectrum of this material showed ligand resonances corresponding to those of  $UO_2(H_2L^{Me})$  but the NH resonance was missing suggesting that degradation was due to the presence of  $DN(SiMe_3)_2$  in the reaction mixture, i.e. the silazide is sufficiently acidic to substitute the Pb from **3(Pb)** by protonolysis. This behaviour contrasts to **1(Pb)** which is stable in the presence of  $HN(SiMe_3)_2$ .

Crystals suitable for X-ray diffraction of both THF and pyridine solvates of **3(Pb)** were grown by diffusion of hexane into pyridine or THF solutions (Figures 1(c) and (d) respectively). The solid state structures of these adducts are different to each other with the Pb cation coordinated by a solvent molecule in **3(Pb)-py** but unsolvated in **3(Pb)-THF**. This results in a significantly longer Pb-N<sub>py</sub> bond (2.58(1) Å) compared to **2(Pb)-py** (2.480(5) Å). It is interesting to note that the Pb...OUO distance in **3-py** of 2.9 Å is very close to those in **1(Pb)** and **2(Pb)** demonstrating the ability of the macrocyclic ligand pockets to distort to optimise metal geometry and separation.

In **3(Pb)-THF** contains a short Pb...O bond and instead of THF coordinating to the Pb cation, an  $\eta^5$ -interaction with the pyrrole  $\pi$ -system of an adjacent molecule exists (Figure 1 and Figure S3). Both structures show bonding with the uranyl endo oxygen although the THF adduct contains a slightly lengthened  $U=O_{endo}$  bond (1.817(9) Å) relative to the pyridine adduct (1.77(1) Å). The solid state IR spectra (Charts S8 and S9) show a  $UO_2$  asymmetric stretch for **3(Pb)-py** at 907  $cm^{-1}$  while that for **3(Pb)-THF** is seen at 896  $cm^{-1}$ . The former is comparable with that for  $(py)UO_2(H_2L^{Me})$  (908  $cm^{-1}$ ) while the latter is closer to that for  $(THF)UO_2Li(HL^{Me})$  (899  $cm^{-1}$ ) which contains a strong  $U=O\cdots Li$  interaction, known to affect the reduction chemistry of the resulting complex.


The treatment of **3(Pb)** with excess pyridine N-oxide resulted in the formation of the py-O adduct **4(Pb)-py**. Single crystals were grown by vapour diffusion of hexane into a concentrated pyridine solution containing excess pyridine N-oxide. The molecular structure (Figure 2) shows the expected coordination of pyridine N-oxide to the uranium centre while the Pb centre is solvated by pyridine.



**Figure 2:** Solid state structure of **4(Pb)-py**. For clarity, all hydrogen atoms are omitted (displacement ellipsoids are drawn at 50% probability). Atom colours: green = uranium; blue = nitrogen; red = oxygen; light grey = lead; dark grey = carbon.


Similarly to **3(Pb)-py** a short bonding interaction (2.612(8) Å) is evident between the Pb and  $O_{endo}$  in **4(Pb)-py** which in this case causes significant elongation of the  $U=O_{endo}$  bond (1.853(8) Å) with little effect on the  $U=O_{exo}$  bond. The former  $U=O$  bond distance is similar to that of reduced uranyl(V) complexes such as  $LiOU^V OLi(HL^{Me})$  ( $U=O_{endo}$  1.834(4) Å)<sup>7</sup> and is slightly longer than the similar bond in  $(THF)_2K[(HO)U^V O_2(H_2L^{Me})]$ <sup>17</sup> (1.821(6) Å). This demonstrates that activation of the linear  $UO_2$  bonds is facilitated by the presence of strong oxo-donors in the equatorial plane as well as by Lewis acidic metal cations coordinating to the axial uranyl oxygen atoms. The elongation of this bond by 0.08 Å is less than seen for other uranyl  $U^V$  complexes in which a strong Lewis acid binds to an oxo group and may reflect the relatively weak Lewis acidity of  $Pb^{II}$ ; the activated  $UO$  bond (1.890(4) Å) in  $OU(OB\{C_6F_5\}_3)(^Ar)acnac)_2$  ( $Ar=3,5-tBu_2C_6H_3$ ) is significantly longer (0.14 Å) than for  $UO_2(^Ar)acnac)_2$  (1.755(5) Å),<sup>18</sup> and in  $OU(OB\{C_6F_5\}_3)(NCN)_2$  ( $NCN = \{Me_3SiN\}CPh\{NSiMe_3\}$ ) at 1.898(3) Å the  $U-OB$  bond is elongated (0.15 Å) relative to 1.750(4) Å in  $UO_2(NCN)_2$ .<sup>19</sup> The IR spectrum of **4(Pb)-py** (Chart S10) shows a  $UO_2$  asymmetric stretch at 893  $cm^{-1}$ , significantly shorter than for **3(Pb)-py** (908  $cm^{-1}$ ) and comparable to  $LiOU^V OLi(HL^{Me})$  (893  $cm^{-1}$ ) and some other reduced  $U^V O_2$  complexes recently synthesised within our group ( $\{(py)R_2AlOU^V OH_2L^{Me}\}$  and  $\{(py)_3MOU^V OH_2L^{Me}\}$ ;  $R = Me, i-Bu, M = Li, Na, K$ )<sup>20</sup> which all appear within the region 891-894  $cm^{-1}$ .

Attempts to oxidise the  $Pb^{II}$  centre with alternative oxidising agents, e.g. o-iodylanisole, p-iodosotoluene, bis-trimethylsilyperoxide, and trimethylamine-N-oxide resulted in the formation of  $UO_2(H_2L^{Me})$  and minor degradation products including a highly symmetrical set of resonances which were assigned as the dinuclear Pb complex **5** (Eq. 1).



The reasons behind the degradation of **3(Pb)** upon reaction with strong oxygen-atom oxidants are unclear; however,  $Pb^{II}$

has a relatively large ionic radius ( $1.2 \text{ \AA}$ )<sup>16</sup> and therefore may be unstable within the cleft of this smaller ligand. We have previously shown that complexes of larger metal cations such as  $\text{Ca}^{\text{II}}$  adopt an alternative, bowl-shaped coordination mode with  $\text{L}^{\text{R}}$  ( $\text{R} = \text{Me}, \text{Et}$ ).<sup>15</sup> As such, **5** was synthesised directly by treatment of  $\text{H}_4\text{L}^{\text{Me}}$  with two equivalents of  $\text{Pb}\{\text{N}(\text{SiMe}_3)_2\}_2$  in pyridine (Eq. 1).



**Figure 3:** Solid state structure of **5**. For clarity, all hydrogen atoms are omitted (displacement ellipsoids are drawn at 50% probability). Atom colours: blue = nitrogen; light grey = lead; dark grey = carbon.

The solid state structure of **5** (Figure 3) shows the expected ‘bowl’ coordination geometry in which the macrocycle folds at the meso-carbon instead of the aryl linker. This provides a larger cleft which can accommodate the two Pb centres without the significant distortion out of the  $\text{N}_4$ -plane seen in the solid state structure of the other, Pacman-shaped Pb complexes in this work. The  $\text{N}_{\text{pyrrole}} \cdots \text{N}_{\text{pyrrole}}$  distance of  $5.25 \text{ \AA}$  is only slightly larger than the equivalent  $\text{N}_{\text{imine}} \cdots \text{N}_{\text{imine}}$  distance for **3(Pb)-py** of  $5.15 \text{ \AA}$  and in fact smaller than that in **1(Pb)-THF** ( $5.27 \text{ \AA}$ ) however the larger  $\text{N}_{\text{pyrrole}}\text{-Pb-N}_{\text{pyrrole}}$  angle for **5** ( $152^\circ$  cf.  $\text{N}_{\text{imine}}\text{-Pb-N}_{\text{imine}}$  angle of  $130^\circ$  for **3(Pb)-py** and  $138^\circ$  for **1(Pb)-THF**) allows more room for the metal to sit within the  $\text{N}_4$ -donor plane and as a result the Pb atoms in **5** sit only  $0.3 \text{ \AA}$  out of their respective  $\text{N}_4$ -pockets. In addition,  $\pi$ -interactions between the lead centre and the aryl ring of the adjacent ligand or the pyridine solvent (See SI Figure S4) add stability to this structure.

The reactions of  $\text{UO}_2(\text{H}_2\text{L}^{\text{Me}})$  in THF or pyridine with various stoichiometries of  $\text{Sn}\{\text{N}(\text{SiMe}_3)_2\}_2$  and under various temperatures yield a complex mixture of diamagnetic and paramagnetic species as evidenced by  $^1\text{H}$  NMR spectroscopy. While slight activation of the  $\text{UO}_2$  unit is suggested by the IR spectrum of **3(Pb)-THF**, it is likely that significantly more interaction occurs with Sn due to its increased Lewis acidity compared to Pb, resulting in the activation of  $\text{UO}_2$  to reduction by a second molecule of  $\text{Sn}\{\text{N}(\text{SiMe}_3)_2\}_2$ . We have studied extensively the reactions of  $\text{UO}_2(\text{H}_2\text{L}^{\text{Me}})$  with electropositive metal amides (e.g. LDA,  $\text{Ln}\{\text{N}(\text{SiMe}_3)_2\}_3$ , and  $\text{Mg}\{\text{N}(\text{SiMe}_3)_2\}_2$ ) and have shown that coordination of these metals within the vacant compartment of the macrocycle results in strong interactions with the endo-oxo of the uranyl and activates this group towards reduction to  $\text{U}^{\text{V}}$ . In contrast, ‘softer’ polarisable metal amides (e.g.  $\text{Fe}\{\text{N}(\text{SiMe}_3)_2\}_2$ ,  $\text{Mn}\{\text{N}(\text{SiMe}_3)_2\}_2$ ) yield the diamagnetic transamination products. In contrast to the above reactions with  $\text{Sn}^{\text{II}}$ , no reaction between  $\text{UO}_2(\text{H}_2\text{L}^{\text{Me}})$  and  $\text{Ge}\{\text{N}(\text{SiMe}_3)_2\}_2$  occurs at ambient temperatures in either THF

or pyridine. Upon heating to  $80 \text{ }^\circ\text{C}$  a number of minor paramagnetic resonances appear suggesting that reduction of the uranyl centre occurs, whilst heating to  $120 \text{ }^\circ\text{C}$  in pyridine results in the formation of the previously reported oxo-silylated complex  $[(\text{Me}_3\text{Si})\text{OUO}(\text{H}_2\text{L}^{\text{Me}})]$  after 24 h as the major product along with multiple minor paramagnetic species. This may indicate that coordination of  $\text{Ge}\{\text{N}(\text{SiMe}_3)_2\}_2$  to the exogenous uranyl oxo may occur, resulting in reductive silylation as seen with  $\text{ZnCl}\{\text{N}(\text{SiMe}_3)_2\}$ .<sup>8</sup>

## Conclusions

Mixed-metal uranyl/Group 14 complexes of two different Schiff base macrocyclic ligands have been synthesised by transamination from  $\text{UO}_2(\text{H}_2\text{L})$  ( $\text{L} = \text{L}^{\text{Me}}$  and  $\text{L}^{\text{A}}$ ). With the larger anthracenyl-linked macrocycle  $\text{Ge}^{\text{II}}$ ,  $\text{Sn}^{\text{II}}$  and  $\text{Pb}^{\text{II}}$  all coordinate in the vacant macrocyclic pocket without reduction of the uranyl. In these complexes the metal ion sits above the  $\text{N}_4$ -donor plane, closer to the uranyl endo-oxygen suggesting a possible bonding interaction. However, it is clear from IR spectroscopy and solid-state structural analysis that only minor activation of the uranyl oxo groups occurs, and so this U-O-M interaction is likely a result of crystal packing forces. In contrast, similar reactions between Group 14 silylamides and the uranyl complex of a smaller aryl-linked macrocycle results in reduction to multiple paramagnetic species for Sn and Ge, likely due to the closer interaction between the metal and  $\text{U}=\text{O}_{\text{endo}}$  enforced by the ligand framework. Only using Pb were we able to isolate the expected mixed-metal uranyl-Pb complex which showed similarly minor activation of the uranyl bonding in the solid state. These data suggest that the proximity of the second metal to uranyl is not an overriding factor in the reduction of uranyl(VI) to uranyl(V) and in the formation of uranyl(VI) oxo-metal bonds. However, the exchange of the equatorial ligand from THF or pyridine in **3(Pb)** with pyridine N-oxide to form **4(Pb)-py** results in significant elongation of the endo-oxo group and the formation of a clear OUO-Pb bonding interaction, similar to those seen in simple uranyl-perfluoroborane Lewis acid-base adducts. This is only the case when using the more constrained macrocycle environment provided by  $\text{L}^{\text{Me}}$ , and supports the premise that the formation of uranyl(VI)-oxo Lewis acid-base interactions requires a ligand environment (i.e. the Pacman macrocycle) that not only defines the approach of the Lewis acid but also the equatorial coordination sphere of the uranyl, in this case the weak-field macrocycle  $\text{N}_4$ -donor set plus the strongly donating pyridine oxide.

## Experimental

### General Details

All manipulations were carried out under a dry, oxygen-free dinitrogen atmosphere using standard Schlenk techniques or in a glove box unless otherwise stated. Solvents (toluene, *n*-hexane, diethyl ether and tetrahydrofuran (THF)) were dried by passage through activated  $4 \text{ \AA}$  molecular sieves or activated

alumina towers and stored over activated 4 Å molecular sieves. Pyridine was distilled from potassium under dinitrogen in a solvent still prior to use. Deuterated solvents were refluxed over potassium, free-pump-thaw degassed three times and vacuum transferred and prior to use.  $^1\text{H}$  NMR spectra were recorded at 298 K unless otherwise stated on either a Bruker AVA400 spectrometer at 399.90 MHz, or AVA500 spectrometer at 500.12 MHz.  $^{13}\text{C}$  NMR spectra were recorded at 298 K on a Bruker AVA500 at 125.77 MHz.  $^{119}\text{Sn}$  and  $^{29}\text{Si}$ -INEPT NMR spectra were run on a Bruker PRO500 spectrometer with a Prodigy cryoprobe at 186 MHz and a Bruker AVA400 spectrometer at 99 MHz respectively.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were referenced internally to residual protio solvent ( $^1\text{H}$ ) or solvent ( $^{13}\text{C}$ ) and are reported relative to tetramethylsilane ( $\delta = 0$  ppm). Chemical shifts are quoted in  $\delta$  (ppm). IR spectra were acquired on a Jasco 410 FT-IR spectrophotometer as  $\text{C}_6\text{H}_6$  solutions, between KBr plates, or as a Nujol mull (w = weak, m = medium, s = strong intensity). Elemental analyses were performed by Mr Stephen Boyer at London Metropolitan University or Pascher Labor, Germany. The compounds  $\text{H}_4\text{L}^{\text{Me}}$ ,<sup>12</sup>  $\text{UO}_2(\text{H}_2\text{L}^{\text{Me}})$ ,<sup>14</sup>  $\text{UO}_2(\text{H}_2\text{L}^{\text{A}})$ ,<sup>11</sup>  $\text{Ge}\{\text{N}(\text{SiMe}_3)_2\}_2$ ,  $\text{Sn}\{\text{N}(\text{SiMe}_3)_2\}_2$  and  $\text{Pb}\{\text{N}(\text{SiMe}_3)_2\}_2$ <sup>21</sup> were synthesised by published methods; all other chemicals were purchased from Sigma Aldrich and used as received.

**1(Ge)-THF** - To a solution of  $\text{UO}_2(\text{H}_2\text{L}^{\text{A}})$  (500 mg, 0.42 mmol) in THF (10 mL) was added a solution of  $\text{Ge}\{\text{N}(\text{SiMe}_3)_2\}_2$  (180 mg, 0.46 mmol) in THF (10 mL) and the solution was stirred for 8 h at 80 °C before the solvent was half removed and hexane added, resulting in the precipitation of microcrystalline **1(Ge)-THF** (340 mg, 0.25 mmol, 61%). Deep red x-ray quality crystals were grown by slow diffusion of hexane into a concentrated solution of **1(Ge)-THF** in THF.

$^1\text{H}$  NMR (500 MHz, THF- $d_6$ ):  $\delta$  9.35 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 9.12 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 8.43 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.99 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.84 (d,  $J = 8.5$  Hz, 2H, 2 x CH), 7.66 (d,  $J = 8.5$  Hz, 2H, 2 x CH), 7.46 (dd,  $J = 8.5$ , 7.0 Hz, 2H, 2 x CH), 7.28 (dd,  $J = 8.5$ , 7.0 Hz, 2H, 2 x CH), 7.20 (d,  $J = 6.9$  Hz, 2H, 2 x CH), 7.17 (d,  $J = 3.6$  Hz, 2H, 2 x CH), 6.62 (d,  $J = 6.9$  Hz, 2H, 2 x CH), 6.60 (d,  $J = 3.6$  Hz, 2H, 2 x CH), 6.55 (d,  $J = 3.6$  Hz, 2H, 2 x CH), 6.02 (d,  $J = 3.6$  Hz, 2H, 2 x CH), 2.27 (qt,  $J = 7.3$ , 3.6 Hz, 4H, 2 x  $\text{CH}_2$ ), 2.09 (q,  $J = 7.3$  Hz, 2H,  $\text{CH}_2$ ), 0.85 (t,  $J = 7.3$  Hz, 3H,  $\text{CH}_3$ ), 0.81 (t,  $J = 7.3$  Hz, 3H,  $\text{CH}_3$ ), 0.80 (t,  $J = 7.3$  Hz, 3H,  $\text{CH}_3$ ), 0.22 (t,  $J = 7.3$  Hz, 3H,  $\text{CH}_3$ ) ppm (one  $\text{CH}_2$  resonance hidden by THF solvent);  $^1\text{H}$  NMR (601 MHz, Pyridine- $d_5$ ):  $\delta$  9.58 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 9.17 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 8.13 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.72 (d,  $J = 8.5$  Hz, 2H, 2 x CH), 7.67 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.50 (d,  $J = 8.4$  Hz, 2H, 2 x CH), 7.34 (d,  $J = 3.7$  Hz, 2H, 2 x CH), 7.25 (m, 4H, 4 x CH), 6.94 (d,  $J = 6.8$  Hz, 2H, 2 x CH), 6.84 (d,  $J = 3.9$  Hz, 2H, 2 x CH), 6.61 (d,  $J = 3.5$  Hz, 2H, 2 x CH), 6.28 (d,  $J = 3.6$  Hz, 2H, 2 x CH), 5.68 (d,  $J = 6.8$  Hz, 2H, 2 x CH), 2.62 – 2.47 (m, 4H, 2 x  $\text{CH}_2$ ), 2.28 (q,  $J = 7.2$  Hz, 2H,  $\text{CH}_2$ ), 1.74 (q,  $J = 7.3$  Hz, 2H,  $\text{CH}_2$ ), 1.25 – 1.04 (m, 12H, 3 x  $\text{CH}_3$ ), 0.67 (t,  $J = 7.3$  Hz, 3H,  $\text{CH}_3$ ) ppm;  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz, THF- $d_6$ ):  $\delta$  162.7( $\text{C}_q$ ), 161.8 (CH), 154.1 (CH), 153.7 ( $\text{C}_q$ ), 153.5 ( $\text{C}_q$ ), 148.2 ( $\text{C}_q$ ), 138.7 ( $\text{C}_q$ ), 137.5 ( $\text{C}_q$ ), 134.0 ( $\text{C}_q$ ), 133.8 ( $\text{C}_q$ ), 128.9 ( $\text{C}_q$ ), 128.8 ( $\text{C}_q$ ), 127.4 (CH), 127.1 (CH), 126.7 (CH), 126.3 (CH), 125.4 (CH), 124.6 (CH), 123.2 (CH), 121.6 (CH), 120.8 (CH),

113.3 (CH), 112.6 (CH), 108.4 (CH), 52.2 ( $\text{C}_q$ ), 46.7 ( $\text{C}_q$ ), 44.6 ( $\text{CH}_2$ ), 40.2 ( $\text{CH}_2$ ), 38.5 ( $\text{CH}_2$ ), 30.4 ( $\text{CH}_2$ ), 11.0 ( $\text{CH}_3$ ), 10.7 ( $\text{CH}_3$ ), 10.6 ( $\text{CH}_3$ ), 10.0 ( $\text{CH}_3$ ) ppm; Analysis Calcd.  $\text{C}_{66}\text{H}_{64}\text{N}_8\text{O}_4\text{GeU}$  (1343.95) requires C % 58.98, H % 4.80, N % 8.34; found C % 58.40, H % 4.37, N % 8.68. FTIR (Nujol mull,  $\text{cm}^{-1}$ ):  $\nu$  1616 (m), 1591 (s), 1553 (m), 1313 (m), 1277 (s), 1266 (s), 1170 (w), 1089 (w), 1054 (m), 1027 (m), 1009 (m), 925 (s,  $\text{UO}_2$  asymmetric stretch), 875 (w), 855 (m), 742 (s), 722 (s); FTIR ( $\text{C}_6\text{H}_6$ ,  $\text{cm}^{-1}$ ):  $\nu$  2969 (m), 2934 (m), 2875 (m), 1595 (vs), 1552 (vs), 1311 (m), 1278 (s), 1255 (m), 1090 (m), 1071 (s), 1058 (s), 1013 (m), 928 (s), 911 (m), 875 (m), 863 (m), 763 (m), 743 (m).

**1(Sn)-THF** - To a solution of  $\text{UO}_2(\text{H}_2\text{L}^{\text{A}})$  (200 mg, 0.16 mmol) in THF (5 mL) was added a solution of  $\text{Sn}\{\text{N}(\text{SiMe}_3)_2\}_2$  (69 mg, 0.16 mmol) in THF (10 mL) and the solution was stirred for 8 h before the solvent was removed and the residue washed with hexane. Recrystallisation from THF/hexane yielded microcrystalline **1(Sn)-THF** (135 mg, 0.12 mmol, 75%).

$^1\text{H}$  NMR (Pyridine- $d_5$ , 500 MHz):  $\delta$  9.62 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 9.38 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 8.15 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 8.12 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.74 (d,  $J = 8.5$  Hz, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.58 (d,  $J = 8.5$  Hz, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.51 (d,  $J = 3.6$  Hz, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 7.33 (dd,  $J = 7.5$  Hz, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.26 (dd,  $J = 7.5$  Hz, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.04 (d,  $J = 6.7$  Hz, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 6.96 (d,  $J = 3.6$  Hz, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 6.61 (d,  $J = 7.0$  Hz, 2H,  $\text{CH}_{\text{Ar}}$ ), 6.56 (d,  $J = 3.8$  Hz, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 6.16 (d,  $J = 3.8$  Hz, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 2.57 (m, 4H, 2 x  $\text{CH}_2$ ), 2.39 (q,  $J = 7.2$  Hz, 2H,  $\text{CH}_2$ ), 1.87 (q,  $J = 7.2$  Hz, 2H,  $\text{CH}_2$ ), 1.28 (t,  $J = 7.2$  Hz,  $\text{CH}_3$ ), 1.15 (t,  $J = 7.2$  Hz,  $\text{CH}_3$ ), 1.12 (t,  $J = 7.2$  Hz,  $\text{CH}_3$ ), 0.79 (t,  $J = 7.2$  Hz,  $\text{CH}_3$ ) ppm;  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz, Pyridine- $d_5$ ):  $\delta$  162.8 ( $\text{C}_q$ ), 161.2 (CH), 154.4 (CH), 153.0 ( $\text{C}_q$ ), 150.9 ( $\text{C}_q$ ), 139.0 ( $\text{C}_q$ ), 138.8 ( $\text{C}_q$ ), 133.2 ( $\text{C}_q$ ), 133.0 ( $\text{C}_q$ ), 128.6 ( $\text{C}_q$ ), 128.1 ( $\text{C}_q$ ), 127.5 (CH), 126.7 (CH), 126.6 (CH), 125.9 (CH), 125.8 (CH), 125.0 (CH), 121.9 (CH), 121.2 (CH), 121.1 (CH), 114.9 (CH), 113.1 (CH), 110.0 (CH), 52.6 ( $\text{C}_q$ ), 46.2 ( $\text{C}_q$ ), 42.1 ( $\text{CH}_2$ ), 40.0 ( $\text{CH}_2$ ), 38.7 ( $\text{CH}_2$ ), 27.2 ( $\text{CH}_2$ ), 11.3 ( $\text{CH}_3$ ), 11.2 ( $\text{CH}_3$ ), 10.7 ( $\text{CH}_3$ ), 10.0 ( $\text{CH}_3$ ) ppm (one  $\text{C}_q$  under pyridine solvent resonances);  $^{119}\text{Sn}$  NMR (186 MHz, pyridine- $d_5$ ):  $\delta$  -459 ppm; Analysis Calcd.  $\text{C}_{46}\text{H}_{64}\text{N}_8\text{O}_4\text{SnU}$  (1390.03) requires C % 57.03, H % 4.64, N % 8.06; found C % 57.20, H % 4.52, N % 8.14; FTIR (Nujol mull,  $\text{cm}^{-1}$ ):  $\nu$  1594 (s), 1554 (m), 1311 (m), 1277 (m), 1262 (m), 1171 (w), 1154 (w), 1088 (m), 1055 (s), 1016 (m), 963 (w), 921 (m,  $\text{UO}_2$  asymmetric stretch), 875 (w), 801 (m), 722 (s); FTIR ( $\text{C}_6\text{H}_6$ ,  $\text{cm}^{-1}$ ):  $\nu$  2968 (m), 2932 (m), 2875 (m), 1599 (vs), 1553 (vs), 1311 (m), 1278 (s), 1091 (m), 1059 (s), 928 (m), 910 (m), 875 (m), 863 (m), 761 (m), 744 (m).

**1(Pb)-THF** - To a solution of  $\text{UO}_2(\text{H}_2\text{L}^{\text{A}})$  (200 mg, 160  $\mu\text{mol}$ ) in THF (5 mL) was added a solution of  $\text{Pb}\{\text{N}(\text{SiMe}_3)_2\}_2$  (87 mg, 160  $\mu\text{mol}$ ) in THF (15 mL) and the solution was stirred for 8 h before the solvent was removed and the residue washed with hexane. Recrystallisation from THF/hexane yielded microcrystalline **1(Pb)-THF** (145 mg, 110  $\mu\text{mol}$ , 64%). Dark red x-ray quality crystals were grown by slow cooling a THF/ $\text{C}_6\text{H}_6$  solution.

$^1\text{H}$  NMR (Pyridine- $d_5$ , 500 MHz):  $\delta$  9.55 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 9.39 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 8.20 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.84 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.75

(d,  $J = 8.5$  Hz, 2H, 2 x CH<sub>Ar</sub>), 7.50 (d,  $J = 8.5$  Hz, 2H, 2 x CH<sub>Ar</sub>), 7.45 (d,  $J = 3.6$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 7.29 (dd,  $J = 7.5$  Hz, 2H, 2 x CH<sub>Ar</sub>), 7.26 (dd,  $J = 7.5$  Hz, 2H, 2 x CH<sub>Ar</sub>), 7.04 (d,  $J = 6.7$  Hz, 2H, 2 x CH<sub>Ar</sub>), 6.88 (d,  $J = 3.6$  Hz, 2H, 2 x CH<sub>Ar</sub>), 6.76 (d,  $J = 3.6$  Hz, 2H, CH<sub>Ar</sub>), 6.53 (d,  $J = 3.6$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 5.79 (d,  $J = 6.7$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 2.61 (q,  $J = 7.3$  Hz, 2H), 2.55 (q,  $J = 7.3$  Hz, 2H), 2.44 (q,  $J = 7.3$  Hz, 2H), 1.97 (q,  $J = 7.3$  Hz, 2H), 1.31 (t,  $J = 7.3$  Hz, 3H), 1.19 (t,  $J = 7.3$  Hz, 3H), 1.10 (t,  $J = 7.3$  Hz, 3H), 0.83 (t,  $J = 7.3$  Hz, 3H) ppm; <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, Pyridine-*d*<sub>5</sub>): δ 163.0 (C<sub>q</sub>), 161.3 (CH), 158.6 (C<sub>q</sub>), 156.1 (CH), 152.7 (C<sub>q</sub>), 151.2 (C<sub>q</sub>), 141.6 (C<sub>q</sub>), 138.7 (C<sub>q</sub>), 133.2 (C<sub>q</sub>), 133.0 (C<sub>q</sub>), 128.8 (C<sub>q</sub>), 128.1 (C<sub>q</sub>), 127.6 (CH), 126.9 (CH), 126.7 (CH), 126.0 (CH), 125.7 (CH), 122.7 (CH), 121.0 (CH), 120.2 (CH), 114.6 (CH), 113.4 (CH), 110.7 (CH), 52.6 (C<sub>q</sub>), 47.3 (C<sub>q</sub>), 41.0 (CH<sub>2</sub>), 39.4 (CH<sub>2</sub>), 38.7 (CH<sub>2</sub>), 26.2 (CH<sub>2</sub>), 11.3 (CH<sub>3</sub>), 11.2 (CH<sub>3</sub>), 10.4 (CH<sub>3</sub>), 9.7 (CH<sub>3</sub>) ppm; Analysis Calcd. C<sub>62</sub>H<sub>56</sub>N<sub>8</sub>O<sub>3</sub>PbU (1406.41) requires C % 52.95, H % 4.01, N % 7.97; found C % 52.85, H % 4.17, N % 7.73; FTIR (Nujol mull, cm<sup>-1</sup>): ν 1597 (s), 1552 (m), 1261 (s), 1090 (m), 1055 (m), 1018 (m), 916 (m, UO<sub>2</sub> asymmetric stretch), 874 (w), 799 (m), 722 (m); FTIR (C<sub>6</sub>H<sub>6</sub>, cm<sup>-1</sup>): ν 2971 (m), 2934 (m), 2876 (m), 1599 (vs), 1402 (w), 1315 (m), 1290 (s), 1277 (s), 1266 (s), 1090 (m), 1058 (s), 916 (m), 905 (s), 874 (s), 862 (s), 761 (m), 744 (m).

**2(Pb)** - To a solution of UO<sub>2</sub>(H<sub>2</sub>L<sup>A</sup>) (100 mg, 80 μmol) in *d*<sub>5</sub>-pyridine (1 mL) was added Pb{N(SiMe<sub>3</sub>)<sub>2</sub>}<sub>2</sub> (43 mg, 80 μmol) followed by pyridine-*N*-oxide (64 mg, 640 μmol, 8 eq.). The solution was filtered and hexane was diffused into the vial over 7 days resulting in the formation of red crystals of **2(Pb)-py** (65 mg, 44 μmol, 54%). **2(Pb)-THF** was synthesised similarly from a THF/C<sub>6</sub>D<sub>6</sub> solution.

<sup>1</sup>H NMR (500 MHz, Pyridine-*d*<sub>5</sub>): δ 10.03 (s, 2H, 2 x CH<sub>im</sub>), 9.25 (s, 2H, 2 x CH<sub>im</sub>), 8.03 (s, 2H, 2 x CH<sub>Ar</sub>), 7.86 (s, 2H, 2 x CH<sub>Ar</sub>), 7.68 (d,  $J = 8.6$  Hz, 2H, 2 x CH), 7.50 (d,  $J = 3.7$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 7.32 – 7.26 (m, 4H, 4 x CH), 7.07 – 7.02 (m, 2H, 2 x CH), 6.88 (d,  $J = 3.7$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 6.79 (d,  $J = 7.0$  Hz, 2H, 2 x CH), 6.71 (d,  $J = 3.5$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 6.46 (d,  $J = 3.5$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 6.26 (br. dd, 1H, CH), 5.85 (d,  $J = 7.0$  Hz, 2H, 2 x CH), 2.66 – 2.54 (m, 4H, 2 x CH<sub>2</sub>), 2.37 (q,  $J = 7.3$  Hz, 2H, CH<sub>2</sub>), 1.81 (q,  $J = 7.3$  Hz, 2H, CH<sub>2</sub>), 1.32 – 1.12 (m, 6H, 2 x CH<sub>3</sub>), 1.07 (t,  $J = 7.3$  Hz, 3H, CH<sub>3</sub>), 0.69 (t,  $J = 7.3$  Hz, 3H, CH<sub>3</sub>) ppm; <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, Pyridine-*d*<sub>5</sub>): δ 162.9 (C<sub>q</sub>), 160.0 (CH), 159.5 (CH), 156.9 (CH), 155.6 (CH), 153.0 (C<sub>q</sub>), 142.2 (C<sub>q</sub>), 133.6 (CH), 133.1 (C<sub>q</sub>), 132.7 (C<sub>q</sub>), 129.2 (C<sub>q</sub>), 128.5 (C<sub>q</sub>), 127.1 (CH), 126.7 (CH), 126.3 (CH), 125.8 (CH), 125.5 (CH), 125.1 (CH), 122.3 (CH), 120.9 (CH), 120.8 (CH), 114.0 (CH), 113.6 (CH), 110.6 (CH), 52.1 (C<sub>q</sub>), 46.9 (C<sub>q</sub>), 41.0 (CH<sub>2</sub>), 33.1 (CH<sub>2</sub>), 25.6 (CH<sub>2</sub>), 11.7 (CH<sub>3</sub>), 10.8 (CH<sub>3</sub>), 10.2 (CH<sub>3</sub>), 9.5 (CH<sub>3</sub>) ppm; Analysis Calcd. C<sub>68</sub>H<sub>58</sub>N<sub>10</sub>O<sub>3</sub>PbU (1508.51) requires C % 54.14, H % 3.88, N % 9.29; found C % 54.23, H % 3.98, N % 9.40; FTIR (Nujol mull, cm<sup>-1</sup>): ν 1588 (s), 1548 (m), 1300 (m), 1276 (s), 1265 (s), 1244 (m), 1170 (m), 1158 (m), 1111 (m), 1090 (m), 1056 (m), 1040 (m), 1017 (m), 953 (m), 902 (m, UO<sub>2</sub> asymmetric stretch), 892 (m), 871 (m), 849 (w), 756 (m), 739 (m), 722 (m).

**3(Pb)-THF** - To a solution of UO<sub>2</sub>(H<sub>2</sub>L<sup>Me</sup>) (100 mg, 93 μmol) in THF (1 mL) was added a solution of Pb{N(SiMe<sub>3</sub>)<sub>2</sub>}<sub>2</sub> (50 mg, 94

μmol) in THF (1 mL). The reaction mixture was heated to reflux for 16 h whereupon the <sup>1</sup>H NMR spectrum indicated the presence of **3(Pb)**. Analytically pure material was isolated by dropwise addition of the reaction mixture to hexane (5 mL) resulting in the precipitation of **3(Pb)-THF** as a fine brown solid (99mg, 82 μmol, 87%) which was filtered and dried under reduced pressure. Slow cooling of the filtrate of this reaction yielded single crystals of **3(Pb)-THF**.

<sup>1</sup>H NMR (400 MHz, THF-*d*<sub>8</sub>): δ 9.07 (s, 2H, 2 x CH<sub>im</sub>), 8.01 (s, 2H, 2 x CH<sub>im</sub>), 7.35 (s, 2H, 2 x CH<sub>Ar</sub>), 6.91 (d,  $J = 3.6$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 6.89 (s, 2H, 2 x CH<sub>Ar</sub>), 6.31 (d,  $J = 3.4$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 6.29 (d,  $J = 3.6$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 5.93 (d,  $J = 3.4$  Hz, 2H, 2 x CH<sub>pyrrole</sub>), 2.40 (s, 6H, 2 x CH<sub>3</sub>), 2.38 (s, 6H, 2 x CH<sub>3</sub>), 1.81 (s, 3H, CH<sub>3</sub>), 1.40 (s, 3H, CH<sub>3</sub>), 1.34 (s, 3H, CH<sub>3</sub>), 1.31 (s, 3H, CH<sub>3</sub>) ppm; <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, THF-*d*<sub>8</sub>): δ 164.6 (C<sub>q</sub>), 161.9 (CH), 157.9 (CH), 157.4 (C<sub>q</sub>), 146.6 (C<sub>q</sub>), 144.8 (C<sub>q</sub>), 139.1 (C<sub>q</sub>), 138.3 (C<sub>q</sub>), 134.7 (C<sub>q</sub>), 132.4 (C<sub>q</sub>), 124.9 (CH), 122.9 (CH), 122.2 (CH), 120.6 (CH), 109.3 (CH), 107.1 (CH), 40.7 (C<sub>q</sub>), 38.7 (C<sub>q</sub>), 34.3 (CH<sub>3</sub>), 33.7 (CH<sub>3</sub>), 30.1 (CH<sub>3</sub>), 27.5 (CH<sub>3</sub>), 18.7 (CH<sub>3</sub>), 18.6 (CH<sub>3</sub>) ppm; Analysis Calcd. C<sub>46</sub>H<sub>50</sub>N<sub>8</sub>O<sub>3</sub>PbU (1208.19) requires C % 45.73, H % 4.17, N % 9.27; found C % 45.72, H % 4.04, N % 9.16; FTIR (Nujol mull, cm<sup>-1</sup>): ν 2725 (m), 2671 (m), 1596 (s), 1573 (m), 1299 (m), 1281 (s), 1272 (s), 1183 (m), 1050 (m), 1019 (m), 896 (m, UO<sub>2</sub> asymmetric stretch), 722 (m); FTIR (C<sub>6</sub>H<sub>6</sub>, cm<sup>-1</sup>): ν 3375 (w), 2958 (m), 2922 (m), 2872 (w), 2862 (w), 2235 (w), 2082 (w), 1596 (vs), 1574 (s), 1357 (w), 1275 (s), 1218 (w), 1181 (w), 898 (s, UO<sub>2</sub> asymmetric stretch) 839 (w), 821 (w), 800 (w), 764 (w), 725 (w).

**3(Pb)-py** - To a solution of UO<sub>2</sub>(H<sub>2</sub>L<sup>Me</sup>) (100 mg, 94 μmol) in *d*<sub>5</sub>-pyridine (0.5 mL) was added a solution of Pb{N(SiMe<sub>3</sub>)<sub>2</sub>}<sub>2</sub> (50 mg, 94 μmol) in *d*<sub>5</sub>-pyridine (0.5 mL). Single crystals were isolated by dropwise addition of the reaction mixture to hexane (5 mL) resulting in the precipitation of **3(Pb)-py** as dark red needles (81 mg, 59 μmol, 63%).

<sup>1</sup>H NMR (500 MHz, Pyridine-*d*<sub>5</sub>): δ 9.23 (s, 2H, 2 x CH<sub>im</sub>), 8.47 (s, 2H, 2 x CH<sub>im</sub>), 7.30 (d,  $J = 3.3$  Hz, 2H), 6.81 (d,  $J = 3.3$  Hz, 2H), 6.66 (s, 2H, CH<sub>Ar</sub>), 6.56 (s, 2H, CH<sub>Ar</sub>), 6.52 (d,  $J = 3.3$  Hz, 2H), 6.22 (d,  $J = 3.3$  Hz, 2H), 2.17 (s, 6H, 2 x CH<sub>3</sub>), 1.93 (s, 3H, CH<sub>3</sub>), 1.84 (s, 6H, 2 x CH<sub>3</sub>), 1.81 (s, 3H, CH<sub>3</sub>), 1.34 (s, 3H, CH<sub>3</sub>), 1.27 (s, 3H, CH<sub>3</sub>) ppm; <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, Pyridine-*d*<sub>5</sub>): δ 164.7 (C<sub>q</sub>), 163.1 (CH), 157.8 (CH), 157.6 (C<sub>q</sub>), 145.1 (C<sub>q</sub>), 144.3 (C<sub>q</sub>), 139.6 (C<sub>q</sub>), 139.1 (C<sub>q</sub>), 134.4 (C<sub>q</sub>), 131.9 (C<sub>q</sub>), 126.9 (CH), 123.0 (CH), 121.6 (CH), 110.7 (CH), 108.9 (CH), 41.6 (C<sub>q</sub>), 39.5 (C<sub>q</sub>), 36.0 (CH<sub>3</sub>), 33.5 (CH<sub>3</sub>), 30.4 (CH<sub>3</sub>), 29.9 (CH<sub>3</sub>), 19.6 (CH<sub>3</sub>), 19.2 (CH<sub>3</sub>) ppm; IR (Nujol Mull, cm<sup>-1</sup>): ν 1600 (s), 1577 (s), 1301 (m), 1271 (s), 1219 (w), 1175 (w), 1113 (w), 1051 (m), 1018 (m), 908 (m, UO<sub>2</sub> asymmetric stretch), 888 (m), 837 (w), 800 (w), 722 (w), 700 (w); FTIR (C<sub>6</sub>H<sub>6</sub>, cm<sup>-1</sup>): ν 2973 (w), 1594 (vs), 1573 (s), 1438 (w), 1278 (s), 1260 (m), 1052 (m), 923 (w), 895 (m, UO<sub>2</sub> asymmetric stretch).

**4(Pb)-py** - To a solution of **3(Pb)-THF** (10 mg, 8.3 μmol) in C<sub>6</sub>D<sub>6</sub> (1 mL) was added pyridine *N*-oxide (1 mg, 10 μmol). Immediate precipitation of a brown precipitate was observed. This was centrifuged and the mother liquor removed before washing the solids with Et<sub>2</sub>O. The isolated precipitate was dried under



reduced pressure to yield a light brown powder (8 mg, 6.5 μmol, 78%). Single crystals were grown from vapour diffusion of hexane into a pyridine solution of **4**(Pb).

$^1\text{H}$  NMR (500 MHz, Pyridine- $d_5$ ):  $\delta$  9.63 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 9.14 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 8.86 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 8.12 (dd,  $J = 7.6$  Hz, 1H,  $\text{CH}_{\text{py}}$ ), 7.99 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 7.74 (s, 2H, 2 x  $\text{CH}_{\text{py}}$ ), 7.71 (dd,  $J = 7.6$  Hz, 2H, 2 x  $\text{CH}_{\text{py}}$ ), 7.62 (s, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 7.36 (s, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 7.08 (s, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 6.65 (s, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 2.55 (s, 6H, 2 x  $\text{CH}_3$ ), 2.49 (s, 3H,  $\text{CH}_3$ ), 2.32 (s, 3H,  $\text{CH}_3$ ), 2.28 (s, 6H, 2 x  $\text{CH}_3$ ), 1.87 (s, 3H,  $\text{CH}_3$ ), 1.58 (s, 3H,  $\text{CH}_3$ ) ppm;  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz, Pyridine- $d_5$ ):  $\delta$  165.0 ( $\text{C}_q$ ), 161.4 ( $\text{C}_q$ ), 161.4 (CH), 154.9 (CH), 147.4 ( $\text{C}_q$ ), 145.9 ( $\text{C}_q$ ), 139.7 ( $\text{C}_q$ ), 139.6 (CH), 135.0 ( $\text{C}_q$ ), 132.7 ( $\text{C}_q$ ), 126.4 (CH), 125.1 (CH), 122.7 (CH), 122.3 (CH), 121.1 (CH), 110.6 (CH), 109.1 (CH), 41.8 ( $\text{C}_q$ ), 39.6 ( $\text{C}_q$ ), 38.5 ( $\text{CH}_3$ ), 37.0 ( $\text{CH}_3$ ), 28.3 ( $\text{CH}_3$ ), 27.4 ( $\text{CH}_3$ ), 19.43 ( $\text{CH}_3$ ), 19.1 ( $\text{CH}_3$ ) ppm; Analysis Calcd.  $\text{C}_{47}\text{H}_{45}\text{N}_9\text{O}_3\text{PbU}$  (1229.17) requires C % 45.93, H % 3.69, N % 10.26; found C % 45.64, H % 3.55, N % 9.99.

**5** - To a solution of  $\text{H}_4\text{L}^{\text{Me}}$  (500 mg, 0.76 mmol, 1 eq.) in THF (20 mL) was added a solution of  $\text{Pb}\{\text{N}(\text{SiMe}_3)_2\}_2$  (800mg, 1.52 mmol, 2 eq.) in THF (10 mL). The solution immediately darkened to yield a deep red solution which was stirred for 8 h before the solvent was removed under reduced pressure and the residue washed with hexane. Recrystallization from THF/hexane yielded **5** as an orange/red powder (735 mg, 0.69mmol, 90%). Single crystals suitable for X-ray diffraction were grown by slow diffusion of hexane into a concentration solution of **5** in pyridine.

$^1\text{H}$  NMR (500 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  8.24 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 8.15 (s, 2H, 2 x  $\text{CH}_{\text{im}}$ ), 6.71 (d,  $J = 3.5$  Hz, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 6.62 (d,  $J = 3.5$  Hz, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 6.48 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 6.37 (d,  $J = 3.5$  Hz, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 6.34 (d,  $J = 3.5$  Hz, 2H, 2 x  $\text{CH}_{\text{pyrrole}}$ ), 6.22 (s, 2H, 2 x  $\text{CH}_{\text{Ar}}$ ), 2.03 (s, 6H, 2 x  $\text{CH}_3$ ), 2.02 (s, 6H, 2 x  $\text{CH}_3$ ), 1.76 (s, 3H,  $\text{CH}_3$ ), 1.73 (s, 3H,  $\text{CH}_3$ ), 1.71 (s, 3H,  $\text{CH}_3$ ), 1.47 (s, 3H,  $\text{CH}_3$ ) ppm;  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  162.8 (CH), 159.2 (CH), 158.9 ( $\text{C}_q$ ), 158.4 ( $\text{C}_q$ ), 142.4 ( $\text{C}_q$ ), 140.9 ( $\text{C}_q$ ), 140.7 ( $\text{C}_q$ ), 140.0 ( $\text{C}_q$ ), 133.1 ( $\text{C}_q$ ), 132.5 ( $\text{C}_q$ ), 123.8 (CH), 123.3 (CH), 122.5 (CH), 120.5 (CH), 110.5 (CH), 110.0 (CH), 39.6 ( $\text{C}_q$ ), 39.4 ( $\text{C}_q$ ), 34.9 ( $\text{CH}_3$ ), 32.3 ( $\text{CH}_3$ ), 24.1 ( $\text{CH}_3$ ), 24.0 ( $\text{CH}_3$ ), 19.5 ( $\text{CH}_3$ ), 19.3 ( $\text{CH}_3$ ) ppm; Analysis Calcd.  $\text{C}_{46}\text{H}_{48}\text{N}_8\text{Pb}_2$  (1143.35) requires C % 48.32, H % 4.23, N % 9.80; found C % 48.50, H % 4.17, N % 9.58.


## Acknowledgements

The authors thank the University of Edinburgh and the EPSRC (Grant numbers EP/M010554/1 and EP/H004823/1) for funding.

## Notes and references

1. G. R. Choppin, *J. Radioanal. Nucl. Chem.*, 2007, **273**, 695-703.
2. P. L. Arnold, G. M. Jones, S. O. Odoh, G. Schreckenbach, N. Magnani and J. B. Love, *Nat. Chem.*, 2012, **4**, 221-227.
3. J. N. Mathur, M. S. Murali and K. L. Nash, *Solvent Extr. Ion Exch.*, 2001, **19**, 357-390; E. P. Horwitz, R. Chiarizia, M. L.

- Dietz and H. Diamond, *Anal. Chim. Acta*, 1993, **281**, 361-372; Y. Sakamura, T. Hijikata, K. Kinoshita, T. Inoue, T. S. Storvick, C. L. Krueger, J. J. Roy, D. L. Grimmer, S. P. Fusselman and R. L. Gay, *J. Alloys Compd.*, 1998, **271**, 592-596.
4. P. L. Arnold, J. B. Love and D. Patel, *Coord. Chem. Rev.*, 2009, **253**, 1973-1978; J. C. Renshaw, L. J. C. Butchins, F. R. Livens, I. May, J. M. Charnock and J. R. Lloyd, *Environ. Sci. Technol.*, 2005, **39**, 5657-5660.
5. M. Bühl and G. Schreckenbach, *Inorg. Chem.*, 2010, **49**, 3821-3827.
6. P. L. Arnold, A.-F. Pecharman and J. B. Love, *Angew. Chem.-Int. Edit.*, 2011, **50**, 9456-9458; P. L. Arnold, E. Hollis, F. J. White, N. Magnani, R. Caciuffo and J. B. Love, *Angew. Chem.-Int. Edit.*, 2011, **50**, 887-890; G. M. Jones, P. L. Arnold and J. B. Love, *Chem. Eur. J.*, 2013, **19**, 10287-10294.
7. P. L. Arnold, A.-F. Pecharman, E. Hollis, A. Yahia, L. Maron, S. Parsons and J. B. Love, *Nat. Chem.*, 2010, **2**, 1056-1061.
8. P. L. Arnold, A.-F. Pecharman, R. M. Lord, G. M. Jones, E. Hollis, G. S. Nichol, L. Maron, J. Fang, T. Davin and J. B. Love, *Inorg. Chem.*, 2015, **54**, 3702-3710.
9. P. L. Arnold, D. Patel, A. J. Blake, C. Wilson and J. B. Love, *J. Am. Chem. Soc.*, 2006, **128**, 9610-9611.
10. P. L. Arnold, G. M. Jones, Q.-J. Pan, G. Schreckenbach and J. B. Love, *Dalton Trans.*, 2012, **41**, 6595-6597.
11. X. J. Zheng, N. L. Bell, C. J. Stevens, Y. X. Zhong, G. Schreckenbach, P. L. Arnold, J. B. Love and Q. J. Pan, *Submitted*, 2016
12. G. Givaja, M. Volpe, J. W. Leeland, M. A. Edwards, T. K. Young, S. B. Darby, S. D. Reid, A. J. Blake, C. Wilson, J. Wolowska, E. J. L. McInnes, M. Schröder and J. B. Love, *Chem. Eur. J.*, 2007, **13**, 3707-3723.
13. F. Allen, *Acta Crystallogr., Sect. B: Struct. Sci.*, 2002, **58**, 380-388.
14. P. L. Arnold, A. J. Blake, C. Wilson and J. B. Love, *Inorg. Chem.*, 2004, **43**, 8206-8208.
15. E. A. Connolly, J. W. Leeland and J. B. Love, *Inorg. Chem.*, 2016, **55**, 840-847.
16. *Handbook of Chemistry and Physics*, 61st edn., 1981.
17. P. L. Arnold, D. Patel, A.-F. Pecharman, C. Wilson and J. B. Love, *Dalton Trans.*, 2010, **39**, 3501-3508.
18. T. W. Hayton and G. Wu, *Inorg. Chem.*, 2009, **48**, 3065-3072.
19. M. J. Sarsfield and M. Helliwell, *J. Am. Chem. Soc.*, 2004, **126**, 1036-1037.
20. M. Zegke, G. S. Nichol, P. L. Arnold and J. B. Love, *Chem. Commun.*, 2015, **51**, 5876-5879.
21. T. Heidemann and S. Mathur, *Eur. J. Inorg. Chem.*, 2014, **2014**, 506-510.



254x133mm (120 x 120 DPI)