

Chemical Science

rsc.li/chemical-science



ISSN 2041-6539



ROYAL SOCIETY
OF CHEMISTRY

Celebrating
IYPT 2019

EDGE ARTICLE

Himanshu Mishra *et al.*

The chemical reactions in electrosprays of water do not always correspond to those at the pristine air–water interface

Cite this: *Chem. Sci.*, 2019, 10, 2566

All publication charges for this article have been paid for by the Royal Society of Chemistry

The chemical reactions in electrosprays of water do not always correspond to those at the pristine air–water interface†

Adair Gallo, Jr.,^{abc} Andreia S. F. Farinha,^{abc} Miguel Dinis,^{ad} Abdul-Hamid Emwas,^{ae} Adriano Santana,^{abc} Robert J. Nielsen,^f William A. Goddard, III^{fg} and Himanshu Mishra^{id*abc}

The recent application of electrosprays to characterize the air–water interface, along with the reports on dramatically accelerated chemical reactions in aqueous electrosprays, have sparked a broad interest. Herein, we report on complementary laboratory and *in silico* experiments tracking the oligomerization of isoprene, an important biogenic gas, in electrosprays and isoprene–water emulsions to differentiate the contributions of interfacial effects from those of high voltages leading to charge-separation and concentration of reactants in the electrosprays. To this end, we employed electrospray ionization mass spectrometry, proton nuclear magnetic resonance, *ab initio* calculations and molecular dynamics simulations. We found that the oligomerization of isoprene in aqueous electrosprays involved minimally hydrated and highly reactive hydronium ions. Those conditions, however, are non-existent at pristine air–water interfaces and oil–water emulsions under normal temperature and pressure. Thus, electrosprays should be complemented with surface-specific platforms and theoretical methods to reliably investigate chemistries at the pristine air–water interface.

Received 11th December 2018
Accepted 21st December 2018

DOI: 10.1039/c8sc05538f

rsc.li/chemical-science

The air–water interface plays a critical role in numerous natural and applied contexts, such as atmospheric chemistries,^{1–3} precipitation,⁴ spray coatings,⁵ and materials synthesis.^{6,7} Indeed, it has been hypothesized that microdroplets generated during the splashing of waves in oceans could have been the chemical reactors leading to the origin of life.^{8–10} Despite its ubiquity and importance, a variety of fundamental phenomena at the air–water interface remain incompletely understood, such as the specific adsorption of ions^{11–13} and chemistries therein.^{10,14–20} The interfacial region, with a typical thickness δ_0 , ≈ 0.5 nm, separates the gas-phase (vapor) from the condensed phase (water), two drastically different regions in terms of hydration – reactions spontaneous in one phase are forbidden in the other.²¹ In fact, the chemical activities of species at the air–water interface can depart significantly from those in the bulk, as has been demonstrated by surface-specific techniques,

including vibrational second harmonic generation and sum frequency generation,^{20,22,23} and polarization-modulated infrared absorption reflection spectroscopy,^{10,13} and indirect approaches, including NMR²⁴ and confocal fluorescence microscopy.²⁵ Even though vibrational spectroscopy-based techniques report directly on thermodynamic properties of the air–water interface, they suffer from interpretational ambiguities and limitations due to low signal-to-noise ratios.^{26–32} Thus, new techniques with higher sensitivity and unambiguous response are needed to help resolve the poorly understood features of the air–water interface while providing benchmarks to judge previous interpretations.³³ In this work, we assess the application of electrospray ionization mass spectrometry (ESIMS) to unravel the thermodynamic properties of pristine air–water interface (Henceforth, we will use the qualifier ‘pristine’ to refer to the air–water interface that is not under the influence of any external sources/agents, such as electrical voltage or a drying gas).

In the recent years, ESIMS, which has been widely used to characterize ionic/molecular species in polar/apolar solvents,³⁴ has been adapted to investigate the pristine air–water interface. In the standard configuration, ESIMS experiments entail the formation of electrosprays by the application of electrical potential and/or pneumatic pressure leading microscale droplets with excess electrical charge; those microdroplets pass through a glass/metallic capillary maintained at elevated

^aKing Abdullah University of Science and Technology (KAUST), Saudi Arabia. E-mail: himanshu.mishra@kaust.edu.sa

^bWater Desalination and Reuse Center (WDRC), Saudi Arabia

^cDivision of Biological and Environmental Sciences (BESE), Saudi Arabia

^dKAUST Catalysis Center (KCC), Saudi Arabia

^eImaging and Characterization Core Laboratory, Thuwal 23955-6900, Saudi Arabia

^fMaterials and Process Simulation Center, California Institute of Technology, Pasadena, CA 91125, USA

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8sc05538f



temperature (~ 473 K) to evaporate the solvent and facilitate the mass spectrometric detection of analytes downstream.^{34–38} In the experiments designed to investigate chemistries at the air–water interface, electrosprays containing one or more reactant(s) are intersected with gases or other electrosprays containing other reactant(s) followed by mass spectrometric detection. For instance, using this platform, thermodynamic properties of the pristine air–water interface have been explored under ambient conditions, including the relative concentrations of interfacial hydronium and hydroxide ions and their activities^{39–43} leading to interpretations that have elicited scientific debate.^{11,12,17–19,44,45} Further, by intersecting electrosprays of pH-adjusted water with gaseous organic acids,⁴⁰ isoprene,⁴¹ and terpenes,^{40,41,46,47} researchers observed instantaneous protonation (<1 ms), and in some cases oligomerization of organics, which led them to conclude that as the bulk acidity of water approaches $\text{pH} \leq 3.6$, the pristine air–water interface behaves as a superacid. While a clear understanding of the emergence of the putative superacidity at the air–water interface is unavailable, we note that in the condensed phase proton-catalyzed oligomerization of isoprene (or olefins in general) requires 60–80% concentrated H_2SO_4 solutions ($\text{pH} < -0.5$).⁴⁸ Similar rate enhancements in aqueous electrosprays have also been observed for the syntheses of abiotic sugar phosphates,^{15,49} the Pomeranz–Fritsch synthesis of isoquinoline,⁵⁰ the reaction between *o*-phthalaldehyde and alanine,⁵¹ and the ozonation of oleic acid,⁵² among others.^{16,53} Herein, we assess the relationships between the chemistries observed in aqueous electrosprays to those at pristine air–water interfaces; we also seek to decouple the factors that contribute to the mechanisms underlying reported dramatic rate enhancements by addressing the following questions:

- (i) Do aqueous electrospray-based platforms report on thermodynamic properties of the pristine air–water interface?
- (ii) Do accelerated reactions in aqueous electrosprays arise only from the significant enhancement of the hydrophobe–water (air–water) interfacial area? If yes, the mechanisms underlying the dramatic rate enhancements therein should be insightful in explaining the accelerated organic reactions in oil–water emulsions also referred to as ‘on-water’ catalysis.^{54–57}
- (iii) Are the rate accelerations in aqueous electrosprays driven solely by the non-equilibrium conditions therein, especially the enhanced concentration of reactants in the micro-/nano-droplets due to the evaporation of water?^{58–61}
- (iv) Are gas-phase reactions implicated in the acceleration of chemical reactions in aqueous electrosprays?^{36,44,50,62–64}

To address those questions, we investigated the oligomerization of isoprene by proton nuclear magnetic resonance ($^1\text{H-NMR}$), a non-invasive technique, as a complementary platform to the ESIMS. Questions (i and ii) were addressed by comparing the effects of enhancing the water–hydrophobe interfacial area in both liquid–vapor and liquid–liquid systems; questions (iii and iv) were addressed by varying the capillary voltages, ionic strengths of the aqueous solutions electrosprayed and intersected with gas-phase isoprene, and $^1\text{H-NMR}$ analysis of condensed vapor from the electrosprays. To highlight the role of hydration in electrosprays, we performed

quantum mechanical calculations employing density functional theory (M06 flavor).

Materials and methods

In our experiments, we used isoprene (99% purity from Sigma-Aldrich), Mili-Q deionized water (18 $\text{M}\Omega$ m resistivity), D_2O (99.9% purity from Sigma Aldrich), ethanol (absolute from Merck Millipore), acetone (HPLC standard from VWR Chemicals), NaCl (>99% purity from Sigma Aldrich), HCl (36.9% concentration from Fisher Scientific), DCl (35% concentration 99% deuterium purity from Sigma Aldrich), and NaOH (>97% purity from Sigma Aldrich) to adjust the pH and ionic strengths.

ESIMS

All experiments were conducted in a commercial Thermo Scientific – LCQ Fleet electrospray ionization mass spectrometer in the positive ion mode, where a DC potential of 6–8 kV was applied to the needle, the tube lens voltage was 75 V, the sheath gas flow rate was 10 arb, the pressure was 1.2 torr at the convection gauge and 0.8×10^{-5} torr at the ion-gauge, the flow rates of analytes were controlled by a calibrated syringe pump and ranged between 1–10 $\mu\text{L min}^{-1}$, the distances from the ion source and the inlet to the mass spectrometer were ~ 2 cm, and the distance between the electrospray and the tube ejecting isoprene was 1 cm.

$^1\text{H-NMR}$

All NMR spectra were acquired using a Bruker 700 AVANAC III spectrometer equipped with a Bruker CP TCI multinuclear CryoProbe (BrukerBioSpin, Rheinstetten, Germany); Bruker Topspin 2.1 software was used to collect and analyze the data. We transferred 100 μL of the (A1) samples into 5 mm NMR tubes, followed by 600 μL of deuterated chloroform (CDCl_3). The $^1\text{H-NMR}$ spectra were recorded at 298 K by collecting 32 scans with a recycle delay of 5 s, using a standard 1D 90° pulse sequence and standard (zg) program from the Bruker pulse library. The chemical shifts were adjusted using tetramethylsilane (TMS) as an internal chemical shift reference. The (A) samples and a sample of as-purchased isoprene (B) were prepared by transferring 100 μL of each to 5 mm NMR tubes, and then adding 550 μL of deuterated water D_2O to the NMR tubes. The $^1\text{H-NMR}$ spectra were recorded by collecting 512 scans with a recycle delay time of 5 s, using an excitation sculpting pulse sequence (zgesgp) program from the Bruker pulse library. The chemical shifts were adjusted using 3-trimethylsilylpropane sulfonic acid (DSS) as an internal chemical shift reference. The free induction decay (FID) data were collected at a spectral width of 16 ppm into 64 k data points. The FID signals were amplified by an exponential line-broadening factor of 1 Hz before Fourier transformation.

Computational methods

To gain molecular-level insights into the protonation and oligomerization of the isoprene in our experiments, we performed (and benchmarked) DFT calculations at the following levels: (A)



geometry and transition state optimization with M06/6-311G* followed by single point calculations with a larger basis set (6-311++G**);⁶⁵ (B) geometry and transition state optimization with M06/6-311++G**; (C) geometry and transition state optimization with M06/6-311++G**, followed by single point calculations with CCSD(T).⁶⁶ We calculated the internal reaction coordinate (IRC) along the potential energy surface connecting adduct (A), transition state (TS), and product (P). Frequency calculations of the optimized geometries yielded Hessians with zero and one imaginary frequency, respectively, for the minima and the transition state structures. The force constants from the frequency calculations on the TS structures were subsequently used to calculate the IRC path along the forward and backward direction of the transition vector, connecting transition state to the minima.^{67,68} The vibrational frequencies from the Hessians were also used to provide the zero-point energies and vibrational contributions to the enthalpies and entropies. The free energies of isoprene at 1 atm were calculated using statistical mechanics for ideal gases.

Molecular simulations

Born–Oppenheimer molecular dynamics (BOMD)⁶⁶ simulations were carried out with the M06 functional and 6-311G(d,p) basis set in the NVT ensemble coupled with a Nose–Hoover thermostat⁶⁹ at 298 K. The total simulation times were 5 ps with 1 fs timesteps.

We used the Gaussian 09 software package⁷⁰ to perform these calculations and simulations.

Results

We investigated chemical reactions between pH-adjusted water and isoprene (C₅H₈, 2-methyl-1,3-butadiene, MW = 68 amu, and solubility in water, $S = 0.7 \text{ g L}^{-1}$ at normal temperature and pressure (NTP): 293 K and 1 atm). We chose to examine reactions of isoprene because (i) we wanted to reproduce previous experimental results to ensure a clear comparison, (ii) isoprene is an important biogenic gas whose fate in the atmosphere is not completely understood,^{1,41,71,72} and (iii) we could investigate chemistries in electrosprays and emulsions by taking advantage of the low boiling point of isoprene ($T_b = 307 \text{ K}$) and the high vapor pressure at NTP ($p = 61 \text{ kPa}$).⁷³

As delineated in Fig. 1 and summarized in Table 1, we report on the following sets of ESIMS (detection limit = $\sim 1 \text{ nM}$) and ¹H-NMR (detection limit = $\sim 10 \text{ }\mu\text{M}$) experiments:

(A) Liquid–liquid collisions

At NTP, we combined liquid isoprene with pH-adjusted H₂O or D₂O, $1 \leq \text{pH} \leq 13$, in a volumetric ratio 1 : 6 : 3 (isoprene : water : air), agitated the emulsions at 1200 rpm in a vortexer for 6, 60, or 360 minutes, and analyzed the organic phases after phase-separation by ESIMS and ¹H-NMR. Since the air in the reaction vessels was saturated with isoprene, those experiments also ensured the presence of the products of reactions between the gas-phase isoprene and pH-adjusted water in the organic phase.



Fig. 1 Summary of the experiments (A), (B), and (C) reported in this work. (A) Liquid–liquid collisions: mixtures of isoprene, pH-adjusted water, and air in the volumetric ratio 1 : 6 : 3 was agitated at 1200 rpm (for 6, 60, 360 minutes) followed by ESIMS analysis of the organic phase. (B) As-purchased liquid isoprene was injected directly in the ESIMS. (C) Gas–liquid collisions: electrospays of water (pH range 1–13) were collided with a stream of air carrying isoprene gas, followed by mass spectrometric detection (Methods).

(A1) Condensed vapors from electrosprays of organic phase from (A). After the liquid–liquid collision reactions (A) were over and the organic phases were electrosprayed in ESIMS for characterization, we condensed the sprays and analyzed them by ¹H-NMR. The ¹H-NMR-based investigation of the reaction products from experiments (A) before and after electrospraying was carried out to pinpoint the effects, if any, of electrospraying on the formation of the products.

(B) Pure components

We analyzed as-purchased isoprene, acetone, and ethanol by ESIMS and ¹H-NMR.

(C) Gas–liquid collisions

We created electrosprays of aqueous solutions with varying ionic strengths and pH, and intersected them with a stream of gas-phase isoprene (0.48 g min^{-1} carried by N₂ gas flowing at 600 mL min^{-1} , *i.e.* isoprene gas concentration was 800 mg L^{-1}) followed by mass spectrometric detection (Methods).

Hereafter, throughout the paper, we will refer to our experiments on the liquid–liquid collisions as (A), condensed vapors from the electrosprays (A1), pure isoprene as (B), and gas–liquid collisions in the ESIMS as (C) (Fig. 1, Table 1).

Intriguingly, the ESIMS spectra from the above-mentioned experiments (A) at pH = 1, (B), and (C) at pH = 1 were nearly identical after normalizing with the maximum intensity (Fig. 2A–C and Section S_a†). The positions of the main peaks in the mass spectra fitted the general formula, [(Isop)_n·H]⁺, which corresponded to covalently bonded oligomers of isoprene with one excess proton. In Section S_b and Fig. S1† we present the evidence proving that the peaks did not correspond to physisorbed clusters. In experiments (A), the mass spectra remained



Table 1 Experimental summary

	(A) Liquid–liquid collisions	(B) Pure components	(C) Gas–liquid collisions
ESIMS	Water(L)–isoprene(L)	Isoprene(L), acetone(L), ethanol(L)	Isoprene(G)–water(L)
	Organics injected	Components injected	Water injected
pH	1–13	—	1–13
pNaCl	—	—	1–9
Shaking time	6, 60, 360 min	—	—
Voltage	6 kV	6 kV	6–8 kV
Capillary temperature	150 °C	30–330 °C	150 °C
¹H-NMR	Organics from (A) and condensed vapors (A1)	Isoprene(L)	—
pH	1.5	—	—
Shaking time	6, 60, 360 min	—	—
Aqueous phase	D ₂ O, H ₂ O	—	—

the same as the duration of mixing varied from 6 min to 6 h (Fig. S2†). We also observed numerous secondary peaks between the primary $[(\text{Isop})_n \cdot \text{H}]^+$ peaks. While a detailed characterization of those peaks falls beyond the scope of this work, we speculate that they are composed of C1–C4 fragments covalently bonded with the $[(\text{Isop})_n \cdot \text{H}]^+$ oligomers, for example $m/z = 233$ (Fig. 2C and S3†).

Next, we investigated the role of water pH on the oligomerization of isoprene in experiments (A) and (C). When the products were characterized by ESIMS, we noticed that the oligomers $[(\text{Isop})_n \cdot \text{H}]^+$ appeared when the aqueous phase had $\text{pH} \leq 3.6$ (Fig. 3). Those observations have been reported

previously^{18,40,41,46,47} and ascribed to the superacidity of the air–water interface at $\text{pH} \leq 3.6$. However, we also found that the ESIMS spectra from both experiments, (A) and (C), yielded oligomers $[(\text{Isop})_n \cdot \text{H}]^+$ for the acidic, basic, and pH-neutral salty solutions (Fig. 3; compare Fig. 2A and C with Fig. S3† panels C1, C2, and C3). We note that for electrosprays produced from $\text{pH} > 7$ water and salty water, counterions, such as Na^+ , could influence the fate of reactions, but we have not investigated those factors.

In experiments (A), after the emulsions comprising liquid isoprene, liquid water at $\text{pH} = 1.5$, and air (containing saturated gaseous isoprene) were vigorously mixed (for 6 min, 60 min, and




Fig. 2 ESIMS spectra for sets of experiments A, B, and C: the dominant peaks correspond to protonated oligomers of isoprene, $[(\text{Isop})_n \cdot \text{H}]^+$, and the secondary peaks correspond to fragments of the isoprene molecules bonded to the primary oligomers. (A) ESIMS spectra of the organic phase from the emulsion of liquid isoprene in water at $\text{pH} = 1$ and air (1 : 6 : 3 v/v/v) that was agitated at 1200 rpm for 360 minutes. (B) ESIMS spectra of as-purchased liquid isoprene. (C) ESIMS spectra of products of gas–liquid collisions between water ($\text{pH} = 1$) and gas-phase isoprene (Methods).






Fig. 4 (A) ^1H -NMR spectra of the organic phase after shaking liquid isoprene with pH 1.5 water and air in a volumetric ratio of 1 : 6 : 3 for 60 minutes. (B) ^1H -NMR spectra of as-purchased isoprene. (A1) ^1H -NMR spectra of the condensed exhaust from the electrospays of the organic phase after the liquid–liquid collision (A) experiments. The nearly identical spectra for experimental sets (A) and (B) demonstrate that there was no detectable oligomerization of isoprene during the vigorous shaking of emulsions comprised of liquid isoprene with pH 1.5 water and air in a volumetric ratio of 1 : 6 : 3 (NMR resolution $\sim 10\ \mu\text{M}$).

water-alcohol mixtures,⁸² among others;^{64,77} these proton-transfer reactions also underlie the basis of the Proton Transfer Reaction Mass Spectrometry (PTRMS) that has been used to detect trace gases in the atmosphere.^{77,83}

In all those mechanisms, the initial ionic strength of water (acidic, basic, or salty) and electrical voltage were crucial for the formation of a stable stream of charged microdroplets – the higher the ionic strength of solutions, the lower the requirement for the electrical voltage.^{35,36,84–86} In fact, due to the electrochemical reactions at the electro spray needle under the influence of high electric fields, the electro sprayed droplets from a positively charged capillary should contain more positive ions than in the bulk^{58,63,85,87,88} (Section S_c, Fig. S5[†]). Interestingly, for pH-adjusted water electro sprayed at 6 kV, we detected oligomers (Fig. 3C) when $\text{pH} \leq 3.6$ or $\text{pH} > 12$, whereas for the NaCl solutions, we observed oligomers at concentrations as low as $10^{-9}\ \text{M}$ ($\text{pNa} = 9$). Yet, the higher intensities of the $[(\text{Isop})_n \cdot \text{H}]^+$ at $\text{pH} \leq 3.6$ in comparison to the salty solutions (Fig. 3C) indicate that the proposed mechanism M_1 is unlikely to play a crucial role in the case of gaseous isoprene interacting with electrospays of water.

Following our logical exclusion of mechanism M_1 , we are left with mechanisms M_2 and M_3 , *i.e.* did the reactions take place on the surface of electro sprayed water droplets or in the gas-phase? Whether or not the electro spray spectra represent

the solvent- or gas-phase chemistries/characteristics is a much-debated matter and case-specific.^{36,38,76,89,90} Obviously, the answer would have a bearing on the questions (ii–iv) outlined above, because the kinetics and thermodynamics of reactions in bulk and gas-phase differ dramatically.²¹ Recently, Silveira and co-workers employed cryogenic ion mobility mass spectrometry to demonstrate the effects of rapid dehydration on the structures of undecapeptide substance during the final stages of electro spray ionization.⁹⁰ To further clarify the role of hydration on the protonation and oligomerization of isoprene in our experiments with mildly acidic water (electrospays and emulsions), we carried out quantum mechanics calculations.

Computational calculations

To gain qualitative molecular-scale insights into our experiments with mildly acidic water, we carried out density functional theory (DFT) calculations (Computational methods). M06, a hybrid meta-generalized gradient approximation (*meta*-GGA) functional, is known to provide an accurate description of the ground-state thermochemistry, thermochemical kinetics, and transition state structures and energies for a wide series of organic reactions.^{91–96} We have previously confirmed that the binding energies of water clusters, $(\text{H}_2\text{O})_n$ (range $n = 2–8, 20$),



along with the hydration and neutralization energies of hydroxide and hydronium ions using DFT functionals (M06, M06-2X, M06-L, B3LYP, X3LYP) are in excellent agreement with the high-level theory (CCSD(T)/aug-cc-p VDZ level), both with and without the basis set superposition error correction.⁹¹ For instance, our *ab initio* predictions of the proton transfer thermodynamics between $\text{H}_3\text{O}^+(\text{g})$ and $\text{Isop}(\text{g})$ was $\Delta G^0 = -30.7 \text{ kcal mol}^{-1}$, in accordance with the difference in the experimental gas-phase basicity (GB) of H_2O ($\text{GB}_{\text{H}_2\text{O}} = -157.7 \text{ kcal mol}^{-1}$) and Isop ($\text{GB}_{\text{ISO}} = -190.6 \text{ kcal mol}^{-1}$), $\Delta \text{GB} = -32.9 \text{ kcal mol}^{-1}$ (Fig. S6†).⁹⁷ Furthermore, the *trans*- or *cis*- $\text{Isop}(\text{g})$ spontaneously added onto ($\text{Isop} \cdot \text{H}^+ + \text{H}_2\text{O}$), leading to cyclic ($\Delta G^0 = -40 \text{ kcal mol}^{-1}$) or acyclic monoterpenes ($\Delta G^0 = -9 \text{ kcal mol}^{-1}$), (Fig. S7 and S8†), as also noted by other researchers.^{48,98} Next, we calculated the kinetic barriers for protonation and oligomerization of Isop on a small water cluster containing an extra proton, $(\text{H}_2\text{O})_3 \cdot \text{H}^+$, as representative of our electrospray experiments and a larger cluster, $(\text{H}_2\text{O})_{36} \cdot \text{H}^+$, representing the pristine air–water interface. The sizes of the clusters were guided, in principle, by previous experimental and theoretical work on the hydration of protons,^{99–101} showing the asymptotic stabilization of a proton with increasing cluster size, and limited by computational expense. The initial geometry of the smaller cluster was obtained from the Cambridge Cluster Database,¹⁰² while the larger cluster geometry was obtained from an SPC/E bulk water box equilibrated at 298 K and at 1 atm pressure yielding the bulk density of 1 g cm^{-3} . We took a cluster of 36 water molecules from the equilibrated water box as a surrogate for the air–water interface. We added a proton to this cluster and applied Born–Oppenheimer Molecular Dynamics (BOMD) simulations

(Computational methods) to obtain various low-energy conformers (Table S2, Section S_d†). We chose a cluster with the proton on the surface that facilitated the subsequent study of the activation barriers through DFT calculations (Computational methods).

In the case of $(\text{H}_2\text{O})_3 \cdot \text{H}^+$, representative of electrosprays, the proton was extremely reactive/acidic due to insufficient hydration – the incipient isoprene molecule (Isop) fell into a shallow potential well forming an adduct (Fig. 5). The free energy barrier for proton transfer from $(\text{H}_2\text{O})_3 \cdot \text{H}^+$ to $\text{Isop}(\text{g})$ was $\Delta G^\ddagger = 6.9 \text{ kcal mol}^{-1}$, and the barrier to the oligomerization with another free $\text{Isop}(\text{g})$ was $\Delta G^\ddagger = 2.1 \text{ kcal mol}^{-1}$ (Fig. 5). On the other hand, the predictions for those barriers for protonation and oligomerization of isoprene with the larger water cluster, $(\text{H}_2\text{O})_{36} \cdot \text{H}^+$, representative of the air–water interface, were $\Delta G^\ddagger = 25.5 \text{ kcal mol}^{-1}$ and $\Delta G^\ddagger = 40.2 \text{ kcal mol}^{-1}$, respectively, which are insurmountable under ambient conditions within a 1 ms time-frame (Fig. 6). The stark differences in the free energy landscapes for the interactions of $\text{Isop}(\text{g})$ with minimally hydrated hydronium ions, available in electrosprays, and larger water clusters can simply be understood in terms of the enthalpies of hydration of gas-phase protons that decrease monotonically with the addition of water molecules;^{99–101} the proton activity also decreases commensurately and, in fact, is expected to be even lower for the air–water interface than in the $(\text{H}_2\text{O})_{36} \cdot \text{H}^+$ cluster.

To test the effects of larger basis-sets on these predictions, we compared the results of our M06/6-311+G*/6-311++G** calculations with M06/6-311++G** for the smaller cluster, and found very similar results (Section S_d, Table S1†). Towards further benchmarking, we carried out single point energy




Fig. 5 Quantum mechanics-based free energy and enthalpy landscapes for protonation and oligomerization of isoprene while interacting with a cluster comprising three water molecules and one excess proton, $(\text{H}_2\text{O})_3 \cdot \text{H}^+$. Due to its incomplete hydration (compared to bulk), the proton exhibited extreme acidity. The free energy barrier for the proton transfer from $(\text{H}_2\text{O})_3 \cdot \text{H}^+$ to isoprene(g) was $\Delta G^\ddagger = 6.9 \text{ kcal mol}^{-1}$ and the barrier to subsequent oligomerization with another free isoprene(g) was $\Delta G^\ddagger = 2.1 \text{ kcal mol}^{-1}$, which are easily surmountable under ambient NTP conditions within the timescale of our experiments ($\sim 1 \text{ ms}$). These model predictions support mechanism M₃.



R: reactants
A-I: first adduct
TS-I: first TS
P-I: first protonated Isop.
A-II: second adduct
TS-II: second TS
P-II: Isop.₂ oligomer



Fig. 6 Quantum mechanics-based free energy and enthalpy landscapes for protonation and oligomerization of isoprene on a cluster consisting of 36 water molecules and an excess proton, $(\text{H}_2\text{O})_{36}\cdot\text{H}^+$, representative of very small water droplets. The kinetic barriers preventing proton transfer to isoprene and its subsequent oligomerization were $\Delta G^\ddagger = 25.5 \text{ kcal mol}^{-1}$ and $\Delta G^\ddagger = 40.2 \text{ kcal mol}^{-1}$ respectively, which were insurmountable under ambient NTP conditions within the timescale of our experiments ($\sim 1 \text{ ms}$). The predictions of this model suggest that the reactions of isoprene in electrosprays cannot involve liquid-phase drops. These predictions also support the proposed mechanisms M_3 by ruling out the possible reactions through M_2 .

calculations on the structures optimized at the M06/6-311++G** level with the CCSD(T)-level theory,⁶⁶ and found that the qualitative trends in the reaction kinetics remained unchanged (Table S1†). We note that we could only perform the benchmarking for the smaller system due to the computational expense. Thus, we expect that the relative differences in the barrier heights for the reactions are unlikely to change significantly between DFT/M06 and CC for our systems. Besides, our ¹H-NMR experiments present unambiguous evidence for these predictions. To summarize, our calculations demonstrate that the kinetic barriers for the protonation and oligomerization of isoprene are easily surmountable in the smaller acidic cluster (available in electrosprays only) and prohibited in larger acidic clusters at 298 K and 1 atm (a surrogate for the pristine air-water interface).

Conclusion and outlook

Based on our experimental investigation of oil-water and air-water interfaces of isoprene with pH-adjusted water, analyzed by ESIMS and ¹H-NMR along with quantum mechanical predictions, we address the questions outlined in the introduction as follows:

(i) Aqueous electrosprays do not always report on the thermodynamic properties of pristine air-water interfaces.

(ii) The observed chemical reactions of isoprene in aqueous electrosprays were not driven by the enhancement in the hydrophobe-water interfacial area, as evidenced by the lack thereof in vigorously mixed emulsions of isoprene and pH-adjusted water. Thus, the mechanisms underlying the ‘on-water’ catalysis^{54–57} must be different from those leading to rate accelerations in the aqueous electrosprays.¹⁰³ Electrosprays of water must facilitate additional chemical pathways, such as reactions with partially hydrated (gas-phase) hydroniums, which are not accessible in vigorously mixed oil-water emulsions or pristine aqueous interfaces.

(iii) Reactions of isoprene in aqueous electrosprays were driven by non-equilibrium conditions therein – most importantly, due to the rapid evaporation of water leading to highly concentrated droplets and partially hydrated hydronium ions (mechanism M_3). It is, thus, crucial to distinguish their contribution from purely ‘interfacial effects’ towards dramatic rate enhancements in aqueous electrosprays.^{50,53,64,78,104,105}

(iv) Gas-phase reactions could play a significant role in the electrosprays – in our experiments, reactions between partially hydrated protons and isoprene led to its protonation and



oligomerization, as recently suggested by Yan & co-workers⁵³ and demonstrated by Jacobs & co-workers.¹⁰⁶

Our experimental and theoretical results demonstrate that chemistries in aqueous electrosprays do not necessarily correspond to those at the pristine air–water interface and oil–water emulsions at NTP. These findings also contradict the previous claims of the superacidity of the pristine air–water interface as the bulk acidity approaches $\text{pH} \leq 3.6$;⁴⁰ the proposal for the mildly acidic environmental surfaces to act as the primary sink for the atmospheric isoprene/terpenes should also be reevaluated.^{41,46,47} While the potential of aqueous electrosprays to produce high-value products appears promising, those reactions are unlikely to be realized at pristine aqueous interfaces because of the seminal role of the non-equilibrium effects, such as the formation of water clusters with minimally hydrated hydronium ions.^{49,78,106} We do note that air–water interfaces could, perhaps, be investigated semi-quantitatively through electrosprays, if the reactants do not participate in gas-phase, acid catalyzed, or redox reactions therein^{51,107–109} and/or the gas-phase reactants do not dissolve in the droplets to re-emerge as interfacial species; a careful case-by-case assessment is needed. The mechanisms underlying the protonation and oligomerization of isoprene gas on electrosprays of $\text{pH} > 7$ water also warrant further investigation. We conclude by stressing on the importance of combining complementary experimental techniques, *ab initio* calculations and molecular dynamics simulations in the quest to unravel phenomena occurring at the pristine air–water interface.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The research reported in this publication was supported by funding from King Abdullah University of Science and Technology (#OSR-2016-CRG5-2992). The authors thank Mr Ivan Gromicho, Scientific Illustrator at KAUST, for preparing Fig. 1. The authors also thank Professor Richard Saykally and Professor Evan Williams (University of California Berkeley), and Dr Manuel Monge Palacios (KAUST) for fruitful discussions. This research used the resources of the Supercomputing Laboratory at King Abdullah University of Science & Technology (KAUST) in Thuwal, Saudi Arabia.

References

- J. H. Seinfeld and S. N. Pandis, *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, Wiley-Interscience, 2edn, 2006.
- J. Zhong, M. Kumar, J. S. Francisco and X. C. Zeng, Insight into Chemistry on Cloud/Aerosol Water Surfaces, *Acc. Chem. Res.*, 2018, **51**(5), 1229–1237.
- M. Kumar and J. S. Francisco, Ion pair particles at the air–water interface, *Proc. Natl. Acad. Sci., India*, 2017, **114**(47), 12401–12406.
- A. V. Shavlov and V. A. Zdzhumandzhi, Metastable states and coalescence of charged water drops inside clouds and fog, *J. Aerosol Sci.*, 2016, **91**, 54–61.
- L. Pawlowski, *The Science and Engineering of Thermal Spray Coatings*, John Wiley & Sons Ltd., Britain, 2008.
- J. H. Bang and K. S. Suslick, Applications of Ultrasound to the Synthesis of Nanostructured Materials, *Adv. Mater.*, 2010, **22**(10), 1039–1059.
- A. Y. Li, Q. J. Luo, S. J. Park and R. G. Cooks, Synthesis and Catalytic Reactions of Nanoparticles formed by Electrospray Ionization of Coinage Metals, *Angew. Chem., Int. Ed.*, 2014, **53**(12), 3147–3150.
- C. M. Dobson, G. B. Ellison, A. F. Tuck and V. Vaida, Atmospheric aerosols as prebiotic chemical reactors, *Proc. Natl. Acad. Sci., India*, 2000, **97**(22), 11864–11868.
- K. Ariga and J. P. Hill, Monolayers at Air–Water Interfaces: From Origins-of-Life to Nanotechnology, *Chem. Rev.*, 2011, **11**(4), 199–211.
- E. C. Griffith and V. Vaida, *In situ* observation of peptide bond formation at the water–air interface, *Proc. Natl. Acad. Sci., India*, 2012, **109**(39), 15697–15701.
- N. Agmon, H. J. Bakker, R. K. Campen, R. H. Henchman, P. Pohl, S. Roke, M. Thämer and A. Hassanali, Protons and Hydroxide Ions in Aqueous Systems, *Chem. Rev.*, 2016, **116**(13), 7642–7672.
- R. J. Saykally, Air/water interface: Two sides of the acid–base story, *Nat. Chem.*, 2013, **5**(2), 82–84.
- E. C. Griffith and V. Vaida, Ionization state of L-phenylalanine at the air–water interface, *J. Am. Chem. Soc.*, 2013, **135**(2), 710–716.
- National Academies of Sciences, Engineering, and Medicine, *The Future of Atmospheric Chemistry Research: Remembering Yesterday, Understanding Today, Anticipating Tomorrow*, The National Academies Press, Washington, DC, 2016, DOI: 10.17226/23573.
- V. Vaida, Prebiotic phosphorylation enabled by microdroplets, *Proc. Natl. Acad. Sci., India*, 2017, **114**(47), 12359–12361.
- S. Banerjee, E. Gnanamani, X. Yan and R. N. Zare, Can all bulk-phase reactions be accelerated in microdroplets?, *Analyst*, 2017, **142**(9), 1399–1402.
- Y. Tabe, N. Kikkawa, H. Takahashi and A. Morita, Surface Acidity of Water Probed by Free Energy Calculation for Trimethylamine Protonation, *J. Phys. Chem. A*, 2013, **118**(2), 977–988.
- A. J. Colussi and S. Enami, Comment on Surface Acidity of Water Probed by Free Energy Calculation for Trimethylamine Protonation, *J. Phys. Chem. C*, 2014, **118**(5), 2894.
- Y. Tabe, N. Kikkawa, H. Takahashi and A. Morita, Reply to Comment on Surface Acidity of Water Probed by Free Energy Calculation for Trimethylamine Protonation, *J. Phys. Chem. C*, 2014, **118**(5), 2895.
- C. Macias-Romero, I. Nahalka, H. I. Okur and S. Roke, Optical imaging of surface chemistry and dynamics in confinement, *Science*, 2017, **357**(6353), 784–787.
- E. M. Arnett, Gas-phase proton transfer- Breakthrough for solution chemistry, *Acc. Chem. Res.*, 1973, **6**(12), 404–409.



- 22 K. B. Eisenthal, Liquid interfaces probed by second-harmonic and sum-frequency spectroscopy, *Chem. Rev.*, 1996, **96**(4), 1343–1360.
- 23 R. K. Lam, S. L. Raj, T. A. Pascal, C. D. Pemmaraju, L. Foglia, A. Simoncig, N. Fabris, P. Miotti, C. J. Hull, A. M. Rizzuto, J. W. Smith, R. Mincigrucci, C. Masciovecchio, A. Gessini, E. Allaria, G. De Ninno, B. Diviacco, E. Roussel, S. Spampinati, G. Penco, S. Di Mitri, M. Trovo, M. Danailov, S. T. Christensen, D. Sokaras, T. C. Weng, M. Coreno, L. Poletto, W. S. Drisdell, D. Prendergast, L. Giannessi, E. Principi, D. Nordlund, R. J. Saykally and C. P. Schwartz, Soft X-Ray Second Harmonic Generation as an Interfacial Probe, *Phys. Rev. Lett.*, 2018, **120**, 023901.
- 24 K. Ariga, T. Nakanishi, J. P. Hill, M. Shirai, M. Okuno, T. Abe and J. Kikuchi, Tunable pK of amino acid residues at the air–water interface gives an L-zyne (Langmuir enzyme), *J. Am. Chem. Soc.*, 2005, **127**(34), 12074–12080.
- 25 H. Y. Yang, Y. Imanishi and A. Harata, Estimating pH at the Air/Water Interface with a Confocal Fluorescence Microscope, *Anal. Sci.*, 2015, **31**(10), 1005–1010.
- 26 J. S. Samson, R. Scheu, N. Smolentsev, S. W. Rick and S. Roke, Sum frequency spectroscopy of the hydrophobic nanodroplet/water interface: Absence of hydroxyl ion and dangling OH bond signatures, *Chem. Phys. Lett.*, 2014, **615**, 124–131.
- 27 C. S. Tian and Y. R. Shen, Structure and charging of hydrophobic material/water interfaces studied by phase-sensitive sum-frequency vibrational spectroscopy, *Proc. Natl. Acad. Sci., India*, 2009, **106**(36), 15148–15153.
- 28 S. Nihonyanagi, R. Kusaka, K.-i. Inoue, A. Adhikari, S. Yamaguchi and T. Tahara, Accurate determination of complex $\chi(2)$ spectrum of the air/water interface, *J. Chem. Phys.*, 2015, **143**(12), 124707.
- 29 S. J. Byrnes, P. L. Geissler and Y. R. Shen, Ambiguities in surface nonlinear spectroscopy calculations, *Chem. Phys. Lett.*, 2011, **516**(4–6), 115–124.
- 30 O. Carrier, E. H. G. Backus, N. Shahidzadeh, J. Franz, M. Wagner, Y. Nagata, M. Bonn and D. Bonn, Oppositely Charged Ions at Water–Air and Water–Oil Interfaces: Contrasting the Molecular Picture with Thermodynamics, *J. Phys. Chem. Lett.*, 2016, **7**(5), 825–830.
- 31 R. K. Lam, J. W. Smith, A. M. Rizzuto, O. Karshioğlu, H. Bluhm and R. J. Saykally, Reversed interfacial fractionation of carbonate and bicarbonate evidenced by X-ray photoemission spectroscopy, *J. Chem. Phys.*, 2017, **146**(9), 094703.
- 32 A. M. Jubb, W. Hua and H. C. Allen, Organization of Water and Atmospherically Relevant Ions and Solutes: Vibrational Sum Frequency Spectroscopy at the Vapor/Liquid and Liquid/Solid Interfaces, *Acc. Chem. Res.*, 2012, **45**(1), 110–119.
- 33 J. R. Platt, Strong Inference – Certain Systematic Methods of Scientific Thinking May Produce Much More Rapid Progress Than Others, *Science*, 1964, **146**(364), 7.
- 34 E. De Hoffmann and V. Stroobant, *Mass spectrometry: principles and applications*, John Wiley & Sons, 2007.
- 35 P. Kebarle and M. Peschke, On the mechanisms by which the charged droplets produced by electrospray lead to gas phase ions, *Anal. Chim. Acta*, 2000, **406**(1), 11–35.
- 36 P. Kebarle and U. H. Verkerk, Electrospray: From Ions in Solution to Ions in the Gas Phase, What We Know Now, *Mass Spectrom. Rev.*, 2009, **28**(6), 898–917.
- 37 S. Banerjee and S. Mazumdar, Electrospray Ionization Mass Spectrometry: A Technique to Access the Information beyond the Molecular Weight of the Analyte, *Int. J. Anal. Chem.*, 2012, **2012**, 40.
- 38 S. Nguyen and J. B. Fenn, Gas-phase ions of solute species from charged droplets of solutions, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**(4), 1111–1117.
- 39 S. Enami, M. R. Hoffmann and A. J. Colussi, Proton Availability at the Air/Water Interface, *J. Phys. Chem. Lett.*, 2010, **1**(10), 1599–1604.
- 40 S. Enami, M. R. Hoffmann and A. J. Colussi, Superacid chemistry on mildly acidic water, *J. Phys. Chem. Lett.*, 2010, **1**(24), 3488–3493.
- 41 S. Enami, H. Mishra, M. R. Hoffmann and A. J. Colussi, Protonation and Oligomerization of Gaseous Isoprene on Mildly Acidic Surfaces: Implications for Atmospheric Chemistry, *J. Phys. Chem. A*, 2012, **116**(24), 6027–6032.
- 42 H. Mishra, S. Enami, R. J. Nielsen, L. A. Stewart, M. R. Hoffmann, W. A. Goddard and A. J. Colussi, Bronsted basicity of the air–water interface, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**(46), 18679–18683.
- 43 H. Mishra, S. Enami, R. J. Nielsen, M. R. Hoffmann, W. A. Goddard and A. J. Colussi, Anions dramatically enhance proton transfer through aqueous interfaces, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**(26), 10228–10232.
- 44 D. Schroder, Ion clustering in electrospray mass spectrometry of brine and other electrolyte solutions, *Phys. Chem. Chem. Phys.*, 2012, **14**(18), 6382–6390.
- 45 X. Yan, M. Delgado, J. Aubry, O. Gribelin, A. Stocco, F. Boisson-Da Cruz, J. Bernard and F. Ganachaud, Central Role of Bicarbonate Anions in Charging Water/Hydrophobic Interfaces, *J. Phys. Chem. Lett.*, 2018, **9**(1), 96–103.
- 46 K. Matsuoka, Y. Sakamoto, T. Hama, Y. Kajii and S. Enami, Reactive Uptake of Gaseous Sesquiterpenes on Aqueous Surfaces, *J. Phys. Chem. A*, 2017, **121**(4), 810–818.
- 47 S. Enami, M. R. Hoffmann and A. J. Colussi, Dry deposition of biogenic terpenes via cationic oligomerization on environmental aqueous surfaces, *J. Phys. Chem. Lett.*, 2012, **3**, 3102–3108.
- 48 V. B. Kazansky, Solvation as a main factor that determines the strength of liquid superacids and the selectivity of the acid-catalyzed reactions of olefins, *Catal. Today*, 2002, **73**(1–2), 127–137.
- 49 I. Nam, H. G. Nam and R. N. Zare, Abiotic synthesis of purine and pyrimidine ribonucleosides in aqueous microdroplets, *Proc. Natl. Acad. Sci., India*, 2018, **115**(1), 36–40.
- 50 J. K. Lee, S. Banerjee, H. G. Nam and R. N. Zare, Acceleration of reaction in charged microdroplets, *Q. Rev. Biophys.*, 2015, **48**(04), 437–444.



- 51 M. I. Jacobs, J. F. Davies, L. Lee, R. D. Davis, F. Houle and K. R. Wilson, Exploring Chemistry in Microcompartments Using Guided Droplet Collisions in a Branched Quadrupole Trap Coupled to a Single Droplet, Paper Spray Mass Spectrometer, *Anal. Chem.*, 2017, **89**(22), 12511–12519.
- 52 D. A. Thomas, L. T. Wang, B. Goh, E. S. Kim and J. L. Beauchamp, Mass Spectrometric Sampling of a Liquid Surface by Nanoliter Droplet Generation from Bursting Bubbles and Focused Acoustic Pulses: Application to Studies of Interfacial Chemistry, *Anal. Chem.*, 2015, **87**(6), 3336–3344.
- 53 X. Yan, R. M. Bain and R. G. Cooks, Organic Reactions in Microdroplets: Reaction Acceleration Revealed by Mass Spectrometry, *Angew. Chem., Int. Ed.*, 2016, **55**(42), 12960–12972.
- 54 S. Narayan, J. Muldoon, M. G. Finn, V. V. Fokin, H. C. Kolb and K. B. Sharpless, On water: unique reactivity of organic compounds in aqueous suspension, *Angew. Chem., Int. Ed.*, 2005, **44**(21), 3275–3279.
- 55 Y. S. Jung and R. A. Marcus, On the theory of organic catalysis on water, *J. Am. Chem. Soc.*, 2007, **129**(17), 5492–5502.
- 56 J. K. Beattie, C. S. P. McErlean and C. B. W. Phippen, The Mechanism of On-Water Catalysis, *Chem.–Eur. J.*, 2010, **16**(30), 8972–8974.
- 57 L. L. Thomas, J. Tirado-Rives and W. L. Jorgensen, Quantum Mechanical/Molecular Mechanical Modeling Finds Diels-Alder Reactions Are Accelerated Less on the Surface of Water Than in Water, *J. Am. Chem. Soc.*, 2010, **132**(9), 3097–3104.
- 58 W. D. Luedtke, U. Landman, Y. H. Chiu, D. J. Levandier, R. A. Dressler, S. Sok and M. S. Gordon, Nanojets, electrospray, and ion field evaporation: Molecular dynamics simulations and laboratory experiments, *J. Phys. Chem. A*, 2008, **112**(40), 9628–9649.
- 59 J. F. de la Mora, The fluid dynamics of Taylor cones, *Annu. Rev. Fluid Mech.*, 2007, **39**, 217–243.
- 60 D. N. Mortensen and E. R. Williams, Ultrafast (1 μ s) Mixing and Fast Protein Folding in Nanodrops Monitored by Mass Spectrometry, *J. Am. Chem. Soc.*, 2016, **138**(10), 3453–3460.
- 61 H. J. Sterling, C. A. Cassou, A. C. Susa and E. R. Williams, Electrothermal Supercharging of Proteins in Native Electrospray Ionization, *Anal. Chem.*, 2012, **84**(8), 3795–3801.
- 62 B. Yuan, A. R. Koss, C. Warneke, M. Coggon, K. Sekimoto and J. A. de Gouw, Proton-Transfer-Reaction Mass Spectrometry: Applications in Atmospheric Sciences, *Chem. Rev.*, 2017, **117**(21), 13187–13229.
- 63 S. Zhou, B. S. Prebyl and K. D. Cook, Profiling pH changes in the electrospray plume, *Anal. Chem.*, 2002, **74**(19), 4885–4888.
- 64 S. L. Zhou and K. D. Cook, Protonation in electrospray mass spectrometry: Wrong-way-round or right-way-round?, *J. Am. Soc. Mass Spectrom.*, 2000, **11**(11), 961–966.
- 65 M. M. Francl, W. J. Pietro, W. J. Hehre, J. S. Binkley, M. S. Gordon, D. J. Defrees and J. A. Pople, Self-consistent molecular-orbital methods. 23. A polarization-type basis set for 2nd -row elements, *J. Chem. Phys.*, 1982, **77**(7), 3654–3665.
- 66 C. J. Cramer, *Essentials of computational chemistry: theories and models*, John Wiley & Sons, 2013.
- 67 K. Fukui, The path of chemical reactions-the IRC approach, *Acc. Chem. Res.*, 1981, **14**(12), 363–368.
- 68 C. Dykstra, G. Frenking, K. Kim and G. Scuseria, *Theory and Applications of Computational Chemistry: the first forty years*, Elsevier, 2011.
- 69 S. Nosé, A molecular dynamics method for simulations in the canonical ensemble, *Mol. Phys.*, 1984, **52**(2), 255–268.
- 70 M. J. Frisch, G. Trucks, H. B. Schlegel, G. Scuseria, M. Robb, J. Cheeseman, G. Scalmani, V. Barone, B. Mennucci and G. Petersson, *Gaussian 09, revision A. 1*, Gaussian Inc., Wallingford CT, 2009, vol. 27, p. 34.
- 71 T. D. Sharkey, A. E. Wiberley and A. R. Donohue, Isoprene emission from plants: Why and how, *Ann. Bot.*, 2008, **101**(1), 5–18.
- 72 A. T. Archibald, J. G. Levine, N. L. Abraham, M. C. Cooke, P. M. Edwards, D. E. Heard, M. E. Jenkin, A. Karunaharan, R. C. Pike, P. S. Monks, D. E. Shallcross, P. J. Telford, L. K. Whalley and J. A. Pyle, Impacts of HO(x) regeneration and recycling in the oxidation of isoprene: Consequences for the composition of past, present and future atmospheres, *Geophys. Res. Lett.*, 2011, **38**(L05804), 6.
- 73 Neutral Thermochemical Data, in *NIST Chemistry WebBook, NIST Standard Reference Database Number 69*, ed. P. J. Linstrom and W. G. Mallard, National Institute of Standards and Technology, Gaithersburg MD, 2017, p. 20899, DOI: 10.18434/T4D303.
- 74 M. Deiana, F. Corongiu, M. Dessi, P. Scano, M. Casu and A. Lai, NMR determination of site-specific deuterium distribution (SNIF-NMR) in squalene from different sources, *Magn. Reson. Chem.*, 2001, **39**(1), 29–32.
- 75 R. M. Bain, C. J. Pulliam and R. G. Cooks, Accelerated Hantzsch electrospray synthesis with temporal control of reaction intermediates, *Chem. Sci.*, 2015, **6**(1), 397–401.
- 76 D. Schroder, M. Budesinsky and J. Roithova, Deprotonation of *p*-Hydroxybenzoic Acid: Does Electrospray Ionization Sample Solution or Gas-Phase Structures?, *J. Am. Chem. Soc.*, 2012, **134**(38), 15897–15905.
- 77 J. F. de la Mora, G. J. Van Berkel, C. G. Enke, R. B. Cole, M. Martinez-Sanchez and J. B. Fenn, Electrochemical processes in electrospray ionization mass spectrometry, *J. Mass Spectrom.*, 2000, **35**(8), 939–952.
- 78 A. J. Ingram, C. L. Boeser and R. N. Zare, Going beyond electrospray: mass spectrometric studies of chemical reactions in and on liquids, *Chem. Sci.*, 2016, **7**(1), 39–55.
- 79 R. M. Bain, C. J. Pulliam, F. They and R. G. Cooks, Accelerated Chemical Reactions and Organic Synthesis in Leidenfrost Droplets, *Angew. Chem., Int. Ed.*, 2016, **55**(35), 10478–10482.
- 80 P. Aussillous and D. Quere, Liquid marbles, *Nature*, 2001, **411**(6840), 924–927.
- 81 E. P. L. Hunter and S. G. Lias, Evaluated Gas Phase Basicities and Proton Affinities of Molecules: An Update, *J. Phys. Chem. Ref. Data*, 1998, **27**(3), 413–656.



