

ChemComm

Chemical Communications

rsc.li/chemcomm



ISSN 1359-7345

COMMUNICATION

Alix C. Harlington *et al.*
Efficient *O*-demethylation of lignin monoaromatics using the
peroxygenase activity of cytochrome P450 enzymes


 Cite this: *Chem. Commun.*, 2022, 58, 13321

 Received 25th August 2022,
Accepted 14th October 2022

DOI: 10.1039/d2cc04698a

rsc.li/chemcomm

Efficient *O*-demethylation of lignin monoaromatics using the peroxygenase activity of cytochrome P450 enzymes†

 Alix C. Harlington,^a Keith E. Shearwin,^a Stephen G. Bell^{*b} and Fiona Whelan^{†a}

A crucial reaction in harnessing renewable carbon from lignin is *O*-demethylation. We demonstrate the selective *O*-demethylation of syringol and guaiacol using different cytochrome P450 enzymes. These can efficiently use hydrogen peroxide which, when compared to nicotinamide cofactor-dependent monooxygenases and synthetic methods, allows for cheap and clean *O*-demethylation of lignin-derived aromatics.

The biological valorisation of lignin has the potential to supply useful chemicals and materials traditionally derived from non-renewable feedstocks.^{1,2} Lignin is comprised of varying proportions of *p*-coumaryl (H-type), coniferyl (G-type), and sinapyl (S-type) alcohol subunits,^{3,4} and its depolymerisation generates a heterogeneous mixture of aromatic compounds (lignin-derived aromatic compounds; LDACs).⁵ The majority of LDACs are derived from methoxylated G and S-type subunits which must be *O*-demethylated to diols before they can be funnelled to high-value ring-opened compounds.^{6,7} However, *O*-demethylation is recognised as a rate-limiting reaction limiting the conversion of LDACs to useful compounds in high atom yield.^{8,9} Thus, efforts have focused on characterisation of demethylase enzymes from bacteria to improve the biological valorisation of lignin.

The cytochrome P450 superfamily of heme-containing enzymes are of interest as *O*-demethylases for lignin valorisation. These enzymes catalyse C–H bond oxidation reactions on a broad range of substrates.^{10,11} Recently, a new family of cytochrome P450s – the CYP255A family – have been characterised that catalyse the aromatic *O*-demethylation of G-type LDACs. The CYP255A2 enzyme (GcoA) catalyses *O*-demethylation of several LDACs, with highest specificity for guaiacol.^{12,13} Similarly, CYP255A1 (AgcA) catalyses

O-demethylation of guaiacol and 4-alkylguaiacols.¹³ Moreover, engineered P450 enzymes, non-heme iron oxygenases and tetrahydrofolate dependent enzymes have been reported that can demethylate guaiacol and syringol derivatives.^{14–16} However, very few S-type LDAC *O*-demethylases have been characterised, making it difficult to establish efficient microbial valorisation of S-type lignin.¹⁷ Accordingly, efforts have been made to expand the substrate range of P450s for S-lignin LDACs.¹⁸ The lack of a P450 *O*-demethylase with native specificity for S-type LDACs limits both our understanding of the enzymatic *O*-demethylation of S-type LDACs and our capacity to engineer *O*-demethylases for efficient lignin valorisation.

Here we report the characterisation of a new cytochrome P450 from *Amycolatopsis thermoflava* N1165 that catalyses the *O*-demethylation of syringol to 3-methoxycatechol (3-MC). Furthermore, we report for the first time the use of H₂O₂-driven reactions to efficiently support the catalytic activity of the CYP255 family.

Several species of bacteria have the capacity to degrade and assimilate S-type LDACs.^{1,8,19,20} However, the catabolic pathways for the degradation of S-type LDACs remain largely unelucidated. We hypothesised that certain bacterial species may encode various cytochrome P450 *O*-demethylases with specificity for different LDACs to support degradation of different lignin monoaromatics. We noticed that *A. thermoflava* N1165 contained two genes that encode CYP255 enzymes. One of these is essentially identical to the GcoA from *Amycolatopsis* sp. strain ATCC 39116, except for a single point mutation (K206R), and therefore could be assigned as a guaiacol *O*-demethylase GcoA. The second gene (WP_037322545) encodes a CYP255 enzyme which shares 47% sequence identity with GcoA and we hypothesised that this enzyme could be a potential S-type LDAC *O*-demethylase (herein referred to as SyoA). Phylogenetic analysis revealed that SyoA shares a common ancestor with the CYP255A and CYP1232 families of *O*-demethylases (Fig. 1).^{12,21}

The SyoA and GcoA enzymes were produced in *E. coli* and purified using affinity and anion exchange chromatography

^a Department of Molecular and Biomedical Science, University of Adelaide, Adelaide, SA 5005, Australia. E-mail: fiona.whelan@adelaide.edu.au

^b Department of Chemistry, University of Adelaide, Adelaide, SA 5005, Australia. E-mail: stephen.bell@adelaide.edu.au

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2cc04698a>






Fig. 3 Active site structures of (a) CYP199A4 bound to 4-methoxybenzoic acid (PDB: 4do1) and (b) GcoA bound to guaiacol (PDB: 5ncb) highlighting the oxygen binding groove of the I-helix. The conserved acid–alcohol pair (D251 and T252) responsible for the monooxygenase activity of most P450s is replaced with an amide and acid pair (Q248 and E249) in GcoA.

peroxygenases that use H_2O_2 contain a residue capable of acid–base chemistry (histidine, aspartate, or glutamate) on the distal side of the heme.^{24–26} Recent reports have demonstrated that mutation of the conserved threonine of the acid–alcohol pair to glutamic acid, confers peroxygenase activity in several members of the P450 superfamily.^{27,28} Given the presence of this glutamic acid residue in the active sites of GcoA and SyoA, we reasoned that these enzymes may be capable of peroxygenase activity using H_2O_2 to *O*-demethylate their substrates.

Addition of H_2O_2 to P450 enzymes can result in destruction of the heme cofactor, leaving the enzyme inactive.²⁹ Heme bleaching assays were used to assess the stability of GcoA and SyoA to the presence of H_2O_2 , and by extension, their potential as peroxygenases. The destruction of heme was determined by monitoring change in absorbance of the Soret heme peak by UV-vis spectroscopy. Addition of 40 mM H_2O_2 to $\sim 1 \mu\text{M}$

substrate-free GcoA and SyoA resulted in destruction of the heme over ~ 15 – 20 min (Fig. S5, ESI[†]). Addition of the preferred substrates improved stability of both enzymes, with the Soret heme peak present after 30 min (Fig. S6, ESI[†]). These experiments demonstrated that compared to many heme enzymes, GcoA and SyoA have some resistance to H_2O_2 .^{29,30} Interestingly, after addition of 40 mM H_2O_2 to guaiacol-bound GcoA and syringol-bound SyoA, the solutions changed colour to red and the UV-vis absorbance spectrum changed over the course of 30 min, indicating a potential catalytic reaction (Fig. S6, ESI[†]).

High-performance liquid chromatography (HPLC) analysis of 10–20 mM H_2O_2 -driven enzyme ($1 \mu\text{M}$) reactions was performed to identify the predicted *O*-demethylated products of guaiacol and syringol. HPLC analysis of the reactions with GcoA and guaiacol showed the formation of a product with a retention time of 7.2 min. This product exhibited a retention time identical to a commercial sample of catechol and it was determined that $\sim 59\%$ ($295 \pm 12 \mu\text{M}$, total turnover number, TTN 295) of guaiacol was *O*-demethylated to catechol by 120 min (Fig. 4a–c). The reaction with GcoA and syringol yielded no products that matched the retention time of 3-MC, indicating syringol could not be efficiently *O*-demethylated by GcoA (Fig. S7, ESI[†]).

The H_2O_2 -driven SyoA reaction with guaiacol yielded a small amount of catechol ($15 \pm 1 \mu\text{M}$, TTN 15) after 120 min (Fig. 4b). In contrast, HPLC analysis of the reaction with SyoA and syringol revealed formation of a peak corresponding to the retention time of an authentic sample of 3-MC. Approximately 27% ($137 \pm 12 \mu\text{M}$, TTN 137) of syringol was converted to 3-MC by SyoA after 60 min (Fig. 4a, b and d). Furthermore, we




Fig. 4 Optimised *in vitro* hydrogen peroxide turnovers demonstrate *O*-demethylase activity of GcoA and SyoA. (a) Scheme showing the *O*-demethylation of guaiacol and syringol by GcoA or SyoA in the presence of hydrogen peroxide. (b) Concentration of demethylated guaiacol and syringol aromatic products produced over 2 hours (guaiacol) and 1 hour (syringol). HPLC analysis of the hydrogen peroxide driven demethylation of (c) guaiacol by GcoA and (d) syringol by SyoA. Error bars indicate the SD from the mean of replicates $n = 3$.



