



Cite this: *Environ. Sci.: Processes Impacts*, 2024, 26, 344

## Contrasted redox-dependent structural control on Fe isotope fractionation during its adsorption onto and assimilation by heterotrophic soil bacteria†

Aridane G. González,<sup>a</sup> Franck Poitrasson,<sup>b</sup> Felix Jiménez-Villacorta,<sup>c</sup> Liudmila S. Shirokova<sup>bd</sup> and Oleg S. Pokrovsky<sup>be</sup>

Despite the importance of structural control on metal stable isotope fractionation in inorganic and abiotic systems, the link between metal structural changes and related isotopic fractionation during reactions with organic surfaces and live cells remains poorly established. We conducted reversible adsorption of Fe(II) and Fe(III) on the surface of exopolysaccharide (EPS)-rich and EPS-poor *Pseudomonas aureofaciens*, and we allowed Fe intracellular uptake by growing cells. We analyzed the Fe isotopic composition of the remaining fluid and cell biomass, and compared the isotopic fractionation during adsorption and assimilation reaction with relative changes in Fe structural status between aqueous solution and bacterial cells, based on available and newly collected X-ray absorption spectroscopy (XAS) observations. Iron(III) adsorption onto *P. aureofaciens* at  $2.8 \leq \text{pH} \leq 6.0$  produced an enrichment of the cell surface in heavier isotopes with  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  ranging from +0.7 to +2.1‰, without a link to pH in EPS-rich cultures. In contrast, the magnitude of isotopic fractionation increased with pH in EPS-poor cultures. Iron(II) adsorption produced an even larger enrichment of the cell surface in heavier isotopes, by up to 3.2‰, tentatively linked to Fe(III) hydroxide precipitation. Intracellular assimilation of Fe(II) favored heavier isotopes and led to  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  of +0.8‰. In addition, Fe(III) cellular uptake produced an enrichment of the bacterial biomass in lighter isotopes with  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  of -1‰. The XAS analyses demonstrated the dominance of Fe(III)-phosphate complexes both at the cell surface and in the cell interior. We suggest that heavier isotope enrichment of the cell surface relative to the aqueous solution is due to strong Fe(III)-phosphoryl surface complexes and Fe complexation to ligands responsible for metal transfer from the surface to the inner cell. In case of Fe(II) adsorption or assimilation, its partial oxidation within the cell compartments may lead to cell enrichment in heavier isotopes. In contrast, loss of symmetry of assimilated Fe(III) relative to the aqueous Fe<sup>3+</sup> ion and longer bonds of intracellular ions relative to aqueous Fe(III)-citrate or hydroxo-complexes could produce an enrichment of cells in lighter isotopes. The versatile nature of Fe(II) and Fe(III) fractionation without a distinct effect of pH and surface exopolysaccharide coverage suggests that, in natural soil and sedimentary environments, Fe isotope fractionation during interaction with heterotrophic bacteria will be primarily governed by Fe complexation with DOM and Fe redox status in the soil pore water.

Received 2nd August 2023  
Accepted 5th December 2023

DOI: 10.1039/d3em00332a

rsc.li/epsi

### Environmental significance

This investigation provides valuable insights into the mechanisms governing Fe isotope fractionation in soil-fluid bacterial systems. Isotopic partitioning during the adsorption of Fe(II) and Fe(III) by a common soil bacterium is dominated by Fe complexation with DOM. The adsorption of Fe(III) results in an enrichment of heavier isotopes on the cell surface that is attributed to the stronger and more symmetrical complexation of Fe(III) with surface phosphoryl moieties compared to hydroxyl complexes in the aqueous solution. In natural environments, the  $\delta^{57}\text{Fe}$  values in solution or microbial biomass may vary from approximately -1 to +3‰, depending on the Fe redox status in soil porewater, and this can occur rapidly over short time scales, independent of pH and the presence of microbial EPS.

<sup>a</sup>Instituto de Oceanografía y Cambio Global, IOCG, Universidad de Las Palmas de Gran Canaria, ULPGC, Spain. E-mail: aridane.gonzalez@ulpgc.es; Tel: +34 928 45 44 51

<sup>b</sup>Géosciences Environnement Toulouse (GET), CNRS UMR 5563, UPS-IRD-CNES 14-16, Avenue Edouard Belin, 31400, Toulouse, France

<sup>c</sup>ESS Bilbao, Parque Tecnológico Bizkaia, Nave 201, 48170 Zamudio, Spain

<sup>d</sup>N.P. Laverov Federal Center for Integrated Arctic Research (FCI Arctic), Russian Academy of Sciences, Arkhangelsk, Russia

<sup>e</sup>BIO-GEO-CLIM Laboratory, National Research Tomsk State University, Tomsk, Russia

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3em00332a>





surface- and intracellularly bound Fe and Fe isotope fractionation between the cell and aqueous solution was conducted considering, in particular, the effect of (i) Fe redox state; (ii) solution pH, and (iii) the presence or not of microbial cell exometabolites. By encompassing a wide variety of soil and aquatic environments, we anticipate that the results can be used to better constrain factors controlling Fe isotopic composition of soil or sediment porewaters and surface waters in the presence of heterotrophic bacteria.

## 2. Materials and methods

### 2.1. Heterotrophic bacterial culture

The bacterial strain *P. aureofaciens* CNMN PsB-03 was obtained from the Laboratory of Plant Mineral Nutrition and Hydric Regime (Institute of Genetics and Plant Physiology, Moldovan Academy of Sciences, Chisinau, Moldova). It was isolated from temperate organic-rich soil and selected due to its capacity of producing EPS on a sucrose-containing medium.<sup>49</sup> The strain of *P. aureofaciens* was maintained at 4 °C in a liquid succinic acid (SA) medium,<sup>50</sup> and cultured either in the sucrose-peptone (SP) media,<sup>51</sup> thus promoting rich EPS synthesis, or in the SA media with poor EPS production. Cultivation was performed at 25 °C during 48–72 h with continuous shaking. The qualitative monosaccharide composition of the EPS produced is different depending on the culture media: the SP-media yields the EPS composed of 76% fructose, 11% glucose and <10% of other sugars, whereas in the SA media, polysaccharides contain 50% glucose, 22% fructose and 14% mannose and other sugars.<sup>52</sup>

Experiments were performed for non-rinsed and rinsed bacterial biomass that produced two sub-samples for both EPS-rich and EPS-poor samples. In the first case, the bacteria were collected from the culture without rinsing, using centrifugation for 20 min at 8000 g. Therefore, for non-rinsed biomass, the EPS content of the cell wall is expected to remain intact. The rinsed biomass was produced by repeated (3 times) centrifugation in an inert electrolyte solution (0.01 M NaNO<sub>3</sub>). Repetitive rinsing of biomass in this electrolyte solution should lead to the removal of part of the EPS from the cell surface. For EPS-rich (SP-media) and EPS-poor (SA-media) cultures, 28% and 6% of the total biomass were removed during centrifugation, respectively. The biomass concentration is given in g<sub>wet</sub> l<sup>-1</sup> (after centrifugation for 20 min at 8000 g). The conversion factor of wet to dry biomass is equal to 3.6 and 5.0 for EPS-rich and EPS-poor cultures, respectively.

### 2.2. Adsorption experiments

Prior to the adsorption experiments, the biomass was collected *via* centrifugation and rinsed with 0.01 M NaNO<sub>3</sub> solution three times *via* centrifugation. The last supernatant was considered as the reference solution used for control experiments. The biomass of *P. aureofaciens* was fixed to 4 g<sub>wet</sub> l<sup>-1</sup> for all the experiments. The pH was adjusted by adding aliquots of 0.01–0.1 M NaOH or HNO<sub>3</sub>. Adsorption experiments were carried out in 0.01 M NaNO<sub>3</sub> under darkness at 25 ± 0.2 °C and continuously shaken bacteria suspension. The initial concentrations of

Fe(II)–ammonium sulfate or Fe(III) chloride (Sigma) solutions were 20 and 3.3 mg l<sup>-1</sup>, respectively (Table ESI-1†). Divalent Fe was added to thoroughly rinse bacterial biomass and maintained in 0.01 M NaNO<sub>3</sub> at pH of 4.9 under continuous nitrogen bubbling and stirring. Iron(III) adsorption was studied on live bacteria as a function of pH (2.8 to 6.0) and with 3 hours of exposure time. Bacteria-free control experiments with Fe(II) and Fe(III) did not yield significant (*i.e.*, >5–10%) dissolved (<0.45 μm) Fe concentration decrease during an exposure time comparable to that of the adsorption experiments with bacteria. This indicated a lack of Fe adsorption on reactor cell walls, Fe(II) oxidation and particulate Fe(III) precipitation.

After sampling, experimental solutions were filtered through 0.45 μm acetate cellulose filters, acidified with ultrapure NaNO<sub>3</sub> and stored at 5 °C pending analysis. The reversibility of adsorption was tested following the method developed by Fowle and Fein.<sup>53</sup> A homogeneous parent bacteria suspension with Fe(III) was adjusted to pH ~ 5, under which 100% of Fe was adsorbed onto bacteria. After 3 hours of interaction, aliquots of this parent suspension solution were collected and adjusted to sequentially lower pH values (to 3 and then to 1.5). The reaction vessels were equilibrated at new pH values for 3 hours and sampled for dissolved Fe. The “desorbed” metal supernatant concentration was used to calculate the level of irreversibly “incorporated” metal. This amount ranged between 5 and 15% of the initially added Fe, which suggests an equilibrium adsorption process with a relatively small quantity of Fe(III) penetrating inside the cell under our experimental adsorption conditions (not shown). For the reversibility of Fe(II) adsorption and oxidation, a similar technique yielded 100 ± 10% reversibility. The average experimental reproducibility of metal adsorption yield, calculated based on 2 to 3 replicates at identical conditions, was between 5 and 10%, as assessed in a series of separate adsorption experiments. For these reasons, the replicates were not measured for isotopic ratios.

### 2.3. Iron assimilation by living cells

Iron incorporation into the cells during microbial growth experiments was performed over 2–3 days after 2 days of cell culturing. This period corresponded to maximal cell viability at the exponential to the beginning of the stationary growth state,<sup>39,40</sup> and was consistent with previous studies on metal isotope fractionation during metal assimilation by bacterial cultures.<sup>38,42</sup> Iron was added in the EDTA-free nutrient medium in the form of either Fe(II) ammonium sulfate, Fe(II) lactate, Fe(III) citrate, or FeCl<sub>3</sub>. Total aqueous Fe concentration ranged from 2 to 100 mg l<sup>-1</sup>. Other nutrient metals (Cu, Zn, Mn, Co, Ni, Mo) were present in trace amounts required for bacteria growth, with concentrations of 1–2 orders of magnitude lower than that of Fe. The main organic ligands controlling Fe speciation in organic-rich nutrient media of *P. aureofaciens*, in addition to cell exometabolites, are soluble proteins, phosphate, and carboxylates which are capable of complexing Fe(II) and Fe(III) in solution. By analogy with Cu<sup>2+</sup> (ref. 38) and according to the Fe-binding capacities, thermodynamic calculations using visual Minteq software demonstrate that >99% of total dissolved Fe is







**Table 1** Isotopic fractionation of iron in the adsorption and assimilation experiments represented as  $\delta^{57}\text{Fe}$  and  $\delta^{56}\text{Fe}$ , including values in the starting stock solutions. The isotopic fractionation factor is computed from eqn (4) in the text

Name	Type of sample	pH	$\delta^{57}\text{Fe}$	2SE	$\delta^{56}\text{Fe}$	2SE	$\Delta^{57}\text{Fe}$ (‰)	2SE
<b>Adsorption experiments (3 h)</b>								
EPS-rich-2	Biomass	4.0	0.678	0.047	0.432	0.061	0.74	0.14
	Solution		−0.062	0.129	−0.062	0.317		
EPS-poor-3	Biomass	4.0	0.963	0.040	0.670	0.099	0.89	0.10
	Solution		0.084	0.091	0.005	0.045		
EPS-rich-4	Biomass	5.5	0.466	0.044	0.279	0.090	0.81	0.11
	Solution		−0.347	0.098	−0.250	0.129		
EPS-poor-5	Biomass	6.0	0.747	0.148	0.515	0.093	1.96	0.22
	Solution		−1.212	0.163	−0.790	0.108		
EPS-poor-1	Biomass	2.0	0.198	0.108	0.129	0.124	0.72	0.15
	Solution		−0.525	0.111	−0.419	0.076		
EPS-poor-2	Biomass	2.8	0.199	0.034	0.122	0.068	0.78	0.13
	Solution		−0.578	0.123	−0.407	0.152		
EPS-rich-3	Solution	4.6	−0.680	0.132	−0.503	0.136	1.29	0.30
EPS-poor-Fe2	Solution	4.9	−0.176	0.191	−0.135	0.121	3.16	1.60
<b>Assimilation experiments (48 h)</b>								
EPS-rich-4	Solution	7.2	0.558	0.058	0.363	0.177	−1.10	0.46
EPS-rich-5	Biomass	7.1	1.003	0.132	0.653	0.124	0.77	0.32
	Solution		0.236	0.291	0.208	0.119		
EPS-poor-6	Biomass	7.5	0.996	0.107	0.709	0.128	0.83	0.66
EPS-poor-8	Biomass	7.1	0.749	0.068	0.490 <sup>a</sup>	0.088	−0.99	0.24
EPS-rich-6	Biomass	7.2	0.172	0.097	0.127	0.063	−1.01	0.51
EPS-rich-7	Biomass	7.2	−0.392	0.095	−0.256	0.056	−1.12	0.23
EPS-poor-Fe2-2	Biomass	7.8	0.749	0.119	0.492	0.093	−0.04	0.19
<b>Starting stock solutions</b>								
FeCl <sub>3</sub>	Stock	—	0.55	0.179	0.349	0.045	—	—
Fe(II)-ammonium sulfate	Stock	—	0.188	0.068	0.104	0.066	—	—
Fe(III)-citrate	Stock	—	0.295	0.098	0.197	0.065	—	—
Fe(II)-lactate	Stock	—	0.788	0.142	0.537	0.106	—	—
BCR482	Stock	—	−0.002	0.109	−0.028	0.034	—	—

<sup>a</sup> Recalculated from  $\delta^{57}\text{Fe}$ .

oxalate, gluconate and phosphate) were measured in transmission mode. The incident beam was monitored with a N<sub>2</sub>-filled gas ionization chamber. XANES spectra were analyzed using linear combination fitting considering several footprint components in agreement with the local environment and geometry around Fe atoms (hydroxide as goethite, carboxylate as ferric citrate and phosphoryl as Fe(III) phosphate). EXAFS signals were analyzed up to a maximum  $k$  value of 12 Å<sup>−1</sup>. Multiple scans of 60 min per scan acquisition time were recorded at different spots of the sample to further avoid beam induced reactions. Data treatment and analysis have been performed using the ATHENA and VIPER software, using a weighted  $k^3 \cdot c(k)$  signal with a window ranging from 2.5 to around 12 Å<sup>−1</sup>. Fourier transforms of the spectra were filtered in the 1–2.3 Å range to isolate contribution from the first coordination sphere. All the biomass samples were analyzed at 5–10 K using a liquid-helium cryostat because the low temperature offers great benefits such as decreasing the thermal disorder, and minimizing the damage in terms of redox sensitive elements. All biomass samples for XAS analysis were prepared on-site at the European Molecular Biology Laboratory (EMBL) in the vicinity of the ESRF and processed at the beamline within several hours after the end of the freeze-drying procedure. This

greatly reduced the possibility of sample degradation, and allowed achievement of the highest Fe concentration for XAS spectra acquisition without increasing the level of initial Fe loading.

## 2.7. Calculation of isotopic fractionation in adsorption and assimilation experiments

The experimental conditions of Fe adsorption and assimilation assays are listed in Table ESI-1.† It presents the data for both biomass and solutions analyzed in this work. For samples from the batch adsorption and assimilation experiments performed in a closed system, the isotopic signature of the bacterial cells ( $\delta^{57}\text{Fe}_{\text{cell}}$ ) was calculated from the mass balance equation using the isotopic ratio in the aqueous phase ( $\delta^{57}\text{Fe}_{\text{solution}}$ ), the proportion of adsorbed Fe ( $A$ , %), and the isotopic composition of the initial solution ( $\delta^{57}\text{Fe}_{\text{initial}}$ ):

$$\delta^{57}\text{Fe}_{\text{cell}} = (100 \times \delta^{57}\text{Fe}_{\text{initial}} - (100 - A) \times \delta^{57}\text{Fe}_{\text{solution}}) / A \quad (3)$$

Similarly, in case if only bacterial cell Fe isotope measurements were performed, the isotopic signature of the fluid phase ( $\delta^{57}\text{Fe}_{\text{solution}}$ ) was calculated *via*:



$$\delta^{57}\text{Fe}_{\text{solution}} = (100 \times \delta^{57}\text{Fe}_{\text{initial}} - A \times \delta^{57}\text{Fe}_{\text{cell}})/(100 - A) \quad (4)$$

In this work, we define the isotopic fractionation factor between Fe in solution and Fe adsorbed onto or assimilated by bacteria as:

$$\Delta^{57}\text{Fe}_{\text{cell-solution}} = \delta^{57}\text{Fe}_{\text{cell}} - \delta^{57}\text{Fe}_{\text{solution}} \quad (5)$$

The values of isotopic signature ( $\delta^{57}\text{Fe}$  and  $\delta^{56}\text{Fe}$ ) and isotopic fractionation factor ( $\Delta^{57}\text{Fe}_{\text{cell-solution}}$ ) and those of the starting products are listed in Table 1. In the adsorption experiments, 6 pairs of samples (cell biomass and corresponding aqueous solution) were processed for isotopic analyses. This allowed straightforward calculation of  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  using eqn (4) (Table 1). In these experiments, we also calculated  $\delta^{57}\text{Fe}_{\text{cell}}$  using eqn (3); the values of  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  obtained from such indirect calculation agreed within propagated analytical uncertainties with directly measured  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  values for 4 out of the 6 experiments. For the remaining two experiments (EPS-poor-1 and 2), it is possible the proportion of adsorbed Fe on the cell surfaces is only known with an uncertainty larger than our 3% estimate. With this caveat in mind, we computed  $\delta^{57}\text{Fe}_{\text{cell}}$  using eqn (3) when only  $\delta^{57}\text{Fe}_{\text{aqueous}}$  was available (exp. EPS-rich-3 and EPS-poor-Fe2 in Table 1). Similarly, in the assimilation experiments, coupling of cell and solution measurements in EPS-rich-5 samples yielded direct  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  ( $+0.767 \pm 0.32\%$  by eqn (4)), which is similar within uncertainty to the calculated value ( $+0.739 \pm 0.25\%$  by eqn (3) and (4)). This allowed using other measured  $\delta^{57}\text{Fe}_{\text{cell}}$  to calculate  $\delta^{57}\text{Fe}_{\text{solution}}$  and  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  via eqn (4) and (5), respectively.

We also obtained  $\delta^{56}\text{Fe}$  values, though slightly less precisely than  $\delta^{57}\text{Fe}$  values on a per amu basis. Given that the relationships between  $\delta^{56}\text{Fe}$  and  $\delta^{57}\text{Fe}$  of both adsorbed and assimilated samples plot on a single mass fractionation line (Fig. ESI-1†), only  $\delta^{57}\text{Fe}$  values are discussed in this paper. The resulting linear regression for  $\delta^{57}\text{Fe}$  vs.  $\delta^{56}\text{Fe}$  (Fig. ESI-1†) was:

$$\delta^{57}\text{Fe} = (0.016 \pm 0.009) + (1.456 \pm 0.022) \times \delta^{56}\text{Fe} \quad (R^2 = 0.996) \quad (6)$$

This is consistent with  $\delta^{57}\text{Fe}$  being equal to  $\sim 3/2$  of  $\delta^{56}\text{Fe}$ . Given the findings reported by Amor *et al.*<sup>64</sup> on mass-independent Fe isotope fractionation during magnetotactic bacteria metabolism, we searched for such mass independent Fe isotope fractionation on our experimental products to see whether such an effect could also occur with other type of bacteria. However, we could not detect any significant mass independent effect outside 0.1‰ uncertainty.

### 3. Results

#### 3.1. Isotope fractionation during Fe adsorption onto and assimilation by the bacteria

In case of  $\text{Fe}^{3+}$  adsorption onto EPS-rich and EPS-poor cells (see Table ESI-1†), the  $\delta^{57}\text{Fe}_{\text{cell}}$  was always higher than  $\delta^{57}\text{Fe}_{\text{solution}}$  (Table 1). However, for the two sets of experiments having the same pH range (close to 4 and 4.6–4.9, respectively), the




Fig. 1  $\delta^{57}\text{Fe}$  values of the initial solution calculated by the mass balance between cell and solution Fe concentrations and isotopic compositions. Asterisks represent the data on Fe isotope fractionation during adsorption onto cyanobacteria, as reported by Mulholland *et al.*, 2015.<sup>32</sup> Note that the isotopic offset during assimilation is strongly dependent on the initial Fe redox status – negative in case of Fe(III) and positive in case of Fe(II) as distinguished by red and blue rectangles, respectively.

$\Delta^{57}\text{Fe}_{\text{cell-solution}}$  is slightly higher for EPS-poor cultures compared to the EPS-rich ones, although the difference is comparable to uncertainties. Iron(III) adsorption onto *P. aureofaciens* at  $2.8 \leq \text{pH} \leq 6.0$  produced an enrichment of the cell surface in the heavier isotope with  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  ranging from +0.72 to +2.0‰ (Table 1 and Fig. 1). The EPS-poor culture exhibited a possible impact of pH on the isotopic offset during adsorption, which increased from +0.72‰ at pH of 2 to +1.96‰ at pH of 6 (Table 1 and Fig. 1).

The isotope fractionation induced by Fe(II) and Fe(III) assimilation during bacterial growth demonstrated the following features. The EPS-poor-8 cells exhibited higher  $\delta^{57}\text{Fe}_{\text{cell}}$  values of assimilated Fe compared to those of the EPS-rich-6 ones for the same pH range (Table 1), even though the starting  $\text{FeCl}_3$  of the former was isotopically lighter than the starting  $\text{FeCl}_3$  of the latter (Table 1). This feature of Fe(III) assimilation is not confirmed for the case of Fe(II) assimilation since the  $\delta^{57}\text{Fe}_{\text{cell}}$  values in EPS-poor-6 and EPS-rich-5 experiments were undistinguishable (Table 1). Intracellular assimilation of Fe(II) led to preferential uptake of heavier isotopes producing  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  of +0.77 and +0.83‰ for EPS-rich-5 and EPS-poor-6 experiments. In contrast, intracellular uptake of Fe(III) yielded an enrichment of the cell in lighter isotopes with  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  close to  $-1\%$  as calculated from EPS-poor-8 and EPS-rich-4,6,7 experiments (Table 1). It is noteworthy that the Fe(III) uptake from both initial  $\text{FeCl}_3$  and Fe-citrate salts yielded similar isotopic offsets (Tables ESI-1† and 1). Considering the propagated uncertainties (see Table 1), the magnitude of isotopic offset ( $\Delta^{57}\text{Fe}_{\text{cell-solution}}$ ) in the EPS-rich cells (from  $-1.01$  to  $-1.12\%$ ) was not distinguishable from that in the EPS-poor cells ( $\Delta^{57}\text{Fe}_{\text{cell-solution}} = -0.99\%$ ) for the experiments using Fe(III) salts (Tables ESI-1† and 1).



### 3.2. XAS characterization of Fe bound to soil bacteria

The structural characterization of Fe adsorbed onto *P. aurifaciens* was based on XAS (EXAFS and XANES) analysis of

samples as was also described in previous work.<sup>40</sup> These results are directly used for testing structural control on adsorbed Fe(III) and Fe(II) from aqueous solution as described in Section 4.1 below. The intracellularly assimilated Fe was not




Fig. 2 Fe K-edge XANES spectra at the Fe K-edge energies (A) and (B), presenting Fe K-edge oscillations (C) and Fourier transform magnitudes (D) of the Fe adsorbed on cells.



Table 2 Summary of Fe structural parameters of Fe(II) and Fe(III) assimilation experiments. Experimental details are listed in Table ESI-1

Sample	pH	Fe(III) phosphate (%)	Goethite (%)	$R$ (Å) ( $\pm 0.05$ Å)	$N$ (atom) ( $\pm 0.5$ )	$\sigma^2$ (Å <sup>2</sup> ) ( $\pm 0.001$ Å <sup>2</sup> )	$R$ -factor
EPS-rich-5	7.1	95	5	2.07	5.9	0.007	0.08
EPS-poor-6	7.8	100		2.07	5.1	0.006	0.13
EPS-rich-4	7.2	100		2.07	6.0	0.007	0.05
EPS-poor-8	7.1	100		2.06	5.6	0.005	0.15

characterized previously and newly obtained results are presented below.

The XANES spectra of assimilated Fe–bacteria samples are illustrated in Fig. 2 and Table 2. When comparing the samples with reference spectra, the analysis clearly shows that the unique organic component in most of the samples (and main component in all cases) is Fe(III)–phosphate, corresponding to Fe binding to the phosphoryl group. The relative proportion of Fe(III)–phosphate and goethite was estimated from linear combination fits (LCF analysis) of XANES spectra and using Fe(III) reference compounds. The choice of the best corresponding standard for each spectrum was made using automated combinational fitting (Athena software).

The EXAFS results allowed extracting the first shell of Fe (Table 2) which exhibited similar features among samples (Fig. 2). Note that the modeling of EXAFS spectra in biological samples is very difficult due to distorted geometries and the presence of different neighbors, like O, N and S, light elements with similar weak cross section values, in the nearest metal coordination shell. In this sense, XANES results may complement the EXAFS *via* identifying the ligands using the edge features.

The filtered Fourier transform curves (EXAFS signals filtered for the first coordination sphere) were fitted using Fe–O bonds. The inelastic loss factor was estimated/set, based on references and the EPS-rich-4 sample, as  $S_0^2 = 0.87$ . Results show a coordination number of around 6 in all cases. This is in good agreement with an octahedral geometry of 6 oxygen atoms surrounding Fe cations, with a Fe–O distance of  $\sim 2.07$  Å. We could not quantify the relative contribution of Fe(III)–phosphoryl or carboxyl complexes that are typical in samples of microorganisms,<sup>38,65,66</sup> although the samples showed an important contribution of Fe(III)–phosphate like groups in the XANES analysis (peak at  $\sim 1.9$ – $2.0$  Å<sup>-1</sup>). Overall, the EXAFS spectra confirmed the XANES results.

## 4. Discussion

### 4.1. Heavier isotope enrichment of cells during Fe(II) and Fe(III) adsorption

The adsorption of Fe onto *P. aureofaciens* allowed quantifying the adsorption constants and surface binding site concentrations. The total amount of proton-binding sites was two times higher in the EPS-rich cells compared to the EPS-poor ones (5.5 and 2.6 mmol g<sub>wet</sub><sup>-1</sup>), whereas the concentration of phosphoryl-like groups was higher in the EPS-poor cells and the concentration of carboxyl-like groups was higher in the EPS-rich cells. The adsorption of Fe(III) started at acidic pH ( $\sim 1.5$ ) and increased with the pH, following the “universal pH-edge” for

metals.<sup>67,68</sup> González *et al.*<sup>40</sup> provided chemical characterization of Fe<sup>3+</sup> adsorbed onto soil bacteria by X-ray absorption spectroscopy (XAS), and demonstrated that Fe(III)–phosphate was the predominant Fe-binding compound on the cell surfaces of the EPS-poor cultures, at pH from 2.0 to 6.0. In addition, the goethite-like component was present in the EPS-rich cells (pH from 3.0 to 4.6). Therefore, EXAFS spectra supported the XANES results.

The present study demonstrated that, during Fe(III) adsorption, the effect of pH was strongly pronounced for the EPS-poor culture ( $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  increased from +0.72‰ at pH = 2.0 to +1.96‰ at pH = 6.0; Table 1 and Fig. 1). The pH effect could not be tested for the EPS-rich culture, for which our two experiments yielded significantly different  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  values ( $+0.74 \pm 0.14$ ‰ and  $+1.29 \pm 0.30$ ‰; Table 1 and Fig. 1) at similar  $4 \leq \text{pH} \leq 4.6$ . The exact reasons for these differences between these EPS-rich and EPS-poor cultures in the circum-neutral pH range (Fig. 1) are not clear. It can be hypothesized that, in the case of EPS-poor strain, the phosphoryl moieties of the cell wall are not protected from solution and adsorption occurs essentially on these high affinity sites. This could produce stronger fractionation at higher pH, where phosphorylated groups are mostly deprotonated. However, this seems to be inconsistent with XAS-based bulk speciation of Fe at the cell surface, because the surface-bound Fe(III) was dominated by Fe–phosphoryl ligands, regardless of pH and the presence or absence of the EPS. At the same time, one should keep in mind that XAS treatment provides dominant (*ca.* 90%) speciation of an element. The remaining 10% are not detectable by spectroscopy but it is not excluded that they can be responsible for observed isotope fractionation, provided that the isotopic offsets of reactions with these minor compounds strongly exceed those of the major binding constituents of Fe. Recently, Oleinikova *et al.*<sup>69</sup> studied the adsorption of organo-ferric colloids onto *P. aureofaciens* and demonstrated an enrichment in  $\delta^{57}\text{Fe}$  (+0.4‰) of cell surfaces compared to the remaining solution. Consistent with the results of Gonzalez *et al.*<sup>40</sup> and of the present study, this could be explained by the dominance of Fe(III)–complexes with phosphoryl groups on cell surfaces. Therefore, the only possible explanation for the observed difference in isotopic offset between the aqueous solution and cell biomass of EPS-rich and EPS-poor cultures is the shielding effect of carboxylate-rich EPS on aqueous Fe ions. These rather inert EPS layers can prevent direct interaction and binding of Fe<sup>2+</sup>(aq.) and Fe<sup>3+</sup>(aq.) with the strong phosphoryl moieties of the external cell membrane.

Globally, the XAS analyses of adsorbed Fe demonstrated a dominance of Fe(III)–phosphate complexes at the cell surface





regardless of the identity of strain and the chemical composition of solution. Therefore, we suggest that the heavier isotope enrichment of the cell surface relative to aqueous solution is due to strong Fe(III)–phosphoryl surface complexes compared to FeOH<sup>2+</sup>, Fe(OH)<sub>2</sub><sup>+</sup> and Fe(III)–organic complexes in aqueous solution. Presumably, these phosphoryl surface complexes exhibit shorter bond lengths and higher symmetry than aqueous Fe(III) inorganic species (FeOH<sup>2+</sup>, Fe(OH)<sub>2</sub><sup>+</sup>, FeCl<sub>2</sub><sup>+</sup>) and Fe(III)–EPS complexes. Note that the isotopic measurements of Fe(III) adsorption rule out the possibility of Fe(OH)<sub>3</sub> precipitation at the cell surface, because the observed isotopic offset ( $\Delta^{57}\text{Fe}_{\text{cell-solution}} = +0.70$  to  $+2.1\%$ ) is opposite to the sign of the Fe<sup>3+</sup><sub>(aq.)</sub> → Fe(OH)<sub>3(solid)</sub> reaction ( $\Delta^{57}\text{Fe}_{\text{Fe(III)hydroxide-Fe}^{3+}\text{(aq.)}}$  from  $-0.2$  to  $-1.5\%$ ), with more fractionated values at slower reaction rates.<sup>70</sup>

In contrast, in case of Fe(II) adsorption on EPS-poor cultures ( $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  of  $+3.16\%$  measured in this study; EPS-poor-F2; Tables ESI-1† and 1, and Fig. 1), in addition to surface complexation with stronger than aqueous ligand, Fe(II) partial oxidation at the cell surface may lead to overall cell enrichment in heavier isotopes. Indeed, it is known that the Fe(II)<sub>aq.</sub> → Fe(III)–hydroxide (solid) reaction exhibits a  $\Delta^{57}\text{Fe}_{\text{solid-solution}}$  of  $+2.3 \pm 0.3\%$ ,<sup>16,19,20</sup> although an isotopic offset as high as  $+4.8\%$  was also reported in case of hematite precipitation.<sup>71</sup>

#### 4.2. Fe assimilation by bacterial cells

Intracellular assimilation of Fe(II) enriched the cells in heavier isotopes producing undistinguishable  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  of  $+0.77 \pm 0.02\%$  and  $+0.83 \pm 0.08\%$  for EPS-rich and EPS-poor cultures, respectively (Tables ESI-1† and 1). Two possible mechanisms can be responsible for this enrichment. First, partial oxidation of Fe(II) on the cell surface (not removed by EDTA treatment) or inside the cells, favorable at circumneutral pH of nutrient media (7.2–7.5), which can yield sizable enrichment in heavier isotopes at the surface, as follows from  $\Delta^{57}\text{Fe}_{\text{solid-solution}}$  of  $+2.3 \pm 0.3\%$  for the Fe(II)<sub>aq.</sub> → Fe(III)–hydroxide (solid).<sup>16,19,20</sup> However, in order for this effect to be pronounced, at least 30% of Fe(II) should be transformed into Fe(III) (15–20% in the case where the isotopic offset of Fe<sup>2+</sup> oxidation equals  $+4.8\%$  as reported by ref. 71). Second, higher symmetry and shorter bonds of Fe(II) in the cells compared to Fe(II)–lactate solutions may be responsible for cell enrichment in heavy isotopes relative to the nutrient media. Unfortunately, we could not resolve the 2nd neighbor symmetry in this type of sample due to the relatively low Fe concentration in the nutrient solution.

In contrast to Fe(II), cellular uptake of Fe(III) produced an enrichment of the cell biomass in lighter isotopes with  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  around  $-1\%$ , regardless of the presence or not of EPS and the form of Fe(III) in the nutrient media (chloride or citrate; Tables ESI-1† and 1, and Fig. 1). Interestingly, the long-term bacterial uptake of Fe(III) oxy(hydr)oxide colloids and organic complexes by *P. aureofaciens* produced  $\Delta^{57}\text{Fe}_{\text{cell-solution}}$  of  $-0.7\%$ ,<sup>69</sup> which is similar to the results of the present study. Iron can be involved in a number of both extra- and intracellular redox reactions, under strong control of organic binding ligands. For example, Fe(II) can be oxidized and Fe(III)

can be reduced within the cells,<sup>47</sup> where both inorganic insoluble Fe, inorganic soluble Fe and organically complexed soluble Fe can be adsorbed onto and transported into the cells. In addition, Fe can be adsorbed onto cell surfaces prior to being reduced to Fe(II) or to being incorporated.<sup>4</sup> The preference of cell intracellular compartments with lighter Fe isotopes may be controlled by two factors. First, Fe(III) in the nutrient media can be reduced to Fe(II). However, the lack of XAS confirmation of this redox transformation and also lack of information on Fe(II) speciation in cell compartments cannot support this possibility. Another mechanism of lighter isotope enrichment of cell biomass could be a loss of symmetry of assimilated Fe(III) relative to octahedrally coordinated aqueous Fe<sup>3+</sup> ions and longer bonds of intracellular ions relative to aqueous Fe(III)–citrate or hydroxo-complexes.

Overall, although the obtained results are fully consistent with available data on Fe isotope fractionation during adsorption and redox transformation, we could not evidence a straightforward structural control on Fe isotope fractionation between aqueous solution and soil bacteria. The versatile nature of Fe(II) and Fe(III) fractionation without the distinct effect of pH and surface exopolysaccharide coverage suggests that, in soil environments, the nature of microorganisms (presence or not of EPS) which adsorb and assimilate Fe and solution pH (between 4 and 6) exhibit a subordinate effect on Fe isotopic signature in the microbial biomass. As such, Fe isotope fractionations during its interaction with heterotrophic bacteria, mineral or organic sediments will be primarily governed by Fe complexation with DOM and Fe redox state in soil or sediment porewater.<sup>72,73</sup>

## 5. Conclusions

With the aim of assessing the role of heterotrophic bacteria in Fe isotope fractionation in aqueous solutions, we measured the isotopic partitioning during Fe(II) and Fe(III) adsorption onto and assimilation by a common soil bacterium. We notably compared the obtained isotopic offsets between the fluid and the biomass with Fe speciation in the nutrient media and Fe chemical status on the cell surface and in the cell interior.

Adsorption of Fe(III) led to cell surface enrichment in heavier isotopes, tentatively linked to the stronger and more symmetrical complex of Fe(III) with surface phosphoryl moieties compared to hydroxyl complexes of Fe(III) in aqueous solution. This was especially pronounced for the EPS-poor culture in circumneutral solutions, when the binding sites of the cell membrane were not protected by the carboxylates of EPS. Adsorption of Fe(II) could yield some Fe(III) hydroxides at the surface, which produced even stronger enrichment of cells (by up to  $3.2\%$ ) in heavier isotopes.

Iron(III) assimilation from nutrient media produced *ca.*  $-1\%$  cell enrichment in lighter isotopes. This isotopic offset was similar for EPS-rich and EPS-poor cultures and Fe(III) citrate or Fe(III) chloride in the initial nutrient media. In contrast, Fe(II) uptake by bacteria yielded an enrichment of cells by *ca.*  $+0.8\%$  in heavier isotopes, following the formation of Fe(III) hydroxides or other minerals inside the cells or due to formation of



stronger bonds with phosphoryl moieties in cell organelles compared to nutrient media.

Taken together, the results obtained indicate that the mechanisms controlling Fe isotope fractionation in the soil fluid–bacterial system are better constrained. In natural settings, the  $\delta^{57}\text{Fe}$  in solution or microbial biomass may vary from *ca.*  $-1$  to  $+3\text{‰}$  depending primarily on the Fe redox status in soil and sediment porewater and these changes may occur over a short time scale (hours to days) regardless of pH and the presence or not of microbial extracellular substances.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

E. E. Emnova (Moldova Academy of Science) is thanked for providing the *P. aureofaciens* strain. F. P. is funded by CNRS. O. P. acknowledges the support from the TSU Development Programme («Priority-2030»).

## References

- 1 K. H. Nealson and D. Saffarini, Iron and manganese in anaerobic respiration: environmental significance, physiology, and regulation, *Annu. Rev. Microbiol.*, 1994, **48**, 311–344.
- 2 D. R. Lovley, Microbial Fe(III) reduction in subsurface environments, *FEMS Microbiol. Rev.*, 1997, **20**, 305–313, DOI: [10.1111/j.1574-6976.1997.tb00316.x](https://doi.org/10.1111/j.1574-6976.1997.tb00316.x).
- 3 B. Thamdrup, in *Advances in microbial ecology*, Springer, 2000, pp. 41–84.
- 4 F. M. M. Morel, A. B. Kustka and Y. Shaked, The role of unchelated Fe in the iron nutrition of phytoplankton, *Limnol. Oceanogr.*, 2008, **53**, 400–404, DOI: [10.4319/lo.2008.53.1.0400](https://doi.org/10.4319/lo.2008.53.1.0400).
- 5 R. M. Cornell and U. Schwertmann, *The iron oxides: structure, properties, reactions, occurrences and uses*, John Wiley & Sons, 2003.
- 6 J. H. Martin, K. H. Coale, K. S. Johnson, S. E. Fitzwater, R. M. Gordon, S. J. Tanner, C. N. Hunter, V. A. Elrod, J. L. Nowicki, T. L. Coley, R. T. Barber, S. Lindley, A. J. Watson, K. Van Scoy, C. S. Law, M. I. Liddicoat, R. Ling, T. Stanton, J. Stockel, C. Collins, A. Anderson, R. Bidigare, M. Ondrusek, M. Latasa, F. J. Millero, K. Lee, W. Yao, J. Z. Zhang, G. Friederich, C. Sakamoto, F. Chavez, K. Buck, Z. Kolber, R. Greene, P. Falkowski, S. W. Chisholm, F. Hoge, R. Swift, J. Yungel, S. Turner, P. Nightingale, A. Hatton, P. Liss and N. W. Tindale, Testing the iron hypothesis in ecosystems of the equatorial Pacific Ocean, *Nature*, 1994, **371**, 123–129, DOI: [10.1038/371123a0](https://doi.org/10.1038/371123a0).
- 7 K. H. Coale, K. S. Johnson, S. E. Fitzwater, R. M. Gordon, S. J. Tanner, F. P. Chavez, L. Ferioli, C. Sakamoto, P. Rogers, F. Millero, P. Steinberg, P. Nightingale, D. Cooper, W. P. Cochlan, M. R. Landry, J. Constantinou, G. Rollwagen, A. Trasvina and R. Kudela, A massive phytoplankton bloom induced by an ecosystem-scale iron fertilization experiment in the equatorial Pacific Ocean, *Nature*, 1996, **383**, 495, DOI: [10.1038/383495a0](https://doi.org/10.1038/383495a0).
- 8 X. Liu and F. J. Millero, The solubility of iron in seawater, *Mar. Chem.*, 2002, **77**, 43–54, DOI: [10.1016/S0304-4203\(01\)00074-3](https://doi.org/10.1016/S0304-4203(01)00074-3).
- 9 E. Murad and W. R. Fischer, in *Iron in soils and clay minerals*, Springer, 1988, pp. 1–18.
- 10 F. J. Millero, S. Sotolongo and M. Izaguirre, The oxidation kinetics of Fe(II) in seawater, *Geochim. Cosmochim. Acta*, 1987, **51**, 793–801, DOI: [10.1016/0016-7037\(87\)90093-7](https://doi.org/10.1016/0016-7037(87)90093-7).
- 11 J. M. Santana-Casiano, M. González-Dávila and F. J. Millero, Oxidation of nanomolar levels of Fe(II) with oxygen in natural waters, *Environ. Sci. Technol.*, 2005, **39**, 2073–2079, DOI: [10.1021/es049748y](https://doi.org/10.1021/es049748y).
- 12 A. G. González, J. M. Santana-Casiano, N. Pérez and M. González-Dávila, Oxidation of Fe(II) in natural waters at high nutrient concentrations, *Environ. Sci. Technol.*, 2010, **44**, 8095–8101, DOI: [10.1021/es1009218](https://doi.org/10.1021/es1009218).
- 13 E. L. Rue and K. W. Bruland, Complexation of iron(III) by natural organic ligands in the Central North Pacific as determined by a new competitive ligand equilibration/adsorptive cathodic stripping, *Mar. Chem.*, 1995, **50**, 117–138, DOI: [10.1016/0304-4203\(95\)00031-L](https://doi.org/10.1016/0304-4203(95)00031-L).
- 14 B. Wu, W. Amelung, Y. Xing, R. Bol and A. E. Berns, Iron cycling and isotope fractionation in terrestrial ecosystems, *Earth-Sci. Rev.*, 2019, **190**, 323–352, DOI: [10.1016/j.earscirev.2018.12.012](https://doi.org/10.1016/j.earscirev.2018.12.012).
- 15 C. Johnson, B. Beard and S. Weyer, *Iron geochemistry: an isotopic perspective*, Springer, 2020.
- 16 T. D. Bullen, A. F. White, C. W. Childs, D. V. Vivit and M. S. Schulz, Demonstration of significant abiotic iron isotope fractionation in nature, *Geology*, 2001, **29**, 699–702, DOI: [10.1130/0091-7613\(2001\)029<0699:DOSAII>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0699:DOSAII>2.0.CO;2).
- 17 A. D. Anbar, Molybdenum stable isotopes: observations, interpretations and directions, *Rev. Mineral. Geochem.*, 2004, **55**, 429–454, DOI: [10.2138/gsrmg.55.1.429](https://doi.org/10.2138/gsrmg.55.1.429).
- 18 A. Gronstal, V. Pearson, A. Kappler, C. Dooris, M. Anand, F. Poitrasson, T. P. Kee and C. S. Cockell, Laboratory experiments on the weathering of iron meteorites and carbonaceous chondrites by iron-oxidizing bacteria, *Meteorit. Planet. Sci.*, 2009, **44**, 233–247, DOI: [10.1111/j.1945-5100.2009.tb00731.x](https://doi.org/10.1111/j.1945-5100.2009.tb00731.x).
- 19 B. L. Beard, R. M. Handler, M. M. Scherer, L. Wu, A. D. Czaja, A. Heimann and C. M. Johnson, Iron isotope fractionation between aqueous ferrous iron and goethite, *Earth Planet. Sci. Lett.*, 2010, **295**, 241–250, DOI: [10.1016/j.epsl.2010.04.006](https://doi.org/10.1016/j.epsl.2010.04.006).
- 20 L. R. Croal, C. M. Johnson, B. L. Beard and D. K. Newman, Iron isotope fractionation by Fe(II)-oxidizing photoautotrophic bacteria, *Geochim. Cosmochim. Acta*, 2004, **68**, 1227–1242, DOI: [10.1016/j.gca.2003.09.011](https://doi.org/10.1016/j.gca.2003.09.011).
- 21 E. D. Swanner, T. Bayer, W. Wu, L. Hao, M. Obst, A. Sundman, J. M. Byrne, F. M. Michel, I. C. Kleinhans, A. Kappler and R. Schoenberg, Iron isotope fractionation during Fe(II) oxidation mediated by the oxygen-producing



- marine cyanobacterium *Synechococcus* PCC 7002, *Environ. Sci. Technol.*, 2017, **51**, 4897–4906, DOI: [10.1021/acs.est.6b05833](https://doi.org/10.1021/acs.est.6b05833).
- 22 G. A. Icopini, A. D. Anbar, S. S. Ruebush, M. Tien and S. L. Brantley, Iron isotope fractionation during microbial reduction of iron: the importance of adsorption, *Geology*, 2004, **32**, 205–208, DOI: [10.1130/G20184.1](https://doi.org/10.1130/G20184.1).
- 23 H. A. Crosby, C. M. Johnson, E. E. Roden and B. L. Beard, Coupled Fe(II)-Fe(III) electron and atom exchange as a mechanism for Fe isotope fractionation during dissimilatory iron oxide reduction, *Environ. Sci. Technol.*, 2005, **39**, 6698–6704, DOI: [10.1021/es0505346](https://doi.org/10.1021/es0505346).
- 24 B. L. Beard, C. M. Johnson, L. Cox, H. Sun, K. H. Nealson and C. Aguilar, Iron isotope biosignatures, *Science*, 1999, **285**, 1889–1892, DOI: [10.1126/science.285.5435.1889](https://doi.org/10.1126/science.285.5435.1889).
- 25 R. A. Wiesli, B. L. Beard and C. M. Johnson, Experimental determination of Fe isotope fractionation between aqueous Fe(II), siderite and 'green rust' in abiotic systems, *Chem. Geol.*, 2004, **211**, 343–362, DOI: [10.1016/j.chemgeo.2004.07.002](https://doi.org/10.1016/j.chemgeo.2004.07.002).
- 26 C. M. Johnson, B. L. Beard, E. E. Roden, D. K. Newman and K. H. Nealson, Isotopic constraints on biogeochemical cycling of Fe, *Rev. Mineral. Geochem.*, 2004, **55**, 359–408, DOI: [10.2138/gsrmg.55.1.359](https://doi.org/10.2138/gsrmg.55.1.359).
- 27 C. M. Johnson, E. E. Roden, S. A. Welch and B. L. Beard, Experimental constraints on Fe isotope fractionation during magnetite and Fe carbonate formation coupled to dissimilatory hydrous ferric oxide reduction, *Geochim. Cosmochim. Acta*, 2005, **69**, 963–993, DOI: [10.1016/j.gca.2004.06.043](https://doi.org/10.1016/j.gca.2004.06.043).
- 28 K. O. Konhauser, W. S. Fyfe, F. G. Ferris and T. J. Beveridge, Metal sorption and mineral precipitation by bacteria in two Amazonian river systems: Rio Solimoes and Rio Negro, Brazil, *Geology*, 1993, **21**, 1103–1106, DOI: [10.1130/0091-7613\(1993\)021<1103:MSAMPB>2.3.CO;2](https://doi.org/10.1130/0091-7613(1993)021<1103:MSAMPB>2.3.CO;2).
- 29 R. J. M. Hudson and F. M. M. Morel, Distinguishing between extra- and intracellular iron in marine phytoplankton, *Limnol. Oceanogr.*, 1989, **34**, 1113–1120, DOI: [10.4319/lo.1989.34.6.1113](https://doi.org/10.4319/lo.1989.34.6.1113).
- 30 N. Yee, L. G. Benning, V. R. Phoenix and F. G. Ferris, Characterization of metal–cyanobacteria sorption reactions: a combined macroscopic and infrared spectroscopic investigation, *Environ. Sci. Technol.*, 2004, **38**, 775–782, DOI: [10.1021/es0346680](https://doi.org/10.1021/es0346680).
- 31 N. Teutsch, U. Von Gunten, D. Porcelli, O. A. Cirpka and A. N. Halliday, Adsorption as a cause for iron isotope fractionation in reduced groundwater, *Geochim. Cosmochim. Acta*, 2005, **69**, 4175–4185, DOI: [10.1016/j.gca.2005.04.007](https://doi.org/10.1016/j.gca.2005.04.007).
- 32 D. S. Mulholland, F. Poitrasson, L. S. Shirokova, A. G. González, O. S. Pokrovsky, G. R. Boaventura and L. C. Vieira, Iron isotope fractionation during Fe(II) and Fe(III) adsorption on cyanobacteria, *Chem. Geol.*, 2015, **400**, 24–33, DOI: [10.1016/j.chemgeo.2015.01.017](https://doi.org/10.1016/j.chemgeo.2015.01.017).
- 33 K. Dideriksen, J. A. Baker and S. L. S. Stipp, Equilibrium Fe isotope fractionation between inorganic aqueous Fe(III) and the siderophore complex, Fe(III)-desferrioxamine B, *Earth Planet. Sci. Lett.*, 2008, **269**, 280–290, DOI: [10.1016/j.epsl.2008.02.022](https://doi.org/10.1016/j.epsl.2008.02.022).
- 34 J. B. Fein, C. J. Daughney, N. Yee and T. A. Davis, A chemical equilibrium model of metal adsorption onto bacterial surfaces, *Geochim. Cosmochim. Acta*, 1997, **61**, 3319–3328, DOI: [10.1016/S0016-7037\(97\)00166-X](https://doi.org/10.1016/S0016-7037(97)00166-X).
- 35 C. J. Daughney, J. B. Fein and N. Yee, A comparison of the thermodynamics of metal adsorption onto two common bacteria, *Chem. Geol.*, 1998, **144**, 161–176, DOI: [10.1016/S0009-2541\(97\)00124-1](https://doi.org/10.1016/S0009-2541(97)00124-1).
- 36 O. S. Pokrovsky, R. E. Martinez, S. V. Golubev, E. I. Kompantseva and L. S. Shirokova, Adsorption of metals and protons on *Gloeocapsa* sp. cyanobacteria: a surface speciation approach, *Appl. Geochem.*, 2008, **23**, 2574–2588, DOI: [10.1016/j.apgeochem.2008.05.007](https://doi.org/10.1016/j.apgeochem.2008.05.007).
- 37 O. S. Pokrovsky, G. S. Pokrovski and A. Feurtet-Mazel, A structural study of cadmium interaction with aquatic microorganisms, *Environ. Sci. Technol.*, 2008, **42**, 5527–5533, DOI: [10.1021/es800521a](https://doi.org/10.1021/es800521a).
- 38 O. S. Pokrovsky, J. Viers, E. E. Emnova, E. I. Kompantseva and R. Freydier, Copper isotope fractionation during its interaction with soil and aquatic microorganisms and metal oxy(hydr)oxides: possible structural control, *Geochim. Cosmochim. Acta*, 2008, **72**, 1742–1757, DOI: [10.1016/j.gca.2008.01.018](https://doi.org/10.1016/j.gca.2008.01.018).
- 39 A. G. González, L. S. Shirokova, O. S. Pokrovsky, E. E. Emnova, R. E. Martínez, J. M. Santana-Casiano, M. González-Dávila and G. S. Pokrovski, Adsorption of copper on *Pseudomonas aureofaciens*: protective role of surface exopolysaccharides, *J. Colloid Interface Sci.*, 2010, **350**, 305–314, DOI: [10.1016/j.jcis.2010.06.020](https://doi.org/10.1016/j.jcis.2010.06.020).
- 40 A. G. González, O. S. S. Pokrovsky, F. Jiménez-Villacorta, L. S. Shirokova, J. M. Santana-Casiano, M. González-Dávila and E. E. Emnova, Iron adsorption onto soil and aquatic bacteria: XAS structural study, *Chem. Geol.*, 2014, **372**, 32–45, DOI: [10.1016/j.chemgeo.2014.02.013](https://doi.org/10.1016/j.chemgeo.2014.02.013).
- 41 A. Coutaud, M. Meheut, J. Viers, J.-L. Rols and O. S. Pokrovsky, Zn isotope fractionation during interaction with phototrophic biofilm, *Chem. Geol.*, 2014, **390**, 46–60, DOI: [10.1016/j.chemgeo.2014.10.004](https://doi.org/10.1016/j.chemgeo.2014.10.004).
- 42 M. Coutaud, M. Méheut, P. Glatzel, G. S. Pokrovski, J. Viers, J.-L. Rols and O. S. Pokrovsky, Small changes in Cu redox state and speciation generate large isotope fractionation during adsorption and incorporation of Cu by a phototrophic biofilm, *Geochim. Cosmochim. Acta*, 2018, **220**, 1–18, DOI: [10.1016/j.gca.2017.09.018](https://doi.org/10.1016/j.gca.2017.09.018).
- 43 F. C. A. Kafantaris and D. M. Borrok, Zinc isotope fractionation during surface adsorption and intracellular incorporation by bacteria, *Chem. Geol.*, 2014, **366**, 42–51, DOI: [10.1016/j.chemgeo.2013.12.007](https://doi.org/10.1016/j.chemgeo.2013.12.007).
- 44 J. H. Jamieson-Hanes, H. K. Shrimpton, H. Veeramani, C. J. Ptacek, A. Lanzirrotti, M. Newville and D. W. Blowes, Evaluating zinc isotope fractionation under sulfate reducing conditions using a flow-through cell and in situ XAS analysis, *Geochim. Cosmochim. Acta*, 2017, **203**, 1–14, DOI: [10.1016/j.gca.2016.12.034](https://doi.org/10.1016/j.gca.2016.12.034).





- 45 F. C. Fang, E. R. Frawley, T. Tapscott and A. Vázquez-Torres, Bacterial stress responses during host infection, *Cell Host Microbe*, 2016, **20**, 133–143, DOI: [10.1016/j.chom.2016.07.009](https://doi.org/10.1016/j.chom.2016.07.009).
- 46 D. J. R. Lane, A. M. Merlot, M.-H. Huang, D.-H. Bae, P. J. Jansson, S. Sahni, D. S. Kalinowski and D. R. Richardson, Cellular iron uptake, trafficking and metabolism: key molecules and mechanisms and their roles in disease, *Biochim. Biophys. Acta, Mol. Cell Res.*, 2015, **1853**, 1130–1144, DOI: [10.1016/j.bbamcr.2015.01.021](https://doi.org/10.1016/j.bbamcr.2015.01.021).
- 47 A. Marchetti and M. T. Maldonado, Iron, *The Physiology of Microalgae*, ed. M. A. Borowitzka, J. Beardall and J. A. Raven Springer International Publishing, Cham, 2016, pp. 233–279.
- 48 G. M. Gadd, Metals, minerals and microbes: geomicrobiology and bioremediation, *Microbiology*, 2010, **156**, 609–643, DOI: [10.1099/mic.0.037143-0](https://doi.org/10.1099/mic.0.037143-0).
- 49 E. E. Emnova, R. III, D. A. Dascualinc, S. Toma, O. Gojnetschi and I. Seniscovscaia, *Soil invertase and levansucrase activities under soil water stress, elevated copper level, and soybean seed inoculation with the levan-producing bacteria Pseudomonas aureofaciens*, 2005, vol. 1–3, pp. 128–136.
- 50 J. M. Meyer and M. A. Abdallah, The fluorescent pigment of *Pseudomonas fluorescens*: biosynthesis, purification and physicochemical properties, *J. Gen. Microbiol.*, 1978, **107**, 319–328, DOI: [10.1099/00221287-107-2-319](https://doi.org/10.1099/00221287-107-2-319).
- 51 U. Behrens and M. Ringpfeil, in *Bericht über das Jahr 1963*, Springer, 1964, pp. 420–429.
- 52 E. E. Emnova, L. D. Varbanets, V. N. Vasiliev, A. G. Ciocarlan, O. S. Brovarskaia, N. J. Caunova, O. G. Ganea and S. I. Toma, Properties of exopolysaccharides from rhizospheric fluorescent bacteria of *Pseudomonas* genus, *Bulletin of Moldovan Academy of Sciences Life Sciences*, 2007, **1**(310), 14–20.
- 53 D. A. Fowle and J. B. Fein, Experimental measurements of the reversibility of metal–bacteria adsorption reactions, *Chem. Geol.*, 2000, **168**, 27–36, DOI: [10.1016/S0009-2541\(00\)00188-1](https://doi.org/10.1016/S0009-2541(00)00188-1).
- 54 K. Knauer, R. Behra and L. Sigg, Adsorption and uptake of copper by the green alga *Scenedesmus subspicatus* (chlorophyta), *J. Phycol.*, 1997, **33**, 596–601, DOI: [10.1111/j.0022-3646.1997.00596.x](https://doi.org/10.1111/j.0022-3646.1997.00596.x).
- 55 S. Le Faucheur, R. Behra and L. Sigg, Thiol and metal contents in periphyton exposed to elevated copper and zinc concentrations: a field and microcosm study, *Environ. Sci. Technol.*, 2005, **39**, 8099–8107, DOI: [10.1021/es050303z](https://doi.org/10.1021/es050303z).
- 56 M. Ma, W. Zhu, Z. Wang and G. J. Witkamp, Accumulation, assimilation and growth inhibition of copper on freshwater alga (*Scenedesmus subspicatus* 86.81 SAG) in the presence of EDTA and fulvic acid, *Aquat. Toxicol.*, 2003, **63**, 221–228, DOI: [10.1016/S0166-445X\(02\)00179-0](https://doi.org/10.1016/S0166-445X(02)00179-0).
- 57 K. A. C. De Schampelaere, J. L. Stauber, K. L. Wilde, S. J. Markich, P. L. Brown, N. M. Franklin, N. M. Creighton and C. R. Janssen, Toward a biotic ligand model for freshwater green algae: surface-bound and internal copper are better predictors of toxicity than free Cu<sup>2+</sup>-ion activity when pH is varied, *Environ. Sci. Technol.*, 2005, **39**, 2067–2072, DOI: [10.1021/es049256l](https://doi.org/10.1021/es049256l).
- 58 F. W. E. Strelow, Improved separation of iron from copper and other elements by anion-exchange chromatography on a 4% cross-linked resin with high concentrations of hydrochloric acid, *Talanta*, 1980, **27**, 727–732, DOI: [10.1016/0039-9140\(80\)80166-4](https://doi.org/10.1016/0039-9140(80)80166-4).
- 59 F. Poitrasson, A. N. Halliday, D.-C. Lee, S. Levasseur and N. Teutsch, Iron isotope differences between Earth, Moon, Mars and Vesta as possible records of contrasted accretion mechanisms, *Earth Planet. Sci. Lett.*, 2004, **223**, 253–266, DOI: [10.1016/j.epsl.2004.04.032](https://doi.org/10.1016/j.epsl.2004.04.032).
- 60 F. Poitrasson and R. Freyrier, Heavy iron isotope composition of granites determined by high resolution MC-ICP-MS, *Chem. Geol.*, 2005, **222**, 132–147, DOI: [10.1016/j.chemgeo.2005.07.005](https://doi.org/10.1016/j.chemgeo.2005.07.005).
- 61 K. Dideriksen, J. A. Baker and S. L. S. Stipp, Fe isotope fractionation between inorganic aqueous Fe(III) and a Fe siderophore complex, *Mineral. Mag.*, 2008, **72**, 313–316, DOI: [10.1180/minmag.2008.072.1.313](https://doi.org/10.1180/minmag.2008.072.1.313).
- 62 M. A. Fehr, P. S. Andersson, U. Hålenius and C.-M. Mörth, Iron isotope variations in holocene sediments of the Gotland Deep, Baltic Sea, *Geochim. Cosmochim. Acta*, 2008, **72**, 807–826, DOI: [10.1016/j.gca.2007.11.033](https://doi.org/10.1016/j.gca.2007.11.033).
- 63 S. Aebischer, C. Cloquet, J. Carignan, C. Maurice and R. Pienitz, Disruption of the geochemical metal cycle during mining: multiple isotope studies of lake sediments from Schefferville, subarctic Québec, *Chem. Geol.*, 2015, **412**, 167–178, DOI: [10.1016/j.chemgeo.2015.07.028](https://doi.org/10.1016/j.chemgeo.2015.07.028).
- 64 M. Amor, V. Busigny, P. Louvat, A. Gélabert, P. Cartigny, M. Durand-Dubief, G. Ona-Nguema, E. Alphandéry, I. Chebbi and F. Guyot, Mass-dependent and -independent signature of Fe isotopes in magnetotactic bacteria, *Science*, 2016, **352**(6286), 705–708, DOI: [10.1126/science.aad7632](https://doi.org/10.1126/science.aad7632).
- 65 M. I. Boyanov, S. D. Kelly, K. M. Kemner, B. A. Bunker, J. B. Fein and D. A. Fowle, Adsorption of cadmium to *Bacillus subtilis* bacterial cell walls: a pH-dependent X-ray absorption fine structure spectroscopy study, *Geochim. Cosmochim. Acta*, 2003, **67**, 3299–3311, DOI: [10.1016/S0016-7037\(02\)01343-1](https://doi.org/10.1016/S0016-7037(02)01343-1).
- 66 O. S. Pokrovsky, G. S. Pokrovski, A. Gélabert, J. Schott and A. Boudou, Speciation of Zn associated with diatoms using X-ray absorption spectroscopy, *Environ. Sci. Technol.*, 2005, **39**, 4490–4498, DOI: [10.1021/es0480419](https://doi.org/10.1021/es0480419).
- 67 A. G. González and O. S. Pokrovsky, Metal adsorption on mosses: toward a universal adsorption model, *J. Colloid Interface Sci.*, 2014, **415**, 169–178, DOI: [10.1016/j.jcis.2013.10.028](https://doi.org/10.1016/j.jcis.2013.10.028).
- 68 N. Yee and J. Fein, Cd adsorption onto bacterial surfaces: a universal adsorption edge?, *Geochim. Cosmochim. Acta*, 2001, **65**, 2037–2042, DOI: [10.1016/S0016-7037\(01\)00587-7](https://doi.org/10.1016/S0016-7037(01)00587-7).
- 69 O. V. Oleinikova, F. Poitrasson, O. Y. Drozdova, L. S. Shirokova, S. A. Lapitskiy and O. S. Pokrovsky, Iron Isotope Fractionation during Bio- and Photodegradation of Organoferric Colloids in Boreal Humic Waters, *Environ. Sci. Technol.*, 2019, **53**, 11183–11194, DOI: [10.1021/acs.est.9b02797](https://doi.org/10.1021/acs.est.9b02797).





- 70 N. Balci, T. D. Bullen, K. Witte-Lien, W. C. Shanks, M. Motelica and K. W. Mandernack, Iron isotope fractionation during microbially stimulated Fe(II) oxidation and Fe(III) precipitation, *Geochim. Cosmochim. Acta*, 2006, **70**, 622–639, DOI: [10.1016/j.gca.2005.09.025](https://doi.org/10.1016/j.gca.2005.09.025).
- 71 L. Wu, B. L. Beard, E. E. Roden and C. M. Johnson, Stable Iron Isotope Fractionation between Aqueous Fe(II) and Hydrated Ferric Oxide, *Environ. Sci. Technol.*, 2011, 1847–1852, DOI: [10.1021/es103171x](https://doi.org/10.1021/es103171x).
- 72 W. B. Homoky, S. Severmann, R. A. Mills, P. J. Statham and G. R. Fones, Pore-fluid Fe isotopes reflect the extent of benthic Fe redox recycling: evidence from continental shelf and deep-sea sediments, *Geology*, 2009, **37**(8), 751–754.
- 73 J. K. Klar, W. B. Homoky, P. J. Statham, A. J. Birchill, E. L. Harris, E. M. S. Woodward, B. Silburn, M. J. Cooper, R. H. James, D. P. Connelly, F. Chever, A. Lichtschlag and C. Graves, Stability of dissolved and soluble Fe(II) in shelf sediment pore waters and release to an oxic water column, *Biogeochemistry*, 2017, **135**, 49–67.

