

Cite this: *Chem. Sci.*, 2024, 15, 19787

All publication charges for this article have been paid for by the Royal Society of Chemistry

Mg²⁺-driven selection of natural phosphatidic acids in primitive membranes†

Krishnakavya Thaipurayil Madanan,^a Yuhan Li,^a Valeria J. Boide-Trujillo,^a David A. Russell^{ab} and Claudia Bonfio^{*ab}

Biological membranes are composed exclusively of phospholipids comprising glycerol-1-phosphate or glycerol-3-phosphate. By contrast, primitive membranes would have likely been composed of heterogeneous mixtures of phospholipids, including non-natural analogues comprising glycerol-2-phosphate, as delivered by prebiotic synthesis. Thus, it is not clear how the selection of natural phospholipids could have come about. Here we show how differences in supramolecular properties, but not molecular properties, could have driven the selection of natural phosphatidic acids in primitive membranes. First, we demonstrate that at the molecular level it is unlikely that any prebiotic synthesis or hydrolysis pathway would have enabled the selection of natural phosphatidic acids. Second, we report that at the supramolecular level, natural phospholipids display a greater tendency to self-assemble in more packed and rigid membranes than non-natural analogues of the same chain length. Finally, taking advantage of these differences, we highlight that Mg²⁺, but not Na⁺, K⁺, Ca²⁺ or Zn²⁺, drives the selective precipitation of non-natural phosphatidic acids from heterogeneous mixtures obtained by prebiotic synthesis, leaving membranes proportionally enriched in natural phosphatidic acids. Our findings delineate a plausible pathway by which the transition towards biological membranes could have occurred under conditions compatible with prebiotic metal-driven processes, such as non-enzymatic RNA polymerization.

Received 10th August 2024
Accepted 18th October 2024

DOI: 10.1039/d4sc05362a

rsc.li/chemical-science

Introduction

Phospholipids are major constituents of biological membranes.¹ Their emergence on early Earth would thus have marked a pivotal step on the path towards primitive cells.² Prebiotic synthesis would have produced heterogeneous mixtures of isomeric phospholipids, but only the natural phospholipids comprising glycerol-1-phosphate (G1P) and glycerol-3-phosphate (G3P) were retained in biological membranes while their non-natural analogues comprising glycerol-2-phosphate (G2P) were presumably lost to chemical evolution. Still, it is not clear how these natural phospholipids were selected.

Prebiotic chemists have sought to identify mechanisms by which many of the natural building blocks of life could have been selected. Yet, investigations have focused almost exclusively on nucleosides and peptides. Canonical RNA and DNA nucleosides could have arisen by selective prebiotic synthesis^{3–5}

or because of their greater hydrolytic and photochemical stability relative to non-canonical isomers.^{6–8} Selection could also have resulted from the increased stability of duplexes comprising natural oligonucleotides, compared to non-natural analogues.⁹ Similarly, the canonical amino acids could have resulted from selective synthesis,¹⁰ the preferential incorporation of natural over non-natural amino acids into peptides,¹¹ or by retention due to the enhanced foldability of oligopeptides containing natural amino acids compared to those containing non-natural analogues.¹² The idea that selection processes could have operated at both the molecular and supramolecular levels is central to every scheme.

Two recent reports have begun to address the selection of natural phospholipids. In work on the emergence of amino alcohol bearing phospholipids, our group¹³ described a general prebiotic synthesis and hydrolysis scheme that is selective for natural phospholipid headgroups comprising G1P over those comprising G2P. Then, in work advancing the prebiotic chemistry of cyclophospholipids, Krishnamurthy and co-workers¹⁴ showed that liposomes formed of glycerol monodecanoate and phospholipid comprising G1P are more tolerant of pH changes and metal ions, such as those required for prebiotic RNA chemistry,¹⁵ than those formed of phospholipid comprising G2P.

^aInstitut de Science et d'Ingénierie Supramoléculaires (ISIS), CNRS UMR 7006, University of Strasbourg, 8 Allée Gaspard Monge, 67000 Strasbourg, France. E-mail: cb2036@cam.ac.uk

^bDepartment of Biochemistry, University of Cambridge, Tennis Court Road, CB2 1GA Cambridge, UK

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc05362a>

These studies raise further questions, particularly in light of the systems chemistry approach in which mixtures of reactants and products must be considered and their properties compared. First, it is not clear if prebiotic synthesis and hydrolysis processes could have led to the selection of natural over non-natural diacyl phospholipids. Prebiotic acylation reactions of individual glycerol phosphates have been reported,^{16,17} but no direct comparison has been made of the reactivity of **G1P** and **G2P** with activated fatty acids, nor of the hydrolytic stability of the corresponding phospholipids. Second, it is not clear if differences in membrane properties could have led to the selection of natural phospholipids from heterogeneous mixtures containing both natural and non-natural forms. Hybrid liposomes often behave differently from those made of individual lipids and so their behavior is difficult to predict.

Here we explore processes operating at the molecular and supramolecular levels that could have led to the selection of natural phosphatidic acids from heterogeneous mixtures on early Earth. We show that hydrolytically stable natural and non-natural phosphatidic acids could have formed on prebiotic acylation of mixtures of glycerol phosphates under a variety of conditions, together with a range of amphiphilic coproducts. We then highlight that natural phosphatidic acids have a greater propensity to self-assemble, compared to their non-natural isomers, and that their liposomes display enhanced rigidity and stability towards changes in pH and metal ion concentrations. Finally, harnessing these differences, we demonstrate that on exposure to Mg^{2+} , but not Na^+ , K^+ , Ca^{2+} or Zn^{2+} , natural phosphatidic acids could have been selected from liposomes composed of heterogeneous mixtures. Overall, our findings offer a plausible rationale for the selection of natural phospholipids, taking advantage of differences in properties that manifest only at the supramolecular level.

Note: throughout our discussion we treat **G1P** and **G3P** as synonymous and for simplicity refer to them with the acronym **G1P**. The stereospecific numbering (*sn*) scheme for glycerophospholipids is used only where appropriate.

Results and discussion

Prebiotic synthesis delivers mixtures of natural and non-natural phosphatidic acids

Synthesis in solution. Several prebiotic routes to phosphatidic acids have been described,^{16–19} and more can be traced by combining steps from multiple reports. For example, glycerol could have been delivered by (photo)reduction of acetaldehyde or dihydroxyacetone,^{20–22} and glycerol-1-phosphate (**G1P**) and glycerol-2-phosphate (**G2P**) could have formed on phosphorylation of glycerol.^{18,20–25} Phosphatidic acids could have then been produced by acylation of glycerol phosphates with activated fatty acids under various conditions. However, previous investigations^{16,17} only addressed the long-chain acylation of single isomers of glycerol phosphate, despite the likely co-occurrence of both **G1P** and **G2P** on early Earth. Thus, it is not clear whether differences in the reactivity of **G1P** and **G2P**

towards acylating agents might explain the selection of natural **G1PAs** over non-natural **G2PAs**.

To probe the acylation of glycerol phosphates in solution, each glycerol phosphate was dissolved in water and the pH of each solution was adjusted to 7.4. A solution of *N*-decanoyl imidazole (NDI) in acetonitrile was then added,¹⁶ and each homogeneous mixture was stirred for 8 days at room temperature (Fig. 1a, b and S1†). The progress of each reaction was



Fig. 1 Prebiotic synthesis of phosphatidic acids. (a) Reactions of **G1P** (10 mM), **G2P** (10 mM), or both (5 mM each) with NDI (100 mM) in solution, or ammonium decanoate (100 mM), cyanamide (100 mM), and imidazole (100 mM) in the dry state, afford natural **G1PAs** and non-natural **G2PAs**. (b) Yields of **G1PAs** and **G2PAs** formed from **G1P**, **G2P**, or both. (c) $^{31}\text{P}\{^1\text{H}\}$ NMR spectra showing the signals for the **G1PAs** and **G2PAs** obtained in reactions described in entries 3 (pH ~7) and 6 (pH ~4 after extraction) of (b). Yields were determined by ^{31}P NMR based on conversion of the respective glycerol phosphate (or sum total glycerol phosphates). The chemical shift of each phosphatidic acid is pH and concentration dependent. *The major positional isomer²⁶ of **MDG1PA** is shown in (a) and combined yields of both isomers of **MDG1PA** are given in (b). See the ESI† for details.

Hydrolysis. Our group recently demonstrated that hydrolysis provides a mechanism for the selection of complex natural phospholipids and their headgroups.¹³ Thinking along similar lines, we wondered if **G2PAs** might be more susceptible to hydrolysis than **G1PAs** and if hydrolysis processes would enable the selection of natural phosphatidic acids. Accordingly, the C10-lipids **DDG1PA** and **DDG2PA** were synthesized^{31,32} and subjected to hydrolysis at room temperature, pH 10, either in homogeneous solution in acetonitrile/carbonate buffer (1 : 1, v/v) or in the form of liposomes in carbonate buffer. The extent of hydrolysis of each lipid was greater in solution than in liposomes at every time point, as observed for other primitive amphiphiles,³³ but differences between the extents of hydrolysis of **DDG1PA** and **DDG2PA** in solution and in liposomes were minimal (Fig. S10†). For example, after 11 days in solution 87% of **DDG1PA** and 87% of **DDG2PA** were hydrolyzed, while in liposomes 35% of **DDG1PA** and 34% of **DDG2PA** were hydrolyzed. Similar hydrolysis experiments were performed at pH 7 (imidazole buffer) and pH 4 (acetate buffer), but slow hydrolysis (data not shown) and precipitation of the lipid (see below) prevented us from observing any differences. These results

Synthesis in the dry state. To probe the acylation of glycerol phosphates in the dry state, each glycerol phosphate was dissolved in water and ammonium decanoate, cyanamide, and imidazole were added before the pH of the mixture was adjusted to 7.4 (Fig. 1a and b).¹⁷ Each mixture was allowed to evaporate at room temperature and the dried mixture was heated at 65 °C for 18 h. The products were extracted into acetonitrile/water/D₂O (5 : 4 : 1, v/v/v), and the formation of phosphatidic acids was verified by ³¹P NMR and TLC as before. In contrast to the results obtained in solution, a broader range of products was produced in the dry state. On acylation of **G1P**, **MDG1PA** and **DDG1PA** were obtained in yields of 15% and 4%, respectively, alongside **G2P** (2%), **MDG2PA** (2%), **DDG2PA** (1%), glycerol-1,2-cyclic phosphate (**G>P**, 16%), and 3-monodecanoyl glycerol-1,2-cyclic phosphate (**MDG>P**, 6%) (Fig. 1b and S5†). The positional isomers, **G2P** and **MDG2PA** probably form on hydrolysis of **G>P** and **MDG>P**, respectively,^{13,29} or by phosphoryl migration under the ultimately acidic conditions of the reaction.²⁶ Similarly, the reaction of **G2P** afforded **MDG2PA** and **DDG2PA** in yields of 14% and 4%, respectively, along with the same array of products as

suggest that, in contrast to the possible emergence of amino alcohol bearing phospholipids,¹³ it is not likely that the selection of natural phosphatidic acids on early Earth resulted from hydrolysis.

Natural and non-natural phosphatidic acids exhibit different supramolecular properties

As **G1PAs** and **G2PAs** form stable liposomes in aqueous buffers,^{14,16} we sought to identify whether differences in their respective supramolecular properties could have led to the selection of natural **G1PAs**. Microscopy observations showed that **DDG1PA** and **DDG2PA** form liposomes of similar size and lamellarity in phosphate buffer (50 mM, pH 7.5) (Fig. 2a).^{14,16} Yet, the biophysical properties of these membranes remain unknown. Previous NMR and neutron diffraction studies on 1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine (**DPG3PC**) and its isomer 1,3-dipalmitoyl-*sn*-glycero-2-phosphocholine (**DPG2PC**) showed that the two phospholipids differ in their gel-to-liquid transition temperature (41.5 °C for **DPG3PC** vs. 37.5 °C for **DPG2PC**), probably due to the different conformations of their glycerol backbones (perpendicular and parallel to the membrane surface for **DPG3PC** and **DPG2PC**, respectively).³⁴ We thus investigated whether natural and non-natural phosphatidic acids also acquire a different spatial organization

within their respective bilayer. Accordingly, liposomes prepared from **DDG1PA** and **DDG2PA** labelled with the fluorescent probe 1,6-diphenyl-1,3,5-hexatriene (DPH) were studied by fluorescence anisotropy. DPH is an apolar dye that localizes in the hydrophobic region of the lipid bilayer and so fluorescence anisotropy measurements provide information on the freedom of motion of the dye within the membrane, the viscosity of the environment in which the dye is embedded, and hence the fluidity of the membrane.³⁵ Our data show that **DDG1PA** and **DDG2PA** form liposomes with membranes as fluid as those composed of unsaturated fatty acids, such as myristoleic acid (**MA**) (Fig. 2b), but that **DDG1PA** membranes are more rigid than **DDG2PA** membranes. Moreover, when **DDG1PA** and **DDG2PA** liposomes were labelled with Laurdan, an amphiphilic dye sensitive to lipid packing and order within the bilayer,³⁶ generalized polarization studies showed that the **DDG1PA** membranes are more ordered and better packed than **DDG2PA** membranes, indicative of stronger hydrophobic interactions between the acyl chains (Fig. 2c and S11†). The fluorescence anisotropy and generalized polarization measurements were made at 25 °C, a temperature expected to be above the gel-to-fluid phase transition temperature of both lipids. These findings confirm that natural and non-natural phosphatidic acids acquire different conformations within their respective bilayers



Fig. 2 Self-assembly and supramolecular properties of **G1PAs** and **G2PAs** in pure or mixed systems. (a) Fluorescence microscopy images show that **DDG1PA** and **DDG2PA** (25 mM, containing 1 mol% NBD-PE) form liposomes of similar size and lamellarity. The scale bar represents 10 μm. (b) DPH fluorescence anisotropy (polarization) of **DDG1PA** and **DDG2PA** membranes compared with those of 1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine (**DPG3PC**) and **MA**. Measurements were made at 25 °C. (c) Laurdan generalized polarization (GP) of **DDG1PA** and **DDG2PA** membranes compared with those of **DPG3PC** and 1,2-dioleoyl-*sn*-glycero-3-phosphocholine (**DOG3PC**). Measurements were made at 25 °C. (d) Structural and conformational differences between **DDG1PA** and **DDG2PA** in membranes. (e) CACs of **G1PAs** and **G2PAs** of different chain length. (f) CACs of mixtures of **DDG1PA** or **DDG2PA** and other C10-amphiphiles. Data show the mean ± SEM, $n \geq 2$. See the ESI† for details.

quantified by ^{31}P NMR (99% loss after addition of 5 mM Mg^{2+}) (Fig. S25†). By contrast, **DDG1PA** liposomes are largely unaffected by Mg^{2+} (16% loss after addition of 5 mM Mg^{2+}) (Fig. S26†). Similar observations were made by epifluorescence microscopy using a higher lipid concentration (25 mM), where precipitation of **DDG2PA** was observed at 7.5 mM MgCl_2 (Fig. 3a). The concentration of Mg^{2+} required to induce precipitation is proportional to the concentration of the lipid.



Fig. 3 Mg^{2+} -driven selection of natural **G1PAs** over non-natural **G2PAs**. (a) Microscopy images showing the relative Mg^{2+} -tolerance of **DDG1PA** and **DDG2PA** liposomes (25 mM). The scale bar represents 10 μm . **DDG1PA** liposomes remain intact while **DDG2PA** liposomes disassemble following aggregation and precipitation of the lipid. The concentration of Mg^{2+} required to induce precipitation is proportional to the concentration of the lipid. (b) $^{31}\text{P}\{^1\text{H}\}$ NMR spectra showing the change in composition of mixed liposomes prepared from **DDG1PA** and **DDG2PA** (1 : 1, 5 mM each) on titration with Mg^{2+} . (c) $^{31}\text{P}\{^1\text{H}\}$ NMR spectra showing the change in composition of mixed liposomes prepared from **G1PAs** and **G2PAs** obtained by prebiotic synthesis on titration with Mg^{2+} . **DDG2PA** precipitates from mixed liposomes on titration with Mg^{2+} , while **DDG1PA** remains in liposomes. See the ESI† for details.

Analogously, **DDG2PA** liposomes were more susceptible to precipitation by CaCl_2 and ZnCl_2 than **DDG1PA** liposomes (Fig. S27–S30†). Finally, by comparison, neither lipid was susceptible to precipitation on addition of up to 100 mM NaCl or KCl , as expected for monovalent cations.⁴³

To probe the greater tolerance of **DDG1PA** over **DDG2PA** to Mg^{2+} , we measured the change in membrane ζ -potential of liposomes made of **DDG1PA** or **DDG2PA** (5 mM each) on titration with MgCl_2 . In the absence of Mg^{2+} , the ζ -potentials of **DDG1PA** and **DDG2PA** liposomes were -56 mV and -74 mV, respectively (Fig. S31†). The more negative surface potential of **DDG2PA** suggests that its phosphate group is less hydrated and hence forms weaker hydrogen bonding interactions in the headgroup space of the bilayer compared to **DDG1PA**. Further, it is possible that the phosphate group of **DDG2PA** is more exposed to the bulk medium than that of **DDG1PA**, as might be expected since **DDG2PA** forms less rigid and less well packed membranes than **DDG1PA** (Fig. 2c). On addition of Mg^{2+} to liposomes composed of each lipid, we observed a gradual increase in ζ -potential, demonstrating the electrostatic interactions of Mg^{2+} with the anionic headgroups and its charge shielding effect (Fig. S31†).⁴³

Intrigued by the lower tolerance of **DDG2PA** over **DDG1PA** to divalent cations, we tested the stability of mixed liposomes made of **DDG1PA** and **DDG2PA** (1 : 1, 5 mM each, prepared in 50 mM phosphate buffer, pH 7.5) in the presence of 5 mM Mg^{2+} . In the event, the sample remained turbid despite the immediate formation of a precipitate.

To determine the composition of the remaining liposomes, the precipitate was removed by filtration, the liposomes were dissolved by addition of 0.2 M Triton X-100,⁴⁴ and the resulting solution was analyzed by ^{31}P NMR (Fig. 3b and S32†). **DDG1PA** was found to be four times more abundant than **DDG2PA** in the filtrate, demonstrating that selective precipitation of the non-natural **DDG2PA** occurred on addition of Mg^{2+} . This finding suggests that a Mg^{2+} -driven selection process could have led to the enrichment of natural phosphatidic acids in mixed liposomes. Moreover, Mg^{2+} appears unique among the metal ions studied as selective precipitation of **DDG2PA** from mixed liposomes could not be effected by titration with Ca^{2+} , Zn^{2+} (Fig. S33 and S34†), Na^+ or K^+ (data not shown).

The mechanism that underpins the Mg^{2+} -induced selection from mixed membranes can be understood in terms of the electrostatic interactions between Mg^{2+} and the anionic surface of the membrane and the differential binding of the isomeric lipids.⁴⁵ Mg^{2+} binding to phosphatidic acids would liberate water of hydration from the membrane, leading to tighter packing of the lipids and ultimately precipitation. Our data show that **DDG2PA** membranes have a more negative surface charge than those of **DDG1PA** and suggest that the headgroups of **DDG2PA** are less well hydrated and so more prone to disruption on dehydration. It is also possible that the binding affinity of Mg^{2+} for **DDG2PA** is greater than that for **DDG1PA** and the difference in binding affinities is enough to enable the selective precipitation of **DDG2PA** in the presence of **DDG1PA**, in contrast to the other metal ions tested. Whether the selective



precipitation of **DDG2PA** from mixed membranes suggests that phase separation into isomeric lipid domains occurs prior to precipitation remains an intriguing possibility.

Finally, to demonstrate the plausibility of the Mg^{2+} -driven selection process on lipid mixtures obtained by prebiotic synthesis, we prepared a mixture of **G1PAs** and **G2PAs** by acylation of **G1P** and **G2P** with NDI, as described above (Fig. 1). The mixture of phospholipids obtained after 8 days (**MDG1PA**, **MDG2PA**, **DDG1PA**, and **DDG2PA**) was lyophilized, then gently hydrated with water at pH 7.0 to give liposomes. The imidazole released on reaction of NDI served as the buffer. On titration with MgCl_2 , **DDG2PA** precipitated and the surviving liposomes became proportionally enriched in **DDG1PA** (^{31}P NMR, Fig. 3c and Fig. S35–S37†). Interestingly, the lysophospholipids **MDG1PA** and **MDG2PA** were unaffected by Mg^{2+} up to a concentration of 5 mM, yet these species could undergo further acylation and thus Mg^{2+} -induced precipitation.

Overall, these results highlight the unique role of Mg^{2+} in driving the selection of **G1PAs** from heterogeneous mixtures of phospholipids under conditions compatible with key metal-driven prebiotic processes like non-enzymatic RNA polymerisation.¹⁵

Conclusion

Due to the heterogeneity of prebiotic building blocks present on early Earth, efficient mechanisms would have been required for the selection of the now canonical biomolecules. In relation to the emergence of phospholipids, recent studies showed that the selection of natural phospholipid headgroups could have resulted from the preferential hydrolysis of their non-natural analogues¹³ and that liposomes composed of non-natural phosphatidic acids are poorly tolerant to pH changes and metal ions.^{14,16} However, no prior study addressed whether processes operating at the molecular level (prebiotic synthesis) or the supramolecular level (membrane properties) could have driven the selection of natural phospholipids from heterogeneous mixtures available on early Earth. Our work addresses these questions.

First, we show that acylation of mixtures of glycerol phosphates in solution affords non-natural **G2PAs** in higher yields than natural **G1PAs**, but acylation in the dry state provides **G1PAs** and **G2PAs** in similar yields. These isomeric phospholipids are comparably stable to hydrolysis. Second, we highlight that natural **G1PAs** possess a greater propensity to self-assemble and to pack in ordered bilayers, compared to non-natural **G2PAs**, and that they form liposomes of enhanced rigidity and broader tolerance to pH and metal ions. Finally, we demonstrate that Mg^{2+} drives the selective precipitation of **G2PAs** and, thus, the accumulation of **G1PAs** in liposomes composed of mixtures of phospholipids obtained by prebiotic synthesis. The effect of Mg^{2+} is unique among the metal ions tested. Taken together, our results suggest that properties that manifest at the supramolecular level can be harnessed for the selection of natural phospholipids, while processes that operate at the molecular level (synthesis and hydrolysis) probably cannot.

In summary, our work outlines prebiotic synthesis and selection pathways that could have driven the transition towards biological phospholipid membranes on early Earth. The phosphatidic acids focused on here are minor constituents of biological membranes but serve as important biosynthetic intermediates in pathways leading to complex lipids, such as phosphatidylethanolamines and phosphatidylcholines. Since the transformation of **G1PAs** into both of these lipids under early Earth conditions has already been described,^{46,47} our study renders these foundational investigations all the more relevant. Importantly, the requirements of Mg^{2+} -driven non-enzymatic RNA replication and the need for Mg^{2+} -stable biological phospholipid liposomes can be harmonized. In fact, Mg^{2+} is essential in driving the selection of metal ion-resistant natural phospholipids. Beyond, the interactions between Mg^{2+} ions and membranes enriched in natural phosphatidic acids could have led to the co-localization of RNA on the membrane and enabled synergistic effects in prebiotic RNA chemistry⁴⁸ as well as RNA-driven recognition and fusion of primitive liposomes.⁴⁹

Data availability

Data are available upon reasonable request from the corresponding author.

Author contributions

K. T. M., Y. L., V. J. B.-T. and D. A. R. carried out the experimental work. D. A. R. and C. B. conceived the project, directed the work and wrote the manuscript with input from all the authors.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The project was funded by the Foundation Jean-Marie Lehn, the University of Strasbourg Institute for Advanced Study (USIAS) and the Agence Nationale de la Recherche (ITI SysChem within the Investissement d'Avenir program ANR-10-IDEX-0002, and AAP IdEx 2022 Attractivité to C. B.), and the CSC Graduate School, funded by the Agence Nationale de la Recherche (CSC-IGS ANR-17-EURE-0016 to K. T. M. and V. J. B. T.). The authors gratefully acknowledge support from Foundation Jean-Marie Lehn, the University of Strasbourg Institute for Advanced Study and the Agence Nationale de la Recherche. The authors thank colleagues at ISIS and lab members for stimulating discussions.

References

- 1 B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts and P. Walter, *Molecular Biology of the Cell*, Garland Science, New York, 2014.



- 2 J. C. Blain and J. W. Szostak, *Annu. Rev. Biochem.*, 2014, **83**, 615–640.
- 3 J. Xu, V. Chmela, N. J. Green, D. A. Russell, M. J. Janicki, R. W. Góra, R. Szabla, A. D. Bond and J. D. Sutherland, *Nature*, 2020, **582**, 60–66.
- 4 J. Xu, N. J. Green, D. A. Russell, Z. Liu and J. D. Sutherland, *J. Am. Chem. Soc.*, 2021, **143**, 14482–14486.
- 5 M. W. Powner, B. Gerland and J. D. Sutherland, *Nature*, 2009, **459**, 239–242.
- 6 A. C. Rios and Y. Tor, *Isr. J. Chem.*, 2013, **53**, 469–483.
- 7 M. W. Powner and J. D. Sutherland, *ChemBioChem*, 2008, **9**, 2386–2387.
- 8 R. A. Sanchez and L. E. Orgel, *J. Mol. Biol.*, 1970, **47**, 531–543.
- 9 S. Bhowmik and R. Krishnamurthy, *Nat. Chem.*, 2019, **11**, 1009–1018.
- 10 K. Ashe, C. Fernández-García, M. K. Corpinot, A. J. Coggins, D.-K. Bučar and M. W. Powner, *Commun. Chem.*, 2019, **2**, 1–7.
- 11 M. Frenkel-Pinter, J. W. Haynes, M. C. A. S. Petrov, B. T. Burcar, R. Krishnamurthy, N. V. Hud, L. J. Leman and L. D. Williams, *Proc. Natl. Acad. Sci. U. S. A.*, 2019, **116**, 16338–16346.
- 12 M. Makarov, A. C. Sanchez Rocha, R. Krystufek, I. Cherepashuk, V. Dzmitruk, T. Charnavets, A. M. Faustino, M. Lebl, K. Fujishima, S. D. Fried and K. Hlouchova, *J. Am. Chem. Soc.*, 2023, **145**, 5320–5329.
- 13 M. Aleksandrova, F. Rahmatova, D. A. Russell and C. Bonfio, *J. Am. Chem. Soc.*, 2023, **145**, 25614–25620.
- 14 S. Pulletikurti, K. S. Veena, M. Yadav, A. A. Deniz and R. Krishnamurthy, *Chem*, 2024, **10**, 1839–1867.
- 15 S. S. Mansy, J. P. Schrum, M. Krishnamurthy, S. Tobé, D. A. Treco and J. W. Szostak, *Nature*, 2008, **454**, 122–125.
- 16 C. Bonfio, C. Caumes, C. D. Duffy, B. H. Patel, C. Percivalle, M. Tsanakopoulou and J. D. Sutherland, *J. Am. Chem. Soc.*, 2019, **141**, 3934–3939.
- 17 D. E. Epps, E. Sherwood, J. Eichberg and J. Oró, *J. Mol. Evol.*, 1978, **11**, 279–292.
- 18 D. Fayolle, E. Altamura, A. D'Onofrio, W. Madanamothoo, B. Fenet, F. Mavelli, R. Buchet, P. Stano, M. Fiore and P. Strazewski, *Sci. Rep.*, 2017, **7**, 18106.
- 19 W. R. Hargreaves, S. J. Mulvihill and D. W. Deamer, *Nature*, 1977, **266**, 78–80.
- 20 B. H. Patel, C. Percivalle, D. J. Ritson, C. D. Duffy and J. D. Sutherland, *Nat. Chem.*, 2015, **7**, 301–307.
- 21 D. J. Ritson and J. D. Sutherland, *Nat. Chem.*, 2023, **15**, 1470–1477.
- 22 D. E. Epps, D. W. Nooner, J. Eichberg, E. Sherwood and J. Oró, *J. Mol. Evol.*, 1979, **14**, 235–241.
- 23 C. Gibbard, S. Bhowmik, M. Karki, E.-K. Kim and R. Krishnamurthy, *Nat. Chem.*, 2018, **10**, 212–217.
- 24 O. R. Maguire, I. B. A. Smokers and W. T. S. Huck, *Nat. Commun.*, 2021, **12**, 5517.
- 25 B. Burcar, M. Pasek, M. Gull, B. J. Cafferty, F. Velasco, N. V. Hud and C. Menor-Salv, *Angew. Chem., Int. Ed.*, 2016, **55**, 13249–13253.
- 26 A. Plueckthun and E. A. Dennis, *Biochemistry*, 1982, **21**, 1743–1750.
- 27 E. K. Ryu and M. MacCoss, *J. Lipid Res.*, 1979, **20**, 561–563.
- 28 L. Kugel and M. Halmann, *J. Am. Chem. Soc.*, 1967, **89**, 4125–4128.
- 29 V. E. Ortuno, S. Pulletikurti, K. S. Veena and R. Krishnamurthy, *Chem. Commun.*, 2022, **58**, 6231–6234.
- 30 H. Bockerhoff and M. Yurkowsky, *Can. J. Biochem.*, 1965, **43**, 1777.
- 31 C. M. Gupta, R. Radhakrishnan and H. G. Khorana, *Proc. Natl. Acad. Sci. U. S. A.*, 1977, **74**, 4315–4319.
- 32 E. G. Bligh and W. J. Dyer, *Can. J. Biochem. Physiol.*, 1959, **37**, 911–917.
- 33 C. Bonfio, D. A. Russell, N. J. Green, A. Mariani and J. D. Sutherland, *Chem. Sci.*, 2020, **11**, 10688–10697.
- 34 G. Buldt and G. H. de Haas, *J. Mol. Biol.*, 1982, **158**, 55–71.
- 35 B. J. Litman and Y. Barenholz, in *Methods in Enzymology*, Academic Press, 1982, vol. 81, pp. 678–685.
- 36 T. Parasassi, G. De Stasio, G. Ravagnan, R. M. Rusch and E. Gratton, *Biophys. J.*, 1991, **60**, 179–189.
- 37 G. S. Harbison and R. G. Griffin, *J. Lipid Res.*, 1984, **25**, 1140–1142.
- 38 A. Blume and H. Eibl, *Biochim. Biophys. Acta*, 1979, **558**, 13–21.
- 39 H. Eibl and A. Blume, *Biochim. Biophys. Acta*, 1979, **553**, 476–488.
- 40 P. Dalai, P. Ustriyana and N. Sahai, *Geochim. Cosmochim. Acta*, 2018, **223**, 216–228.
- 41 C. Bonfio, E. Godino, M. Corsini, F. Fabrizi de Biani, G. Guella and S. S. Mansy, *Nat. Catal.*, 2018, **1**, 616–623.
- 42 K. B. Muchowska, S. J. Varma and J. Moran, *Nature*, 2019, **569**, 104–107.
- 43 H. Träuble and H. Eibl, *Proc. Natl. Acad. Sci. U.S.A.*, 1974, **71**, 214–219.
- 44 E. London and G. W. Feigenson, *J. Lipid Res.*, 1979, **20**, 408–412.
- 45 H. Hauser, *Chem. Phys. Lipids*, 1991, **57**, 309–325.
- 46 M. Rao, J. Eichberg and J. Oró, *J. Mol. Evol.*, 1982, **18**, 196–202.
- 47 M. Rao, J. Eichberg and J. Oró, *J. Mol. Evol.*, 1987, **25**, 1–6.
- 48 T. Czerniak and J. P. Saenz, *Proc. Natl. Acad. Sci. U. S. A.*, 2022, **119**, e2119235119.
- 49 J. A. Peruzzi, M. L. Jacobs, T. Q. Vu, K. S. Wang and N. P. Kamat, *Angew. Chem., Int. Ed.*, 2019, **58**, 18683–18690.

